University of Dundee

Microbial Dysbiosis in Bronchiectasis and Cystic Fibrosis

Dicker, Alison J.; Chalmers, James D.

Published in:
Archivos de Bronconeumologia

DOI:
[10.1016/j.arbres.2016.11.018](https://doi.org/10.1016/j.arbres.2016.11.018)

Publication date:
2017

Licence:
CC BY-NC-ND

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Dicker, A. J., & Chalmers, J. D. (2017). Microbial Dysbiosis in Bronchiectasis and Cystic Fibrosis. *Archivos de Bronconeumologia*, 53(9), 471-472. <https://doi.org/10.1016/j.arbres.2016.11.018>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Editorial: Microbial Dysbiosis in Bronchiectasis and Cystic Fibrosis

A. J. Dicker and J. D. Chalmers*

Scottish Centre for Respiratory Research, University of Dundee, Ninewells Hospital & Medical School, Dundee, U.K.

*Corresponding Author: James D Chalmers, Scottish Centre for Respiratory Research, University of Dundee, Ninewells Hospital & Medical School, Dundee DD1 9SY, UK, jchalmers@dundee.ac.uk, phone 01382383642

Identification and treatment of bacterial infection is central to our management of bronchiectasis, whether caused by cystic fibrosis or by other underlying disorders.(1) The pathophysiology of these diseases is understood to be a vicious cycle of infection, inflammation and tissue damage which we attempt to break by eradicating or suppressing pathogenic bacteria using inhaled, oral and intravenous antibiotics.(1, 2)

Our understanding of this pathophysiology is being changed, however, by developments in molecular microbiology and genomics. Non-culture based methods of identifying bacteria, viruses and fungi in the lungs of CF and bronchiectasis patients is a rapidly expanding field of research. By utilising high-throughput sequencing platforms to target conserved regions of DNA, such as the bacterial 16S rRNA gene, identification of the bacteria present in sputum, bronchoalveolar lavage (BAL) or upper airway swabs is possible without the need for microbial culture.(3) There is potential to identify potential pathogens more sensitively, to identify non-culturable bacteria, to understand “phenotypes” of disease based on microbiota profiles, and to better understand the impact of treatment on microbes and the host response.

Cystic fibrosis is perhaps the most intensively studied condition with respect to the lung microbiome. Recent data shows that the upper airway microbiome starts to change in CF infants in the first year of life with overgrowth of *Staphylococcus* even in children without antibiotic treatment.(4, 5) As disease progresses, the proteobacteria increase in relative abundance with the emergence of “traditional” CF pathogens such as *Pseudomonas spp.* Longitudinal studies have identified that the lung microbiota is stable over time, at least in the short term, and does not vary significantly during exacerbations.(6) One of the largest studies in CF patients to date (n=269) confirmed patterns of microbiota diversity observed in other smaller-scale studies; dysbiosis (a reduction in microbiota diversity with dominance of one or a few genera) is associated with more severe disease, age and increasing antibiotic use.(7) A subset of genera, namely *Veillonella*, *Burkholderia*, *Prevotella*, *Streptococcus*, *Rothia* and *Actinomyces*, appear to form a core CF microbiota whilst genera like *Pseudomonas* and *Staphylococcus* become more dominant as disease progresses. Very few studies have extensive longitudinal follow up of patients or have data on the impact of therapies such as inhaled antibiotics or CFTR modulators; this will need to be addressed before the utility of microbiota data in clinical settings is fully realised.

The microbiota in bronchiectasis has been less well studied than in CF; small studies have shown that it is diverse and relatively stable over time, with a reduction in diversity identified at exacerbations and also associated with increasing disease severity.(8) Like CF, there are common shared genera such as *Veillonella*, *Streptococcus* and *Prevotella* with dominance of pathogens such as *Pseudomonas* or *Haemophilus* associated with increasing disease severity.(9) Intriguingly, a randomized control trial

(RCT) of erythromycin suggested that long term macrolide therapy, a common treatment strategy, may be associated with an increase in relative abundance of *Pseudomonas spp.*(9) This is important because of the well-known association between *Pseudomonas aeruginosa* infection and poor clinical outcomes.(10) Dominance of *Pseudomonas spp.* by sequencing has also been associated with more exacerbations and increased levels of inflammation such as matrix metalloproteinases.(9, 11)

While there are increasing numbers of bronchiectasis microbiota studies, a significant percentage have arisen from a single RCT in Australia that included only patients with a history of 2 or more exacerbations in the previous year among other exclusion criteria.(9) It has been shown that 52-93% of bronchiectasis patients in real-life are ineligible for RCTs.(12) Larger studies from diverse populations are now needed.

Studies show that individuals microbiota profiles vary significantly from each other; what is normal for one patient during disease stability may not be for another. The key to being able to use microbiota data clinically will be a better understanding of how an individual patients microbiota changes over time, and how this is associated with altered lung inflammatory profiles, clinical phenotypes and clinical outcomes. In both CF and bronchiectasis it appears that this change towards dysbiosis alongside changes in clinical and inflammatory profiles is clinically important and useful to stratify patients. What is now needed is to determine if microbiota profiles can add to existing stratification methods such as the Bronchiectasis Severity Index, by providing additional prediction of exacerbations or treatment response.(13, 14)

A barrier to effective implantation of microbiota data into practice is a lack of agreed technical approaches. Sampling techniques and DNA extraction methods vary greatly. Different regions of the 16S rRNA gene have been sequenced and numerous different sequence analysis pipelines exist. All of these can cause slight variation in the proportions of genera identified. Contamination of samples or laboratory reagents with can cause misleading results if not identified and dealt with, particularly in low biomass samples. These technical and bioinformatic challenges need to be overcome and methods standardised if microbiota data is to become useful in clinical care.

The bacterial microbiota alone will not explain all the phenotypic diversity seen in CF and bronchiectasis patients. Viruses, fungi and mycobacteria also contribute to the lung microbiome in the healthy lung, in CF and bronchiectasis disease stability and exacerbation. Indeed for bronchiectasis no 'virome' or 'mycobiota' studies have been published to date, yet it is acknowledged that mycosis is becoming an increasing problem.(15) In both CF and bronchiectasis, therapeutic and prophylactic use of antibiotics are key parts of the physicians' arsenal, yet whilst the treatments may improve disease symptoms, the effect they have on the fungi, bacteria and viruses inhabiting the

airway is still not known. A better understanding how antibiotic usage in bronchiectasis and CF may contribute towards the development of a dysbiotic microbiome could lead towards better antibiotic stewardship in early disease to slow the development of dysbiosis and drug resistance.

Microbial sequencing technologies are a powerful tool to understand these diseases as never before. Application of novel tools is not, however, an end in itself. The challenge for the pulmonary research community is to now apply this technology to answer clinically important questions and allow microbiome science to have a meaningful impact on patient care.

1. Burgel PR, Bellis G, Olesen HV, Viviani L, Zolin A, Blasi F, Elborn JS. Future trends in cystic fibrosis demography in 34 european countries. *Eur Respir J* 2015;46:133-141.
2. Chalmers JD, Aliberti S, Blasi F. Management of bronchiectasis in adults. *Eur Respir J* 2015;45:1446-1462.
3. Wang Z, Bafadhel M, Haldar K, Spivak A, Mayhew D, Miller BE, Tal-Singer R, Johnston SL, Ramsheh MY, Barer MR, Brightling CE, Brown JR. Lung microbiome dynamics in copd exacerbations. *Eur Respir J* 2016;47:1082-1092.
4. Prevaes SM, de Winter-de Groot KM, Janssens HM, de Steenhuijsen Piters WA, Tramper-Stranders GA, Wyllie AL, Hasrat R, Tiddens HA, van Westreenen M, van der Ent CK, Sanders EA, Bogaert D. Development of the nasopharyngeal microbiota in infants with cystic fibrosis. *Am J Respir Crit Care Med* 2016;193:504-515.
5. Mika M, Korten I, Qi W, Regamey N, Frey U, Casaulta C, Latzin P, Hilty M. The nasal microbiota in infants with cystic fibrosis in the first year of life: A prospective cohort study. *Lancet Respir Med* 2016.
6. Fodor AA, Klem ER, Gilpin DF, Elborn JS, Boucher RC, Tunney MM, Wolfgang MC. The adult cystic fibrosis airway microbiota is stable over time and infection type, and highly resilient to antibiotic treatment of exacerbations. *PLoS One* 2012;7:e45001.
7. Coburn B, Wang PW, Diaz Caballero J, Clark ST, Brahma V, Donaldson S, Zhang Y, Surendra A, Gong Y, Elizabeth Tullis D, Yau YC, Waters VJ, Hwang DM, Guttman DS. Lung microbiota across age and disease stage in cystic fibrosis. *Sci Rep* 2015;5:10241.
8. Tunney MM, Einarsson GG, Wei L, Drain M, Klem ER, Cardwell C, Ennis M, Boucher RC, Wolfgang MC, Elborn JS. Lung microbiota and bacterial abundance in patients with bronchiectasis when clinically stable and during exacerbation. *Am J Respir Crit Care Med* 2013;187:1118-1126.
9. Rogers GB, Bruce KD, Martin ML, Burr LD, Serisier DJ. The effect of long-term macrolide treatment on respiratory microbiota composition in non-cystic fibrosis bronchiectasis: An analysis from the randomised, double-blind, placebo-controlled trial. *The Lancet Respiratory Medicine* 2014;2:988-996.
10. Finch S, McDonnell MJ, Abo-Leyah H, Aliberti S, Chalmers JD. A comprehensive analysis of the impact of pseudomonas aeruginosa colonization on prognosis in adult bronchiectasis. *Annals of the American Thoracic Society* 2015;12:1602-1611.
11. Taylor SL, Rogers GB, Chen AC, Burr LD, McGuckin MA, Serisier DJ. Matrix metalloproteinases vary with airway microbiota composition and lung function in non-cystic fibrosis bronchiectasis. *Annals of the American Thoracic Society* 2015;12:701-707.
12. Chalmers JD, McDonnell MJ, Rutherford R, Davidson J, Finch S, Crichton M, Dupont L, Hill AT, Fardon TC, De Soyza A, Aliberti S, Goeminne P. The generalizability of bronchiectasis randomized controlled trials: A multicentre cohort study. *Respir Med* 2016;112:51-58.

13. Chalmers JD, Goeminne P, Aliberti S, McDonnell MJ, Lonni S, Davidson J, Poppelwell L, Salih W, Pesci A, Dupont LJ, Fardon TC, De Soyza A, Hill AT. The bronchiectasis severity index. An international derivation and validation study. *Am J Respir Crit Care Med* 2014;189:576-585.
14. Rogers GB, Zain NM, Bruce KD, Burr LD, Chen AC, Rivett DW, McGuckin MA, Serisier DJ. A novel microbiota stratification system predicts future exacerbations in bronchiectasis. *Annals of the American Thoracic Society* 2014;11:496-503.
15. Nguyen LD, Viscogliosi E, Delhaes L. The lung mycobiome: An emerging field of the human respiratory microbiome. *Frontiers in microbiology* 2015;6:89.