University of Dundee

Pain and the global burden of disease

Rice, Andrew S. C.; Smith, Blair H.; Blyth, Fiona M.

Published in:
Pain

DOI:
[10.1097/j.pain.0000000000000454](https://doi.org/10.1097/j.pain.0000000000000454)

Publication date:
2016

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Rice, A. S. C., Smith, B. H., & Blyth, F. M. (2016). Pain and the global burden of disease. *Pain*, 157(4), 791-796.
<https://doi.org/10.1097/j.pain.0000000000000454>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

PAIN

Pain and the global burden of disease

--Manuscript Draft--

Manuscript Number:	PAIN-D-14-14328R1
Article Type:	Commentary (INVITED ONLY)
Section/Category:	Clinical Epidemiology
Keywords:	Global burden of disease; Pain; epidemiology
Corresponding Author:	Andrew SC Rice, MB BS MD FRCA FFPMRCA Imperial College London London, UNITED KINGDOM
First Author:	Andrew SC Rice, MB BS MD FRCA FFPMRCA
Order of Authors:	Andrew SC Rice, MB BS MD FRCA FFPMRCA
	Blair H Smith, MB ChB, MD, MEd, FRCGP, FRCP Edin
	Fiona M Blyth, MB BS, MPH, FAFPHM, PhD

This is the accepted, peer reviewed version of an article published in *PAIN*. The final published version is available online: <http://dx.doi.org/10.1097/j.pain.0000000000000454>

Pain and the Global Burden of Disease

Andrew S.C. Rice¹, Blair H. Smith², Fiona M. Blyth³

1. Andrew SC Rice MB BS, MD, FRCA, FFPMRCA, Professor of Pain Research, Department of Surgery and Cancer, Imperial College, Chelsea and Westminster Hospital campus, 369 Fulham Road, London SW10 9NH, United Kingdom

2. Blair H. Smith MB ChB, MD, MEd, FRCGP, FRCP Edin. Professor of Population Health Science, University of Dundee, Ninewells Hospital and Medical School, Kirsty Semple Way, Dundee DD2 4BF, United Kingdom.

3. Fiona M Blyth, MB BS, MPH, FAFPHM, PhD, University of Sydney Centre for Education and Research on Ageing, Concord Repatriation General Hospital, Hospital Road, Concord NSW 2139, Australia.

1. Introduction

From time to time a major global research publication appears which has profound implications for the pain research and clinical management community. The purpose of this article is to draw to the attention of PAIN's readership one such article, The Global Burden of Disease (GBD) Study 2013 which was recently published in the *Lancet* [26]. It comprises a comprehensive and extensive assessment of the global burden of disease down to country level. The GBD Study also documents changes in this burden over a 23 year period ending in 2013. Looking forward, there are plans for these data to be updated on an annual basis.

Here, we summarise, interpret and contextualise the key aspects of the GBD Study 2013 pertaining to pain. What is striking, and indeed highlighted in an accompanying commentary [4], about the morbidity and disability burden data is the high prominence of pain, and diseases associated with pain, as a global cause of disability in both the developed and developing world. Furthermore, an important implication is that whilst some diseases have decreased in prevalence since 1990 (most notably some associated with acute infectious diseases), those burdens associated with chronic pain are in many cases increasing. This is at least partly because of powerful demographic forces such as the aging of populations in the developed world and similarly increased survival into middle and old age in the developing world, resulting both in growing population size and rising prevalence of chronic diseases. In GBD 2010 40% of the overall disease burden in developing countries was due to communicable diseases, 49% to non-communicable diseases (NCDs), and 11% to injuries, and the poorest people in the world were affected at younger ages (40 and under) [27]. In developing countries populations are ageing at a much faster rate than occurred in developed countries: in the UK it took 45 years for the proportion of the population aged over 65 to double from 7% to 14% while it is predicted to take only 14 years in Columbia. To quote Alex Kalache [14] "The developed world became rich before it became old. In developing countries, people are becoming old before they become rich".

2. What did they measure, and how?

The GBD Study 2013 was a major project [26], the paper having 679 co-authors, 58 pages and 1,868 pages of online appendix. Data on 301 diseases and injuries, and 2,337 of their sequelae were drawn from 188 countries, providing systematic evidence on disease burden at global, regional and national levels.

For acute conditions (<3 months' duration), the incidence was presented (number of new cases in 2013 per 100,000 population); for chronic conditions, prevalence was presented (total number of individuals with the condition in 2013, and the number of individuals per 100,000 population). In addition, the impact of the diseases and their sequelae was calculated as 'Years Lived with Disability' (YLDs) – the prevalence multiplied by a rigorously derived disability weighting factor that represents the public's assessment of the severity of health loss.

The methods are clearly described in the paper and were developed from those used in the 2010 GBD study [27], and results were compared with the re-analysed results of the first GBD study, which was in 1990.

3. The global burden of disease – a painful picture

The GBD Study showed that only a small fraction of the world's population (4.3%) was free of disease, injury or sequelae. Because of increases in world population and life expectancy, the total number of individuals suffering almost every condition increased, and the prevalence of multi-morbidity also increased. Furthermore, mortality was found to be declining faster than disease prevalence and disability, with a resultant growth in the overall burden of disease. In other words, while life expectancy has increased, healthy life expectancy has not kept pace.

The commonest acute conditions, by far, were upper respiratory infections with in excess of 18.7 billion cases in 2013, increased by 38% in number since 1990, and 6.5% in age-standardised rate. Most of the commonest acute conditions were infectious diseases and short-term injuries, but tooth pain due to permanent caries was the 5th commonest acute condition, with more than 200 million cases estimated in 2013 (an absolute increase of 36% since 1990, and a rate increase of 1.4%).

A total of 59 chronic conditions were estimated each to have a global prevalence in excess of 1%, with eight conditions affecting more than 10% of the global population. The commonest of these was asymptomatic dental caries (2.4 billion individuals, rising from 1.7 billion in 1990). Second commonest (and therefore the commonest symptomatic chronic condition) was recurrent tension-type headaches (1.6 billion in 2013, up from 1.1 billion in 1990); the age-standardised rate of this condition was unchanged in the comparison period. Seven of the commonest chronic conditions were found to be primary pain conditions – migraines were the 7th commonest, low back pain was 12th, neck pain was 19th, 'other musculoskeletal conditions' were 32nd, osteoarthritis 33rd, and urolithiasis was 57th. Although these all demonstrated large increases in total numbers of individuals affected, the age-standardised rates of chronic conditions were mostly either unchanged or only slightly reduced in comparison with 1990. Only five conditions demonstrated an increase in rate, including 'other musculoskeletal conditions', for which a 6.6% increase in rate was estimated, and urolithiasis (20.1% increase).

Probably, though, the most important picture of disease burden is represented by the YLDs, which take both the prevalence and severity into account. YLDs describe the impact of conditions on individuals, and by implication on their communities and societies, and the health and social services which must provide for them. Many of the most prevalent conditions (e.g. asymptomatic dental caries) are found to contribute less to YLDs, because of relatively low disability weightings. It is therefore in examining causes and rates of YLDs around the world that we find chronic pain to be the most important condition – globally, regionally, and in every individual country. It should be noted that for some regions (especially developing country areas such as Sub-Saharan Africa) there were no or scant primary data and the YLDs were estimated from the population structure and

weighted prevalence rates from other regions. There is still a great need for primary data from these areas to improve burden estimation.

Globally, the single greatest cause of YLDs was chronic low back pain, which was estimated to cause more than 146 million YLDs in 2013, an increase of 61% since 1990 (with no significant change in age-standardised rate). This compares with the second greatest cause – major depressive disorder – which contributed in excess of 51 million YLDs (increased in number by 53% since 1990, and in rate by 4.7%). Chronic neck pain was the 4th greatest cause (>34 million YLDs); migraine was 6th (>28 million YLDs); ‘other musculoskeletal disorders’ was 10th (>22 million YLDs); osteoarthritis was 13th (>12 million YLDs); and medication overuse headache was 18th (>9 million YLDs). The main four chronic pain conditions featured in the top ten causes of YLDs both in developed countries and in developing countries.

Indeed, chronic back pain and chronic neck pain featured in the top ten causes of YLDs in every country examined, and migraine was also ranked in the top ten everywhere other than Sub-Saharan Africa. Musculoskeletal disorders together with fractures and soft tissue injuries accounted for 20.8% of all YLDs globally in 2013. Mental health conditions were also important causes of YLD, with major depressive disorder appearing in every country’s list of top ten causes, and anxiety featuring prominently; both of these are important co-morbidities with chronic pain. Regions with notable variation from the most important global causes of YLDs were Africa and the Middle East. In North Africa and the Middle East, opioid use disorder was listed as the 8th most important cause (and the most important in Qatar). Data for every country and condition assessed are available as supplementary tables in the paper’s extensive online appendix.

The study was designed to assess the global burden of disease, and cannot provide definitive explanations for the findings. However, the authors postulate that the main reasons for the increase in YLDs caused by these chronic pain conditions are related to ageing populations (rates of YLDs are declining much more slowly than mortality rates), increasing obesity and reducing levels of physical activity. Rates of obesity, road traffic accidents, other types of injury are all increasing in the developing world. They further note that there is still little policy discussion around their prevention and management, despite their burden being highlighted as larger than expected in the GBD 2010 Study [27].

4. The burden of “hidden” pain

The preceding section documents how chronic pain is clearly the most important current and future cause of morbidity and disability across the world, with large recent increases in both the number of individuals affected and years lost to disability, coupled with static or rising prevalence rates. However, even this picture actually under-estimates the total burden imposed by chronic pain. Here we discuss a number of disparate exemplar conditions where pain can be a major feature, although it was not possible to ascertain the exact contribution of pain to the overall YLD from the published GBD data.

The diabetes pandemic is well documented and diabetes was ranked as the 7th most important cause of YLDs globally, rising from 10th in the 1990 survey – a percentage change of 136% and now accounting for 29.5 million YLD. Up to 26% of type 2 diabetics suffer from painful diabetic neuropathy, representing a population prevalence of 0.8% [25].

With respect to infectious diseases, in Sub-Saharan Africa, diseases arising from HIV/AIDS (other than mycobacterial infection) were listed as the 4th most important cause of YLDs, and were top of the list in six countries. The number of YLDs caused by HIV-related diseases, standing at >4 million in 2013, had increased by more than 300%, with an age-standardised rate increase of 172% since 1990. HIV/AIDS is associated with many causes of pain, but one of the most relevant, especially as the trend towards world-wide access to antiretroviral drugs continues, is HIV-associated sensory neuropathy (HIV-SN). The anticipated decline in HIV-SN with withdrawal of neurotoxic antiretroviral drugs is not yet evident and this peripheral neuropathy still afflicts between 27% and 57% of people living with HIV of whom between 38% and 90% experience neuropathic pain [8; 11; 21]. Therefore, HIV-SN is one of the most frequent manifestations of HIV in people who are adequately treated with antiretroviral drugs. The burden of neuropathic pain in HIV-SN is further increased as it is peculiarly resistant to analgesic therapies which are effective in other painful neuropathies [9; 12; 22; 23].

Varicella and herpes zoster were globally associated with 198,000 YLDs, having increased by 48% since 1990, although there was a slight decline when the age-standardised change was taken into account. However, it is not possible to ascertain to what extent this is attributable to the main chronic pain complication of zoster, postherpetic neuralgia. Whilst postherpetic neuralgia would be expected to become more prevalent with an ageing population, this may well be mitigated by the ongoing introduction of zoster vaccination in a number of countries [20].

Although now a comparably rare disease globally, leprosy is still prevalent in parts of the developing world and accounted for some 40,000 YLDs, these having increased by 73% since 1990 (age standardised increase 6%). Some of this burden is probably attributable to neuropathic pain, which chronically afflicts some 20-25% of leprosy patients even long after mycobacteriological cure, because of persisting nerve damage and new immune mediated neuritis reactions [13; 18; 24].

Overall cancer accounted for 6.8 million YLDs, an increase of 83% (age adjusted 8.5 million YLDs). Neuropathic or mixed pain has been found to affect 20-40% of people with cancer [5]. These findings are importantly contextualised by considering ongoing gains in cancer survival for many common cancers.

We are increasingly aware of the importance of neuropathic pain in neurological disorders and it is notable, for example, that multiple sclerosis and Parkinson's disease accounted for 755,000 and 695,000 YLDs respectively, both having increased since 1990.

Sickle cell disorders and trait accounted for 342,000 and 1.4 million YLDs respectively.

Disability following war and conflict were ranked as the top causes of YLDs in Cambodia, Nicaragua, and Rwanda and second in Vietnam. Modern munitions dictated that limb trauma and amputation (and therefore post amputation pain) were one of the major survivable injuries of the First World War [10] and this continues to be a major feature of modern day conflicts, including those inflicted on civilians by landmines long after active conflicts have finished [3; 7; 16; 17; 19].

Finally, the contribution of harms associated with analgesic treatments to the global burden of disease cannot be ignored, although it is difficult to ascertain the exact impact of prescribed, as opposed to misused, therapies. Overall the adverse effects of medical treatment globally accounted for 201,000 YLDs. Nevertheless, it is sobering to note in passing that opioid use disorders accounted for 5.8 million YLDs and medication overuse headache for 9.8 million, with both having risen since 1990. Cannabis is often advocated for the treatment of pain, particularly neuropathic pain, despite a distinct lack of evidence for efficacy and unknown risks of harm [12] – cannabis use disorders accounted for 396,000 YLDs.

5. Implications for pain research, practice and policy.

GBD 2010 [27] and 2013 [26] changed our understanding of the huge global burden of disease that is driven by pain conditions, and also highlighted areas where the pain community can contribute to further improvements in how this burden is accurately assessed and, ultimately, reduced. This global challenge requires an agenda and action plan that must span research, policy and practice, and actively address the problems pertaining both to developed and to developing countries.

The over-arching priority is global recognition of chronic pain as a distinct clinical entity, rather than merely as a symptom of other conditions, with funding specifically identified and allocated to address this most disabling of conditions across a diverse range of diseases and settings.

5.1 Research

Pain research has historically been – and continues to be - a low priority for research funders; for example in 2007 only 0.61% of the United States National Institutes of Health budget was allocated to pain research [1]. Funding for clinical services has been similarly problematic.

Experience to date with estimating the global burden of pain has identified significant gaps in our basic understanding of how pain affects populations, ranging from the prevalence of common pain conditions to the cultural dimensions of pain burden (such as stigma and shame) in different societies.

We need population studies that measure burden of pain (occurrence and impact) in countries where no primary studies have been done, most notably in developing countries and regions such as Sub-Saharan Africa. This will first require the development of standardised and robust measures for use in population studies of common pain-associated conditions such as neck pain. Then we need better characterisation of the natural history and main trajectories of common pain conditions so that incidence and remission rates, as well as prevalence rates and the proportions who experience pain-related disability at all levels of severity, can be better understood. There is a need for better measurement of neuropathic and disease-associated pain, and a more consistent approach to characterising pain as a sequel of injury (for example, spinal cord injury), surgery or disease.

New population-level studies are also needed to assess how the cross-cutting risk factors for chronic diseases – such as poverty, diet, obesity, physical activity levels and other common exposures associated with chronic diseases – interplay with pain within an explicit public health framework [6]. Linked to this will be the identification and implementation of policy and political approaches to reducing the burden of chronic pain. These are at least as important as clinical interventions, given the association of chronic pain with poverty and secular demographic trends in the societies affected.

5.2 Practice

A number of areas of clinical practice require scrutiny in the light of the high visibility of pain in the GBD picture. However, those of us involved in clinical pain management should also recognise that our efforts are likely to be more effective if they are conducted not in isolation, but rather imbedded as part of an overall multi-disciplinary rehabilitation programme aimed at returning individuals to functioning as productive members of their societies.

To directly address the burden of disease, research must continue to identify and evaluate new clinical interventions for chronic pain, including its prevention. These will include a range of modalities and/or be multi-modal, including pharmacological, physical, psychological, social and

1 surgical. Nevertheless, for effective translation of new and existing evidence on prevention and
2 treatment into clinical practice it is arguably time for the international pain community to become
3 more active in adopting emerging key factors in successful translation and implementation from the
4 rapidly developing field of implementation science.

5
6 However, changing practice is not just about introduction of novel interventions, but also,
7 importantly, implementation of existing evidence-based effective, and cost-effective, treatments for
8 chronic pain throughout community, primary care and specialist care settings (see for example:
9 Scottish Intercollegiate Guideline Network guidelines (SIGN) [2]. These include many interventions
10 that are relatively inexpensive (e.g. self-management; physical activity) and that are suitable for
11 delivery to large numbers of people in low resource settings. Associated with this is the development
12 of “minimum” and “optimum” standards of pain service provision appropriate to each country,
13 including staffing levels and increased local availability of appropriate drugs on the WHO Essential
14 Medicines Lists [15]. International review and comparisons of current service provision in
15 comparison with such standards will be essential, as will strategies to bridge the gaps.

16
17 Finally, as ever, appropriate education of healthcare professionals and managers, patients, and
18 society in the mechanisms, prevention and management of chronic pain is vital.

19 20 21 22 *5.3. Implications for policy*

23
24 We need to articulate a coherent and integrated framework for pain conditions, to be used within
25 the GBD and other policy contexts. This should to replace the somewhat fragmented and
26 conceptually clouded approach to date in which some, but not all, common regional pain conditions
27 are included, where pain is not explicitly included as an important sequel of other conditions, and
28 the role of pain as a risk factor for other conditions is not well-conceptualised. Despite the current
29 approach, enough is known about the global pain burden to warrant significant responses from
30 global and national agencies. The WHO’s focus on non-communicable diseases remains on
31 cardiovascular and respiratory diseases, diabetes and cancer. This is noteworthy, as there is little to
32 suggest to date that there has been significant institutional response to the GBD evidence about
33 pain burden. It is incumbent on the pain community to engage with the ongoing GBD initiative, and
34 to advocate globally, for action and progress on the universal burden of pain.

35
36 The difficulties of undertaking research and interventions in resource-poor countries are substantial
37 and are likely to require active engagement with the WHO, the United Nations, and potentially
38 partnering with other professional associations. For example the Bone and Joint Decade
39 (www.bjd.org.au) notably played a key role in supporting the work of the expert reference group for
40 musculoskeletal conditions in GBD 2010. Much of this research is likely to be done by researchers
41 whose primary expertise is not in pain research (for example, studies where pain is the sequel of
42 other conditions), again pointing to the importance of collaborations and partnerships. Another
43 form of engagement would be for members of the international pain community with the relevant
44 expertise to have direct input through the GBD collaborators network (www.healthdata.org/gbd).

45
46 In conclusion, GBD 2013 [26] has changed our understanding of the huge burden of disease and
47 disability that is driven by pain-associated conditions and places chronic pain firmly at the crux of
48 healthcare provision - this has major societal implications for research, planning, delivery and
49 resource allocation. Looking forward and empowered by this knowledge, it is incumbent on the
50 global pain research and clinical communities to recognise and address this challenge, interact with
51 and influence the GBD programme and its methods, and advocate for a better and more detailed
52 understanding of pain as the major component of global disease burden. *Carpe diem!*

References

- [1] Relieving pain in America: a blueprint for transforming prevention, care, education, and research.: Institute of Medicine, 2011.
- [2] SIGN 136: Management of Chronic Pain. Scottish Intercollegiate Guidelines Network. Edinburgh: Scottish Intercollegiate Guidelines Network, 2013.
<http://www.sign.ac.uk/guidelines/fulltext/136/index.html>
- [3] Aldington D, Small C, Edwards D, Ralph J, Woods P, Jagdish S, Moore RA. A survey of post-amputation pains in serving military personnel. *Journal of the Royal Army Medical Corps* 2014;160(1):38-41.
- [4] Atun R. Transitioning health systems for multimorbidity. *The Lancet* 2015;386(9995):721-722.
- [5] Bennett MI, Rayment C, Hjermstad M, Aass N, Caraceni A, Kaasa S. Prevalence and aetiology of neuropathic pain in cancer patients: a systematic review. *Pain* 2012;153(2):359-365.
- [6] Blyth FM, Van Der Windt DA, Croft PR. Chronic Disabling Pain: A Significant Public Health Problem. *Am J Prev Med* 2015;49(1):98-101.
- [7] Carlen PL, Wall PD, Nadvorna H, Steinbach T. Phantom limbs and related phenomena in recent traumatic amputations. *Neurology* 1978;28(3):211-217.
- [8] Cherry CL, Kamerman PR, Bennett DLH, Rice ASC. HIV-associated sensory neuropathy: still a problem in the post-stavudine era? *Future Virology* 2012;7(9):1-6.
- [9] Dinat N, Marinda E, Moch S, Rice AS, Kamerman PR. Randomized, Double-Blind, Crossover Trial of Amitriptyline for Analgesia in Painful HIV-Associated Sensory Neuropathy. *PLoS One* 2015;10(5):e0126297.
- [10] Edwards DS, Mayhew ER, Rice ASC. "Doomed to go in company with miserable pain": surgical recognition and treatment of amputation-related pain on the Western Front during World War 1. *The Lancet* 2014;384(9955):1715-1719.
- [11] Ellis RJ, Rosario D, Clifford DB, McArthur JC, Simpson D, Alexander T, Gelman BB, Vaida F, Collier A, Marra CM, Ances B, Atkinson JH, Dworkin RH, Morgello S, Grant I. Continued high prevalence and adverse clinical impact of human immunodeficiency virus-associated sensory neuropathy in the era of combination antiretroviral therapy: the CHARTER Study. *ArchNeurol* 2010;67(5):552-558.
- [12] Finnerup NB, Attal N, Haroutounian S, McNicol E, Baron R, Dworkin RH, Gilron I, Haanpää M, Hansson P, Jensen TS, Kamerman PR, Lund K, Moore A, Raja SN, Rice ASC, Rowbotham M, Sena E, Siddall P, Smith BH, Wallace M. Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis. *The Lancet Neurology* 2015;14(2):162-173.
- [13] Haroun OMO, Hietaharju A, Bizuneh E, Tesfaye F, Brandsma JW, Haanpää M, Rice ASC, Lockwood DNJ. Investigation of neuropathic pain in treated leprosy patients in Ethiopia: A cross-sectional study. *Pain* 2012;153(8):1620-1624.
- [14] Kalache A. Rural aging: reduction of existing inequities. Keynote address, June 10, 2000. *J Rural Health* 2001;17(4):312-313.
- [15] Kamerman PR, Wadley AL, Davis KD, Hietaharju A, Jain P, Kopf A, Meyer A-C, Raja SN, Rice ASC, Smith BH, Treede R-D, Wiffen PJ. World Health Organization essential medicines lists: where are the drugs to treat neuropathic pain? *PAIN* 2015;156(5):793-797.
- [16] Lacoux P, Ford N. Treatment of neuropathic pain in Sierra Leone. *Lancet Neurol* 2002;1(3):190-195.
- [17] Lacoux PA, Crombie IK, Macrae WA. Pain in traumatic upper limb amputees in Sierra Leone. *Pain* 2002;99(1-2):309-312.

- [18] Lasry-Levy E, Hietaharju A, Pai V, Ganapati R, Rice ASC, Haanpaa M, Lockwood DNJ. Neuropathic Pain and Psychological Morbidity in Patients with Treated Leprosy: A Cross-Sectional Prevalence Study in Mumbai. *Plos Neglected Tropical Diseases* 2011;5(3).
- [19] Le Feuvre P, Aldington D. Know pain know gain: proposing a treatment approach for phantom limb pain. *Journal of the Royal Army Medical Corps* 2014;160(1):16-21.
- [20] Oxman MN, Levin MJ, Johnson GR, Schmader KE, Straus SE, Gelb LD, Arbeit RD, Simberkoff MS, Gershon AA, Davis LE, Weinberg A, Boardman KD, Williams HM, Zhang JH, Peduzzi PN, Beisel CE, Morrison VA, Guatelli JC, Brooks PA, Kauffman CA, Pachucki CT, Neuzil KM, Betts RF, Wright PF, Griffin MR, Brunell P, Soto NE, Marques AR, Keay SK, Goodman RP, Cotton DJ, Gnann JW, Jr., Loutit J, Holodniy M, Keitel WA, Crawford GE, Yeh SS, Lobo Z, Toney JF, Greenberg RN, Keller PM, Harbecke R, Hayward AR, Irwin MR, Kyriakides TC, Chan CY, Chan ISF, Wang WWB, Annunziato PW, Silber JL, the Shingles Prevention Study Group. A Vaccine to Prevent Herpes Zoster and Postherpetic Neuralgia in Older Adults. *The New England Journal of Medicine* 2005;352(22):2271-2284.
- [21] Phillips TJC, Brown M, Ramirez JD, Perkins J, Woldeamanuel YW, Williams ACdC, Orengo C, Bennett DLH, Bodi I, Cox S, Maier C, Krumova EK, Rice ASC. Sensory, psychological, and metabolic dysfunction in HIV-associated peripheral neuropathy: A cross-sectional deep profiling study. *Pain* 2014;155(9):1846-1860.
- [22] Phillips TJC, Cherry CL, Cox S, Marshall SJ, Rice ASC. Pharmacological Treatment of Painful HIV-Associated Sensory Neuropathy: A Systematic Review and Meta-Analysis of Randomised Controlled Trials. *PLoS ONE* 2010;5(12):e14433.
- [23] Simpson DM, Rice AS, Emir B, Landen J, Semel D, Chew ML, Sporn J. A randomized, double-blind, placebo-controlled trial and open-label extension study to evaluate the efficacy and safety of pregabalin in the treatment of neuropathic pain associated with human immunodeficiency virus neuropathy. *Pain* 2014;155(10):1943-1954.
- [24] Thakur S, Dworkin RH, Haroun OM, Lockwood DN, Rice AS. Acute and chronic pain associated with leprosy. *Pain* 2015;156(6):998-1002.
- [25] van Hecke O, Austin SK, Khan RA, Smith BH, Torrance N. Neuropathic pain in the general population: a systematic review of epidemiological studies. *Pain* 2014;155(4):654-662.
- [26] Vos T, Barber RM, Bell B, Bertozzi-Villa A, Biryukov S, Bolliger I, Charlson F, Davis A, Degenhardt L, Dicker D, Duan L, Erskine H, Feigin VL, Ferrari AJ, Fitzmaurice C, Fleming T, Graetz N, Guinovart C, Haagsma J, Hansen GM, Hanson SW, Heuton KR, Higashi H, Kassebaum N, Kyu H, Laurie E, Liang X, Lofgren K, Lozano R, MacIntyre MF, Moradi-Lakeh M, Naghavi M, Nguyen G, Odell S, Ortblad K, Roberts DA, Roth GA, Sandar L, Serina PT, Stanaway JD, Steiner C, Thomas B, Vollset SE, Whiteford H, Wolock TM, Ye P, Zhou M, Ávila MA, Aasvang GM, Abbafati C, Ozgoren AA, Abd-Allah F, Aziz MIA, Abera SF, Aboyans V, Abraham JP, Abraham B, Abubakar I, Abu-Raddad LJ, Abu-Rmeileh NME, Aburto TC, Achoki T, Ackerman IN, Adelekan A, Ademi Z, Adou AK, Adsuar JC, Arnlov J, Agardh EE, Al Khabouri MJ, Alam SS, Alasfoor D, Albittar MI, Alegretti MA, Aleman AV, Alemu ZA, Alfonso-Cristancho R, Alhabib S, Ali R, Alla F, Allebeck P, Allen PJ, AlMazroa MA, Alsharif U, Alvarez E, Alvis-Guzman N, Ameli O, Amini H, Ammar W, Anderson BO, Anderson HR, Antonio CAT, Anwari P, Apfel H, Arsenijevic VSA, Artaman A, Asghar RJ, Assadi R, Atkins LS, Atkinson C, Badawi A, Bahit MC, Bakfalouni T, Balakrishnan K, Balalla S, Banerjee A, Barker-Collo SL, Barquera S, Barregard L, Barrero LH, Basu S, Basu A, Baxter A, Beardsley J, Bedi N, Beghi E, Bekele T, Bell ML, Benjet C, Bennett DA, Bensenor IM, Benzian H, Bernabe E, Beyene TJ, Bhala N, Bhalla A, Bhutta Z, Bienhoff K, Bikbov B, Abdulhak AB, Blore JD, Blyth FM, Bohensky MA, Basara BB, Borges G, Bornstein NM, Bose D, Boufous S, Bourne RR, Boyers LN, Brainin M, Brauer M, Brayne CEG, Brazinova A, Breitborde NJK, Brenner H, Briggs ADM, Brooks PM, Brown J, Brughra TS, Buchbinder R, Buckle GC, Bukhman G, Bulloch AG, Burch M, Burnett R, Cardenas R, Cabral NL, Nonato IRC, Campuzano JC, Carapetis JR, Carpenter DO, Caso V, Castaneda-Orjuela CA, Catala-Lopez F, Chadha VK, Chang J-C, Chen H, Chen W, Chiang PP, Chimed-Ochir O,

Chowdhury R, Christensen H, Christophi CA, Chugh SS, Cirillo M, Coggeshall M, Cohen A,
 Colistro V, Colquhoun SM, Contreras AG, Cooper LT, Cooper C, Cooperrider K, Coresh J,
 Cortinovis M, Criqui MH, Crump JA, Cuevas-Nasu L, Dandona R, Dandona L, Dansereau E,
 Dantes HG, Dargan PI, Davey G, Davitoiu DV, Dayama A, De la Cruz-Gongora V, de la Vega SF,
 De Leo D, del Pozo-Cruz B, Dellavalle RP, Deribe K, Derrett S, Des Jarlais DC, Dessalegn M,
 deVeber GA, Dharmaratne SD, Diaz-Torne C, Ding EL, Dokova K, Dorsey ER, Driscoll TR,
 Duber H, Durrani AM, Edmond KM, Ellenbogen RG, Endres M, Ermakov SP, Eshrati B,
 Esteghamati A, Estep K, Fahimi S, Farzadfar F, Fay DFJ, Felson DT, Fereshtehnejad S-M,
 Fernandes JG, Ferri CP, Flaxman A, Foigt N, Foreman KJ, Fowkes FGR, Franklin RC, Furst T,
 Futran ND, Gabbe BJ, Gankpe FG, Garcia-Guerra FA, Geleijnse JM, Gessner BD, Gibney KB,
 Gillum RF, Ginawi IA, Giroud M, Giussani G, Goenka S, Goginashvili K, Gona P, de Cosio TG,
 Gosselin RA, Gotay CC, Goto A, Gouda HN, Guerrant RI, Gughani HC, Gunnell D, Gupta R,
 Gupta R, Gutierrez RA, Hafezi-Nejad N, Hagan H, Halasa Y, Hamadeh RR, Hamavid H,
 Hammami M, Hankey GJ, Hao Y, Harb HL, Haro JM, Havmoeller R, Hay RJ, Hay S, Hedayati
 MT, Pi IBH, Heydarpour P, Hajar M, Hoek HW, Hoffman HJ, Hornberger JC, Hosgood HD,
 Hossain M, Hotez PJ, Hoy DG, Hsairi M, Hu H, Hu G, Huang JJ, Huang C, Huiart L, Hussein A,
 Iannarone M, Iburg KM, Innos K, Inoue M, Jacobsen KH, Jassal SK, Jeemon P, Jensen PN, Jha
 V, Jiang G, Jiang Y, Jonas JB, Joseph J, Juel K, Kan H, Karch A, Karimkhani C, Karthikeyan G,
 Katz R, Kaul A, Kawakami N, Kazi DS, Kemp AH, Kengne AP, Khader YS, Khalifa SEAH, Khan EA,
 Khan G, Khang Y-H, Khonelidze I, Kielsing C, Kim D, Kim S, Kimokoti RW, Kinfu Y, Kinge JM,
 Kissela BM, Kivipelto M, Knibbs L, Knudsen AK, Kokubo Y, Kosen S, Kramer A, Kravchenko M,
 Krishnamurthi RV, Krishnaswami S, Defo BK, Bicer BK, Kuipers EJ, Kulkarni VS, Kumar K,
 Kumar GA, Kwan GF, Lai T, Laloo R, Lam H, Lan Q, Lansingh VC, Larson H, Larsson A,
 Lawryniewicz AEB, Leasher JL, Lee J-T, Leigh J, Leung R, Levi M, Li B, Li Y, Li Y, Liang J, Lim S, Lin
 H-H, Lind M, Lindsay MP, Lipshultz SE, Liu S, Lloyd BK, Ohno SL, Logroscino G, Looker KJ,
 Lopez AD, Lopez-Olmedo N, Lortet-Tieulent J, Lotufo PA, Low N, Lucas RM, Lunevicius R,
 Lyons RA, Ma J, Ma S, Mackay MT, Majdan M, Malekzadeh R, Mapoma CC, Marcenes W,
 March LM, Margono C, Marks GB, Marzan MB, Masci JR, Mason-Jones AJ, Matzopoulos RG,
 Mayosi BM, Mazorodze TT, McGill NW, McGrath JJ, McKee M, McLain A, McMahon BJ,
 Meaney PA, Mehndiratta MM, Mejia-Rodriguez F, Mekonnen W, Melaku YA, Meltzer M,
 Memish ZA, Mensah G, Meretoja A, Mhimbira FA, Micha R, Miller TR, Mills EJ, Mitchell PB,
 Mock CN, Moffitt TE, Ibrahim NM, Mohammad KA, Mokdad AH, Mola GL, Monasta L,
 Montico M, Montine TJ, Moore AR, Moran AE, Morawska L, Mori R, Moschandreas J, Moturi
 WN, Moyer M, Mozaffarian D, Mueller UO, Mukaigawara M, Murdoch ME, Murray J, Murthy
 KS, Naghavi P, Nahas Z, Naheed A, Naidoo KS, Naldi L, Nand D, Nangia V, Narayan KMV, Nash
 D, Nejjari C, Neupane SP, Newman LM, Newton CR, Ng M, Ngalesoni FN, Nhung NT, Nisar MI,
 Nolte S, Norheim OF, Norman RE, Norrving B, Nyakarahuka L, Oh IH, Ohkubo T, Omer SB,
 Opio JN, Ortiz A, Pandian JD, Pabello CIA, Papachristou C, Park E-K, Parry CD, Caicedo AJP,
 Patten SB, Paul VK, Pavlin BI, Pearce N, Pedraza LS, Pellegrini CA, Pereira DM, Perez-Ruiz FP,
 Perico N, Pervaiz A, Pesudovs K, Peterson CB, Petzold M, Phillips MR, Phillips D, Phillips B,
 Piel FB, Plass D, Poenaru D, Polanczyk GV, Polinder S, Pope CA, Popova S, Poulton RG,
 Pourmalek F, Prabhakaran D, Prasad NM, Qato D, Quistberg DA, Rafay A, Rahimi K, Rahimi-
 Movaghar V, Rahman Su, Raju M, Rakovac I, Rana SM, Razavi H, Refaat A, Rehm J, Remuzzi
 G, Resnikoff S, Ribeiro AL, Riccio PM, Richardson L, Richardus JH, Riederer AM, Robinson M,
 Roca A, Rodriguez A, Rojas-Rueda D, Ronfani L, Rothenbacher D, Roy N, Ruhago GM, Sabin
 N, Sacco RL, Ksoreide K, Saha S, Sahathevan R, Sahraian MA, Sampson U, Sanabria JR,
 Sanchez-Riera L, Santos IS, Satpathy M, Saunders JE, Sawhney M, Saylan MI, Scarborough P,
 Schoettker B, Schneider IJC, Schwebel DC, Scott JG, Seedat S, Sepanlou SG, Serdar B, Servan-
 Mori EE, Shackelford K, Shaheen A, Shahraz S, Levy TS, Shangguan S, She J, Sheikhbahaei S,
 Shepard DS, Shi P, Shibuya K, Shinohara Y, Shiri R, Shishani K, Shiue I, Shrimme MG,
 Sigfusdottir ID, Silberberg DH, Simard EP, Sindi S, Singh JA, Singh L, Skirbekk V, Sliwa K, Soljak

M, Soneji S, Soshnikov SS, Speyer P, Sposato LA, Sreeramareddy CT, Stoeckl H, Stathopoulou VK, Steckling N, Stein MB, Stein DJ, Steiner TJ, Stewart A, Stork E, Stovner LJ, Stroumpoulis K, Sturua L, Sunguya BF, Swaroop M, Sykes BL, Tabb KM, Takahashi K, Tan F, Tandon N, Tanne D, Tanner M, Tavakkoli M, Taylor HR, Te Ao BJ, Temesgen AM, Have MT, Tenkorang EY, Terkawi AS, Theadom AM, Thomas E, Thorne-Lyman AL, Thrift AG, Tleyjeh IM, Tonelli M, Topouzis F, Towbin JA, Toyoshima H, Traebert J, Tran BX, Trasande L, Trillini M, Truelsen T, Trujillo U, Tsilimbaris M, Tuzcu EM, Ukwaja KN, Undurraga EA, Uzun SB, van Brakel WH, van de Vijver S, Dingenen RV, van Gool CH, Varakin YY, Vasankari TJ, Vavilala MS, Veerman LJ, Velasquez-Melendez G, Venketasubramanian N, Vijayakumar L, Villalpando S, Violante FS, Vlassov VV, Waller S, Wallin MT, Wan X, Wang L, Wang J, Wang Y, Warouw TS, Weichenthal S, Weiderpass E, Weintraub RG, Werdecker A, Wessells KRR, Westerman R, Wilkinson JD, Williams HC, Williams TN, Woldeyohannes SM, Wolfe CDA, Wong JQ, Wong H, Woolf AD, Wright JL, Wurtz B, Xu G, Yang G, Yano Y, Yenesew MA, Yentur GK, Yip P, Yonemoto N, Yoon S-J, Younis M, Yu C, Kim KY, Zaki MES, Zhang Y, Zhao Z, Zhao Y, Zhu J, Zonies D, Zunt JR, Salomon JA, Murray CJL. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *The Lancet* 2015;386(9995):743-800.

- [27] Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M, Shibuya K, Salomon JA, Abdalla S, Aboyans V, Abraham J, Ackerman I, Aggarwal R, Ahn SY, Ali MK, Alvarado M, Anderson HR, Anderson LM, Andrews KG, Atkinson C, Baddour LM, Bahalim AN, Barker-Collo S, Barrero LH, Bartels DH, Basanez MG, Baxter A, Bell ML, Benjamin EJ, Bennett D, Bernabe E, Bhalla K, Bhandari B, Bikbov B, Bin Abdulhak A, Birbeck G, Black JA, Blencowe H, Blore JD, Blyth F, Bolliger I, Bonaventure A, Boufous S, Bourne R, Boussinesq M, Braithwaite T, Brayne C, Bridgett L, Brooker S, Brooks P, Brugha TS, Bryan-Hancock C, Bucello C, Buchbinder R, Buckle G, Budke CM, Burch M, Burney P, Burstein R, Calabria B, Campbell B, Canter CE, Carabin H, Carapetis J, Carmona L, Cella C, Charlson F, Chen H, Cheng AT, Chou D, Chugh SS, Coffeng LE, Colan SD, Colquhoun S, Colson KE, Condon J, Connor MD, Cooper LT, Corriere M, Cortinovis M, de Vaccaro KC, Couser W, Cowie BC, Criqui MH, Cross M, Dabhadkar KC, Dahiya M, Dahodwala N, Damsere-Derry J, Danaei G, Davis A, De Leo D, Degenhardt L, Dellavalle R, Delossantos A, Denenberg J, Derrett S, Des Jarlais DC, Dharmaratne SD, Dherani M, Diaz-Torne C, Dolk H, Dorsey ER, Driscoll T, Duber H, Ebel B, Edmond K, Elbaz A, Ali SE, Erskine H, Erwin PJ, Espindola P, Ewoigbokhan SE, Farzadfar F, Feigin V, Felson DT, Ferrari A, Ferri CP, Fevre EM, Finucane MM, Flaxman S, Flood L, Foreman K, Forouzanfar MH, Fowkes FG, Franklin R, Fransen M, Freeman MK, Gabbe BJ, Gabriel SE, Gakidou E, Ganatra HA, Garcia B, Gaspari F, Gillum RF, Gmel G, Gosselin R, Grainger R, Groeger J, Guillemin F, Gunnell D, Gupta R, Haagsma J, Hagan H, Halasa YA, Hall W, Haring D, Haro JM, Harrison JE, Havmoeller R, Hay RJ, Higashi H, Hill C, Hoen B, Hoffman H, Hotez PJ, Hoy D, Huang JJ, Ibeanusi SE, Jacobsen KH, James SL, Jarvis D, Jasrasaria R, Jayaraman S, Johns N, Jonas JB, Karthikeyan G, Kassebaum N, Kawakami N, Keren A, Khoo JP, King CH, Knowlton LM, Kobusingye O, Koranteng A, Krishnamurthi R, Laloo R, Laslett LL, Lathlean T, Leasher JL, Lee YY, Leigh J, Lim SS, Limb E, Lin JK, Lipnick M, Lipshultz SE, Liu W, Loane M, Ohno SL, Lyons R, Ma J, Mabweijano J, MacIntyre MF, Malekzadeh R, Mallinger L, Manivannan S, Marcenes W, March L, Margolis DJ, Marks GB, Marks R, Matsumori A, Matzopoulos R, Mayosi BM, McAnulty JH, McDermott MM, McGill N, McGrath J, Medina-Mora ME, Meltzer M, Mensah GA, Merriman TR, Meyer AC, Miglioli V, Miller M, Miller TR, Mitchell PB, Mocumbi AO, Moffitt TE, Mokdad AA, Monasta L, Montico M, Moradi-Lakeh M, Moran A, Morawska L, Mori R, Murdoch ME, Mwaniki MK, Naidoo K, Nair MN, Naldi L, Narayan KM, Nelson PK, Nelson RG, Nevitt MC, Newton CR, Nolte S, Norman P, Norman R, O'Donnell M, O'Hanlon S, Olives C, Omer SB, Ortblad K, Osborne R, Ozgediz D, Page A, Pahari B, Pandian JD, Rivero AP, Patten SB, Pearce N, Padilla RP, Perez-Ruiz F, Perico N, Pesudovs K, Phillips D, Phillips MR,

Pierce K, Pion S, Polanczyk GV, Polinder S, Pope CA, 3rd, Popova S, Porrini E, Pourmalek F, Prince M, Pullan RL, Ramaiah KD, Ranganathan D, Razavi H, Regan M, Rehm JT, Rein DB, Remuzzi G, Richardson K, Rivara FP, Roberts T, Robinson C, De Leon FR, Ronfani L, Room R, Rosenfeld LC, Rushton L, Sacco RL, Saha S, Sampson U, Sanchez-Riera L, Sanman E, Schwebel DC, Scott JG, Segui-Gomez M, Shahraz S, Shepard DS, Shin H, ShivaKoti R, Singh D, Singh GM, Singh JA, Singleton J, Sleet DA, Sliwa K, Smith E, Smith JL, Stapelberg NJ, Steer A, Steiner T, Stolk WA, Stovner LJ, Sudfeld C, Syed S, Tamburlini G, Tavakkoli M, Taylor HR, Taylor JA, Taylor WJ, Thomas B, Thomson WM, Thurston GD, Tleyjeh IM, Tonelli M, Towbin JA, Truelsen T, Tsilimbaris MK, Ubeda C, Undurraga EA, van der Werf MJ, van Os J, Vavilala MS, Venketasubramanian N, Wang M, Wang W, Watt K, Weatherall DJ, Weinstock MA, Weintraub R, Weisskopf MG, Weissman MM, White RA, Whiteford H, Wiersma ST, Wilkinson JD, Williams HC, Williams SR, Witt E, Wolfe F, Woolf AD, Wulf S, Yeh PH, Zaidi AK, Zheng ZJ, Zonies D, Lopez AD, Murray CJ, AlMazroa MA, Memish ZA. Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. Lancet 2012;380(9859):2163-2196.

Conflict of interest statement: ASCR and BHS have no conflict to declare. FMB is an author of the GBD study