University of Dundee

The Geomycology of Elemental Cycling and Transformations in the Environment

Gadd, Geoffrey Michael

Published in:
Microbiology Spectrum

DOI:
[10.1128/microbiolspec.FUNK-0010-2016](https://doi.org/10.1128/microbiolspec.FUNK-0010-2016)

Publication date:
2017

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Gadd, G. M. (2017). The Geomycology of Elemental Cycling and Transformations in the Environment. *Microbiology Spectrum*, 5(1), 1-16. <https://doi.org/10.1128/microbiolspec.FUNK-0010-2016>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **The geomycology of elemental cycling and transformations in the environment**

3 Geoffrey Michael Gadd^{1,2}

5 ¹Geomicrobiology Group

6 School of Life Sciences

7 University of Dundee

8 Dundee

9 DD1 5EH

10 Scotland

11 UK

13 ²Laboratory of Environmental Pollution and Bioremediation

14 Xinjiang Institute of Ecology and Geography

15 Chinese Academy of Sciences

16 Urumqi 830011

17 People's Republic of China

19 Tel: +44 (0)1382 384767

20 e-mail: g.m.gadd@dundee.ac.uk

22 **Running Title: Geomycology of elemental cycling**

23 **Keywords:** Geomycology, geomicrobiology, elemental cycling, metal-mineral
24 transformations, bioremediation, biomineralization

SUMMARY

Geomicrobiology addresses the roles of microorganisms in geological and geochemical processes and geomycology is a part of this topic focussing on the fungi. Geoactive roles of fungi include organic and inorganic transformations important in nutrient and element cycling, rock and mineral bioweathering, mycogenic biomineral formation, and metal-fungal interactions. Lichens and mycorrhizas are significant geoactive agents. Organic matter decomposition is important for cycling of major biomass-associated elements, e.g. C, H, N, O, P and S as well as all other elements found in lower concentrations. Transformations of metals and minerals are central to geomicrobiology, and fungi effect changes in metal speciation, as well as mediate mineral formation or dissolution. Such mechanisms are components of biogeochemical cycles for metals as well as associated elements in biomass, soil, rocks and minerals, e.g. S, P, and metalloids. Fungi may have the greatest geochemical influence within the terrestrial environment. However, they are also important in the aquatic environment and significant components of the deep subsurface, extreme environments and habitats polluted by xenobiotics, metals and radionuclides. Applications of geomycology include metal and radionuclide bioleaching, biorecovery, detoxification, and bioremediation, and the production of biominerals or metal(loid) elements with catalytic or other properties. Adverse effects include biodeterioration of natural and synthetic materials, rock and mineral-based building materials (e.g. concrete), cultural heritage, metals, alloys and related substances, and adverse effects on radionuclide mobility and containment. The ubiquity and importance of fungi in the biosphere underlines the importance of geomycology as a conceptual framework encompassing the environmental activities of fungi.

INTRODUCTION

The significance of fungi in natural environments is extensive and profound. Their most obvious roles are as decomposers of organic materials, and as animal and plant pathogens and symbionts. It is therefore obvious that they are of major importance in the global carbon cycle through such activities and as important determinants of plant growth and productivity. However, their importance in terms of nutrient and element cycling greatly extends beyond this core activity and they are involved in the biogeochemical cycling of many other elements and substances, as well as many other related processes of environmental significance. The growing discipline of geomicrobiology addresses the roles of microorganisms in geological and geochemical processes (1,2), and geomycology can be considered to be a part of this topic that focusses on the fungi (3,4). The often clear demarcation between mycological and bacteriological research has ensured that the geoactive properties and significance of fungi have been unappreciated in wider geomicrobiological contexts. The range of prokaryotic metabolic diversity found in archaea and bacteria, including their abilities to use a variety of different terminal electron acceptors in respiration and effect redox transformations of many metal species (5,6), has also contributed to a narrow overall view of the significance of eukaryotic organisms in important biosphere processes. A recent collection of geomicrobiology review articles managed to completely exclude fungi (as well as algae), even to the extent of defining “microbes” as being only bacteria and archaea (7). Nevertheless, appreciation of fungi as agents of geochemical change is growing, and their significance is being discovered even in locations not usually regarded as prime fungal habitats, e.g. rocks, acid mine drainage, deep aquatic sediments, hydrothermal vents and the igneous oceanic crust (8-11). Their significance as bioweathering agents of rocks and minerals is probably better understood than bacterial roles (12), and this ability is of prime importance in the weathering of human structures in the built environment and cultural heritage (13-15). On the positive side, the geoactive properties of fungi can be used for

human benefit and several aspects may contribute to providing solutions to several important global challenges. Geomycology is relevant to reclamation and revegetation of polluted habitats, bioremediation, nuclear decommissioning and radionuclide containment, biorecovery of important elements, and the production of novel biomaterials. This chapter outlines important geoactive properties of fungi in relation to important environmental processes, their positive and negative applications and impact on human society.

THE FUNGAL HABITAT

Fungi are ubiquitous components of the microbial communities of any terrestrial environment, including such hostile habitats as the Arctic, hot deserts, and metal-rich and hypersaline soils (16). They are significant inhabitants of the aquatic environment as decomposers of organic matter but are also involved in other elemental cycles, e.g. Manganese oxidation (17). Fungi are ubiquitous in habitats polluted by xenobiotics, toxic metals and radionuclides, both terrestrial and aquatic, as well as leachates and other solid or liquid wastes (18). In such habitats, fungi may exhibit a variety of mechanisms that determine tolerance and survival, and which are also components of elemental cycles for pollutant elements (18). These “extreme” locations may also act as a reservoir of novel organisms with unusual properties. For example, acid mine drainage is now known to harbor fungal communities, the preponderance of earlier research on this habitat being devoted to prokaryotes, and some isolates possess unusual element bioaccumulation properties. New species include *Fodinomyces uranophilus* and *Coniochaeta fodinicola* from uranium mine locations that can bind mobile uranium (10,19) and a *Penidiella* sp. from an acidic abandoned mine location that was capable of accumulating rare earth elements such as dysprosium (Dy) (20). A global fungal role in biogeochemical cycling of the elements (e.g. C, H, N, O, P, S, metals, metalloids) is therefore obvious and interlinked with the ability to adopt a variety of

growth, metabolic and morphological strategies, adaptive capabilities to environmental extremes and, their symbiotic associations with animals, plants, algae and cyanobacteria (16,21,22). The ability of many fungi to grow oligotrophically by scavenging nutrients from the air and rainwater helps them survive on stone and rock surfaces which are considered to be an inhospitable environment (9,23). In addition, organic and inorganic residues on mineral surfaces or within cracks and fissures can act as nutrient sources in the sub-aerial rock environment (24). One of the most successful means for fungi to survive in the extreme sub-aerial environment is underpinned by their symbiotic associations with algae and cyanobacteria as lichens where the phototrophs provide a source of carbon and protection from light and irradiation (24,25). Lichens enable colonization of a wide range of environments including those at climatic extremes such as the Arctic and Antarctic, exposed rock surfaces and dry deserts.

In the deep subsurface, the research emphasis is mostly on prokaryotes, but the presence of fungi is now well known (11, 26-28). Fungi occur in abundance and high diversity in such varied environments as deep-sea sediments (29), hydrothermal vents (30,31), and methane cold-seeps (29,32). They are now also known as abundant inhabitants of the igneous oceanic crust which has consequently been described as the largest fungal habitat on Earth (11). Fungi seem to play an important ecological role in the igneous oceanic crust as they exist in symbiosis with chemolithotrophic prokaryotes, decompose organic matter, dissolve and form minerals, and therefore are involved in the cycling of elements (11,33,34). Fossilized microorganisms have been observed in drilled cores and dredged samples from the ocean floor with a majority of these findings representing fungi (34,35). These fungi existed in a close symbiotic-like relationship with two types of prokaryotes, which appeared to use the structural framework of the mycelium for their growth (34). It therefore seems clear that

geomycological processes are significant in a wide range of biosphere habitats, including those traditionally thought to be inimical to fungal growth and development (36).

ORGANIC MATTER DECOMPOSITION AND BIOGEOCHEMICAL CYCLING OF COMPONENT ELEMENTS

Organic matter decomposition is the attribute most commonly associated with fungi, and is a major contribution to global biogeochemistry as well as the spoilage of foodstuffs and organic materials (21,36). In fact, fungal processes represent a potential control point in the global carbon cycle (37). To some extent, research on this aspect of chemoorganotrophic metabolism has obscured the wider global implications of decomposition in terms of the cycling of other elements and nutrients, and also contributed to a lack of attention to fungal roles in wider geochemical cycles.

Most biogeochemical attention on fungi has been given to carbon and nitrogen cycles, and the ability of fungi to utilize a wide spectrum of organic compounds is well known. Simple compounds such as sugars, organic acids, and amino acids can easily be transported into the cell while more complex molecules are first broken down to smaller molecules by extracellular enzymes before cellular entry. Such compounds include natural substances such as cellulose, pectin, lignin, lignocellulose, chitin and starch to anthropogenic products like hydrocarbons, pesticides, and other xenobiotics (21,37). Organometals (compounds with at least one metal-carbon bond) can also be attacked by fungi. Degradation of organometallic compounds can be carried out by fungi either by direct enzymic action or by facilitating abiotic degradation, e.g. by alteration of external pH through metabolite excretion. Tributyltin oxide and tributyltin naphthenate may be degraded to mono- and dibutyltins by fungal action, inorganic Sn(II) being the final degradation product (38). Organomercury compounds may be

detoxified by conversion to Hg(II) by fungal organomercury lyase, the Hg(II) being subsequently reduced to Hg(0) by mercuric reductase, a system analogous to that found in mercury-resistant bacteria.

Some fungi have remarkable degradative properties and ligninolytic fungi, like the white rot *Phanerochaete chrysosporium*, can degrade many xenobiotics including aromatic hydrocarbons, chlorinated organics, polychlorinated biphenyls, nitrogen-containing aromatics and many other pesticides, dyes and xenobiotics (39,40). Such activities are of importance in polluted habitats and have been applied in bioremediation where ligninolytic fungi have been used to treat soil contaminated with substances like pentachlorophenol (PCP) and polynuclear aromatic hydrocarbons (PAHs) (21, 41-43). In many cases, xenobiotic-transforming fungi need additional utilizable carbon sources for co-metabolism because although capable of degradation, they cannot adequately utilize these substrates as an energy source. Inexpensive utilizable lignicellulosic wastes such as corn cobs, straw and sawdust can therefore be used as nutrient additions for enhanced pollutant degradation. Wood-rotting and other fungi have also received considerable attention for the bleaching of dyes and industrial effluents, and biotreatment of various agricultural wastes such as forestry, pulp and paper by-products, sugar cane bagasse, coffee pulp, sugar beet pulp, apple and tomato pulp, and cyanide (42).

As mentioned, fungi are highly important in the degradation of naturally-occurring complex molecules in terrestrial and aquatic habitats. Since around 95% of plant tissue is composed of carbon, hydrogen, oxygen, nitrogen, phosphorus and sulfur, decomposition activities of fungi are clearly important in relation to redistribution of these elements between organisms and environmental compartments. As well as C, H, O, N, P, and S, another 15 elements are typically found in living plant tissues - K, Ca, Mg, B, Cl, Fe, Mn, Zn, Cu, Mo, Ni, Co, Se,

Na, Si. However, all 90 or so naturally-occurring elements may be found in plants, mostly at low concentrations although this may be highly dependent on environmental conditions. These include toxic and inessential metals and metalloids including As, Hg, Pb and U. Some plants accumulate relatively high concentrations of metals like Ni and Cd. Plant metal concentrations may reflect environmental conditions and provide a bioindicator of toxic metal pollution or a metalliferous substrate. Metal-accumulating plants are also receiving attention in bioremediation (=phytoremediation).

Similar concepts of element cycling also relate to animal and microbial biomass. Animals also contain multiple elements in varying amounts. The human body (like other organisms) is mostly water and around 99% of the mass comprises oxygen, carbon, hydrogen, nitrogen, calcium and phosphorus. However, many other elements are present in lower amounts including substances taken up as contaminants in food and water. It follows that decomposition, degradative and pathogenic activities of fungi are linked to the redistribution and cycling of all these constituent elements on local and global scales. There must be a fungal component therefore in the biogeochemical cycles of virtually all biomass-associated elements (3,21). The release of elements and nutrient moieties through degradation makes them available to other microorganisms and plants, and also chemical interaction with the environmental pool of different chemical species.

FUNGAL INTERACTIONS WITH THE INORGANIC ENVIRONMENT: ROCKS, MINERALS AND METALS

Bioweathering

Rocks and minerals represent a vast reservoir of elements, many essential for life, and such elements must be released in forms that may be assimilated by the biota. These include essential metals as well as elements like S and P (3,44). Many important microbial processes are influenced by minerals including nutrient acquisition, cell adhesion and biofilm formation (45). Essential nutrients and environmental contaminants sorbed to mineral surfaces can be acquired or removed by organisms including metals and organic compounds (46,47). Other elements and nutrients may be released from minerals as a result of bioweathering, and fungi have notable activities in this context (3,16,24,48).

Bioweathering can be defined as the erosion, decay and decomposition of rocks and minerals mediated by living organisms. Fungi are well suited as geoactive weathering agents since they possess a variety of growth, metabolic and morphological strategies and can be resistant to a range of environmental extremes such as metal toxicity, UV radiation, and desiccation. Their mutualistic associations with plants (mycorrhizas), algae and cyanobacteria (lichens) are particularly significant geoactive agents (3,16,24,48). The ability of fungi to translocate water, ions and nutrients within the mycelial network is another important feature for exploiting heterogeneous environments (49-51).

Fungi appear to be ubiquitous components of the microbiota of all rocks, building stone and concrete, and have been reported from a wide range of rock types, e.g. limestone, marble, granite, sandstone, basalt, gneiss, dolerite and quartz, even from the most extreme environments (9,16,48). Rock surfaces may be subject to moisture deficit and nutrient limitation although many species can tolerate extremes of UV irradiation, salinity, pH, and water potential (16,24,25,48,52). Nutrients can be scavenged from the atmosphere and rainwater, and they also use organic and inorganic residues on surfaces or within cracks and

fissures, waste products of other microorganisms, decaying plants and insects, dust particles, aerosols and animal faeces as nutrient sources (24). Fungi may receive protection from environmental extremes by the presence of melanin pigments and mycosporines in their cell walls, and by the production of mucilaginous exopolymeric substances that may entrap inorganic particulates, e.g. clay minerals, providing further protection (9,53). Fungal interactions with rock-mineral substrates can result in dissolution and biodeterioration but also the formation of patinas, films, varnishes, and crusts (3,9). In soil, fungus-mineral interactions are also an integral component of environmental cycling of elements and nutrients (4,21).

Biomechanical deterioration of rocks and minerals can occur through hyphal penetration and burrowing into decaying material and along crystal planes in, e.g. calcitic and dolomitic rocks (3,24,54). Intracellular turgor pressure may be a significant factor in biomechanical disruption (55,56). Spatial exploration of the environment to locate and exploit new substrates is facilitated by a range of sensory responses that determine the direction of hyphal growth such as thigmotropism (or contact guidance) (57). Biochemical weathering of rocks and minerals can occur through excretion of geoactive metabolites (58,59), and this is believed to be a more significant process than mechanical degradation although a combination of mechanisms is often likely. This can result in pitting and etching of surfaces to complete dissolution of mineral grains (60-62). Bioweathering is a highly significant process and has direct consequences not only for rock and mineral dissolution, but the mobilization and immobilization of metals, nutrient release, and the formation of secondary minerals (2,3).

Metal mobilization

Metal mobilization from rocks, minerals, soil and other substrates can be a consequence of protonolysis, carbonic acid formation from respiratory CO₂, complexation by Fe(III)-binding

siderophores and other excreted metabolites, e.g. amino acids, phenolic compounds, and organic acids, and methylation (for e.g. Hg, Se and As) which can result in volatilization. Fungal-excreted carboxylic acids can attack mineral surfaces providing protons as well as a metal-chelating anion, e.g. citrate (58). Oxalic acid can leach metals that form soluble oxalate complexes, e.g. Al and Fe (63). Solubilization mechanisms can result in metal mobilization from toxic metal containing minerals, e.g. pyromorphite ($\text{Pb}_5(\text{PO}_4)_3\text{Cl}$), contaminated soil and other solid wastes (64-66). Fungi may also mobilize metals and attack mineral surfaces by redox transformations: Fe(III) and Mn(IV) solubility is increased by reduction to Fe(II) and Mn(II) respectively. Fungal reduction of Hg(II) to volatile elemental Hg(0) has also been recorded (67). As discussed earlier, metals may be mobilized from organic substrates during decomposition (21).

Metal immobilization

Fungi are effective accumulators of metals and related substances. Important mechanisms include biosorption to cell walls, pigments and exopolymers, intracellular transport, accumulation and sequestration, or bioprecipitation on and/or around hyphae (3,53,68-76). Living or dead fungal biomass are effective biosorbents for a variety of metals including Ni, Zn, Ag, Cu, Cd and Pb as well as actinides, e.g. U, Th, with a variety of functional groups being involved (24,53,76). The presence of chitin, and pigments like melanin, may enhance the ability of fungal biomass to act as a biosorbent. Fungal biomineralization processes lead to metal immobilization as biominerals or elemental forms, as described below (3).

Biomineralization

Biomineralization refers to the processes by which organisms form minerals. Biologically-induced mineralization (BIM) is where the organism modifies the local microenvironment

creating conditions amenable for extracellular chemical precipitation of mineral phases. The organism does not appear to control the biomineralization process in BIM while a great degree of control over biomineralization is exerted in biologically-controlled biomineralization (BCM), e.g. the complex cellular biomineral structures found in certain other eukaryotes like diatoms (77). Fungal biomineralization therefore usually refers to biologically-induced mineralization. This can result from the bioweathering mechanisms discussed previously such as redox transformations and metabolite excretion (78,79) and organic matter decomposition where released substances re-precipitate with metals in the microenvironment, and *vice versa* (4,21,80-82). As mentioned, fungal surfaces provide many reactive sites for sorption ($\equiv$ biosorption) and this can also lead to the formation of mineral precipitates (2,71,83).

Common Mineral and Biomineral Transformations by Fungi

Fungi are involved in many environmental mineral transformations at differing scales (84-86). They are clearly a very important group of geoactive organisms especially when considering their ubiquity and capacity for production of mineral-transforming metabolites, their symbiotic associations, and the aforementioned consequences of their major significance in organic matter decomposition (4,5).

Carbonates Insoluble carbonates may be broken down by fungal attack, usually the result of acid formation (87-89). Such activity is particularly evident on limestones and marble used in building construction, but can also occur in natural limestone (88,90). Fungal attack on carbonates (dolomites and limestones) can result in transformation of these substrates to dolomite ($\text{CaMg}(\text{CO}_3)_2$), glushinskite ($\text{MgC}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$), weddellite ($\text{CaC}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$), whewellite ($\text{CaC}_2\text{O}_4 \cdot \text{H}_2\text{O}$), and possibly struvite ($\text{NH}_4\text{MgPO}_4 \cdot 6\text{H}_2\text{O}$) (91).

Certain fungi can deposit calcium carbonate extracellularly (92-95). Calcite (CaCO_3) and calcium oxalate monohydrate (whewellite; $\text{CaC}_2\text{O}_4 \cdot \text{H}_2\text{O}$) was precipitated on hyphae of *Serpula himantoides* when grown in simulated limestone microcosms (93). Urease-positive fungi degrade urea liberating carbonate (96). This process results in the precipitation of metal-containing carbonates which provides a means of metal immobilization and biorecovery (94). Incubation of *Neurospora crassa* in urea-containing media resulted in the formation of calcite, as well as carbonates containing other metals. When a carbonate-laden *N. crassa* culture supernatant was mixed with CdCl_2 , the Cd was precipitated in the form of highly pure otavite (CdCO_3) (94). After incubation in media containing urea and CaCl_2 and/or SrCl_2 , *Pestalotiopsis* sp. and *Xepiculopsis graminea* (syn. *Myrothecium gramineum*), isolated from calcareous soil, precipitated calcite (CaCO_3), strontianite (SrCO_3), vaterite in different forms [CaCO_3 , $(\text{Ca}_x\text{Sr}_{1-x})\text{CO}_3$] and olekminskite [$\text{Sr}(\text{Sr,Ca})(\text{CO}_3)_2$] suggesting that urease-positive fungi could play an important role in the environmental fate, bioremediation or biorecovery of Sr or other metals and radionuclides that form insoluble carbonates (95). *Paecilomyces javanicus* mediated the formation of an unknown lead mineral phase after incubation in liquid media with lead shot. After 2 weeks incubation, precipitated mineral phase particles were found to contain plumbonacrite ($\text{Pb}_{10}(\text{CO}_3)_6\text{O}(\text{OH})_6$). However, after 4 weeks incubation, the lead particles that accumulated inside the fungal pellets were transformed into a white mineral phase composed of lead oxalate (PbC_2O_4), hydrocerussite ($\text{Pb}_3(\text{CO}_3)_2(\text{OH})_2$) and a new lead hydroxycarbonate mineral species, thus revealing novel steps in lead carbonation by fungi (97).

Oxalates Calcium oxalate is the most common form of oxalate in the environment, occurring as the dihydrate ($\text{CaC}_2\text{O}_4 \cdot 3\text{H}_2\text{O}$, weddellite) or the more stable monohydrate ($\text{CaC}_2\text{O}_4 \cdot \text{H}_2\text{O}$, whewellite) (59,79). Calcium oxalate can be associated with free-living, pathogenic and plant symbiotic fungi, and lichens, and is formed by precipitation of soluble calcium as the oxalate

(59,61,87,98,99). Fungal calcium oxalate can exhibit a variety of crystalline forms (tetragonal, bipyramidal, plate-like, rhombohedral or needles) (100). Calcium oxalate has an important influence on soil biogeochemistry, acting as a calcium reservoir, and can also influence phosphate availability. Fungi can produce many other metal oxalates on interacting with a variety of different metals and metal-bearing minerals, e.g. Ca, Cd, Co, Cu, Mg, Mn, Sr, Zn, Ni and Pb (3,59,64,79,101-103). The formation of toxic metal oxalates may contribute to fungal metal tolerance (53,102). In many arid and semi-arid regions, calcareous soils and near surface limestones (calcretes) are secondarily cemented with calcite (CaCO_3) and whewellite (calcium oxalate monohydrate, $\text{CaC}_2\text{O}_4\cdot\text{H}_2\text{O}$) and the presence of fungal filaments biomineralized with these substances has been reported (52). Calcium oxalate can also be degraded to calcium carbonate, and this may again cement pre-existing limestones (104). Other experimental work has demonstrated fungal precipitation of secondary calcite, whewellite, and glushkinskite ($\text{MgC}_2\text{O}_4\cdot 2\text{H}_2\text{O}$) (3,16,48,93). Fungal attack on a dolomitic and seawater substrate resulted in the formation of Ca-oxalates (weddellite, $\text{CaC}_2\text{O}_4\cdot 2\text{H}_2\text{O}$; whewellite, $\text{CaC}_2\text{O}_4\cdot\text{H}_2\text{O}$) and glushinskite ($\text{MgC}_2\text{O}_4\cdot 2\text{H}_2\text{O}$) (105).

Oxides Several fungi can oxidize Mn(II) to Mn(IV) O_2 including *Acremonium* spp. (17,106,107). Fungal oxidation is probably non-enzymatic in many cases although involvement of laccase and/or multicopper oxidases have been shown in ascomycetes (17,106). Non-enzymatic microbial Mn^{2+} oxidation may be effected through production of organic acids such as citrate, lactate, malate, gluconate, or tartrate. Some fungi can oxidize Mn(II) and Fe(II) in metal-bearing minerals such as siderite (FeCO_3) and rhodochrosite (MnCO_3) resulting in their precipitation as oxides (108). Manganese and iron oxides are major components (20–30%) along with clay (~60%) and various trace elements in desert varnish (9,108). Oxidation of Fe(II) and Mn(II) by fungi can lead to the formation of dark patinas on glass surfaces (109).

Manganese-reducing microbes may mobilize oxidized manganese, releasing it into the aqueous phase. Most of those fungi that reduce Mn(IV) oxides reduce them indirectly (non-enzymatically) with the likely mechanism being the production of metabolic products that act as reductants for Mn(IV) such as oxalate (1,103).

Phosphates Phosphorus occurs primarily as organic phosphate esters and inorganic forms, e.g. calcium, aluminium, and iron phosphates. Organic phosphates are broken down by phosphatases which liberate orthophosphate during the microbial decomposition of organic material. Fungi also mobilize orthophosphate from insoluble inorganic phosphates by producing acids or chelators, e.g. gluconate, citrate, oxalate, and lactate, which complex the metal resulting in dissociation. Phosphate-solubilization is very important in the plant mycorrhizosphere (110). Microbes can also play a role in the formation of phosphate minerals such as vivianite ($\text{Fe}_3(\text{PO}_4)_2 \cdot 8\text{H}_2\text{O}$), strengite ($\text{FePO}_4 \cdot 2\text{H}_2\text{O}$), and variscite ($\text{AlPO}_4 \cdot 2\text{H}_2\text{O}$). The orthophosphate may be derived from organic phosphate degradation while Fe or Al may arise from solubilization of other minerals. Such formation of phosphate minerals is probably most common in soil (1). Fungal biodeterioration of metallic lead can result in pyromorphite ($\text{Pb}_5[\text{PO}_4]_3\text{X}$ [X= F, Cl or OH]) formation (111-113). Many fungi can solubilize uranium oxides and depleted uranium and reprecipitate secondary uranium phosphate minerals, uramphite and/or chernikovite, which can encrust fungal hyphae to high accumulation values (73,74,114). These minerals appear capable of long-term U retention (73,74,114,115). *Aspergillus niger* and *Paecilomyces javanicus* precipitated U-containing phosphate biominerals when grown with an organic P source with the hyphal matrix acting to localize the uranium minerals. The uranyl phosphates identified included potassium uranyl phosphate hydrate ($\text{KPUO}_6 \cdot 3\text{H}_2\text{O}$), meta-ankoleite [$(\text{K}_{1.7}\text{Ba}_{0.2})(\text{UO}_2)_2(\text{PO}_4)_2 \cdot 6\text{H}_2\text{O}$], uranyl phosphate hydrate [$(\text{UO}_2)_3(\text{PO}_4)_2 \cdot 4\text{H}_2\text{O}$], meta-ankoleite ($\text{K}(\text{UO}_2)(\text{PO}_4) \cdot 3\text{H}_2\text{O}$), uramphite

($\text{NH}_4\text{UO}_2\text{PO}_4 \cdot 3\text{H}_2\text{O}$) and chernikovite [$(\text{H}_3\text{O})_2(\text{UO}_2)_2(\text{PO}_4)_2 \cdot 6\text{H}_2\text{O}$] (80). These organisms could also mediate lead bioprecipitation during growth on organic P substrates (81). These minerals were identified as pyromorphite ($\text{Pb}_5(\text{PO}_4)_3\text{Cl}$) which was only produced by *P. javanicus*, and lead oxalate (PbC_2O_4), which was produced by *A. niger* and *P. javanicus* (81). Several yeasts could also mediate lead bioprecipitation when utilizing an organic phosphorus-containing substrate (glycerol 2-phosphate, phytic acid) as sole phosphorus source. The minerals precipitated here included lead phosphate ($\text{Pb}_3(\text{PO}_4)_2$), pyromorphite ($\text{Pb}_5(\text{PO}_4)_3\text{Cl}$), anglesite (PbSO_4), and the lead oxides massicot and litharge (PbO). All yeasts examined produced pyromorphite, and most produced anglesite (82).

Silicates Silicates comprise 30% of all minerals and about 90% of the Earth's crust (116) (1,60,116). Many species of fungi play a role in the dissolution of silicates and therefore in the formation of clay minerals, and in soil and sediment formation (54, 87, 117-122). The presence of clay minerals can be a typical symptom of rock bioweathering by lichens and ectomycorrhizas (118,119). Bioweathering is mainly indirect, through the production of metabolites together with biomechanical effects (123,124). Geoactive metabolites may be excreted into the bulk phase but may also be produced by adhering organisms on silicate surfaces resulting in etching (125,126). After colonization of sheets of muscovite, a phyllosilicate mineral, by *Aspergillus niger*, dissolution was evident by a network of fungal "footprints" that reflected coverage by the mycelium (126). New biominerals resulted from fungal interactions with both zinc silicate and zinc sulfide, largely resulting from organic acid excretion. Zinc oxalate dihydrate was formed and mineral surfaces showed varying patterns of bioweathering and biomineral formation (127). Silicate dissolution may release limiting nutrients like bound P and Fe. In lichen bioweathering of silicates, calcium, potassium, iron, clay minerals and nanocrystalline aluminous iron oxyhydroxides become mixed with fungal

organic polymers (118), while biotite $(\text{K}(\text{Mg},\text{Fe}(\text{II}))_3\text{AlSi}_3\text{O}_{10}(\text{OH},\text{O},\text{F})_2)$ was penetrated by fungal hyphae along cleavages, partially converting it to vermiculite $((\text{Mg},\text{Fe}(\text{II}),\text{Al})_3(\text{Al},\text{Si})_4\text{O}_{10}(\text{OH})_2 \cdot 4\text{H}_2\text{O})$ (117). The fungal partner has also been reported to be involved in formation of secondary silicates, such as opal $(\text{SiO}_2 \cdot n\text{H}_2\text{O})$ and forsterite $(\text{Mg}_2\text{SiO}_4)$, in lichen thalli (128). The transformation rate of mica and chlorite to clay minerals was pronounced in ectomycorrhizosphere soil and probably a result production of organic acids and direct extraction of K^+ and Mg^{2+} by fungal hyphae (119). Fungal-clay mineral interactions also play an important role in soil development, aggregation and stabilization (16,129). Interactions between clay minerals and fungal biomass alters the sorptive properties of both clay minerals and fungal hyphae (130,131) and also affect the size, shape and structure of mycelial pellets (132).

Reduction or oxidation of metals and metalloids Many fungi can precipitate reduced forms of metals and metalloids, e.g. Ag(I) reduction to elemental silver Ag(0); selenate $[\text{Se}(\text{VI})]$ and selenite $[\text{Se}(\text{IV})]$ to elemental selenium $[\text{Se}(0)]$; tellurite $[\text{Te}(\text{IV})]$ to elemental tellurium $[\text{Te}(0)]$ (133-135). Reduction of Hg(II) to volatile Hg(0) can also be mediated by fungi (67,68). Increased arsenate reduction contributed to tolerance in an *Aspergillus* sp. (136,137). Mn oxidation/reduction has been described above.

Other mycogenic minerals A range of minerals other than those mentioned above have been found in association with fungi (2,3,73,74,77,80-82,114). Mycogenic secondary minerals associated with fungal hyphae and lichen thalli include desert varnish (MnO and FeO), ferrihydrite $(5\text{Fe}_2\text{O}_3 \cdot 9\text{H}_2\text{O})$, iron gluconate, calcium formate, forsterite, goethite $(\alpha\text{-Fe}^{3+}\text{O}(\text{OH}))$, moolooite $(\text{Cu}(\text{C}_2\text{O}_4) \cdot 0.4\text{H}_2\text{O})$, halloysite $(\text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4)$, and hydrocerussite $(\text{Pb}_3(\text{CO}_3)_2(\text{OH})_2)$ (16,48,52,108,120,128,138). Another biogenic mineral (tepius) has been

identified in association with a lichen carpet occurring in high mountain ranges in Venezuela (128).

Halide transformations Several fungi have the ability to produce a variety of atmospheric methyl halides. This ability is widespread in both free-living and symbiotic fungi, and is dependent on substrate concentration and community composition (139,140). The production of chloromethane (CH_3Cl) by wood-rotting fungi, e.g. *Phellinus* spp., may be particularly significant with one estimate of annual global input to the atmosphere from this source being 160 000 t of which 75% is released from tropical and subtropical forests (139). Filamentous fungi may also contribute to the global circulation of stable iodine and also the long-lived radioiodine, ^{129}I (half-life: 1.6×10^7 years), released from nuclear facilities (141).

FUNGAL SYMBIOSES IN GEOMYCOLOGY

Many fungi form partnerships with plants (mycorrhizas) and algae or cyanobacteria (lichens) that are significant geoactive agents. In general terms, the mycobiont is provided with carbon by the photobionts, while the mycobiont may protect the symbiosis from harsh environmental conditions (e.g., desiccation, metal toxicity), and provide increased access to inorganic nutrients such as phosphate and essential metals.

Lichens are fungi that exist in facultative or obligate symbioses with one or more photosynthesizing partners occurring in almost all surface terrestrial environments (142). Lichens play important roles in retention and distribution of nutrient (e.g. C, N) and trace elements, in soil formation, and rock bioweathering (54,87,143). Lichens can accumulate metals such as lead (Pb) and copper (Cu), and many other elements, including radionuclides (144). They also form a variety of metal-organic biominerals, e.g. oxalates, especially during

growth on metal-rich substrates (98,143). On copper sulfide bearing rocks, precipitation of copper oxalate (moolooite) can occur within lichen thalli (145,146).

The majority of terrestrial plants depend on symbiotic mycorrhizal fungi (147,148). Mycorrhizal fungi can mediate metal and phosphate solubilization from mineral sources, extracellular precipitation of metal oxalates, and immobilize metals within biomass (65,66,149-157). Such activities lead to changes in the physico-chemical characteristics of the root environment and enhanced bioweathering of soil minerals (55,157,158). Furthermore, ectomycorrhizal mycelia may respond to different soil silicate and phosphate minerals (e.g. apatite, quartz, potassium feldspar) by regulating growth and metabolic activity (159,160).

Mycorrhizal fungi often excrete bioweathering agents such as low molecular weight carboxylic acids and siderophores (65,161). Ectomycorrhizal fungi can also form narrow pores in weatherable minerals in podzol E horizons, probably by dissolution of Al silicates (162,163). Such excretions can also release elements from apatite and wood ash (K, Ca, Ti, Mn, Pb) (164). Ericoid mycorrhizal and ectomycorrhizal fungi can dissolve several cadmium, copper, zinc and lead-bearing minerals including metal phosphates (65,66,152,161,165). Mobilization of phosphorus from inorganic and organic phosphorus sources is generally regarded as one of the most important functions of mycorrhizal fungi, and this can also result in redistribution of incorporated metals, and the formation of other secondary minerals including other metal phosphates. The ericoid mycorrhiza *Oidiodendron maius* can solubilize zinc oxide and phosphate (161). Many ericoid mycorrhizal and ectomycorrhizal fungi are able to solubilize zinc, cadmium, copper phosphates and lead chlorophosphate (pyromorphite) releasing phosphate and component metals (65,152). An association of arbuscular mycorrhizal fungi (AMF) with *Lindenbergia philippensis*, sampled

from a Zn-contaminated settling pond at a zinc smelter, enhanced Zn accumulation in Zn-loaded rhizosphere sediment compared to treatments that suppressed AMF colonization. A significant proportion of Zn was present as crystalline and other solid materials that were associated with the root mucilaginous sheath (166). Such results may indicate a role for AMF in enhancing Zn immobilization in the rhizosphere of plants that successfully colonize Zn mining and smelting disposal sites (153,166,168).

ENVIRONMENTAL AND APPLIED SIGNIFICANCE OF GEOMYCOLOGY

The kinds of processes detailed previously can impact upon human society not only through their environmental significance and biotechnological applications, but also in deleterious contexts such as biodeterioration and biocorrosion. The biodeterioration of stone and mineral artefacts represents a loss of cultural heritage (13,14). Materials used to stabilize building blocks (mortar) and to coat surfaces prior to painting (plaster or stucco) are also susceptible to biodeterioration (13). Highly deteriorated stone surfaces provide a “proto-soil” for colonization by mosses, ferns and higher plants (14). Mechanisms of stone deterioration are complex and include most of the direct and indirect mechanisms previously discussed for mineral dissolution (13,169). Extracellular polymeric substances (EPS) are also capable of metal complexation and weakening of mineral lattices through wetting and drying cycles, as well as the production of efflorescences, i.e. secondary minerals produced through reaction of anions from excreted acids with cations from the stone (170). Physical damage may be caused by hyphal penetration of weakened areas (88,138). Lichens cause damage due to penetration by their rhizines, composed of fungal filaments, and expansion/contraction of the thallus on wetting/drying (171). “Lichen acids”, mainly oxalic acid, cause damage at the stone/lichen interface, and lichen thalli may accumulate up to 50% calcium oxalate, depending on the substrate (172,173). In addition, carbonic acid formed in the lichen thallus

can solubilize calcium and magnesium carbonates in calcareous stone (174). Fungal biodeterioration of ancient ivory (natural apatite; walrus tusk) was accompanied by widespread etching and tunneling by hyphae and extensive formation of calcium oxalate monohydrate, whewellite (175). Concrete and cement can be biodeteriorated and in some environments, fungi dominate the concrete-deteriorating microbiota (13,14,176-178). Microbial attack on concrete is mediated by protons, inorganic and organic acids and the production of hydrophilic slimes leading to biochemical and biomechanical deterioration (13,75,169). Several species of microfungi were able to colonize samples of the concrete used as radioactive waste barrier in the Chernobyl reactor and leached iron, aluminium, silicon and calcium, and re-precipitated silicon and calcium oxalate (75).

Mineral and metal solubilization mechanisms enable metal removal from industrial wastes, low-grade ores, and metal-bearing minerals. This may have application in bioremediation, metal biorecovery and recycling (58,68,179,180). Metals can be solubilized from fly ash (originating from municipal solid waste incineration), contaminated soil, electronic scrap and other waste materials by fungal activity (179,181). Although fungal systems cannot compare with the efficiency of bacterial bioleaching, they may be more suited to specific bioreactor applications (58). A variety of fungal mechanisms result in metal immobilization such as biosorption, bioaccumulation and bioprecipitation. Biosorption is a physico-chemical process, and is a property of both living and dead organisms (and their components), and fungi are effective agents for removal of metals, radionuclides and other substances from solution (69,70,71,76,130,182-193). Urease-positive fungi can be used to precipitate metal-containing carbonates, some in nanoscale dimensions, thus providing a means of metal biorecovery as well as potentially useful nanoscale biomineral products (94,95). Similarly, the formation of other insoluble metal compounds by fungi or their metabolites could also be

considered as a means to biorecover metals, metalloids and radionuclides, e.g. oxalates, oxides, oxalates, and phosphates, as well as the production of elemental metal or metalloid forms (2,78). Some biomineral and elemental products, including those of nanoscale dimensions, are of relevance to the production of novel advanced biomaterials with applications in metal and radionuclide bioremediation, antimicrobial treatments (e.g. nano-silver), solar energy and electrical battery applications, and microelectronics (194). In a novel approach, urease-positive *Neurospora crassa* was used to precipitate manganese carbonate. After thermal treatment at 300°C, the carbonized biomass-manganese oxide composite material was used in lithium-ion batteries (LiB) and supercapacitors where it was found to exhibit excellent electrochemical properties. In LiB, around 90% charge capacity was retained after 200 charge-discharge cycles (195).

The ability of fungi and bacteria to transform metalloids has been successfully used for bioremediation of contaminated land and water. Selenium methylation results in volatilization and this has been used to remove selenium from the San Joaquin Valley and Kesterson Reservoir, California (196). Mycorrhizal associations may have application in phytoremediation (197,198), the use of plants to remove or detoxify environmental pollutants (199), by metal phytoextraction or by acting as a biological barrier (200-202). Glomalin, an insoluble glycoprotein, is produced in copious amounts on hyphae of arbuscular mycorrhizal fungi and can sequester metals such as Cu, Cd and Pb (203). Arbuscular mycorrhizal fungi can also decrease U translocation from plant roots to shoot (204-206). For ericaceous mycorrhizas, the fungus prevents translocation of Cu and Zn to host plant shoots (147,207,208). The development of stress-tolerant plant-mycorrhizal associations may be a promising strategy for phytoremediation and soil amelioration (161,209,210).

Some of the geomycological processes detailed previously may have consequences for abiotic soil treatment processes, notably the immobilization of toxic metals by phosphate formation. Apatite $\text{Ca}_5(\text{PO}_4)_3(\text{F},\text{Cl},\text{OH})$, pyromorphite $\text{Pb}_5(\text{PO}_4)_3\text{Cl}$, mimetite $\text{Pb}_5(\text{AsO}_4)_3\text{Cl}$ and vanadinite $\text{Pb}_5(\text{VO}_4)_3\text{Cl}$ are the most common prototypes of the apatite mineral family. Such minerals hold promise for stabilization and recycling of industrial and nuclear waste and have been explored for treatment of lead-contaminated soils and waters (211-216). The stability of these minerals is therefore of interest in any soil remediation strategy seeking to reduce the effects of potentially-toxic elements, like Pb, V and As. For example, pyromorphite is a highly insoluble lead phosphate mineral under a wide range of geochemical conditions and has often been suggested as a means to reduce Pb bioavailability. However, solubilization of pyromorphite and formation of lead oxalate by several free-living and symbiotic fungi demonstrates that pyromorphite may not be as effective at immobilizing lead as some previous studies have suggested (64,65). Similarly, despite the insolubility of vanadinite, fungi exerted both biochemical and biophysical effects on the mineral including etching, penetration and the formation of new biominerals (217). Lead oxalate was precipitated by *Aspergillus niger* during the bioleaching of vanadinite and mimetite which implies a general fungal mechanism for the transformation of lead-containing apatite group minerals (e.g. vanadinite, pyromorphite, mimetite) (217,218). This pattern of fungal bioweathering of lead apatites could be extended to other metal apatites, such as calcium apatite $[\text{Ca}_5(\text{PO}_4)_3(\text{OH},\text{F},\text{Cl})]$. Here, the formation of monohydrated (whewellite) and dihydrated (weddelite) calcium oxalate can be accomplished by many different fungal species (79,93,99,175,219,220). The ability of free-living and mycorrhizal fungi to transform toxic metal-containing minerals should therefore be taken into account in risk assessments of the long-term environmental consequences of *in situ* chemical remediation techniques, revegetation strategies or natural attenuation of contaminated sites. The bioweathering

potential of fungi has been suggested as a possible means for the bioremediation of asbestos rich soils. Several fungi could extract iron from asbestos mineral fibres (e.g. 7.3% from crocidolite and 33.6% from chrysotile by a *Verticillium* sp.), thereby removing the reactive iron ions responsible for DNA damage (221).

CONCLUSIONS

The geoactive roles of fungi have often received scant attention in geomicrobiological contexts but they are of clear importance in several key areas. These include a variety of organic and inorganic transformations important in nutrient and element cycling, rock and mineral bioweathering, mycogenic biomineral formation, and metal-fungal interactions. Lichens and mycorrhizas are of special significance as geoactive agents. Organic matter decomposition is important for the cycling of major biomass-associated elements, e.g. C, H, N, O, P and S as well as all other elements that may be found in lower concentrations. Transformations of metals and minerals are central to many geomicrobial processes, and fungi can effect changes in metal speciation, toxicity and mobility, as well as mediate mineral formation or dissolution. Such mechanisms are important components of natural biogeochemical cycles for metals as well as associated elements in biomass, soil, rocks and minerals, e.g. S and P, and metalloids, actinides and metal radionuclides. It is within the terrestrial environment where fungi have the greatest abundance and geochemical influence. However, they are also important in aquatic habitats and are now recognized as significant components of aquatic sediments and the deep subsurface. Geomycological processes can have beneficial or detrimental consequences in a human context. Beneficial applications in environmental biotechnology include metal and radionuclide bioleaching, biorecovery, detoxification, and bioremediation, and in the production of biominerals or metal(loid) elements with catalytic or other properties in nanoparticle, crystalline or colloidal forms. The

latter may be relevant to the development of novel biomaterials. Adverse effects include biodeterioration and destruction of natural and synthetic materials, rock and mineral-based building materials (e.g. concrete), cultural heritage, biocorrosion of metals, alloys and related substances, and adverse effects on radionuclide speciation, mobility and containment. The ubiquity and importance of fungi in biosphere processes underlines the importance of geomycology as a conceptual framework encompassing the environmental activities of fungi, their impact, and their applied significance.

ACKNOWLEDGEMENTS

The author gratefully acknowledges research support from the Natural Environment Research Council, the Biotechnology and Biological Sciences Research Council, the Royal Societies of London and Edinburgh, CCLRC Daresbury SRS, British Nuclear Fuels plc, the National Nuclear Laboratory and the Nuclear Decommissioning Agency. Financial support for some of the research described was received from the Natural Environment Research Council (NE/M010910/1 (TeaSe); NE/M011275/1 (COG3)) which is gratefully acknowledged, as well as the National Natural Science Foundation of China (U1503281). G. M. Gadd gratefully acknowledges an award under the 1000 Talents Plan with the Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Urumqi, China.

REFERENCES

- Ehrlich HL, Newman DK.** 2009. *Geomicrobiology*, 5th edition. CRC Press/Taylor and Francis Group, Boca Raton, FL.
- Gadd GM.** 2010. Metals, minerals and microbes: geomicrobiology and bioremediation. *Microbiol* **156**: 609 – 643.

- 622 **3. Gadd GM.** 2007. Geomycology: biogeochemical transformations of rocks, minerals,
623 metals and radionuclides by fungi, bioweathering and bioremediation. *Mycol Res* **111**:
624 3-49.
- 625 **4. Gadd GM.** 2008. Fungi and their role in the biosphere, p 1709-1717. *In* Jorgensen
626 SE, B. Fath B ed., *Encyclopedia of Ecology*. Elsevier, Amsterdam.
- 627 **5. Gadd GM.** 2008. Bacterial and fungal geomicrobiology: a problem with
628 communities? *Geobiol* **6**: 278–284.
- 629 **6. Kim BH, Gadd GM.** 2008. *Bacterial Physiology and Metabolism*. Cambridge
630 University Press, Cambridge.
- 631 **7. Druschel GK, Kappler A.** 2015. Geomicrobiology and microbial geochemistry.
632 *Elements* **11**: 389-394.
- 633 **8. Reitner J, Schumann G, Pedersen K.** 2006. Fungi in subterranean environments, p
634 377-403. *In* Gadd GM ed., *Fungi in Biogeochemical Cycles*. Cambridge University
635 Press, Cambridge.
- 636 **9. Gorbushina AA.** 2007. Life on the rocks. *Environ Microbiol* **9**: 1613-1631.
- 637 **10. Vázquez-Campos X, Kinsela AS, Waite TD, Collins R.N, Neilan BA.** 2014.
638 *Fodinomyces uranophilus* gen. nov. sp. nov. and *Coniochaeta fodinicola* sp. nov., two
639 uranium mine inhabiting Ascomycota fungi from northern Australia. *Mycologia* **106**:
640 1073–1089.
- 641 **11. Ivarsson M, Bengtson S, Neubeck A.** 2016. The igneous oceanic crust – Earth’s
642 largest fungal habitat? *Fungal Ecol* **20**: 249-255.
- 643 **12. Uroz S, Calvaruso C, Turpault M-P, Frey-Klett P.** 2009. Mineral weathering by
644 bacteria: ecology, actors and mechanisms. *Trends Microbiol* **17**: 378-387.
- 645 **13. Scheerer S, Ortega-Morales O, Gaylarde C.** 2009. Microbial deterioration of stone
646 monuments: an updated overview. *Adv Appl Microbiol* **66**: 97-139.

- 647 **14. Cutler N, Viles H.** 2010. Eukaryotic microorganisms and stone biodeterioration.
648 *Geomicrobiol J* 27: 630–646.
- 649 **15. Gadd GM.** 2016. Geomicrobiology of the built environment. *Nature Microbiol* in
650 press.
- 651 **16. Burford EP, Fomina M, Gadd GM.** 2003. Fungal involvement in bioweathering and
652 biotransformation of rocks and minerals. *Mineral Mag* **67**: 1127-1155.
- 653 **17. Miyata N, Tani Y, Iwahori K, Soma M.** 2004. Enzymatic formation of manganese
654 oxides by an *Acremonium*-like hyphomycete fungus, strain KR21-2. *FEMS Microbiol*
655 *Ecol* **47**: 101-109.
- 656 **18. Fomina M, Burford EP, Gadd GM.** 2005. Toxic metals and fungal communities, p
657 733-758. In Dighton J, White JF, Oudemans P ed., *The Fungal Community. Its*
658 *Organization and Role in the Ecosystem*. CRC Press, Boca Raton, FL.
- 659 **19. Vázquez-Campos X, Kinsela AS, Collins RN, NeilaBA, Aoyagi N, Waite TD.**
660 2015. Uranium binding mechanisms of the acid-tolerant fungus *Coniochaeta*
661 *fodinicola*. *Environ Sci Technol* **49**: 8487–8496.
- 662 **20. Horiike T, Yamashita M.** 2015. A new fungal isolate, *Penidiella* sp. strain T9,
663 accumulates the rare earth element dysprosium. *Appl Environ Microbiol* **81**: 3062–
664 3068.
- 665 **21. Gadd GM.** 2004. Mycotransformation of organic and inorganic substrates.
666 *Mycologist* **18**: 60-70.
- 667 **22. Braissant O, Cailleau G, Aragno M, Verrecchia EP.** 2004. Biologically induced
668 mineralization in the tree *Milicia excelsa* Moraceae.: its causes and consequences to
669 the environment. *Geobiol* **2**: 59-66.
- 670 **23. Wainwright M, Tasnee AA, Barakah F.** 1993. A review of the role of oligotrophic
671 microorganisms in biodeterioration. *Int Biodet Biodeg* **31**: 1-13.

24. **Sterflinger K.** 2000. Fungi as geologic agents. *Geomicrobiol J* **17**: 97-124.
25. **Gorbushina AA, Krumbein WE, Hamann R, Panina L, Soucharjevsky S, Wollenzien U.** 1993. On the role of black fungi in colour change and biodeterioration of antique marbles. *Geomicrobiol J* **11**: 205-221.
26. **Ivarsson M.** 2012. The seafloor basalts as fungal habitats. *Biogeosci* **9**: 3625-3635.
27. **Nagano Y, Nagahama T.** 2012. Fungal diversity in deep-sea extreme environments. *Fungal Ecol* **5**: 463-471.
28. **Orsi WD, Biddle JF, Edgcomb VD.** 2013. Deep sequencing of seafloor eukaryotic rRNA reveals active fungi across marine subsurface provinces. *PLoS One* **8**: 1-10.
29. **Nagano Y, Nagahama T, Hatada Y, Nunoura T, Takami H, Miyazaki J, Takai K, Horikoshi K.** 2010. Fungal diversity in deep-sea sediments and the presence of novel fungal groups. *Fungal Ecol* **3**: 316-325.
30. **Le Calvez T, Burgaud G, Mahe S, Barbier G, P.** 2009. Fungal diversity in deep sea hydrothermal ecosystems. *Appl Environ Microbiol* **75**: 6415-6421.
31. **Connell L, Barrett A, Templeton A, Staudigel H.** 2009. Fungal diversity associated with an active deep sea volcano: Vailulu'u Seamount, Samoa. *Geomicrobiol J* **26**: 597-605.
32. **Nagahama T, Takahashi E, Nagano Y, Abdel-Wahab MA, Miyazaki M.** 2011. Molecular evidence that deep branching fungi are major fungal components in deep-sea methane cold-seep sediments. *Environ Microbiol* **13**: 2359-2370.
33. **Ivarsson M, Bengtson S, Skogby H, Lazor P, Broman C, Belivanova V, Marone F.** 2015. A fungal-prokaryotic consortium at the basalt-zeolite interface in seafloor

igneous crust. *PLoS One* **10**10.:

e014016. <http://dx.doi.org/10.1371/journal.pone.0140106>.

- 34. Bengtson S, Ivarsson M, Astolfo A, Belivanova V, Broman C, Marone F, Stampanoni M.** 2014. Deep-biosphere consortium of fungi and prokaryotes in Eocene sub-seafloor basalts. *Geobiol* **12**: 489-496.
- 35. Schumann G, Manz W, Reitner J, Lustrino M.** 2004. Ancient fungal life in North Pacific Eocene oceanic crust. *Geomicrobiol J* **21**: 241-246.
- 36. Gadd GM. ed..** 2006. *Fungi in Biogeochemical Cycles*. Cambridge University Press, Cambridge.
- 37. Boddy L.** 2016. Fungi, ecosystems, and global change, p 361-400. *In* Watkinson SC, Boddy L, Money NP ed., *The Fungi*. Elsevier, Amsterdam.
- 38. Gadd GM.** 2000. Microbial interactions with tributyltin compounds: detoxification, accumulation, and environmental fate. *Sci Total Environ* **258**: 119-127.
- 39. Cerniglia CE, Sutherland JB.** 2001. Bioremediation of polycyclic aromatic hydrocarbons by ligninolytic and non-ligninolytic fungi, p 136-187. *In* Gadd GM ed., *Fungi in Bioremediation*. Cambridge University Press, Cambridge.
- 40. Cerniglia CE, Sutherland JB.** 2006. Relative roles of bacteria and fungi in polycyclic aromatic hydrocarbon biodegradation and bioremediation of contaminated soils, p 182-211. *In* Gadd GM ed., *Fungi in Biogeochemical Cycles*. Cambridge University Press, Cambridge.
- 41. Singleton I.** 2001. Fungal remediation of soils contaminated with persistent organic pollutants, p 79-96. *In* Gadd GM ed., *Fungi in Bioremediation*. Cambridge University Press, Cambridge.
- 42. Gadd GM. ed..** 2001. *Fungi in Bioremediation*. Cambridge University Press, Cambridge.

- 721 **43. Sutherland JB.** 2004. Degradation of hydrocarbons by yeasts and filamentous fungi,
722 p 443-455. *In* Arora DK ed., *Fungal Biotechnology in Agricultural, Food, and*
723 *Environmental Applications*. Marcel Dekker, New York.
- 724 **44. Gadd GM, Burford EP, Fomina M, Melville K.** 2007. Mineral transformations and
725 biogeochemical cycles: a geomycological perspective, p 78-111. *In* Gadd GM, Dyer
726 P, Watkinson S ed., *Fungi in the Environment*. Cambridge University Press,
727 Cambridge.
- 728 **45. Hochella MF.** 2002. Sustaining Earth: thoughts on the present and future roles in
729 mineralogy in environmental science. *Mineral Mag* **66**: 627-652.
- 730 **46. Kraemer SM, Cheah SF, Zapf R, Xu JD, Raymond KN, Sposito G.** 1999. Effect
731 of hydroxamate siderophores on Fe release and PbII. adsorption by goethite. *Geochim*
732 *Cosmochim Acta* **63**: 3003-3008.
- 733 **47. Vaughan DJ, Pattrick RAD, Wogelius RA.** 2002. Minerals, metals and molecules:
734 ore and environmental mineralogy in the new millennium. *Mineral Mag* **66**: 653-676.
- 735 **48. Burford EP, Kierans M, Gadd GM.** 2003. Geomycology: fungal growth in mineral
736 substrata. *Mycologist* **17**: 98-107.
- 737 **49. Boswell GP, Jacobs H, Davidson FA, Gadd GM, Ritz K.** 2002. Functional
738 consequences of nutrient translocation in mycelial fungi. *J Theor Biol* **217**: 459-477.
- 739 **50. Boswell GP, Jacobs H, Davidson FA, Gadd GM, Ritz K.** 2003. Growth and function
740 of fungal mycelia in heterogeneous environments. *Bull Math Biol* **65**: 447-477.
- 741 **51. Boswell GP, Jacobs H, Ritz K, Gadd GM, Davidson FA.** 2007. The development
742 of fungal networks in complex environments. *Bull Math Biol* **69**: 605-634.
- 743 **52. Verrecchia EP.** 2000. Fungi and sediments, p 69–75. *In* Riding RE, Awramik SM
744 ed., *Microbial Sediments*. Springer-Verlag, Berlin.
- 745 **53. Gadd GM.** 1993. Interactions of fungi with toxic metals. *New Phytol* **124**: 25-60.

- 746 **54. Banfield JP, Barker WW, Welch SA, Taunton A.** 1999. Biological impact on
747 mineral dissolution: application of the lichen model to understanding mineral
748 weathering in the rhizosphere. *Proc Nat Acad Sci USA* **96**: 3404–3411.
- 749 **55. Bonneville S, Smits MM, Brown A, Harrington J, Leake JR, Brydson R, Benning**
750 **LG.** 2009. Plant-driven fungal weathering: early stages of mineral alteration at the
751 nanometer scale. *Geology* **37**: 615-618.
- 752 **56. Li L, Liu L, Chen J, Teng HH.** 2016. Cellular dissolution at hypha- and spore-
753 mineral interfaces revealing unrecognized mechanisms and scales of fungal
754 weathering. *Geol Soc Amer Data, Repository Item* 2016094.
- 755 **57. Bowen AD, Davidson FA, Keatch R, Gadd GM.** 2007. Induction of contour
756 sensing in *Aspergillus niger* by stress and its relevance to fungal growth mechanics
757 and hyphal tip structure. *Fungal Gen Biol* **44**: 484-491.
- 758 **58. Burgstaller W, Schinner F.** 1993. Leaching of metals with fungi. *J Biotechnol* **27**:
759 91-116.
- 760 **59. Gadd GM.** 1999. Fungal production of citric and oxalic acid: importance in metal
761 speciation, physiology and biogeochemical processes. *Adv Microb Physiol* **41**: 47-92.
- 762 **60. Ehrlich H L.** 1998. Geomicrobiology: its significance for geology. *Earth Sci Rev* **45**:
763 45-60.
- 764 **61. Gharieb MM, Sayer JA, Gadd GM.** 1998. Solubilization of natural gypsum
765 $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$. and the formation of calcium oxalate by *Aspergillus niger* and *Serpula*
766 *himantoides*. *Mycol Res* **102**: 825-830.
- 767 **62. Adeyemi AO, Gadd GM.** 2005. Fungal degradation of calcium-, lead- and silicon-
768 bearing minerals. *Biometals* **18**: 269-281.
- 769 **63. Strasser H, Burgstaller W, Schinner F.** 1994. High yield production of oxalic acid
770 for metal leaching purposes by *Aspergillus niger*. *FEMS Microbiol Lett* **119**: 365-370.

- 771 **64. Sayer JA, Cotter-Howells JD, Watson C, Hillier S, Gadd GM.** 1999. Lead mineral
772 transformation by fungi. *Curr Biol* **9**: 691-694.
- 773 **65. Fomina MA, Alexander IJ, Hillier S, Gadd GM.** 2004. Zinc phosphate and
774 pyromorphite solubilization by soil plant-symbiotic fungi. *Geomicrobiol J* **21**: 351-
775 366.
- 776 **66. Fomina MA, Alexander IJ, Colpaert JV, Gadd GM.** 2005. Solubilization of toxic
777 metal minerals and metal tolerance of mycorrhizal fungi. *Soil Biol Biochem* **37**: 851-
778 866.
- 779 **67. Gadd GM.** 1993. Microbial formation and transformation of organometallic and
780 organometalloid compounds. *FEMS Microbiol Rev* **11**: 297-316.
- 781 **68. Gadd GM.** 2000. Bioremedial potential of microbial mechanisms of metal mobilization
782 and immobilization. *Curr Opinion Biotechnol* **11**: 271-279.
- 783 **69. Gadd GM.** 2001. Accumulation and transformation of metals by microorganisms, p
784 225-264. In Rehm H-J, Reed G, Puhler A, Stadler P ed., *Biotechnology, a Multi-volume*
785 *Comprehensive Treatise, Volume 10: Special Processes*. Wiley-VCH Verlag GmbH,
786 Weinheim, Germany.
- 787 **70. Gadd GM.** 2001. Metal transformations, p 359-382. In Gadd GM ed., *Fungi in*
788 *Bioremediation*. Cambridge University Press, Cambridge.
- 789 **71. Gadd GM.** 2009. Biosorption: critical review of scientific rationale, environmental
790 importance and significance for pollution treatment. *J Chem Technol Biotechnol* **84**:
791 13-28.
- 792 **72. Baldrian P.** 2003. Interaction of heavy metals with white-rot fungi. *Enzyme Microb*
793 *Technol* **32**: 78-91.
- 794 **73. Fomina M, Charnock JM, Hillier S, Alvarez R, Gadd GM.** 2007. Fungal
795 transformations of uranium oxides. *Environ Microbiol* **9**: 1696-1710.

74. **Fomina M, Charnock J, Bowen AD, Gadd GM.** 2007. X-ray absorption spectroscopy XAS. of toxic metal mineral transformations by fungi. *Environ Microbiol* **9**: 308-321.
75. **Fomina M, Podgorsky VS, Olishevskaya SV, Kadoshnikov VM, Pisanska IR, Hillier S, Gadd GM.** 2007. Fungal deterioration of barrier concrete used in nuclear waste disposal. *Geomicrobiol J* **24**: 643-653.
76. **Fomina M, Gadd GM.** 2014. Biosorption: current perspectives on concept, definition and application. *Biores Technol* **160**: 3-14.
77. **Gadd GM, Raven JA.** 2010. Geomicrobiology of eukaryotic microorganisms. *Geomicrobiol J* **27**: 491-519.
78. **Gadd GM, Rhee YJ, Stephenson K, Wei Z.** 2012. Geomycology: metals, actinides and biominerals. *Environ Microbiol Rep* **4**: 270–296.
79. **Gadd GM, Bahri-Esfahani J, Li Q, Rhee YJ, Wei Z, Fomina M, Liang X.** 2014. Oxalate production by fungi: significance in geomycology, biodeterioration and bioremediation. *Fungal Biol Rev* **28**: 36–55.
80. **Liang X, Hillier S, Pendrowski H, Gray N, Ceci A, Gadd GM.** 2015. Uranium phosphate biomineralization by fungi. *Environ Microbiol* **17**: 2064-2075.
81. **Liang X, Kierans M, Ceci A, Hillier S, Gadd GM.** 2016. Phosphatase-mediated bioprecipitation of lead by soil fungi. *Environ Microbiol* **18**: 219–231.
82. **Liang X, Csetenyi L, Gadd GM.** 2016. Lead bioprecipitation by yeasts utilizing organic phosphorus substrates. *Geomicrobiol J* **33**: 294-307.
83. **Lloyd JR, Pearce CI, Coker VS, Pattrick RADP, van der Laan G, Cutting, R, Vaughan DV, Paterson-Beedle M, Mikheenko IP, Yong P, Macaskie LE.** 2008. Biomineralization: linking the fossil record to the production of high value functional materials. *Geobiol* **6**: 285-297.

- 821 **84. Rosling A, Roose T, Herrmann AM, Davidson FA, Finlay RD, Gadd GM.** 2009.
822 Approaches to modelling mineral weathering by fungi. *Fungal Biol Rev* **23**: 1-7.
- 823 **85. Smits MM.** 2009. Scale matters? Exploring the effect of scale on fungal-mineral
824 interactions. *Fungal Biol Rev* **23**: 132-137.
- 825 **86. Hutchens E.** 2009. Microbial selectivity on mineral surfaces: possible implications
826 for weathering processes. *Fungal Biol Rev* **23**: 115-121.
- 827 **87. Adamo P, Violante P.** 2000. Weathering of rocks and neogenesis of minerals
828 associated with lichen activity. *Appl Clay Sci* **16**: 229-256.
- 829 **88. Cockell CS, Herrera A.** 2008. Why are some microorganisms boring? *Trends*
830 *Microbiol* **16**: 101-106.
- 831 **89. Lian B, Wang B, Pan M, Liu C, Teng HH.** 2008. Microbial release of potassium
832 from K-bearing minerals by thermophilic fungus *Aspergillus fumigatus*. *Geochim*
833 *Cosmochim Acta* **72**: 87-98.
- 834 **90. Golubic S, Radtke G, Le Campion-Alsumard T.** 2005. Endolithic fungi in marine
835 ecosystems. *Trends Microbiol* **13**: 229–235.
- 836 **91. Kolo K, Keppens E, Preat A, Claeys P.** 2007. Experimental observations on fungal
837 diagenesis of carbonate substrates. *J Geophys Res* **112**: 1-20.
- 838 **92. Verrecchia E.P, Dumont JL, Rolko KE.** 1990. Do fungi building limestones exist in
839 semi-arid regions? *Naturwissenschaften* **77**: 584-586.
- 840 **93. Burford EP, Hillier S, Gadd GM.** 2006. Biomineralization of fungal hyphae with
841 calcite CaCO₃. and calcium oxalate mono- and dihydrate in carboniferous limestone
842 microcosms. *Geomicrobiol J* **23**: 599-611.
- 843 **94. Li Q, Csetenyi L, Gadd GM.** 2014. Biomineralization of metal carbonates by
844 *Neurospora crassa*. *Environ Sci Technol* **48**: 14409-14416.

- 845 **95. Li Q, Csetenyi L, Paton GI, Gadd GM.** 2015. CaCO₃ and SrCO₃ bioprecipitation
846 by fungi isolated from calcareous soil. *Environ Microbiol* **17**: 3082-3097.
- 847 **96. Kumari D, Qian X-Y, Pan X, Achal V, Li Q, Gadd GM.** 2016. Microbially-
848 induced carbonate precipitation for immobilization of toxic metals. *Adv Appl*
849 *Microbiol* **94**: 79-108.
- 850 **97. Rhee YJ, Hillier S, Gadd GM.** 2016. A new lead hydroxycarbonate produced during
851 transformation of lead metal by the soil fungus *Paecilomyces javanicus*.
852 *Geomicrobiol J* in press.. Any page numbers yet?
- 853 **98. Adamo P, Vingiani S, Violante P.** 2002. Lichen-rock interactions and bioformation
854 of minerals. *Dev Soil Sci* **28B**: 377-391.
- 855 **99. Pinzari F, Zotti M, De Mico A, Calvini P.** 2010. Biodegradation of inorganic
856 components in paper documents: formation of calcium oxalate crystals as a
857 consequence of *Aspergillus terreus* Thom growth. *Int Biodet Biodegrad* **64**:499–505.
- 858 **100. Arnott HJ.** 1995. Calcium oxalate in fungi, p 73-111. *In* Khan SR ed., *Calcium*
859 *Oxalate in Biological Systems*. CRC Press, Boca Raton, FL.
- 860 **101. Sayer JA, Gadd GM.** 1997. Solubilization and transformation of insoluble metal
861 compounds to insoluble metal oxalates by *Aspergillus niger*. *Mycol Res* **101**: 653-661.
- 862 **102. Jarosz-Wilkolazka A, Gadd GM.** 2003. Oxalate production by wood-rotting fungi
863 growing in toxic metal-amended medium. *Chemosphere* **52**: 541-547.
- 864 **103. Wei Z, Hillier S, Gadd GM.** 2012. Biotransformation of manganese oxides by fungi:
865 solubilization and production of manganese oxalate biominerals. *Environ Microbiol*
866 **14**: 1744-1753.
- 867 **104. Verrecchia EP, Braissant O, Cailleau G.** 2006. The oxalate-carbonate pathway in
868 soil carbon storage: the role of fungi and oxalotrophic bacteria, p 289-310. *In* Gadd
869 GM ed., *Fungi in Biogeochemical Cycles*. Cambridge University Press, Cambridge.

- 870 **105. Kolo K, Claeys P.** 2005. *In vitro* formation of Ca-oxalates and the mineral
871 glushinskite by fungal interaction with carbonate substrates and seawater.
872 *Biogeosciences* **2**: 277-293.
- 873 **106. Miyata M, Tani Y, Sakata M, Iwahori K.** 2007. Microbial manganese oxide
874 formation and interaction with toxic metal ions. *J Biosci Bioeng* **104**: 1-8.
- 875 **107. Saratovsky I, Gurr SJ, Hayward MA.** 2009. The structure of manganese oxide
876 formed by the fungus *Acremonium* sp. strain KR21-2. *Geochim Cosmochim Acta* **73**:
877 3291-3300.
- 878 **108. Grote G, Krumbein WE.** 1992. Microbial precipitation of manganese by bacteria
879 and fungi from desert rock and rock varnish. *Geomicrobiol J* **10**: 49-57.
- 880 **109. Eckhardt FEW.** 1985. Solubilisation, transport, and deposition of mineral cations by
881 microorganisms-efficient rock-weathering agents, p 161-173. In Drever J ed., *The*
882 *Chemistry of Weathering*, vol. 149. Nato Asi Ser C.
- 883 **110. Whitelaw MA, Harden TJ, Helyar KR.** 1999. Phosphate solubilization in solution
884 culture by the soil fungus *Penicillium radicum*. *Soil Biol Biochem* **31**: 655-665.
- 885 **111. Rhee YJ, Hillier S, Gadd GM.** 2012. Lead transformation to pyromorphite by fungi.
886 *Curr Biol* **22**: 237-241.
- 887 **112. Rhee YJ, Hillier S, Pendrowski H, Gadd GM.** 2014. Pyromorphite formation in a
888 fungal biofilm community growing on lead metal. *Environ Microbiol* **16**: 1441-1451.
- 889 **113. Rhee YJ, Hillier S, Pendrowski H, Gadd GM.** 2014. Fungal transformation of
890 metallic lead to pyromorphite in liquid medium. *Chemosphere* **113**: 17-21.
- 891 **114. Fomina M, Charnock JM, Hillier S, Alvarez R, Livens F, Gadd GM.** 2008. Role
892 of fungi in the biogeochemical fate of depleted uranium. *Curr Biol* **18**: 375-377.
- 893 **115. Gadd GM, Fomina M.** 2011. Uranium and fungi. *Geomicrobiol J* **28**: 471-482.

- 894 **116. Brehm U, Gorbushina A, Mottershead D.** 2005. The role of microorganisms and
895 biofilms in the breakdown and dissolution of quartz and glass. *Palaeogeo Palaeoclim*
896 *Palaeoecol* **219**: 117-129.
- 897 **117. Barker WW, Banfield JF.** 1996. Biologically versus inorganically mediated
898 weathering reactions: relationships between minerals and extracellular microbial
899 polymers in lithobiotic communities. *Chem Geol* **132**: 55-69.
- 900 **118. Barker WW, Banfield JF.** 1998. Zones of chemical and physical interaction at
901 interfaces between microbial communities and minerals: a model. *Geomicrobiol J* **15**:
902 223-244.
- 903 **119. Arocena JM, Glowa KR, Massicotte HB, Lavkulich L.** 1999. Chemical and
904 mineral composition of ectomycorrhizosphere soils of subalpine fir *Abies lasiocarpa*
905 Hook.. Nutt.. in the AE horizon of a Luvisol. *Can J Soil Sci* **79**: 25-35.
- 906 **120. Arocena JM, Zhu LP, Hall K.** 2003. Mineral accumulations induced by biological
907 activity on granitic rocks in Qinghai Plateau, China. *Earth Surf Proc Landforms* **28**:
908 1429-1437.
- 909 **121. Tazaki K.** 2006. Clays, microorganisms, and biomineralization, p 477-497. In
910 Bergaya F, Theng BKG, Lagaly G ed., *Handbook of Clay Science, Developments in*
911 *Clay Science*, Vol. 1. Elsevier, Amsterdam.
- 912 **122. Theng BKG, Yuan G.** 2008. Nanoparticles in the soil environment. *Elements* **4**: 395-
913 399.
- 914 **123. Cromack K, Solkins P, Graustein WC, Speidel K, Todd AW, Spycher G, Li CY,**
915 **Todd RL.** 1979. Calcium oxalate accumulation and soil weathering in mats of the
916 hypogeous fungus *Hysterangium crassum*. *Soil Biol Biochem* **11**: 463-468.

- 124. Delatorre MA, Gomez-Alarcon G, Vizcaino C, Garcia MT.** 1993. Biochemical mechanisms of stone alteration carried out by filamentous fungi living on monuments. *Biogeochem* **19**: 129-147.
- 125. Bennett PC, Rogers JA, Hiebert FK, Choi WJ.** 2001. Silicates, silicate weathering, and microbial ecology. *Geomicrobiol J* **18**: 3-19.
- 126. Wei Z, Kierans M, Gadd GM.** 2012. A model sheet mineral system to study fungal bioweathering of mica. *Geomicrobiol J* **29**: 323-331.
- 127. Wei Z, Liang X, Pendlowski H, Hillier S, Suntornvongsagul K, Sihanonth P, Gadd GM.** 2013. Fungal biotransformation of zinc silicate and sulfide mineral ores. *Environ Microbiol* **15**: 2173-2186.
- 128. Gorbushina AA, Boettcher M, Brumsack HJ, Krumbein WE, Vendrell-Saz M.** 2001. Biogenic forsterite and opal as a product of biodeterioration and lichen stromatolite formation in table mountain systems tepuis. of Venezuela. *Geomicrobiol J* **18**: 117-132.
- 129. Ritz K, Young IM.** 2004. Interaction between soil structure and fungi. *Mycologist* **18**: 52-59.
- 130. Fomina M, Gadd GM.** 2002. Metal sorption by biomass of melanin-producing fungi grown in clay-containing medium. *J Chem Technol Biotechnol* **78**: 23-34.
- 131. Morley GF, Gadd GM.** 1995. Sorption of toxic metals by fungi and clay minerals. *Mycol Res* **99**: 1429-1438.
- 132. Fomina M, Gadd GM.** 2002. Influence of clay minerals on the morphology of fungal pellets. *Mycol Res* **106**: 107-117.
- 133. Kierans M, Staines AM, Bennett H, Gadd GM.** 1991. Silver tolerance and accumulation in yeasts. *Biol Metals* **4**: 100-106.

- 134. Gharieb MM, Wilkinson SC, Gadd GM.** 1995. Reduction of selenium oxyanions by unicellular, polymorphic and filamentous fungi: cellular location of reduced selenium and implications for tolerance. *J Industrial Microbiol* **14**: 300-311.
- 135. Gharieb MM, Kierans M, Gadd GM.** 1999. Transformation and tolerance of tellurite by filamentous fungi: accumulation, reduction and volatilization. *Mycol Res* **103**: 299-305.
- 136. Canovas D, Duran C, Rodriguez N, Amils R, de Lorenzo V.** 2003. Testing the limits of biological tolerance to arsenic in a fungus isolated from the River Tinto. *Environ Microbiol* **5**: 133-138.
- 137. Canovas D, Mukhopadhyay R, Rosen BP, de Lorenzo V.** 2003. Arsenate transport and reduction in the hyper-tolerant fungus *Aspergillus* sp P37. *Environ Microbiol* **5**: 1087-1093.
- 138. Hirsch P, Eckhardt FEW, Palmer R J, Jr.** 1995. Methods for the study of rock inhabiting microorganisms – a mini review. *J Microbiol Meth* **23**: 143–167.
- 139. Watling R, Harper DB,** 1998. Chloromethane production by wood-rotting fungi and an estimate of the global flux to the atmosphere. *Mycol Res* **102**: 769-787.
- 140. Redeker KR, Treseder KK, Allen MF.** 2004. Ectomycorrhizal fungi: a new source of atmospheric methyl halides? *Global Change Biol* **10**: 1009-1016.
- 141. Ban-nai T, Muramatsu Y, Amachi S.** 2006. Rate of iodine volatilization and accumulation by filamentous fungi through laboratory cultures. *Chemosphere* **65**: 2216-2222.
- 142. Haas JR, Purvis OW.** 2006. Lichen biogeochemistry, p 344-276. In Gadd GM ed., *Fungi in Biogeochemical Cycles*. Cambridge University Press, Cambridge.
- 143. Chen J, Blume H-P, Beyer L.** 2000. Weathering of rocks induced by lichen colonization – a review. *Catena* **39**: 121-146.

- 144. Purvis OW, Pawlik-Skowronska B.** 2008. Lichens and metals, p 175-200. *In* Avery SV, Stratford M, van West P ed., *Stress in Yeasts and Filamentous Fungi*. Elsevier, Amsterdam.
- 145. Purvis OW.** 1996. Interactions of lichens with metals. *Science Prog* **79**: 283-309.
- 146. Purvis OW, Halls C.** 1996. A review of lichens in metal-enriched environments. *Lichenologist* **28**: 571-601.
- 147. Smith SE, Read DJ.** 1997. *Mycorrhizal Symbiosis*, 2nd edn. Academic Press, San Diego.
- 148. Wang B, Qiu, Y-L.** 2006. Phylogenetic distribution and evolution of mycorrhizas in land plants. *Mycorrhiza* **16**: 299-363.
- 149. Lapeyrie F., Picatto C., Gerard J, Dexheimer J.** 1990. TEM Study of intracellular and extracellular calcium oxalate accumulation by ectomycorrhizal fungi in pure culture or in association with *Eucalyptus* seedlings. *Symbiosis* **9**: 163-166.
- 150. Lapeyrie F, Ranger J, Vairelles D.** 1991. Phosphate-solubilizing activity of ectomycorrhizal fungi *in vitro*. *Can J Bot* **69**: 342-346.
- 151. Blaudez D, Botton B, Chalot M.** 2000. Cadmium uptake and subcellular compartmentation in the ectomycorrhizal fungus *Paxillus involutus*. *Microbiol* **146**: 1109-1117.
- 152. Fomina M, Charnock JM, Hillier S, Alexander IJ, Gadd GM.** 2006. Zinc phosphate transformations by the *Paxillus involutus*/pine ectomycorrhizal association. *Microbial Ecol* **52**: 322-333.
- 153. Christie P, Li XL, Chen BD.** 2004. Arbuscular mycorrhiza can depress translocation of zinc to shoots of host plants in soils moderately polluted with zinc. *Plant and Soil* **261**: 209-217.

- 154. Bellion M, Courbot M, Jacob C, Blaudez D, Chalot M.** 2006. Extracellular and cellular mechanisms sustaining metal tolerance in ectomycorrhizal fungi. *FEMS Microbiol Lett* **254**: 173-181.
- 155. Finlay R, Wallander H, Smits M, Holmstrom S, Van Hees P, Lian B, Rosling A.** 2009. The role of fungi in biogenic weathering in boreal forest soils. *Fungal Biol Rev* **23**: 101-106.
- 156. Smits MM, Bonneville S, Benning LG, Banwart SA, Leake JR.** 2012. Plant-driven weathering of apatite – the role of an ectomycorrhizal fungus. *Geobiol* **10**: 445-456.
- 157. McMaster TJ.** 2012. Atomic force microscopy of the fungi-mineral interface: applications in mineral dissolution, weathering and biogeochemistry. *Curr Opinion Biotechnol* **23**: 562-569.
- 158. Bonneville S, Morgan DJ, Schmalenberger A, Bray A, Brown A, Banwart SA, Benning LG.** 2011. Tree-mycorrhiza symbiosis accelerate mineral weathering: evidences from nanometer-scale elemental fluxes at the hypha-mineral interface. *Geochim Cosmochim Acta* **75**: 6988-7005.
- 159. Rosling A, Lindahl BD, Taylor AFS, Finlay RD.** 2004. Mycelial growth and substrate acidification of ectomycorrhizal fungi in response to different minerals. *FEMS Microbiol Ecol* **47**: 31-37.
- 160. Rosling A, Lindahl BD, Finlay RD.** 2004. Carbon allocation to ectomycorrhizal roots and mycelium colonising different mineral substrates. *New Phytol* **162**: 795-802.
- 161. Martino E, Perotto S, Parsons R, Gadd GM.** 2003. Solubilization of insoluble inorganic zinc compounds by ericoid mycorrhizal fungi derived from heavy metal polluted sites. *Soil Biol Biochem* **35**: 133-141.

- 162. Jongmans AG, van Breemen N, Lundstrom US, van Hees PAW, Finlay RD, Srinivasan M, Unestam T, Giesler R, Melkerud P-A, Olsson M.** 1997. Rock-eating fungi. *Nature* **389**: 682–683.
- 163. Van Breemen N, Lundstrom US, Jongmans AG.** 2000. Do plants drive podzolization via rock-eating mycorrhizal fungi? *Geoderma* **94**: 163-171.
- 164. Wallander H, Mahmood S, Hagerberg D, Johansson L, Pallon J.** 2003. Elemental composition of ectomycorrhizal mycelia identified by PCR-RFLP analysis and grown in contact with apatite or wood ash in forest soil. *FEMS Microbiol Ecol* **44**: 57-65.
- 165. Leyval C, Joner EJ.** 2001. Bioavailability of heavy metals in the mycorrhizosphere, p 165-185. In Gobran GR, Wenzel WW, Lombi E ed., *Trace Elements in the Rhizosphere*. CRC Press, Boca Raton, FL.
- 166. Kangwankraiphaisan T, Suntornvongsagul K, Sihanonth P, Klysubun W, Gadd GM.** 2013. Influence of arbuscular mycorrhizal fungi AMF. on zinc biogeochemistry in the rhizosphere of *Lindenbergia philippensis* growing in zinc-contaminated sediment. *Biometals* **26**: 489–505.
- 167. Christie P, Li XL, Chen BD.** 2004. Arbuscular mycorrhiza can depress translocation of zinc to shoots of host plants in soils moderately polluted with zinc. *Plant and Soil* **261**: 209-217.
- 168. Turnau K, Gawroński S, Ryszka P, Zook D.** 2012. Mycorrhizal-based phytostabilization of Zn–Pb tailings: lessons from the Trzebionka mining works Southern Poland., p 327-348. In Kothe E, Varma A ed., *Bio-geo Interactions in Metal-Contaminated Soils*. Springer-Verlag, Berlin.
- 169. Sand W.** 1997. Microbial mechanisms of deterioration of inorganic substrates: a general mechanistic overview. *Int Biodeter Biodeg* **40**: 183-190.

- 170. Wright JS.** 2002. Geomorphology and stone conservation: sandstone decay in Stoke-on-Trent. *Struct Surv* **20**: 50–61.
- 171. Gaylarde C, Morton G.** 2002. Biodeterioration of mineral materials, p 516–528. In Bitton G ed., *Environmental Microbiology*, Vol. 1. Wiley, New York.
- 172. Seaward MRD.** 2003. Lichens, agents of monumental destruction. *Microbiol Today* **30**, 110–112.
- 173. Lisci L, Monte M, and Pacini E.** 2003. Lichens and higher plants on stone: a review. *Int Biodet Biodegrad* **51**: 1–17.
- 174. Tiano P.** 2002. Biodegradation of cultural heritage: decay, mechanisms and control methods. *Seminar article, New University of Lisbon, Department of Conservation and Restoration*, 7-12 January 2002. http://www.arcchip.cz/w09/w09_tiano.pdf.
- 175. Pinzari F, Tate J, Bicchieri M, Rhee YJ, Gadd GM.** 2013. Biodegradation of ivory natural apatite.: possible involvement of fungal activity in biodeterioration of the Lewis Chessmen. *Environ Microbiol* **15**: 1050-1062.
- 176. Gu JD, Ford TE, Berke NS, Mitchell R.** 1998. Biodeterioration of concrete by the fungus *Fusarium*. *Int Biodet Biodegrad* **41**: 101-109.
- 177. Nica D, Davis JL, Kirby L, Zuo G, Roberts DJ.** 2000. Isolation and characterization of microorganisms involved in the biodeterioration of concrete in sewers. *Int Biodet Biodegrad* **46**: 61-68.
- 178. Gu JD.** 2009. Corrosion, microbial, p 259-269. In Schaechter M ed., *Encyclopedia of Microbiology*. Elsevier, Amsterdam.
- 179. Brandl H.** 2001. Heterotrophic leaching, p 383-423. In Gadd GM ed., *Fungi in Bioremediation*. Cambridge University Press, Cambridge.
- 180. Santhiya D, Ting YP.** 2005. Bioleaching of spent refinery processing catalyst using *Aspergillus niger* with high-yield oxalic acid. *J Biotechnol* **116**: 171-184.

- 1062 **181. Brandl H, Faramarzi MA.** 2006. Microbe-metal-interactions for the
1063 biotechnological treatment of metal-containing solid waste. *China Partic* **4**: 93-97.
- 1064 **182. De Rome L, Gadd GM.** 1987. Copper adsorption by *Rhizopus arrhizus*,
1065 *Cladosporium resinae* and *Penicillium italicum*. *Appl Microbiol Biotechnol* **26**: 84-90.
- 1066 **183. Gadd GM, Mowll JL.** 1985. Copper uptake by yeast-like cells, hyphae and
1067 chlamydospores of *Aureobasidium pullulans*. *Exp Mycol* **9**: 230-240.
- 1068 **184. Gadd GM, De Rome L.** 1988. Biosorption of copper by fungal melanin. *Appl*
1069 *Microbiol Biotechnol* **29**: 610-617.
- 1070 **185. Volesky B.** 1990. *Biosorption of Heavy Metals*. CRC Press, Boca Raton, FL.
- 1071 **186. Volesky B.** 2007. Biosorption and me. *Wat Res* **41**: 4017-4029.
- 1072 **187. Gadd GM, White C.** 1989. The removal of thorium from simulated acid process
1073 streams by fungal biomass. *Biotechnol Bioeng* **33**: 592-597.
- 1074 **188. Gadd GM, White C.** 1990. Biosorption of radionuclides by yeast and fungal biomass.
1075 *J Chem Technol Biotechnol* **49**: 331-343.
- 1076 **189. Gadd GM, White C.** 1992. Removal of thorium from simulated acid process streams
1077 by fungal biomass: potential for thorium desorption and reuse of biomass and desorbent.
1078 *J Chem Technol Biotechnol* **55**: 39-44.
- 1079 **190. Gadd GM, White C.** 1993. Microbial treatment of metal pollution - a working
1080 biotechnology? *Trends Biotechnol* **11**: 353-359
- 1081 **191. White C, Wilkinson SC, Gadd G.M.** 1995. The role of microorganisms in biosorption
1082 of toxic metals and radionuclides. *Int Biodeter Biodegrad* **35**: 17-40.
- 1083 **192. Wang HL, Chen C.** 2006. Biosorption of heavy metals by *Saccharomyces cerevisiae*:
1084 a review. *Biotechnol Adv* **24**: 427-451.
- 1085 **193. Wang J, Chen C.** 2009. Biosorbents for heavy metals removal and their future.
1086 *Biotechnol Adv* **27** page numbers???*??*

- 1087 **194. Dameron CT, Reese RN, Mehra RK, Kortan AR, Carroll PJ, Steigerwald ML,**
1088 **Brus LE, Winge DR.** 1989. Biosynthesis of cadmium sulphide quantum
1089 semiconductor crystallites. *Nature* **338**: 596-597.
- 1090 **195. Li Q, Liu D, Jia Z, Csetenyi L, Gadd GM.** 2016. Fungal biomineralization of
1091 manganese as a novel source of electrochemical materials. *Curr Biol* **26**: 950-955.
- 1092 **196. Thomson-Eagle ET, Frankenberger WT.** 1992. Bioremediation of soils
1093 contaminated with selenium, p 261-309. In Lal R, Stewart BA ed., *Advances in Soil*
1094 *Science*. Springer, New York.
- 1095 **197. Rosen K, Zhong WL, Martensson A.** 2005. Arbuscular mycorrhizal fungi mediated
1096 uptake of Cs-137 in leek and ryegrass. *Sci Total Environ* **338**: 283-290.
- 1097 **198. Gohre V, Paszkowski U.** 2006. Contribution of the arbuscular mycorrhizal symbiosis
1098 to heavy metal phytoremediation. *Planta* **223**: 1115-1122.
- 1099 **199. Salt DE, Smith RD, Raskin I.** 1998. Phytoremediation. *Ann Rev Plant Physiol Plant*
1100 *Mol Biol* **49**: 643-668.
- 1101 **200. Leyval C, Turnau K, Haselwandter K.** 1997. Effect of heavy metal pollution on
1102 mycorrhizal colonization and function: physiological, ecological and applied aspects.
1103 *Mycorrhiza* **7**: 139-153.
- 1104 **201. Krupa P, Kozdroj J.** 2004. Accumulation of heavy metals by ectomycorrhizal fungi
1105 colonizing birch trees growing in an industrial desert soil. *World J Microbiol*
1106 *Biotechnol* **20**: 427-430.
- 1107 **202. Adriaensen K, Vralstad T, Noben JP, Vangronsveld J, Colpaert JV.** 2005.
1108 Copper-adapted *Suillus luteus*, a symbiotic solution for pines colonizing Cu mine
1109 spoils. *Appl Environ Microbiol* **71**: 7279-7284.

- 1110 **203. Gonzalez-Chavez MC, Carrillo-Gonzalez R, Wright SF, Nichols KA.** 2004. The
1111 role of glomalin, a protein produced by arbuscular mycorrhizal fungi, in sequestering
1112 potentially toxic elements. *Environ Poll* **130**: 317-323.
- 1113 **204. Rufyikiri G, Huysmans L, Wannijn J, Van Hees M, Leyval C, Jakobsen I.** 2004.
1114 Arbuscular mycorrhizal fungi can decrease the uptake of uranium by subterranean
1115 clover grown at high levels of uranium in soil. *Environ Poll* **130**: 427-436.
- 1116 **205. Chen BD, Jakobsen I, Roos P, Zhu YG.** 2005. Effects of the mycorrhizal fungus
1117 *Glomus intraradices* on uranium uptake and accumulation by *Medicago truncatula* L.
1118 from uranium-contaminated soil. *Plant Soil* **275**: 349-359.
- 1119 **206. Chen BD, Zhu YG, Zhang XH, Jakobsen I.** 2005. The influence of mycorrhiza on
1120 uranium and phosphorus uptake by barley plants from a field-contaminated soil.
1121 *Environ Sci Poll Res* **12**: 325-331.
- 1122 **207. Bradley R, Burt AJ, Read DJ.** 1981. Mycorrhizal infection and resistance to heavy
1123 metals. *Nature* **292**: 335-337.
- 1124 **208. Bradley B, Burt AJ, Read DJ.** 1982. The biology of mycorrhiza in the Ericaceae.
1125 VIII. The role of mycorrhizal infection in heavy metal resistance. *New Phytol* **91**:
1126 197-209.
- 1127 **209. Schutzendubel A, Polle A.** 2002. Plant responses to abiotic stresses: heavy metal-
1128 induced oxidative stress and protection by mycorrhization. *J Exp Bot* **53**: 1351-1365.
- 1129 **210. Cairney JWG, Meharg AA.** 2003. Ericoid mycorrhiza: a partnership that exploits
1130 harsh edaphic conditions. *Eur J Soil Sci* **54**: 735-740.
- 1131 **211. Ruby MV, Davis A, Nicholson A.** 1994. In situ formation of lead phosphates in soils
1132 as a method to immobilize lead. *Environ Sci Technol* **28**: 646-654.
- 1133 **212. Cotter-Howells J.** 1996. Lead phosphate formation in soils. *Environ Pollut* **93**: 9-16.

- 1134 **213. Cotter-Howells J, Caporn S.** 1996. Remediation of contaminated land by formation
1135 of heavy metal phosphates. *Appl Geochem* **11**: 335-342.
- 1136 **214. Ioannidis TA, Zouboulis AI.** 2003. Detoxification of a highly toxic lead-loaded
1137 industrial solid waste by stabilization using apatites. *J Hazard Mater* **97**: 173-191.
- 1138 **215. Manning DAC.** 2008. Phosphate minerals, environmental pollution and sustainable
1139 agriculture. *Elements* **4**: 105-108.
- 1140 **216. Oelkers EH, Montel J-M.** 2008. Phosphates and nuclear waste storage. *Elements* **4**:
1141 113-116.
- 1142 **217. Ceci A, Rhee YJ, Kierans M, Hillier S, Pendlowski H, Gray N, Persiani AM,**
1143 **Gadd GM.** 2015. Transformation of vanadinite $Pb_5VO_{4.3}Cl$ by fungi. *Environ*
1144 *Microbiol* **17**: 2018-2034.
- 1145 **218. Ceci A, Kierans M, Hillier S, Persiani AM, Gadd GM.** 2015. Fungal bioweathering
1146 of mimetite and a general geomycological model for lead apatite mineral
1147 biotransformations. *Appl Environ Microbiol* **81**: 4955-4964.
- 1148 **219. Guggiari M, Bloque R, Aragno M, Verrecchia E, Job D, Junier P.** 2011.
1149 Experimental calcium-oxalate crystal production and dissolution by selected wood-rot
1150 fungi. *Int Biodet Biodegrad* **65**:803–809.
- 1151 **220. Fomina M, Hillier S, Charnock JM, Melville K, Alexander I.J, Gadd GM.** 2005.
1152 Role of oxalic acid over-excretion in toxic metal mineral transformations by
1153 *Beauveria caledonica*. *Appl Environ Microbiol* **71**: 371-381.
- 1154 **221. Daghino S, Turci F, Tomatis M, Favier A, Perotto S, Douki T, Fubini, B.** 2006.
1155 Soil fungi reduce the iron content and the DNA damaging effects of asbestos fibers.
1156 *Environ Sci Technol* **40**: 5793-5798.