University of Dundee

The Ax21 protein influences virulence and biofilm formation in the nosocomial pathogen *Stenotrophomonas maltophilia*

An, Shi-q; Tang, Ji-Liang

Published in:
Archives of Microbiology

DOI:
[10.1007/s00203-017-1433-7](https://doi.org/10.1007/s00203-017-1433-7)

Publication date:
2018

Licence:
CC BY

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

An, S., & Tang, J-L. (2018). The Ax21 protein influences virulence and biofilm formation in the nosocomial pathogen *Stenotrophomonas maltophilia*. *Archives of Microbiology*, 200(1), 183-187.
<https://doi.org/10.1007/s00203-017-1433-7>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The Ax21 protein influences virulence and biofilm formation in *Stenotrophomonas maltophilia*

Shi-qi An¹ · Ji-liang Tang²

Received: 22 August 2017 / Revised: 7 September 2017 / Accepted: 19 September 2017
© The Author(s) 2017. This article is an open access publication

Abstract *Stenotrophomonas maltophilia* is an antibiotic-resistant Gram-negative pathogen, which is associated with hospital-acquired infection. The genome encodes a protein highly related to the Ax21 protein of *Xanthomonas oryzae* that is implicated in interactions of this plant pathogen with rice. Here, we report on the pleiotropic nature of *ax21* mutation in *S. maltophilia* and the effects of addition of the Ax21 protein on the restoration of the wild-type phenotype. We show that loss by mutation of Ax21 leads to reduced motility, reduced biofilm formation, reduced tolerance to the antibiotic tobramycin and reduced virulence to larvae of *Galleria mellonella*, as well as alteration in the expression of specific genes associated with virulence or antibiotic resistance. Addition of the Ax21 protein restored motility and the level of gene expression towards wild type. These findings are consistent with the notion that the Ax21 protein is involved in intraspecies communication, although other interpretations cannot be discounted.

Keywords *Stenotrophomonas* · Antibiotic resistance · Motility · Biofilm formation · Virulence

Introduction

Ax21 is an outer membrane protein that is extensively conserved in plant pathogenic *Xanthomonas* and the associated genera *Xylella* and *Stenotrophomonas*, including *S. maltophilia*, some strains of which are hospital-acquired human pathogens (Ryan et al. 2009). Studies on Ax21 in *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) have shown that the protein can be sulphated and that it is secreted into the bacterial medium in association with the outer membrane vesicles (Han et al. 2012; Bahar et al. 2014). Although a sulphated peptide derived from the N-terminus of Ax21 was originally thought to be a specific trigger for XA21-dependent innate immunity in rice, subsequent work has shown that this is not the case. Nevertheless, this peptide does induce defence-related responses in plants (Danna et al. 2011). A second proposed role for Ax21 is as a diffusible signal that controls the gene expression in *Xoo* as a response to bacterial cell density (Bahar et al. 2014). These observations led us to examine the potential role(s) of Ax21 in *S. maltophilia*. In 2011, it was reported that the mutation of *ax21* had effects on different phenotypes in *S. maltophilia* (McCarthy et al. 2011). However, this paper was recently retracted due to errors in data presentation of Fig. 2 (McCarthy et al. 2017). Here, we report on the outcomes of repeated key experiments that indicate the pleiotropic nature of *ax21* mutation and the effects of addition of the Ax21 protein on the restoration of the wild-type phenotype. Here, our aim was to repeat key experiments that indicate the pleiotropic nature of *ax21* mutation and the effects of the addition of the Ax21 protein on the restoration of the wild-type phenotype.

Communicated by Jorge Membrillo-Hernández.

✉ Shi-qi An
s.an@dundee.ac.uk

✉ Ji-liang Tang
jltang@gxu.edu.cn

¹ Division of Molecular Microbiology, School of Life Sciences, University of Dundee, Dundee, UK

² College of Life Science and Technology, Guangxi University, 100 Daxue Road, Nanning, Guangxi 530004, China

Materials and methods

Bacterial strains and growth conditions

The wild-type *S. maltophilia* was strain K279a (Crossman et al. 2008). Unmarked deletion of *smlt0387* (designed as *ax21*) was constructed in *S. maltophilia* K279a strain using the pEX18Gm suicide vector, which uses the *sacB*-based counterselection method (Hoang et al. 1998). To delete *ax21*, the following procedure was used: a PCR fragment containing 150 nucleotides upstream of the *ax21* start site and 150 nucleotides downstream were amplified and then inserted into pEX18Gm that had been digested with the same enzymes and linked to a *Bam*HI–*Hind*III-digested PCR fragment. The pEX18Gm–*ax21* construct was sequenced to confirm that PCR did not introduce a mutation before the construct was mobilized from *E. coli* into *S. maltophilia* K279a. The transconjugants carrying the integrated plasmid on the chromosome were selected on L-agar plates containing 10% (wt/vol) sucrose (Oxoid, UK) and 25 µg/mL gentamicin (Fisher, UK). The resistant colonies were screened using colony PCR to identify mutants. For complementation studies, the *smlt0387* gene was cloned into pBBR1MCS (Kovach et al. 1995). Strains and plasmids used during this study are detailed in Table 1. For the majority of the experiments, the NYGB medium was used as growth medium for *S. maltophilia* strains, which comprises 20 g/L glycerol (Oxoid, UK), 3 g/L yeast extract (Difco, UK) and 5 g/L bacteriological peptone (Oxoid, UK). The assessment of bacterial clumping or biofilm formation was carried out in L medium, which comprises sodium chloride, 5 g/L; yeast extract, 5 g/L; Bactotryptone (Difco, UK), 10 g/L and D-glucose (Fisher, UK), 1 g/L. Peptides Ax21 (Smlt0387) and Ax21Y (Smlt0387 with Y altered to A) used in the experiments were generated by Cambridge Peptides (<http://www.cambridgepeptides.com>) and at 500 nM unless otherwise stated.

RNA extraction and qRT-PCR

For RNA extractions, *S. maltophilia* strains were cultivated at 30 °C in NYGB broth (without antibiotic) to logarithmic phase ($OD_{600} \approx 0.8$). A volume of 800 µL of RNA protect (Qiagen, UK) was added to 400 µL of culture and incubated at room temperature for 5 min. These suspensions were centrifuged and the resulting pellets were stored at –80 °C after removal of the supernatant. Following the manufacturer's instructions, total RNA was isolated from cells after thawing, using the RNeasy Mini Kit (Qiagen, UK) and then treated with DNase (Ambion, UK). PCR was used to confirm the removal of DNA contamination. Specific RT-PCR primers were used to amplify the central fragments of approximately 200 bp in length from *smlt1112* (Frd-AGG ACCCCTGGAACGTTTG; Rev-CACATCCGGCACCAC ATAGG), *smlt1390* (Frd-AGTTGGGCATCAACACCGAT; Rev-GGGTTGCCTTCTTGCTCTGA), *smlt2175* (Frd-AGC CAGAAGGAAACCACCTG; Rev-GCGGTCATAGGTCTG CACTT) and *smlt3949* (Frd-TTCCAGTTCGATAACGCC GC; Rev-CTCAGGCGACCCACATACAA). Quantification of gene expression was assessed using a Rotor-Gene Q (Qiagen, UK) and QuantiFast SYBR Green PCR Kit (Qiagen, UK). Furthermore, the expression of genes encoding the RNA polymerase sigma factor RpoN (Smlt1112), a putative outer membrane surface haemagglutinin (Smlt1390), putative TonB receptor (Smlt2175) and putative two-component regulator TctD (Smlt3949) were analysed because of their known role in virulence and antibiotic resistance in *Stenotrophomonas* (Devos et al. 2015; Ferrer-Navarro et al. 2016).

Motility assays

Bacterial motility assays were carried out on NYGB medium that was solidified using 0.6% Eiken agar (Eiken Chemical, Tokyo). A sterile 200-µl tip was used to inoculate *S. maltophilia* strains to the centre of the plate. The plates were visualized after incubation at 30 °C for 48 h.

Table 1 Bacterial strains and plasmids used in this work

Strain or plasmid	Relevant characteristics	Source or references
<i>Stenotrophomonas maltophilia</i>		
K279a	Clinical isolate	Crossman et al. (2008)
K279a <i>ax21</i>	<i>smlt0387</i> mutant of K279a	This study
<i>ax21</i> (pSmlt0387)	<i>ax21</i> mutant complemented with <i>smlt0387</i> using pBBR1MCS	This study
Plasmids		
pEX18Gm	Broad-host-range allelic exchange vector, Gm ^r	Hoang et al. (1998)
pBBR1MCS	Broad-host-range cloning vector, Cm ^r	Kovach et al. (1995)

Biofilm formation assay

Biofilm development was assessed on glass by crystal violet staining as described in O'Toole and Kolter (1998). *Stenotrophomonas maltophilia* strains were cultivated to logarithmic phase and then diluted to an OD at 600 nm of 0.1 in L medium. A volume of 5 mL of culture was incubated at 30 °C for 24 h in static glass tubes (14 mL). Once the medium and unattached cells were removed, adherent bacteria were washed twice with sterile water and then stained with 0.1% (w/v) crystal violet (Fisher, UK). Water was used to remove all unbound dye. The bound crystal violet was quantified by solubilizing in ethanol and reading at 595 nm.

Antibiotic killing curves

Killing curves were carried out at 30 °C as previously described by Macfarlane et al. (1999). *Stenotrophomonas maltophilia* strains were grown to mid-log phase on NYGB and then diluted to 10^6 in 100 mL of pre-warmed PBS containing the aminoglycoside tobramycin (Sigma, UK) at 100 µg/mL. Similarly, an antibiotic-free control was inoculated. At 0, 10, 20, 30, 50, 100, 120 and 180 min after antibiotic exposure, 0.1 mL volumes were removed, diluted in PBS and inoculated onto NYG agar plates. To determine viable CFU, these plates were incubated for 24 h at 30 °C.

Virulence assay

Galleria mellonella larvae were stored at 4 °C in wood shavings. For experiments, live versus dead larvae were observed after 24 h post-infection. *Galleria mellonella* were injected with 10 µL of successively diluted bacteria (1×10^6 CFU). Infected *G. mellonella* were placed on Whatman paper-lined Petri dishes and incubated at 37 °C. The *G. mellonella* were monitored for their survival after a 24-h period. Three separate tests were conducted consisting of ten larvae for each strain. The control groups for each experiment consisted of *G. mellonella* injected with PBS alone and a group of uninfected *G. mellonella*.

Results and discussion

The predicted proteome of *S. maltophilia* K279a, a clinical isolate (Crossman et al. 2008), contains two proteins that are homologous to Ax21 of *Xoo* strain PXO99^A: Smlt0387 (BLASTP probability score e-77) and Smlt0184 (e-62). Interestingly, Ax21 homologue Smlt0387 was recently identified in *S. maltophilia* to be secreted in association with outer membrane vesicles, which play roles in both infection and antibiotic resistance (Devos et al. 2015; Ferrer-Navarro et al. 2016).

The possible role of Ax21 in *S. maltophilia* was initially assessed by examination of the effect of deletion of *smlt0387* on a number of phenotypes. Deletion of *smlt0387* had a pleiotropic effect, leading to reduced motility on 0.6% Eiken agar (Fig. 1a), reduced biofilm formation on a glass surface (Fig. 1b), reduced tolerance to the aminoglycoside antibiotic

Fig. 1 Mutation of *ax21* has pleiotropic effects in *S. maltophilia* K279a. **a** The *ax21* mutant shows reduced motility in 0.6% Eiken agar, and complementation with *smlt0387* in *trans* restores motility

to wild type. **b** The *ax21* mutant shows reduced biofilm formation on glass as quantified by crystal violet staining. Complementation with *smlt0387* in *trans* restores motility to wild type

Fig. 2 Mutation of *ax21* has effects on antibiotic tolerance and virulence in *S. maltophilia* K279a. **a** The *ax21* mutant shows reduced tolerance to the aminoglycoside tobramycin at 100 µg/mL as revealed by a killing curve. **b** The *ax21* mutant shows reduced virulence in the *G.*

mellonella larva infection model. These mutant phenotypes could be restored to wild-type levels in all cases through complementation by *in trans* expression of a wild-type copy of the gene

tobramycin (Fig. 2a) and reduced virulence to larvae of *Galleria mellonella* (Fig. 2b). *In trans* complementation restored these altered phenotypes to the wild-type phenotype (Figs. 1 and 2). The pleiotropic effects of loss by mutation *Ax21* are consistent with previous observations in different pathovars of *Xanthomonas oryzae*, where *ax21* mutants have altered biofilm formation, extracellular polysaccharide synthesis and virulence (Qian et al. 2013; Park et al. 2014). Furthermore, a correlation has been shown between the abundance of *Ax21* in different strains of *S. maltophilia* and the mortality rate when those strains were tested in a Zebrafish model of infection (Ferrer-Navarro et al. 2013, 2016). Curiously, analysis of a strain with mutation in *smlt0184* did not show any alteration in the phenotypes tested.

For a number of bacterial cell-to-cell signalling systems, the phenotypic effects caused by deletion of the gene encoding the signal synthetase can be reversed by exogenous addition of the signal molecule (Papenfort and Bassler 2016). To test this potential role for *Ax21*, we repeated the motility tests in the presence of a synthetic *Ax21* protein. Addition of the protein at 500 nM restored wild-type motility (Fig. 3a). A variant *Ax21* protein (*Ax21Y*) in which the tyrosine residue that is sulphated in *Ax21* of *Xoo* was replaced by an alanine residue also restored motility (Fig. 3a). This is intriguing, since *S. maltophilia* lacks homologues of *RaxST* believed to be responsible for the sulphation of *Ax21* in *Xoo*.

Quantitative reverse transcription polymerase chain reaction (qRT-PCR) was then used to establish if the phenotypic

Fig. 3 Exogenous *Ax21* and the non-sulphatable Y22A variant form of *Ax21* (here designated *AX21Y*) are active in the regulation in *S. maltophilia*. **a** Exogenous addition of either *AX21* or *AX21Y* to the medium restored motility to an *ax21* mutant. **b** Exogenous addition

of either *AX21* or *AX21Y* to the *ax21* mutant restored the level of expression of *smlt1112*, *smlt1390*, *smlt2175* and *smlt3949* towards wild type as measured by qRT-PCR

effects of *smlt0387* deletion were associated with specific changes in gene expression. The findings (Fig. 3b) showed that loss of *ax21* led to elevated expression of *Smlt1112* and *Smlt1390*, but decrease in the expression of *Smlt2175* and *Smlt3949*. Addition of Ax21 or Ax21Y restored the expression of all of these genes towards wild type.

The effects of Ax21 on *S. maltophilia* thus appear to extend beyond changes that may influence the production or degradation of the molecule, consistent with the notion that the Ax21 protein is a signal involved in intraspecies communication (Winzer et al. 2002). However, other interpretations of the findings cannot be discounted. For example, pleiotropic effects may occur if loss of Ax21 causes dysfunction of the outer membrane leading to cell stress. Furthermore, work in *Xanthomonas* and *Stenotrophomonas* has indicated an influence of the DSF (Diffusible Signal Factor) cell-to-cell signal on the synthesis or secretion of Ax21 (Qian et al. 2013; Devos et al. 2015), raising the possibility that Ax21 acts indirectly through an influence on DSF signalling.

Conclusions

Ax21 influences a diverse range of functions in the nosocomial pathogen *S. maltophilia* leading to altered virulence, tobramycin tolerance and biofilm formation. Further work is needed to establish whether Ax21 is truly a cell–cell signal.

Acknowledgements We thank Robert Ryan, Max Dow and Delphine Caly for initial data, helpful discussions and critical reading of the manuscript. This work was supported by the Ba Gui Scholar Program of Guangxi Zhuang Autonomous region of China (No. 2014A002).

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Bahar O, Pruitt R, Luu DD, Schwessinger B, Daudi A, Liu F, Ruan R, Fontaine-Bodin L, Koebnik R, Ronald P (2014) The *Xanthomonas* Ax21 protein is processed by the general secretory system and is secreted in association with outer membrane vesicles. *Peer J* 2:e242. doi:10.7717/peerj.242
- Crossman LC, Gould VC, Dow JM, Vernikos GS, Okazaki A, Sebahia M, Saunders D, Arrowsmith C, Carver T, Peters N, Adlem E, Kerhornou A, Lord A, Murphy L, Seeger K, Squares R, Rutter S, Quail MA, Rajandream MA, Harris D, Churcher C, Bentley SD, Parkhill J, Thomson NR, Avison MB (2008) The complete genome, comparative and functional analysis of *Stenotrophomonas maltophilia* reveals an organism heavily shielded by drug resistance determinants. *Genome Biol* 9:R74. doi:10.1186/gb-2008-9-4-r74
- Danna CH, Millet YA, Koller T, Han SW, Bent AF, Ronald PC, Ausubel FM (2011) The *Arabidopsis* flagellin receptor FLS2 mediates the perception of *Xanthomonas* Ax21 secreted peptides. *Proc Natl Acad Sci USA* 108:9286–9291. doi:10.1073/pnas.1106366108
- Devos S, Van Oudenhoove L, Stremersch S, Van Putte W, De Rycke R, Van Driessche G, Vitse J, Raemdonck K, Devreese B (2015) The effect of imipenem and diffusible signaling factors on the secretion of outer membrane vesicles and associated Ax21 proteins in *Stenotrophomonas maltophilia*. *Front Microbiol* 6:298. doi:10.3389/fmicb.2015.00298
- Ferrer-Navarro M, Planell R, Yero D, Mongiardini E, Torrent G, Huedo P, Martínez P, Roher N, Mackenzie S, Gibert I, Daura X (2013) Abundance of the quorum-sensing factor Ax21 in four strains of *Stenotrophomonas maltophilia* correlates with mortality rate in a new zebrafish model of infection. *PLoS One* 8:e67207. doi:10.1371/journal.pone.0067207
- Ferrer-Navarro M, Torrent G, Mongiardini E, Conchillo-Solé O, Gibert I, Daura X (2016) Proteomic analysis of outer membrane proteins and vesicles of a clinical isolate and a collection strain of *Stenotrophomonas maltophilia*. *J Proteom* 142:122–129. doi:10.1016/j.jprot.2016.05.001
- Han SW, Lee SW, Bahar O, Schwessinger B, Robinson MR, Shaw JB, Madsen JA, Brodbelt JS, Ronald PC (2012) Tyrosine sulfation in a Gram-negative bacterium. *Nat Commun* 3:1153. doi:10.1038/ncomms2157
- Hoang TT, Karkhoff-Schweizer RR, Kutchma AJ, Schweizer HP (1998) A broad-host-range Flp-FRT recombination system for site-specific excision of chromosomally-located DNA sequences: application for isolation of unmarked *Pseudomonas aeruginosa* mutants. *Gene* 212:77–86
- Kovach ME, Elzer PH, Hill DS, Robertson GT, Farris MA, Roop RM 2nd, Peterson KM (1995) Four new derivatives of the broad-host-range cloning vector pBBR1MCS, carrying different antibiotic-resistance cassettes. *Gene* 166:175–176
- Macfarlane EL, Kwasnicka A, Ochs MM, Hancock RE (1999) PhoP-PhoQ homologues in *Pseudomonas aeruginosa* regulate expression of the outer-membrane protein OprH and polymyxin B resistance. *Mol Microbiol* 34:305–316
- McCarthy Y, Dow JM, Ryan RP (2011) The Ax21 protein is a cell–cell signal that regulates virulence in the nosocomial pathogen *Stenotrophomonas maltophilia*. *J Bacteriol* 193:6375–6378. doi:10.1128/JB.05949-11
- McCarthy Y, Dow JM, Ryan RP (2017) Retraction: the Ax21 protein is a cell–cell signal that regulates virulence in the nosocomial pathogen *Stenotrophomonas maltophilia*. *J Bacteriol* 99(10):e00156–e00157. doi:10.1128/JB.00156-17
- O'Toole GA, Kolter R (1998) Initiation of biofilm formation in *Pseudomonas fluorescens* WCS365 proceeds via multiple, convergent signalling pathways: a genetic analysis. *Mol Microbiol* 28:449–461
- Papenfort K, Bassler BL (2016) Quorum sensing signal-response systems in Gram-negative bacteria. *Nat Rev Microbiol* 14:576–588. doi:10.1038/nrmicro.2016.89
- Park HJ, Lee SW, Han SW (2014) Proteomic and functional analyses of a novel porin-like protein in *Xanthomonas oryzae* pv. *oryzae*. *J Microbiol* 52:1030–1035. doi:10.1007/s12275-014-4442-0
- Qian G, Zhou Y, Zhao Y, Song Z, Wang S, Fan J, Hu B, Venturi V, Liu F (2013) Proteomic analysis reveals novel extracellular virulence-associated proteins and functions regulated by the diffusible signal factor (DSF) in *Xanthomonas oryzae* pv. *oryzicola*. *J Proteome Res* 12:3327–3341. doi:10.1021/pr4001543
- Ryan RP, Monchy S, Cardinale M, Taghavi S, Crossman L, Avison MB, Berg G, van der Lelie D, Dow JM (2009) The versatility and adaptation of bacteria from the genus *Stenotrophomonas*. *Nat Rev Microbiol* 7:514–525. doi:10.1038/nrmicro2163
- Winzer K, Hardie KR, Williams P (2002) Bacterial cell-to-cell communication: sorry, can't talk now—gone to lunch! *Curr Opin Microbiol* 5:216–222