University of Dundee

Irradiance, as well as body site and timing of readings, is important in determining ultraviolet A minimal erythema dose. (Response to Gambichler et al. July BJD)

Eadie, E.; Ibbotson, S. H.; Dawe, R. S.

Published in:
British Journal of Dermatology

DOI:
[10.1111/bjd.16005](https://doi.org/10.1111/bjd.16005)

Publication date:
2018

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Eadie, E., Ibbotson, S. H., & Dawe, R. S. (2018). Irradiance, as well as body site and timing of readings, is important in determining ultraviolet A minimal erythema dose. (Response to Gambichler et al. July BJD). *British Journal of Dermatology*, 178(1), 297-298. <https://doi.org/10.1111/bjd.16005>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Irradiance, as well as body site and timing of readings, is important in determining ultraviolet A minimal erythematol dose. (Response to Gamblicher et al. July BJD)

Journal:	<i>British Journal of Dermatology</i>
Manuscript ID	Draft
Manuscript Type:	Letter to the Editor
Date Submitted by the Author:	n/a
Complete List of Authors:	Eadie, Ewan; Photobiology Unit, Department of Medical Physics, Ninewells Hospital and Medical School Ibbotson, Sally; Photobiology Unit, University Department of Dermatology Dawe, Robert; Ninewells Hospital and Medical School, Photobiology Unit, Department of Dermatology
Keywords:	minimal erythematol dose, UVA-1, irradiance

SCHOLARONE™
Manuscripts

This is the peer reviewed version of the following article: 'Irradiance, as well as body site and timing of readings, is important in determining ultraviolet A minimal erythematol dose. (Response to Gamblicher et al. July BJD)', *British Journal of Dermatology*, which has been published in final form at <http://dx.doi.org/10.1111/bjd.16005>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Irradiance, as well as body site and timing of readings, is important in determining ultraviolet A minimal erythema dose.

Word count: 350

Tables: 0

Figures: 0

Authors: E. Eadie¹, S. H. Ibbotson², R. S. Dawe³

¹Photobiology Unit, Department of Medical Physics, Ninewells Hospital and Medical School, Dundee, DD1 9SY.

²Photobiology Unit, University of Dundee, Department of Dermatology, Ninewells Hospital and Medical School, Dundee, DD1 9SY

³Photobiology Unit, Department of Dermatology, Ninewells Hospital and Medical School, Dundee, DD1 9SY

Corresponding Author: Ewan Eadie, Photobiology Unit, Ninewells Hospital and Medical School, Dundee, DD1 9SY. Ewan.eadie@nhs.net

Funding sources: None

Conflict of Interest: None declared

Dear Sir,

Irradiance, as well as body site and timing of readings, is important in determining ultraviolet A minimal erythema dose. (Response to Gambichler *et al.* July BJD)

Gambichler *et al.* demonstrated that, in their population, using a 25 mWcm^{-2} ultraviolet A-1 (UVA-1) source the median 24-hour delayed minimal erythema dose (MED) on the *inner forearm* was $> 130 \text{ Jcm}^{-2}$.¹ This differs from the 20 Jcm^{-2} to 28 Jcm^{-2} median MED reported from our centre.² The authors suggested the disparity might be explained by different methodologies. We agree and wish to expand on this point. Rather than being contradictory, the studies by Gambichler *et al.* and Beattie *et al.* are in excellent agreement.

The first main difference between the studies was the time when the MED was determined. Beattie *et al.* demonstrated that UVA-1 erythema peaked between 4 and 8 hours (h) with the MED being approximately half that at 24 hours.²

A second difference between the two studies was the site of testing. The inner forearm was tested in the Gambichler study whilst both back and inner forearm were tested in the Dundee study. Our study demonstrated that the back is around twice as sensitive to UVA1 as the inner forearm.

In recognition of this, in their discussion Gambichler *et al.* noted that their result at 24 h on the inner forearm should be compared with the Beattie result at the same time point and in the same body location, that is a median MED of 68 Jcm^{-2} .

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Critical, however, is a third difference between the two studies. In a study by Kagetsu *et al.* it was demonstrated that UVA-induced erythema is irradiance dependent,³ at 24 hour observations. They showed that a higher irradiance gave a lower MED.

The irradiance of the Gambichler *et al.* study was 25 mWcm⁻² and the Beattie study was 70 – 77 mWcm⁻². Taking the median MED from the recent study and correcting for time of observation, site of testing and irradiance of light source results in a reduction in MED from >130 Jcm⁻² to >24 Jcm⁻², similar to the median MED of 20 Jcm⁻² – 28 Jcm² reported by Beattie *et al.*

This highlights that variables including body site, time point *and irradiance*, must be considered when interpreting ultraviolet threshold erythema dose characteristics.

References:

1 Gambichler T, Majert J, Pljakic A *et al.* Determination of the minimal erythema dose for ultraviolet A1 radiation. *Br J Dermatol* 2017; **177**: 238-44.

2 Beattie PE, Dawe RS, Ferguson J *et al.* Dose-response and time-course characteristics of UV-A1 erythema. *Arch Dermatol* 2005; **141**: 1549-55.

3 Kagetsu N, Gange RW, Parrish JA. UVA-induced erythema, pigmentation, and skin surface temperature changes are irradiance dependent. *J Invest Dermatol* 1985; **85**: 445-7.