University of Dundee

Antibiotics and antibiotic resistance genes in global lakes

Yang, Yuyi; Song, Wenjuan; Lin, Hui; Wang, Weibo; Du, Linna; Xing, Wei

Published in:
Environment International

DOI:
[10.1016/j.envint.2018.04.011](https://doi.org/10.1016/j.envint.2018.04.011)

Publication date:
2018

Licence:
CC BY-NC-ND

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Yang, Y., Song, W., Lin, H., Wang, W., Du, L., & Xing, W. (2018). Antibiotics and antibiotic resistance genes in global lakes: A review and meta-analysis. *Environment International*, 116, 60-73.
<https://doi.org/10.1016/j.envint.2018.04.011>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

© 2018. This manuscript version is made available under the CC-BY-NC-ND 4.0
license <http://creativecommons.org/licenses/by-nc-nd/4.0/>

1 Antibiotics and antibiotic resistance genes in global lakes: A review and meta-analysis

2

3 Yuyi Yang^{a,b,*,1}, Wenjuan Song^{c,1}, Hui Lin^d, Weibo Wang^a, Linna Du^e, Wei Xing^{a,*}

4

5 ^a Key Laboratory of Aquatic Botany and Watershed Ecology, Wuhan Botanical

6 Garden, Chinese Academy of Sciences, Wuhan, 430074, China

7 ^b School of Life Sciences, University of Dundee, Dundee, DD1 5EH, Scotland, UK

8 ^c Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences,

9 Urumqi, 830011, China

10 ^d Institute of Environmental Resources and Soil Fertilizers, Zhejiang Academy of

11 Agricultural Sciences, Hangzhou, 310021, China

12 ^e Department of Agriculture and Biotechnology, Wenzhou Vocational College of

13 Science and Technology, Wenzhou, 325006, China

14

Corresponding author: Yuyi Yang, School of Life Sciences, University of Dundee, DD1 5EH, Scotland, UK, Email: ytyang@dundee.ac.uk; Wei Xing, Wuhan Botanical Garden, Chinese Academy of Sciences, Wuhan, 430074, China, Email: xingwei108@wbgcas.cn.

¹The authors contributed equally to this paper

Abstract

Lakes are an important source of freshwater, containing nearly 90% of the liquid surface fresh water worldwide. Long retention times in lakes mean pollutants from discharges slowly circulate around the lakes and may lead to high ecological risk for ecosystem and human health. In recent decades, antibiotics and antibiotic resistance genes (ARGs) have been regarded as emerging pollutants. The occurrence and distribution of antibiotics and ARGs in global freshwater lakes are summarized to show the pollution level of antibiotics and ARGs and to identify some of the potential risks to ecosystem and human health. Fifty-seven antibiotics were reported at least once in the studied lakes. Our meta-analysis shows that sulfamethoxazole, sulfamerazine, sulfameter, tetracycline, oxytetracycline, erythromycin, and roxithromycin were found at high concentrations in both lake water and lake sediment. There is no significant difference in the concentration of sulfonamides in lake water from China and that from other countries worldwide; however, there was a significant difference in quinolones. Erythromycin had the lowest predicted hazardous concentration for five percent of the species (HC₅) and the highest ecological risk in lakes. There was no significant difference in the concentration of sulfonamide resistance genes (*sul1* and *sul2*) in lake water and river water. There is surprisingly limited research on the role of aquatic biota in propagation of ARGs in freshwater lakes. As an environment that is susceptible to cumulative build-up of pollutants, lakes provide an important environment to study the fate of antibiotics and transport of ARGs with a broad range of niches including bacterial community, aquatic plants

37 and animals.

38 **Keywords:** species sensitivity distributions; niche; aquatic biota; antibiotics;

39 antibiotic resistance genes; meta-analysis

1. Introduction

The discovery of penicillin by Sir Alexander Fleming in 1928 opened the modern era of antibiotic innovation and development (Fleming 1929). Today, antibiotics are not only used as medicine for humans but are also widely used in animal husbandry and aquaculture. However, antibiotics that are unintentionally discharged into the environment pose a great threat to ecosystems and human health. These antibiotics can accumulate in food webs and, even more alarming, antibiotic resistance genes (ARGs) can be transferred between environmental bacteria and human pathogens (Bengtsson-Palme and Larsson 2015; Du and Liu 2012; Li et al. 2015; Martinez et al. 2015; Van Boeckel et al. 2015). Antibiotic resistance is not a new phenomenon (D'Costa et al. 2011; Wright and Poinar 2012), but the rapid and widespread increase of ARGs has been accelerated in recent years with the increase in discharge of antibiotics and other pollutants (e.g., heavy metals) into the environment (Bengtsson-Palme et al. 2014; Czekalski et al. 2014; Yang et al. 2017b; Yin et al. 2013). In fact, ARGs have recently been regarded as an emerging pollutant (Pruden et al. 2006). As such, antibiotics and their effects on the environment (ARGs, antibiotic resistant bacteria (ARB), etc.) have become an important theme in environmental science.

Aquatic environments are major pools for antibiotics and ARGs. Effluents from wastewater treatment plants, industry, hospitals and pig farms, for example, will all eventually reach some water source (Lavilla Lerma et al. 2014; Liu et al. 2012; Zhang et al. 2009a; Zhu et al. 2013). The effects of antibiotics on aquatic micro-organisms,

the nitrogen cycle and natural ecosystems have been summarized (Grenni et al. 2018; Roose-Amsaleg and Laverman 2016; Vălitalo et al. 2017). The distribution and environmental behaviour (e.g., adsorption and degradation) of antibiotics in aquatic environments have also been reviewed, (Bu et al. 2013; Kümmerer 2009a; b; Liu and Wong 2013). However, some aquatic environments, specifically lakes and rivers, behave differently due to their different hydraulic characteristics. In rivers, the concentration of pollutants in sediments gradually decreases downstream of a source due to hydraulic characteristics (Pruden et al. 2012; Reuther 2009). Rivers seem to have received most attention among aquatic environments, likely due to their rapid transport of antibiotics and ARGs and the obvious identification of pollution sources and landscapes in different reaches (Chen et al. 2013; Pruden et al. 2012; Rodriguez-Mozaz et al. 2015; Storteboom et al. 2010). In lakes, the residence time of contaminants increases because of the long water retention time relative to rivers. This means that pollutants from discharges slowly circulate around the lakes and makes pollution control in lake basins especially critical (Lyandres 2012; Reuther 2009). Due to these characteristics, lakes are predicted to have the potential to store and accumulate ARGs to a greater extent than rivers (Czekalski et al. 2015). Lakes serve as an important drinking source of freshwater, containing nearly 90% of the liquid surface fresh water worldwide; in contrast, rivers contain only 2% (McConnell and Abel 2013).

Until now, no comprehensive review of antibiotics and ARGs in lakes has been conducted. In this review, we collated the occurrence, distribution and risk of

antibiotics and ARGs in lakes. The objectives were to (1) identify antibiotics with high concentrations through a meta-analysis and prioritize antibiotics with potential ecological risk in lakes; (2) examine the difference in antibiotic pollution between lakes in China and in other countries worldwide, as well as the temporal trend of antibiotics in the most-studied lake, Taihu Lake; and (3) reveal the ARG pollution status in lakes. Factors influencing ARG propagation, potential risks of ARGs to human health, and the bioaccumulation of antibiotics in lakes were also discussed.

2. Methodology

2.1 Data collection

To survey antibiotics and ARGs in lakes worldwide, we used the Web of Knowledge database (<http://apps.webofknowledge.com>) to retrieve publications. Our search terms included [lake AND antibiotic] AND/OR [lake OR river AND/OR antibiotic resistance genes] AND/OR [lake OR river AND/OR antimicrobial resistance genes]. The database was searched for studies published up to 18 January 2018. Publications were then checked individually to eliminate any duplicates or irrelevant articles. Ultimately, 45 relevant papers reporting antibiotics in global lakes met our criteria. The name, country, antibiotic and/or ARG concentrations, and sampling dates are provided in Tables S1-S5, Supplementary material I. Seventeen and forty-seven studies respectively reported ARG analysis using culture-independent methods in lakes and rivers (Tables S1-S3, Supplementary material II). Trimethoprim/sulfamethoxazole (TMP/SMX), also known as co-trimoxazole, is used to treat a variety of bacterial infections, and TMP/SMX concentrations were included

with sulfonamides. It should be noted that several anonymous lakes in India that receive effluent from treatment plants had concentrations of quinolone antibiotics at $\mu\text{g L}^{-1}$ or mg L^{-1} levels (Fick et al. 2009), concentrations that are 10^3 to 10^4 times higher than those detected in other lakes (Table S3, Supplementary material I). Therefore, these lakes were not included in the meta-analysis. Toxicology data for antibiotics on aquatic organisms were retrieved from the United States Environmental Protection Agency (USEPA) ECOTOX database (<https://cfpub.epa.gov/ecotox/>). The search criteria used to gather the appropriate toxicity data included the following: freshwater medium, endpoint of lethal concentration required to kill 50% of the population (LC_{50}) or half maximal effective concentration (EC_{50}), exposure duration of <10 days, and test location of the laboratory. Toxicity data for samples without concentration ranges were not used in this study.

2.2 Statistical analysis

2.2.1 Meta-analysis

The mean concentrations of antibiotics were either provided in the publications or calculated using the original data. Concentrations that were “not detected” or “below detection limit” were entered as zero values, similar to a recent meta-analysis on organic contaminants (Meng et al. 2016). Antibiotic concentrations from the same lake but from different sampling years were considered as different data entries; however, antibiotic concentrations from the same lake but from different seasons in the same year were used to obtain a single mean value. The concentrations of antibiotics in the water and sediment were recorded in ng L^{-1} and ng g^{-1} , respectively.

Only antibiotics with at least two values were included in the meta-analysis. The median values and their confidence intervals for the mean concentrations of antibiotics in lakes across the Earth were calculated with the `wilcox.test` function in R software (R Foundation for Statistical Computing, Vienna, Austria). Comparisons of the concentrations of antibiotics in lakes from China and other countries worldwide were examined by a Mann-Whitney U test in R software. The temporal trend of antibiotics in Taihu Lake, China was analysed with a Pearson correlation analysis. Mann-Whitney U tests were used to compare the maximum relative abundances of *sul1* and *sul2* between lakes and rivers, and the absolute abundances of *sul1* between lakes and rivers.

2.2.2 Ecological risk assessment of antibiotics

Information on the toxicological impacts of many potentially harmful antibiotics is lacking (Välitalo et al. 2017). Therefore, as shown by Staples et al. (2008), antibiotics that had EC₅₀ or LC₅₀ data for at least four aquatic species within the ECOTOX database were analysed for species sensitivity distribution (SSD) analysis. The hazardous concentrations for 5% of the species (HC₅) derived from SSD were accompanied by a confidence limit that conveys information about the shape of the statistical distribution of toxicity values and their variance (Belanger et al. 2016). The values of HC₅ for antibiotics were calculated with R software. HC₅ values were then used to calculate the predicted no effect concentrations (PNEC) with an application factor of 5. The ecological risk of antibiotics in lake water was analysed with a hazard quotient (HQ), which is the ratio of the measured environmental concentration (MEC)

and PNEC.

3. Antibiotics in lakes

3.1 Overview of antibiotics detected in global lakes

The detection of antibiotics in lakes began in 2002 in Lake Ontario with the use of liquid chromatography-mass spectrometry (LC-MS) and liquid chromatography fluorescence detection (LC-FLD) (Nakata et al. 2005). Most later studies used the LC/MS/MS system, which is more efficient than only using the parent ions as in LC-MS analysis (Tables S1-S5, Supplementary material I). There were 57 antibiotics reported at least once in the water and sediments (Tables S1-S5, Supplementary material I). Antibiotics detected in lakes were classified into the following categories: sulfonamides, tetracyclines, quinolones, macrolides, lincosamides, and others (β -lactam, quinoxalines, and polyether and amphenicol antibiotics) (Table 1). The detection frequency may be related with the degradation and adsorption behaviour of antibiotics in lake water and sediments. Tetracyclines, fluoroquinolones and macrolides have a strong adsorption on particles and sediments (Huang et al. 2011; Kümmerer 2009a). Partitioning in the sediment could be one of the main reasons for the frequent detection of tetracyclines and fluoroquinolones in Chinese rivers (Zhou et al. 2011). β -Lactam antibiotics are easily hydrolysed in weakly acidic or alkaline conditions (Braschi et al. 2013) and could be effectively removed by ozonation, Fenton/photo-Fenton, or semiconductor photocatalysis (Homem and Santos 2011). For these reasons, β -lactam antibiotics may have a low detectability in some environments. The number of studies and the investigated antibiotics are summarized

in Table 1. In general, it can be concluded from Table 1 that (1) sulfonamides and quinolones are the most commonly investigated groups in lakes worldwide; (2) sulfonamides and tetracycline antibiotics are only found in lake sediments from China; (3) water was the main target for antibiotic analysis in freshwater lakes, rather than sediment; (4) China was the primary country for the studies on lake antibiotics, although 11 other countries also had studies (including Switzerland, Sweden, Turkey, Greece, Germany, Spain, Italy, the United States of America, Canada, India, and Bolivia) (Fig. S1 in Supplementary material I). No reports about antibiotics in lakes were recorded in Africa and Oceania, which are needed in further studies to understand the antibiotic pollution status in lakes worldwide.

3.2 Levels of antibiotics in the water and sediments of global lakes

3.2.1 Antibiotics in the bulk water of lakes

The average concentrations of 40 antibiotics in lake waters across the globe were retrieved, and the median concentration value and confidence intervals were calculated for each antibiotic (Fig. 1). Seven of the antibiotics had median values that were less than 1 ng L⁻¹ or at the ppt level, including sulfapyridine, sulfathiazole, lomefloxacin, danofloxacin, marbofloxacin, josamycin, and spiramycin. The median values for thirteen of the antibiotics were between 1 and 10 ng L⁻¹, including 5 sulfonamides, 1 tetracycline, 4 quinolones, 2 macrolides, and 1 lincosamide. Twenty of the antibiotics had median concentrations that were above 10 ng L⁻¹, including 6 sulfonamides and trimethoprim, 3 tetracyclines, 3 quinolones, 5 macrolides and lincosamides, and 2 other antibiotics.

Sulfonamides

Sulfamethoxazole (SMX), sulfameter (SME), trimethoprim (TMP) and sulfamerazine (SMR) each had median concentrations that were higher than 11 ng L⁻¹ (Fig. 1). The median concentration of SMX in the lakes was 26.75 ng L⁻¹ (Fig. 1), which is similar to concentrations detected in several rivers, including the Yangtze River, China (from <5 to 23 ng L⁻¹) (Chang et al. 2010) and the Seine River, France (from “not detected” to 40 ng L⁻¹) (Tamtam et al.). The concentration of SMX in the lakes was also lower than concentrations from the Nairobi River Basin in Kenya (from “below the limit of quantification” to 13,800 ng L⁻¹) (Ngumba et al. 2016) and reservoirs in China (1.0 to 585 ng L⁻¹) (Peng et al. 2014). High SMX concentrations were observed in the bulk water of Baiyangdian Lake, China (from “not detected” to 940 ng L⁻¹) (Li et al. 2012) and Lake Lead, United States of America (0.28 to 1200 ng L⁻¹) (Park et al. 2014). The median of the average SME concentration values was 16.95 ng L⁻¹ (Fig. 1), which was higher than concentrations from the Yellow Sea (from “not detected” to 1.2 ng L⁻¹) (Na et al. 2011) and the Daliao River, China (Jia et al. 2011). The highest SME concentrations were observed in Honghu Lake, China, which had an average value of 54.9 ng L⁻¹ (Wang et al. 2017c). The mean concentration of TMP in lake water samples from across the Earth was 12.60 ng L⁻¹ (Fig. 1). This concentration is comparable to that seen in waters from the Jialing Jiang River, China (5 to 7 ng L⁻¹) (Chang et al. 2010) and much lower than those detected in the Pearl River, China (56.4 to 605 ng L⁻¹) (Yang et al. 2011) and Haihe River, China (<LOQ-230 ng L⁻¹) (Luo et al. 2011). High TMP concentrations were found in Lake Ontario (from “not

detected” to 63.30 ng L⁻¹) in Canada (Li et al. 2010) and Lake Lead, United States of America (0.0015 to 120 ng L⁻¹) (Park et al. 2014).

Tetracyclines

Tetracycline (TC), oxytetracycline (OTC) and chlortetracycline (CTC) in lake water samples had median values of average concentrations exceeding 17 ng L⁻¹ (Fig. 1). TC concentrations in the bulk water of most lakes were close to those of Bohai Bay (from “below the limit of quantification” to 30 ng L⁻¹) (Zou et al. 2011), lower than those in the Huangpujiang River (from “not detected” to 114 ng L⁻¹) (Jiang et al. 2011) and rivers in Hong Kong (30 to 497 ng L⁻¹) (Selvam et al. 2017). The OTC concentrations in most lakes were close to that of the Yangtze River (ND-22.5 ng L⁻¹) (Yan et al. 2013) and lower than Huangpujiang River (from “not detected” to 219.8 ng L⁻¹) (Chen and Zhou 2014) and rivers in Hong Kong (7-104 ng L⁻¹) (Selvam et al. 2017). High OTC concentrations (>50 ng L⁻¹) were found in Taihu Lake (Xu et al. 2014), Baiyangdian Lake (Cheng et al. 2014; Cheng et al. 2017) and Honghu Lake (Wang et al. 2017c). High CTC concentrations were found in the bulk water of Taihu Lake with different sampling times from May 2010 to December 2015 (Hu et al. 2017a; Xu et al. 2014; Zhou et al. 2016) and Honghu Lake (Wang et al. 2017c).

Quinolones

Fourteen types of quinolone antibiotics have been detected in lake waters (Table S3, Supplementary material I). Ofloxacin (OFL), norfloxacin (NOR) and ciprofloxacin (CIP) were the most frequent. NOR and OFL had median concentrations of more than 10 ng L⁻¹ (Fig. 1), which were concentrations that are comparable to that of the

238 Jiulongjiang River (Zheng et al. 2011) and Yangtze River (Chang et al. 2010). High
239 concentrations of NOR (57.04 ng L⁻¹) and OFL (376.39 ng L⁻¹) were also found in
240 Baiyangdian Lake (Cheng et al. 2017).

241 *Macrolides and lincosamides*

242 Erythromycin (ETM), azithromycin, roxithromycin, lincomycin and clindamycin had
243 median concentrations higher than 10 ng L⁻¹. The ETM concentrations in most studies
244 were comparable to that of the Yangtze River, China (8 to 24 ng L⁻¹) (Chang et al.
245 2010) and lower than those of the Pearl River, China (40 to 560 ng L⁻¹) (Peng et al.
246 2011), Ume River, Sweden (80.3 to 160.5 ng L⁻¹) and Dal River, Sweden (347.4 to
247 694.8 ng L⁻¹) (Lindim et al. 2016). High ETM concentrations were observed in
248 Baiyangdian Lake (Li et al. 2012) and Taihu Lake (Xu et al. 2014), both in China.
249 High ROX concentrations were found in Baiyangdian Lake (from “not detected” to
250 155 ng L⁻¹) (Li et al. 2012) and Taihu Lake (from “not detected” to 218.3 ng L⁻¹) (Xu
251 et al. 2014). The ROX concentrations in most lakes were comparable to that found in
252 the Han River, South Korea (3 to 14 ng L⁻¹) (Choi et al. 2008b), as well as the Haihe
253 River (from “not detected” to 12 ng L⁻¹) (Zou et al. 2011) and Songhuajiang River
254 (0.2 to 11.5 ng L⁻¹) (Wang et al. 2017b) in China. High concentrations of lincomycin
255 (357 ng L⁻¹) and clindamycin (503 ng L⁻¹) were detected in Taihu Lake, China (Wu et
256 al. 2014).

257 *Others*

258 β -lactam (amoxicillin, oxacillin, ceftriaxone and cefazolin), quinoxaline (carbadox),
259 polyether (narasin) and amphenicol (florfenicol, chloramphenicol and thiamphenicol)

antibiotics were seldom reported, in contrast to other groups of antibiotics (Table S5, Supplementary material I). However, the median values for florfenicol and chloramphenicol exceeded 10 ng L^{-1} (Fig. 1). The mean concentrations of chloramphenicol in Taihu Lake, China reached 32.5 ng L^{-1} (Zhou et al. 2016), which were similar to those observed in the Han River, Korea (31 ng L^{-1}) (Choi et al. 2008a).

3.2.2 Antibiotics in lake sediments

The average concentrations of thirty-five antibiotics in lake sediments from across the globe were collected and their median values and confidence intervals calculated (Fig. 2). Six of the antibiotics had median values that were less than 1 ng g^{-1} , including 3 sulfonamides and 3 quinolones. The median concentrations of the 21 antibiotics were in the range of 1 to 10 ng g^{-1} . Only 8 antibiotics had median values that were higher than 10 ng g^{-1} (Fig. 2). These results indicate that the average concentrations of most antibiotics in lake sediments are often detected at concentrations less than 10 ng g^{-1} . SMX is a very well-studied sulfonamide antibiotic and has been recovered from the sediments from many lakes, including high concentrations in sediments from Honghu Lake, China (57.32 ng g^{-1}) and East Dongting Lake, China (30.57 ng g^{-1}) (Yang et al. 2016). TC had the highest median concentrations among the tetracycline antibiotics recovered from the sediment samples. High concentrations of TC were also found in lake sediments from Taihu Lake (Xu et al. 2014), Honghu Lake (Yang et al. 2016), and Baiyangdian Lake (Cheng et al. 2014), all in China. Baiyangdian Lake also had high ROX concentrations (from “not detected” to 302 ng g^{-1}) (Li et al. 2012).

3.2.3 Partition coefficient of antibiotics in lakes

A Mann-Whitney test shows that the median concentrations of the antibiotics and ARGs recovered from global lake water and lake sediment are significantly different ($p < 0.1$ level). Partitioning coefficients are important to understand the exchange of antibiotics between the water and sediment. The partitioning coefficients of antibiotics (K_d) between lake water and sediment have only been studied in two Chinese lakes, Bosten Lake (Lei et al. 2015) and Taihu Lake (Xu et al. 2014). Several K_d values were obtained or calculated from the data available from previous studies (Cheng et al. 2014; Zhang et al. 2015a; Zhang et al. 2018; Zhang et al. 2016; Zhou et al. 2016) (Table S6, Supplementary material I). A significant correlation was found between the log K_d values of the antibiotics and the log values of their octanol/water partition coefficients (K_{ow}) in Bosten Lake ($p < 0.1$); however, this correlation was not found for Taihu Lake (Fig. S2, Supplementary material I).

The normalized organic carbon distribution coefficients (K_{oc}) of antibiotics were shown to be positively correlated with their K_{ow} values in Chinese rivers (Li et al. 2018). However, K_{oc} is not a suitable measure for antibiotics in Taihu Lake since antibiotics may bind to mineral surfaces more strongly than to organic matter (Xu et al. 2014). Strong sorption of tetracycline and oxytetracycline to sediments of Taihu Lake is due to cation exchange and complexation reactions between antibiotics and the charged and polar sites of sorbents (Ji et al. 2016). Different partition coefficients of multi-phase distributions of antibiotics discovered in Baiyangdian Lake further showed that colloid-bounding influenced the environmental behaviour of antibiotics in lakes (Cheng et al. 2017). Clearly, in contrast to the partitioning coefficient data in

rivers (Li et al. 2018), more research is needed to understand the behaviour and fate of antibiotics in lakes.

3.2.4 Spatial difference and temporal trend of antibiotics in lakes

Due to the limited data available, only four antibiotics were used to compare lakes from China and lakes from other countries worldwide. The results of the Mann-Whitney test showed that there was no significant difference between SMX and SMZ in lake water from China and lake water from other countries (Fig. 3). However, the Chinese lakes showed significantly higher levels of NOR and OFL relative to levels observed for the other countries ($p < 0.05$). High concentrations of antibiotics frequently occurred in lakes in China (Fig. 3). These higher concentrations are perhaps not entirely unexpected, because the country has the highest production and usage of antibiotics (Zhang et al. 2015b).

Taihu Lake is one of the most heavily investigated lakes for antibiotic pollution (Ji et al. 2016; Xie et al. 2015; Xie et al. 2017; Xu et al. 2014; Zhang et al. 2013; Zhong et al. 2013; Zhou et al. 2016; Zhu et al. 2015). The mean concentrations of antibiotics recovered from Taihu Lake have been recorded for over four years. Figure 4 shows a possible temporal trend for antibiotic pollution. It is clear that the concentrations of SMX and OFL in the water decreased from 2010 to 2015 ($p < 0.05$ and $p < 0.1$ levels, respectively). The concentration of tetracyclines and SMZ, however, varied across this time interval (Fig. S3, Supplementary material I). Within sediment samples, the concentrations of SMX, sulfathiazole, SMZ, TC, OTC, CTC, OFL and NOR were fairly similar between 2013 and 2014 (Zhou et al. 2016), and lower than those in 2010

(Xu et al. 2014) (Tables S1-S3, Supplementary material I). Although it is difficult to determine a precise trend within the antibiotic pollution data available from Taihu Lake, it does appear as though the pollution in the lake is lessening.

3.2.5 Occurrence of antibiotics in aquatic biota

Antibiotic pollution within aquatic biota from lakes is rarely reported and has only been observed in Baiyangdian Lake (Li et al. 2012) and Taihu Lake (Xie et al. 2015; Xie et al. 2017). Only 13 antibiotics have been detected in these aquatic biota, with fewer antibiotics recovered from water than sediment samples (Table 2). Quinolones, especially CIPs, play a particularly important role in the accumulation of antibiotics among the aquatic plants of Baiyangdian Lake (8.37 to 6532 $\mu\text{g kg}^{-1}$) (Li et al. 2012). Quinolones (17.8 to 167 $\mu\text{g kg}^{-1}$) and macrolides (“below the limit of detection” to 182 $\mu\text{g kg}^{-1}$), particularly ROX, NOR, and OFL, were frequently detected within aquatic animals and birds (Li et al. 2012). The bioaccumulation factor (BAF) of antibiotics in aquatic biota from Baiyangdian Lake reached upwards of 17 000 L kg^{-1} (Li et al. 2012), while lower BAF values were found in Taihu Lake (Xie et al. 2015; Xie et al. 2017). The BAF values for aquatic biotas from other lakes and several rivers are summarized in Table 2 (Bai et al. 2014; Gao et al. 2012). BAFs were significantly different for the various antibiotics. These differences could be due to lipid solubility and the octanol-water partition coefficient (K_{ow}) of antibiotics, as well as sampling issues among the aquatic biotas (Li et al. 2012). pH is also an important factor influencing BAFs and has previously been shown to influence the bioaccumulation of antibiotics within *Daphnia* (Ding et al. 2016). “Bioaccumulative” chemicals are those

with BAFs greater than 5000 in aquatic organisms, and “potential bioaccumulative” chemicals are those with BAFs ranging from 2000 to 5000 (Gao et al. 2012). ROX, ENR and CIP are “bioaccumulative” chemicals, whereas SMZ, ETM, NOR and fleroxacin (FLE) are “potential bioaccumulative” chemicals. ENR and CIP showed higher BAFs than other reported antibiotics, and future studies should investigate their transport through the food chain.

3.3 Ecological risk of antibiotics

The inhibitory effects of antibiotics on aquatic biota depend on the type and dose of antibiotic, as well as the organisms sampled (Liu et al. 2014; Magdaleno et al. 2015). The HC₅ values of 17 antibiotics on aquatic biota ranged over 3 orders of magnitude (Fig. 5a and Table S7, Supplementary material I). The HC₅ of ETM was 11.73 µg L⁻¹, the lowest value among the antibiotics analysed. The vast majority of antibiotics had HC₅ values less than 100 µg L⁻¹, and only TMP was higher than 1500 µg L⁻¹. To better elucidate the ecological risk of antibiotics in lake water, the HQ values of mean concentrations of antibiotics in the lakes were analysed. HQ values ≤ 0.1 indicated an insignificant ecological risk, whereas HQ values < 0.1 and ≤ 1 represented a low ecological risk, values from 1 < and ≤ 10 were a moderate ecological risk, and HQ > 10 represented a high ecological risk (Ågerstrand and Rudén 2010). Notably, all the HQ values of the antibiotics examined here were below 0.1 (Fig. 5b); hence, there does not appear to be any significant ecological risk to aquatic biotas from antibiotics. ETM had the highest median HQ values, followed by clarithromycin (CLA) (Fig. 5b). ETM has previously been identified as a potential ecological risk to aquatic biotas in

rivers of China (Bu et al. 2013; Chen et al. 2017). Additionally, ETM, CLA, and azithromycin were included on a watch list of substances that could pose a significant ecological risk for aquatic environments in countries within the European Union (Välitalo et al. 2017). These results indicate that the macrolides in lakes worldwide may need special attention due to the ecological risk they pose. These values could also be used as references to evaluate the ecological risk of antibiotics in freshwater systems.

4. Antibiotic resistance genes in lakes

4.1 ARGs in lakes using culture-dependent methods

Antibiotic resistant bacteria (ARB) have been regarded as a threat to human health since the 1970s (Grabow et al. 1974). They have been isolated from lakes since the 1980s (Jones et al. 1986). *Escherichia coli*, *Enterococcus* and *Pseudomonas* are among the most frequency isolated, but, more recently, ARGs within ARBs have been discovered. ARGs found most frequently in *E. coli* from Hamilton Harbour, Lake Ontario, Canada were *tetA*, *tetB*, *bla_{TEM}*, *aadA1* and *sulII* (Hamelin et al. 2006). In organic-rich bacteria-contaminated sediments of Lake Geneva, Switzerland, 22% - 48% of *E. coli* and 16% - 37% of *Enterococcus* (ENT) carried *bla_{TEM}* (Thevenon et al. 2012). The most abundant resistance profile was seen in the resistance clones of 29 ARGs from Jinshan Lake, China; collectively, 88% of TC-resistant, 94% of SMX/TMP-resistant, 90% of ampicillin-resistant, 78% of streptomycin-resistant, and 72% of quinolone-resistant clones can be described by their corresponding ARGs (Wang et al. 2013). Among the extended spectrum beta-lactamase genes in bacteria

from Taihu Lake, China, *bla_{TEM}* was the most abundant, followed by *bla_{SHV}*, and *bla_{CTM-X}* was the least abundant (Yin et al. 2013). Most of the bacteria isolated from urban lakes in China carried at least two different functional ARGs; among these bacteria, gram-negative isolates had a higher detection frequency of ARGs than gram-positive isolates (Zhang et al. 2016).

4.2 ARGs in lakes based on qPCR and metagenomic methods

A total of 17 studies used culture-independent methods to quantify the abundance of ARGs in lakes (Tables S1-S2, Supplementary material II). The lakes assessed in the 17 included studies are located in the United States of America, Switzerland, Italy, India, Sweden, and China. Notably, nearly 50 published reports studied ARG abundance in rivers (Tables S1-S3, Supplementary material II). Additionally, 45 identified studies assessed the abundance of antibiotics in lakes.

Among the studies that assessed rivers, 34, 10, and 3 used qPCR, metagenomics, and high-throughput PCR (HT-PCR), respectively. Among the studies that assessed lakes, 14 used qPCR and 3 used metagenomics. HT-PCR has been used for ARG analysis of samples from swine farms (Zhu et al. 2013), rivers (Ouyang et al. 2015; Zheng et al. 2017; Zhou et al. 2017), and drinking water treatment plants (Xu et al. 2016). Although different issues arose in the studies that employed HT-PCR, this method still appears to be very promising, particularly because it is capable of processing a large number of samples simultaneously and detecting a large number of ARGs per sample (Lamas et al. 2016). HT-PCR may be an important alternative tool to conventional qPCR for ARG analysis in lakes.

414 A metagenomic analysis showed that Kazipally Lake, India was contaminated by
 415 fluoroquinolones and had a higher level of ARG abundance than the non-polluted
 416 Nydalasjön Lake, Sweden (Bengtsson-Palme et al. 2014). The abundances of ARGs in
 417 sediments of Lake Michigan, USA (Chu et al. 2017) and pristine Lake Namco and
 418 Qiangyangyong Glacier Lake, China (Chen et al. 2016) were also lower in
 419 comparison with effluents from sewage treatment plants. The *sul2* and *qnrD* genes
 420 were the most common ARGs in Kazipally Lake (Bengtsson-Palme et al. 2014), while
 421 the *bacA* gene was the most prevalent and dominant in pristine lakes on the Tibetan
 422 Plateau (Chen et al. 2016).

423 Sulfonamide resistance genes (*sul*) and tetracycline resistance genes (*tet*) were the
 424 most studied ARGs in lakes and rivers (Tables S1-S3, Supplementary material II). In
 425 lake water, the abundance of *sul1* is usually higher than that of *sul2* ($p < 0.1$ level).
 426 However, there is no significant difference between the relative abundances of *sul1*
 427 and *sul2* from lake and river water (Fig. 6). For the *sul1* gene, abundances in urban
 428 lakes in China (Yang et al. 2017b), 21 Swiss lakes (Czekalski et al. 2015), and Lake
 429 Geneva (Czekalski et al. 2012) were detected at 10^{-3} to 10^{-2} copies per 16S rRNA
 430 level. These levels are higher than those from Bosten Lake (10^{-5} to 10^{-3} copies per 16S
 431 rRNA) (Zhou et al. 2014) and similar to those from the Liuxi River (Xiong et al.
 432 2014). The abundance of the *tetA* gene reached upwards of 10^6 copies per mL in Lake
 433 Maggiore (Di Cesare et al. 2015) and 10^5 copies per mL in Taihu Lake (Zhang et al.
 434 2009c). These abundance levels were higher than those in the Huangpujiang River
 435 (10^3 copies per mL) (Jiang et al. 2013) and Lake Superior (LaPara et al. 2011).

All absolute abundances of *sulI* in lake sediments were in the range 10^{-6} to 10^{-10} copies per gram of sediment, which showed no difference compared with those in rivers ($p = 0.84$) (Fig. 6). Lake Geneva had a comparatively high abundance of *sulI*, with a value of 2.2×10^9 copies per gram of sediment (Czekalski et al. 2014). The abundance of *tet* genes in sediments of Taihu Lake (Zhang et al. 2009c) and Lake Geneva (Czekalski et al. 2014) reached upwards of 10^6 copies per gram, which is comparable to those in the Almendares River (Knapp et al. 2012). The abundance of *sul* genes was higher than that of *tet* genes in Lake Geneva (Czekalski et al. 2014) and 15 lakes along the Yangtze River (Yang et al. 2017a). Similar results were found in the sediments of the Haihe River (Dang et al. 2017) and Xiangjiang River (Xu et al. 2017). For other types of ARGs, the abundances of ARGs in Taihu Lake were found in the following order: *strB* > *qnrB* > *strA* > *qnrS* (Han et al. 2013). The relative abundances of *bla_{TEM}*, *aadA*, *bla_{CTX-M}*, and *bla_{SHV}* were higher than that of *bla_{NDM}* in Lake Geneva sediments (Devarajan et al. 2015).

4.3 The role of microbial community and mobile genetic elements in ARG propagation

Microbial community adaptation to pollutants or wastewater can result in changes in the ARG profile through several independent mechanisms, including (i) selective enrichment of a population (clonal amplification of ARB), (ii) advantageous mutations or gene rearrangements and duplication, and (iii) horizontal transfer of ecologically important genes, such as ARGs (Kondrashov 2012; Oh et al. 2013). Increases in ARG abundances are related to an increased number of possible ARBs in

the environment; this phenomenon was observed for the ARG abundance and number of *Enterobacteriaceae* in lakes in Brazilian zoos (de Faria et al. 2016) and *Bacteroidales* in rivers (Jia et al. 2014). Increased levels of ARGs and significant changes in the microbial community were adapted to the wastewater effluent in Lake Michigan (Chu et al. 2017) and antibiotic pollution in Kazipally Lake (Bengtsson-Palme et al. 2014). The bacterial community was identified as a contributing factor driving ARG composition other than mobile genetic elements (MGEs) in Tiaoxi River (Zheng et al. 2017) and Wen-Rui Tang River (Zhou et al. 2017).

The horizontal gene transfer (HGT) frequency of ARGs in aquatic bacteria was higher than that in terrestrial bacteria (Hu et al. 2016). Transformation, transduction, and conjugative transfer are the main mechanisms of HGT (Thomas and Nielsen 2005). Mobile genetic elements, such as plasmids, transposons, insertion sequences and integrons, are the important carriers in HGT of ARGs (Stokes and Gillings 2011). In Lake Michigan, high proportions of plasmid-associated ARGs were found in sediments, ranging from 32% to 100% of the total identified ARGs (Chu et al. 2017). Even in the pristine sediments from Lake Namco and Qiangyong Glacier Lake, a variety of integrons and plasmids existed, among which pGMI1000MP plasmid and CH34 megaplasmid were some of the most abundant (Chen et al. 2016). In an antibiotic-polluted Indian lake, 26 known and 21 putative novel plasmids were recovered in the lake metagenome (Bengtsson-Palme et al. 2014). A drastic increase in the abundance of integron integrases and ISCR elements was also found in the Ter

River, Spain after the river received some treated wastewater discharge (Lekunberri et al. 2018).

The class 1 integron (*intI1*) is one of the main vectors for HGT of ARGs in the environment. Nearly 40% of the isolated bacteria in Taihu Lake had *intI1* (Yin et al. 2013), which may facilitate the spread of ARGs between different bacteria. *intI1* has a significantly positive relationship with the abundance of ARGs in the aquatic environment, such as sediment samples from 15 lakes along the Yangtze River (Yang et al. 2017a), water from urban lakes (Yang et al. 2017b), the Haihe River (Dang et al. 2017) and the Tiaoxi River (Zheng et al. 2017). Interestingly, the relative abundances of *intI1* in lakes and rivers showed no significant difference in comparison with those in wastewater treatment plant sludges and treatment plant effluent via meta-analysis (Gillings 2018). The lack of a difference seems to further confirm that integrons play an important role in the HGT of ARGs in environment. Identification of the transfer of the mobile resistome in bacteria by bioinformatic analysis suggests that the transfer is mainly controlled by bacterial community composition (Hu et al. 2017b; Hu et al. 2016). Therefore, ARG propagation under selection pressure in lakes may mainly occur through enrichment of specific populations (ARB) and HGT of ARGs between different bacterial taxa.

4.4 Factors influencing ARG distribution in lakes

Chemical pollution

Chemical pollution discharged into lakes, such as antibiotics and heavy metals, may exert selection pressure on ARGs. The effects of antibiotics on ARG distribution

varied in each study. The positive pressure of antibiotics on ARG distribution in field surveys has been observed and/or analysed in East Dongting Lake, Honghu Lake (Yang et al. 2016), and Kazipally Lake (Bengtsson-Palme et al. 2014). Similar results were also observed in rivers in Pakistan (Khan et al. 2013) and China, including the Haihe River (Luo et al. 2010), Pearl River (Chen et al. 2015) and Beijiang River (Zou et al. 2009). However, a significant pressure of antibiotics on ARG distribution was not seen in rivers in South Africa (Suzuki et al. 2015) and Vietnam (Nakayama et al. 2017). OTC levels were even negatively correlated with the abundance of the *tetL* gene in Drweca River, Poland (Harnisz et al. 2015). In experimental settings, no observable effect of antibiotics on ARG abundances was found in lake sediment microcosms (Berglund et al. 2014). Increased selection on tetracycline resistance genes was apparent at exposure levels as low as 20 µg L⁻¹ OTC in some aquatic systems (Knapp et al. 2008). A recent study also showed that *in situ* degradation of antibiotics prevails at high concentrations of sulfonamides (5000 ng L⁻¹), whereas physiological adaptation by ARG spread would be more important under relatively lower concentrations of sulfonamides (1000 ng L⁻¹) (Vila-Costa et al. 2017). Therefore, studies that assess the selection pressure induced by the presence of antibiotics on the persistence and evolution of resistance genes in lakes, especially the long-term impact on antibiotics at environmental concentrations, are clearly required to help understand this relationship.

Heavy metals are another important factor influencing ARG distribution. The level of heavy metal contamination is several orders of magnitude greater than antibiotic

pollution (Stepanauskas et al. 2006) and may exert co-selection on ARGs (Baker-Austin et al. 2006; Di Cesare et al. 2016; Seiler and Berendonk 2012). Heavy metals have been regarded as a main cause of ARG contamination in sediments of Lake Geneva, Switzerland (Czekalski et al. 2014; Devarajan et al. 2015). Similar results were also found for the effect of heavy metals at sub-lethal levels on ARG distribution in the Xiangjiang River, China (Xu et al. 2017). Antibiotics and co-selection with heavy metals further accounted for the variation of ARGs in the bulk water of urban lakes (Yang et al. 2017b). In contrast, no relationship between ARGs and heavy metals was found in sediments from Chaohu Lake, China (Wu et al. 2016).

In addition, other pollutants have also been found to enrich ARGs in certain environments, including biocides and chemical preservatives (Romero et al. 2017). Biocide (benzyltrimethylammonium chloride) at environmental concentrations ($10 \mu\text{g L}^{-1}$) enriches specific ARGs, such as *bla*_{CTX-M} and *tetA*, in natural water from eutrophic lakes (Yang and Wang 2018). Even persistent organic pollutants, such as polycyclic aromatic hydrocarbons, unexpectedly accelerated the propagation of ARGs in coastal water (Wang et al. 2017a). Nanoparticles also helped shape the microbial community and increased the abundance of ARGs (Metch et al. 2018). Based on these findings, more comprehensive studies are needed to understand the individual and co-selection role of chemical pollution at environmental levels on the propagation of ARGs in lakes.

Physicochemical factors

The organic matter in the sediment was an important factor influencing ARG distribution in Lake Geneva (Devarajan et al. 2015) and Chaohu Lake (Wu et al. 2016). In Lake Maggiore, on the Italian-Swiss border, the abundance of *tetA* and *sul2* in bulk water was positively correlated with dissolved oxygen and negatively correlated with chlorophyll *a* (Di Cesare et al. 2015). Total organic carbon and total dissolved nitrogen were also positively correlated with the absolute abundance of total ARGs in urban rivers (Zhou et al. 2017). Conductivity, salinity, and dissolved oxygen concentration in Warta River, Poland were positively correlated with the abundance of sulfonamide resistance genes (Koczura et al. 2016). Unlike most living organisms, dissolved oxygen in lakes is not necessary for some bacteria to survive and shape the bacterial community (Baatar et al. 2016). As suggested by Di Cesare et al. (2015), resistant microbes inhabit specific niches in lakes or the aquatic environment.

Human activities

Human activities result in the input of ARGs into lakes through several ways, including effluent from hospitals and wastewater treatment plants (Guo et al. 2017; Rizzo et al. 2013; Zhang et al. 2009b) and runoff or discharge from agriculture, aquaculture, and animal husbandry (Heuer et al. 2011; Seiler and Berendonk 2012; Zhu et al. 2013) (Fig. 7). The parameters for evaluating the effect of human activities on ARGs in lakes are usually the number or capacity of wastewater treatment plants (WWTPs), distance to the point source (e.g., WWTP, hospital, aquaculture sites), and landscape (e.g., urban area, agriculture area), which can also be an indicator for pollution levels. Swiss lakes with a greater number and capacity of WWTPs in the

catchment usually exhibited high abundance of *sulI* (Czekalski et al. 2015). Similar results were also observed in the South Platte River Basin, USA where *sulI* abundances in river sediments were correlated with upstream capacities of animal feeding operations and wastewater treatment plants (Pruden et al. 2012). A high relative abundance of ARGs was also found in sites close to the sewage discharge point in Lake Geneva (Czekalski et al. 2014). A high abundance of *sul* and *tet* genes was additionally found in lakes along the Yangtze River, particularly near lakes with a high proportion of built-up land (Yang et al. 2017a). Similarly, Nanhu Lake and Shahu Lake, which are located in a high-density building area of China with elevated human activities, had a high relative abundance of total ARGs (Yang et al. 2017b).

4.5 The role of aquatic biota in the spread of ARGs.

Aquatic biotas play an important role in the propagation of ARGs but few studies have focused on this topic. *Daphnia* has been shown to be a refuge for ARGs in freshwater communities, where bacterial taxa associated with *Daphnia* were potential carriers of *tetA* (Eckert et al. 2016). Bacteriophages in sewage and rivers have also been uncovered as reservoirs for ARGs and could accelerate the spread of ARGs (Colombo et al. 2017; Colomer-Lluch et al. 2014; Colomer-Lluch et al. 2011). In the non-aquaculture freshwater Lake Mogan, Turkey, antibiotic resistance is common among the phenotypes of the fish-mucus-dwelling bacterial populations (Ozaktas et al. 2012). In fish farms, fish faeces contributed to ARG enrichment in sediments despite a lack of contemporaneous antibiotic treatments at the farms and different resistome compositions in the intestinal contents of individual farmed fish (Muziasari et al.

2017). The hindgut of zebra fish was the most important intestinal region supporting ARG dissemination and helped promote the spread of ARG into the water environment (Fu et al. 2017). Shrimp in Iskenderun Bay, Turkey also harbour multi-resistant bacteria that may propagate dissemination of ARGs in the environment (Matyar et al. 2008). Different aquatic plant species and tissues contained vastly different bacterial communities (Crump and Koch 2008), which may affect the composition and transport behaviours of ARGs. Clearly, aquatic biotas are an important carrier and/or refuge for ARGs and a useful media for the exchange of ARGs between aquatic biota and the environment (Fig. 7). A broad range of niches, not only bacteria but also phages, aquatic plants and animals, therefore, should be considered when the enrichment, fate and transport of ARGs in lakes are evaluated.

4.6 Potential risk of ARGs in lakes to humans

Unlike antibiotics, there is no available method to calculate the potential risks to humans based on the presence of ARGs in lakes. Risk assessment frameworks of antibiotic resistance have been proposed and include hazard identification, hazard assessment/characteristics, exposure assessment, and risk assessment (Chereau et al. 2017; Manaia 2017). However, there are many factors hindering the risk assessment of ARGs in lakes for human health, such as the uncertain affiliation between bacterial taxa and ARGs (Manaia 2017), and the unclear doses of ARB for various health outcomes and pathways (Ashbolt et al. 2013). The exposure pathways of ARGs in lakes to humans can be classified into three categories: (1) ingestion of water containing ARGs, (2) ingestion of aquatic products, and (3) direct contact with the

lake, such as recreation and swimming (Chereau et al. 2017; Eckert et al. 2018) (Fig. 7). Regarding the exposure frequency, the ingestion of lake water containing ARGs (i.e., drinking water) is the main risk to humans. Many lakes are important drinking water sources for humans, including Lake Michigan, Lake Geneva, and Lake Zurich. Recent studies have shown that bacteria and ARGs were still present in tap water after treatment, though significantly reduced (Su et al. 2018). In one case, the relative abundance of ARGs in drinking water even increased after chlorination (Jia et al. 2015). Chlorine disinfection is not able to destroy antibiotic-resistance genes, but it can inactivate and decrease bacterial counts of ARBs (Furukawa et al. 2017). ARGs in inactive bacteria or naked DNA persist and may propagate through HGT to elevate the risk to humans. Therefore, regarding the risk of ARGs to humans, both ARGs in active bacteria and naked DNA must be considered. Comprehensive analyses of the persistence of ARGs in environment and their resistance to conventional drinking water treatment methods are also important to develop a priority list of harmful ARGs to humans.

5. Perspectives of study on antibiotics and ARGs in lakes

5.1 Priority lists of harmful antibiotics in lakes

Fifty-seven kinds of antibiotics were detected in lakes worldwide; however, many antibiotics, such as oxazolidinones and ansamycins, have not yet been investigated in lake environments and were not included here. Additionally, metabolites of antibiotics, again not studied here, may be harmful to the aquatic environment and have not yet received sufficient study (Baumann et al. 2015). The non-targeted, large screening of

over a thousand emerging contaminants is possible through ultra-high-performance liquid chromatography coupled with quadrupole time-of-flight mass spectrometry (UHPLC-QTOF MS) (Hernández et al. 2015). This technique could be a powerful tool for the analysis of antibiotics and their metabolites in lakes. This information could be combined with BAFs and ecotoxicology data to develop a priority list of harmful antibiotics in the aquatic environment and assist in lake management.

5.2 The relationship between chemical pollution and ARGs

Some of the antibiotics detected in the lakes studied here were sulfonamides, tetracyclines, quinolones and macrolides. The most common types of ARGs detected in lakes were *sul* and *tet* genes. Additional types of ARGs were discovered using the metagenomic method and HT-qPCR, such as ARGs based on the antibiotics to which they conferred resistance: aminoglycosides, β -lactams, chloramphenicol, macrolide-lincosamide-streptogramin B (MLSB), and vancomycin (Zheng et al. 2017; Zhu et al. 2017). However, few studies have conducted a combined analysis of antibiotics and ARGs. The large-scale screening of organic pollutants, including antibiotics and heavy metals, and ARG profiles should be applied to develop a more comprehensive understanding of the relationship between chemical pollution and ARGs in freshwater lakes. In addition, a good understanding of the lake microcosm or macrocosm is needed to evaluate the individual and co-selection role of chemical pollution on the propagation of ARGs in lakes at their environmental level. This approach is useful to identify the main pollutants exerting selection pressure on ARG propagation and influencing the persistence of ARGs in lakes.

5.3 ARGs and bacterial communities in lakes

The antibiotic resistance of *Escherichia coli*, *Enterococcus* and *Pseudomonas* present in lakes has been well studied and provides important information about the transfer of ARGs in the bacterial community. Furthermore, there is still a significant difference between the profile of ARB via metagenomic analysis and culture-dependent ARB communities (Narciso-da-Rocha and Manaia 2016). Moreover, ARGs and their affiliated taxa are still very unclear. Hence, culture-independent methods should be used with culture-based approaches to identify major sources of ARB profiles. This would help to understand the composition of ARB in the bacterial community in freshwater lakes and assess the transfer risk between lake ARB and humans. It would also be beneficial to study the evolution of ARGs in different bacterial taxonomic groups based on isolated bacteria and their ARG sequences.

5.4 Transport of antibiotics and ARGs in lakes

Specific hydraulic conditions in lakes, such as long retention of water exchange, may result in special characteristics in the antibiotic distribution and ARG spread in the lake ecosystem. Furthermore, lake sediments can be used as natural archives of perturbation histories, including acidification, eutrophication, and pollution; lake sediments can also provide important information on the historical trend of chemical pollution and ARGs (Anderson 1995; Eckert et al. 2018). Lakes are clearly an important environment to investigate the fate of antibiotics and spread of ARGs (Fig. 7). Once chemical pollutants (e.g., antibiotics and heavy metals) are discharged into lakes via runoff or rivers, partitioning between water and sediment can occur, and, via

disturbance, sediment could be a secondary pollution source. Antibiotics can also accumulate in aquatic biota through the food chain. ARGs discharged into the environment will be enriched, selected, or transferred between different bacterial taxa under the selection pressure of chemical pollution. ARG carriers, such as bacteria, phages, and even extracellular DNA, can be adsorbed on the surface of aquatic biota or inhaled into the alimentary canal of aquatic animals. The faeces from these aquatic animals or their decomposing bodies release ARGs into the sediment, resulting in changes in the ARG composition of sediments. Antibiotics and ARGs in water, sediment, and aquatic biota should be studied to give a more complete understanding of their relationship and to better assess the fate and risk of antibiotics, ARGs and ARB in freshwater lakes.

6. Conclusion

This paper systematically reviewed the distribution of antibiotics and ARGs in global lakes. A total of 57 antibiotics reported at least once in the water and sediments of global lakes were assessed. The concentrations of these antibiotics in lake water were higher than those in lake sediment. As seen from our meta-analysis, sulfamethoxazole, sulfamerazine, sulfameter, tetracycline, oxytetracycline, erythromycin, and roxithromycin were found at high concentrations in both the lake water and sediment. There was also no significant difference in sulfamethoxazole and sulfamethazine concentrations in lake waters from China and the lake water from other countries worldwide; however, there was a significant difference in ofloxacin and norfloxacin concentrations between lake waters from China and from the rest of the world. The

concentration of sulfamethoxazole and ofloxacin in bulk water from Taihu Lake significantly decreased between 2010 and 2015. ROX, ENR and CIP are “bioaccumulative” chemicals in aquatic organisms, whereas ETM had the highest ecological risk based on hazard quotient. There was also no significant difference in the concentration of *sul* genes in rivers and lakes. Chemical pollution, including antibiotics and heavy metals, influenced the distribution of ARGs in lakes through selection pressure. The bacterial community is the principle force in shaping the resistome, followed by mobile genetic elements.

Unfortunately, the continuous discharge of chemical pollutants into lakes will not be readily solved in the near future, and selection pressure on ARGs in lakes will continue. It is therefore urgent that we (1) develop priority lists of antibiotics and their metabolites in lakes; (2) identify some of the key pollutants influencing ARG propagation for proper lake management; (3) identify the persistence of ARGs in the environment and their resistance to conventional drinking water treatment methods; (4) understand the long-term ARG propagation exerted by individual or combined pollutants at environmental concentrations; and (5) understand the role of aquatic biota in the transport of antibiotics and ARG propagation.

Declaration of interest

None

Acknowledgements

The authors thank the anonymous reviewers and editor Prof. Michael Gillings for their helpful and constructive comments that greatly contributed to improving the

722 final version of the paper. They would also like to thank Prof. Stephen Christopher
723 Maberly from The Centre for Ecology & Hydrology, United Kingdom for his
724 comments and improving the English level of this manuscript.

725 This project was supported in part by National Natural Science Foundation of China
726 [grant number 31400113, 31370479], Youth Innovation Promotion Association of
727 Chinese Academy of Sciences [grant number 2015282], Wenzhou Science and
728 Technology Major Project [grant number 2016N0031], Basic public welfare research
729 projects in Zhejiang province [grant number LGF18E090007], and State Key
730 Laboratory Breeding Base for Zhejiang Sustainable Pest and Disease Control [grant
731 number 2010DS700124-ZZ1802].

References

- Ågerstrand, M., Rudén, C., 2010. Evaluation of the accuracy and consistency of the Swedish environmental classification and information system for pharmaceuticals. *Sci. Total. Environ.* 408, 2327-2339.
- Anderson, N.J., 1995. Using the past to predict the future: lake sediments and the modelling of limnological disturbance. *Ecol. Model.* 78, 149-172.
- Ashbolt, N.J., Amezcuita, A., Backhaus, T., Borriello, P., Brandt, K.K., Collignon, P., Coors, A., Finley, R., Gaze, W.H., Heberer, T., Lawrence, J.R., Larsson, D.G.J., McEwen, S.A., Ryan, J.J., Schonfeld, J., Silley, P., Snape, J.R., Van den Eede, C., Topp, E., 2013. Human health risk assessment (HHRA) for environmental development and transfer of antibiotic resistance. *Environ. Health Persp.* 121, 993-1001.
- Baatar, B., Chiang, P.W., Rogozin, D.Y., Wu, Y.T., Tseng, C.H., Yang, C.Y., Chiu, H.H., Oyuntsetseg, B., Degermendzhy, A.G., Tang, S.L., 2016. Bacterial communities of three saline meromictic lakes in Central Asia. *Plos One* 11, e0150847.
- Bai, Y., Meng, W., Xu, J., Zhang, Y., Guo, C., 2014. Occurrence, distribution and bioaccumulation of antibiotics in the Liao River Basin in China. *Environ. Sci. Process. Impacts* 16, 586-593.
- Baker-Austin, C., Wright, M.S., Stepanauskas, R., McArthur, J., 2006. Co-selection of antibiotic and metal resistance. *Trends Microbiol.* 14, 176-182.
- Baumann, M., Weiss, K., Maletzki, D., Schüssler, W., Schudoma, D., Kopf, W., Kühnen, U., 2015. Aquatic toxicity of the macrolide antibiotic clarithromycin and its metabolites. *Chemosphere* 120, 192-198.
- Belanger, S., Barron, M., Craig, P., Dyer, S., Galay-Burgos, M., Hamer, M., Marshall, S., Posthuma, L., Raimondo, S., Whitehouse, P., 2016. Future needs and recommendations in the development of species sensitivity distributions: Estimating toxicity thresholds for aquatic ecological communities and assessing impacts of chemical exposures. *Integr Environ Assess Manag* 13, 664-674.
- Bengtsson-Palme, J., Boulund, F., Fick, J., Kristiansson, E., Larsson, D.G.J., 2014. Shotgun metagenomics reveals a wide array of antibiotic resistance genes and mobile elements in a polluted lake in India. *Front. Microbiol.* 5, e648.
- Bengtsson-Palme, J., Larsson, D.G.J., 2015. Antibiotic resistance genes in the environment: prioritizing risks. *Nat. Rev. Microbiol.* 13, 396-396.
- Berglund, B., Khan, G.A., Lindberg, R., Fick, J., Lindgren, P.E., 2014. Abundance and dynamics of antibiotic resistance genes and integrons in lake sediment microcosms. *Plos One* 9, e10815.
- Braschi, I., Blasioli, S., Fellet, C., Lorenzini, R., Garelli, A., Pori, M., Giacomini, D., 2013. Persistence and degradation of new β -lactam antibiotics in the soil and water environment. *Chemosphere* 93, 152-159.
- Bu, Q., Wang, B., Huang, J., Deng, S., Yu, G., 2013. Pharmaceuticals and personal care products in the aquatic environment in China: A review. *J. Hazard. Mater.*

775 262, 189-211.

776 Chang, X., Meyer, M.T., Liu, X., Zhao, Q., Chen, H., Chen, J.-a., Qiu, Z., Yang, L.,
777 Cao, J., Shu, W., 2010. Determination of antibiotics in sewage from hospitals,
778 nursery and slaughter house, wastewater treatment plant and source water in
779 Chongqing region of Three Gorge Reservoir in China. *Environ. Pollut.* 158,
780 1444-1450.

781 Chen, B.W., Liang, X.M., Huang, X.P., Zhang, T., Li, X.D., 2013. Differentiating
782 anthropogenic impacts on ARGs in the Pearl River Estuary by using suitable
783 gene indicators. *Water Res.* 47, 2811-2820.

784 Chen, B.W., Liang, X.M., Nie, X.P., Huang, X.P., Zou, S.C., Li, X.D., 2015. The role
785 of class I integrons in the dissemination of sulfonamide resistance genes in the
786 Pearl River and Pearl River Estuary, South China. *J. Hazard. Mater.* 282,
787 61-67.

788 Chen, B.W., Yuan, K., Chen, X., Yang, Y., Zhang, T., Wang, Y.W., Luan, T.G., Zou,
789 S.C., Li, X.D., 2016. Metagenomic analysis revealing antibiotic resistance
790 genes (ARGs) and their genetic compartments in the Tibetan environment.
791 *Environ. Sci. Technol.* 50, 6670-6679.

792 Chen, H., Jing, L., Teng, Y., Wang, J., 2017. Characterization of antibiotics in a
793 large-scale river system of China: Occurrence pattern, spatiotemporal
794 distribution and environmental risks. *Sci. Total Environ.* 618, 409-418.

795 Chen, K., Zhou, J.L., 2014. Occurrence and behavior of antibiotics in water and
796 sediments from the Huangpu River, Shanghai, China. *Chemosphere* 95,
797 604-612.

798 Cheng, D.M., Liu, X.H., Wang, L., Gong, W.W., Liu, G.N., Fu, W.J., Cheng, M., 2014.
799 Seasonal variation and sediment-water exchange of antibiotics in a shallower
800 large lake in North China. *Sci. Total. Environ.* 476, 266-275.

801 Cheng, D.M., Liu, X.H., Zhao, S.N., Cui, B.S., Bai, J.H., Li, Z.J., 2017. Influence of
802 the natural colloids on the multi-phase distributions of antibiotics in the
803 surface water from the largest lake in North China. *Sci. Total. Environ.* 578,
804 649-659.

805 Chereau, F., Opatowski, L., Tourdjman, M., Vong, S., 2017. Risk assessment for
806 antibiotic resistance in South East Asia. *BMJ* 5, 358, doi: 10.1136/bmj.j3393.

807 Choi, K., Kim, Y., Jung, J., Kim, M.H., Kim, C.S., Kim, N.H., Park, J., 2008a.
808 Occurrences and ecological risks of roxithromycin, trimethoprim, and
809 chloramphenicol in the Han River, Korea. *Environ. Toxicol. Chem* 27,
810 711-719.

811 Choi, K., Kim, Y., Park, J., Park, C.K., Kim, M., Kim, H.S., Kim, P., 2008b. Seasonal
812 variations of several pharmaceutical residues in surface water and sewage
813 treatment plants of Han River, Korea. *Sci. Total. Environ.* 405, 120-128.

814 Chu, B.T.T., Petrovich, M.L., Chaudhary, A., Wright, D., Murphy, B., Wells, G.,
815 Poretsky, R., 2017. Metagenomic analysis reveals the impact of wastewater
816 treatment plants on the dispersal of microorganisms and genes in aquatic
817 sediments. *Appl. Environ. Microbiol.* doi: 10.1128/AEM.02168-17

818 Colombo, S., Arioli, S., Neri, E., Della Scala, G., Gargari, G., Mora, D., 2017.

819 Viromes as genetic reservoir for the microbial communities in aquatic
820 environments: A focus on antimicrobial-resistance genes. *Front. Microbiol.* 8,
821 e1095.

822 Colomer-Lluch, M., Calero-Caceres, W., Jebri, S., Hmaied, F., Muniesa, M., Jofre, J.,
823 2014. Antibiotic resistance genes in bacterial and bacteriophage fractions of
824 Tunisian and Spanish wastewaters as markers to compare the antibiotic
825 resistance patterns in each population. *Environ. Int.* 73, 167-175.

826 Colomer-Lluch, M., Jofre, J., Muniesa, M., 2011. Antibiotic resistance genes in the
827 bacteriophage DNA fraction of environmental samples. *Plos One* 6, e17549.

828 Crump, B.C., Koch, E.W., 2008. Attached bacterial populations shared by four species
829 of aquatic angiosperms. *Appl. Environ. Microb.* 74, 5948-5957.

830 Czekalski, N., Berthold, T., Caucci, S., Egli, A., Buergermann, H., 2012. Increased
831 levels of multiresistant bacteria and resistance genes after wastewater
832 treatment and their dissemination into Lake Geneva, Switzerland. *Front.*
833 *Microbiol.* 3, e106.

834 Czekalski, N., Diez, E.G., Buergermann, H., 2014. Wastewater as a point source of
835 antibiotic-resistance genes in the sediment of a freshwater lake. *ISME J.* 8,
836 1381-1390.

837 Czekalski, N., Sigdel, R., Birtel, J., Matthews, B., Buergermann, H., 2015. Does human
838 activity impact the natural antibiotic resistance background? Abundance of
839 antibiotic resistance genes in 21 Swiss lakes. *Environ. Int.* 81, 45-55.

840 D'Costa, V.M., King, C.E., Kalan, L., Morar, M., Sung, W.W.L., Schwarz, C., Froese,
841 D., Zazula, G., Calmels, F., Debruyne, R., Golding, G.B., Poinar, H.N., Wright,
842 G.D., 2011. Antibiotic resistance is ancient. *Nature* 477, 457-461.

843 Dang, B.J., Mao, D.Q., Xu, Y., Luo, Y., 2017. Conjugative multi-resistant plasmids in
844 Haihe River and their impacts on the abundance and spatial distribution of
845 antibiotic resistance genes. *Water Res.* 111, 81-91.

846 de Faria, A.C.S., de Godoy, I., Sanches, A.A.A., Iglesias, G.A., Candido, S.L., da Paz,
847 R.C.R., Nakazato, L., Dutra, V., 2016. Detection of resistance genes and
848 evaluation of water quality at zoo lakes in Brazil. *Ciencia Rural* 46, 860-866.

849 Devarajan, N., Laffite, A., Graham, N.D., Meijer, M., Prabakar, K., Mubedi, J.I.,
850 Elongo, V., Mpiana, P.T., Ibelings, B.W., Wildi, W., Pote, J., 2015.
851 Accumulation of clinically relevant antibiotic-resistance genes, bacterial load,
852 and metals in freshwater lake sediments in Central Europe. *Environ. Sci.*
853 *Technol.* 49, 6528-6537.

854 Di Cesare, A., Eckert, E.M., Corno, G., 2016. Co-selection of antibiotic and heavy
855 metal resistance in freshwater bacteria. *J. Limnol.* 75, 59-66.

856 Di Cesare, A., Eckert, E.M., Teruggi, A., Fontaneto, D., Bertoni, R., Callieri, C.,
857 Corno, G., 2015. Constitutive presence of antibiotic resistance genes within the
858 bacterial community of a large subalpine lake. *Mol. Ecol.* 24, 3888-3900.

859 Ding, J., Lu, G., Liu, J., Yang, H., Li, Y., 2016. Uptake, depuration, and
860 bioconcentration of two pharmaceuticals, roxithromycin and propranolol, in
861 *Daphnia magna*. *Ecotox. Environ. Safe.* 126, 85-93.

862 Du, L.F., Liu, W.K., 2012. Occurrence, fate, and ecotoxicity of antibiotics in

863 agro-ecosystems. A review. *Agron. Sustain. Dev.* 32, 309-327.

864 Eckert, E.M., Di Cesare, A., Coci, M., Corno, G., 2018. Persistence of antibiotic
865 resistance genes in large subalpine lakes: the role of anthropogenic pollution
866 and ecological interactions. *Hydrobiologia*,
867 <https://doi.org/10.1007/s10750-017-3480-0>

868 Eckert, E.M., Di Cesare, A., Stenzel, B., Fontaneto, D., Corno, G., 2016. *Daphnia* as a
869 refuge for an antibiotic resistance gene in an experimental freshwater
870 community. *Sci. Total. Environ.* 571, 77-81.

871 Fick, J., Soderstrom, H., Lindberg, R.H., Phan, C., Tysklind, M., Larsson, D.G.J.,
872 2009. Contamination of surface, ground, and drinking water from
873 pharmaceutical production. *Environ. Toxicol. Chem* 28, 2522-2527.

874 Fleming, A., 1929. On the antibacterial action of cultures of a *Penicillium*, with
875 special reference to their use in the isolation of *B. influenzae*. *Brit. J. Exp.*
876 *Pathol.* 10, 226-236.

877 Fu, J., Yang, D., Jin, M., Liu, W., Zhao, X., Li, C., Zhao, T., Wang, J., Gao, Z., Shen,
878 Z., Qiu, Z., Li, J.W., 2017. Aquatic animals promote antibiotic resistance gene
879 dissemination in water via conjugation: Role of different regions within the
880 zebra fish intestinal tract, and impact on fish intestinal microbiota. *Mol. Ecol.*
881 26, 5318-5333.

882 Furukawa, T., Jikumaru, A., Ueno, T., Sei, K., 2017. Inactivation effect of
883 antibiotic-resistant gene using chlorine disinfection. *Water* 9, 547.
884 doi:10.3390/w9070547

885 Gao, L., Shi, Y., Li, W., Liu, J., Cai, Y., 2012. Occurrence, distribution and
886 bioaccumulation of antibiotics in the Haihe River in China. *J. Environ.*
887 *Monitor.* 14, 1247-1254.

888 Gillings, M.R., 2018. DNA as a Pollutant: the Clinical Class 1 Integron. *Curr. Pollut.*
889 *Rep.* <https://doi.org/10.1007/s40726-018-0076-x>

890 Grabow, W.O.K., Prozesky, O.W., Smith, L.S., 1974. Drug resistant coliforms call for
891 review of water quality standards. *Water Res.* 8, 1-9.

892 Grenni, P., Ancona, V., Barra Caracciolo, A., 2018. Ecological effects of antibiotics on
893 natural ecosystems: A review. *Microchem. J.* 136, 25-39.

894 Guo, J.H., Li, J., Chen, H., Bond, P.L., Yuan, Z.G., 2017. Metagenomic analysis
895 reveals wastewater treatment plants as hotspots of antibiotic resistance genes
896 and mobile genetic elements. *Water Res.* 123, 468-478.

897 Hamelin, K., Bruant, G., El-Shaarawi, A., Hill, S., Edge, T.A., Bekal, S., Fairbrother,
898 J.M., Harel, J., Maynard, C., Masson, L., Brousseau, R., 2006. A virulence and
899 antimicrobial resistance DNA microarray detects a high frequency of virulence
900 genes in *Escherichia coli* isolates from great lakes recreational waters. *Appl.*
901 *Environ. Microb.* 72, 4200-4206.

902 Han, N.N., Zhang, S.H., Wang, P.F., Wang, C., 2013. Characterization of antibiotic
903 resistance *E. coli* and antibiotic resistance genes in aquatic environment of
904 Taihu Lake, China. *Progress in Environmental Protection and Processing of*
905 *Resource*, Pts 1-4 295-298, 630-634.

906 Harnisz, M., Korzeniewska, E., Golas, I., 2015. The impact of a freshwater fish farm

907 on the community of tetracycline-resistant bacteria and the structure of
 908 tetracycline resistance genes in river water. *Chemosphere* 128, 134-141.
 909 Hernández, F., Ibáñez, M., Botero-Coy, A.-M., Bade, R., Bustos-López, M.C.,
 910 Rincón, J., Moncayo, A., Bijlsma, L., 2015. LC-QTOF MS screening of more
 911 than 1,000 licit and illicit drugs and their metabolites in wastewater and
 912 surface waters from the area of Bogotá, Colombia. *Anal. and Bioanal. Chem.*
 913 407, 6405-6416.
 914 Heuer, H., Schmitt, H., Smalla, K., 2011. Antibiotic resistance gene spread due to
 915 manure application on agricultural fields. *Curr. Opin. Microbiol.* 14, 236-243.
 916 Homem, V., Santos, L., 2011. Degradation and removal methods of antibiotics from
 917 aqueous matrices – A review. *J. Environ. Manage.* 92, 2304-2347.
 918 Hu, X.L., Bao, Y.F., Hu, J.J., Liu, Y.Y., Yin, D.Q., 2017a. Occurrence of 25
 919 pharmaceuticals in Taihu Lake and their removal from two urban drinking
 920 water treatment plants and a constructed wetland. *Environ. Sci. Pollut. Res.*
 921 24, 14889-14902.
 922 Hu, Y., Gao, G.F., Zhu, B., 2017b. The antibiotic resistome: gene flow in
 923 environments, animals and human beings. *Frontiers of Medicine* 11, 161-168.
 924 Hu, Y., Yang, X., Li, J., Lv, N., Liu, F., Wu, J., Lin, I.Y.C., Wu, N., Weimer, B.C.,
 925 Gao, G.F., Liu, Y., Zhu, B., 2016. The transfer network connecting the animal
 926 and human microbiomes. *Appl. Environ. Microbiol.* 82, 6672-6681.
 927 Huang, C.H., Renew, J.E., Smeby, K.L., Pinkston, K., Sedlak, D.L., 2011. Assessment
 928 of potential antibiotic contaminants in water and preliminary occurrence
 929 analysis. *J. Contemp. Water Res. Educ.* 120, 30-40.
 930 Ji, L.L., Bai, Z.T., Deng, L.P., Ashraf, M.A., 2016. Sorption of tetracycline,
 931 oxytetracycline and tylosin to eight surface sediments of Taihu Lake. *J.*
 932 *Environ. Biol.* 37, 1087-1095.
 933 Jia, A., Hu, J., Wu, X., Peng, H., Wu, S., Dong, Z., 2011. Occurrence and source
 934 apportionment of sulfonamides and their metabolites in Liaodong Bay and the
 935 adjacent Liao River basin, North China. *Environ. Toxicol. Chem* 30,
 936 1252-1260.
 937 Jia, S., Shi, P., Hu, Q., Li, B., Zhang, T., Zhang, X.X., 2015. Bacterial community
 938 shift drives antibiotic resistance promotion during drinking water chlorination.
 939 *Environ. Sci. Technol.* 49, 2271-2279.
 940 Jia, S.Y., He, X.W., Bu, Y.Q., Shi, P., Miao, Y., Zhou, H.P., Shan, Z.J., Zhang, X.X.,
 941 2014. Environmental fate of tetracycline resistance genes originating from
 942 swine feedlots in river water. *J. Environ. Sci. Health Part. B* 49, 624-631.
 943 Jiang, L., Hu, X., Yin, D., Zhang, H., Yu, Z., 2011. Occurrence, distribution and
 944 seasonal variation of antibiotics in the Huangpu River, Shanghai, China.
 945 *Chemosphere* 82, 822-828.
 946 Jiang, L., Hu, X.L., Xu, T., Zhang, H.C., Sheng, D., Yin, D.Q., 2013. Prevalence of
 947 antibiotic resistance genes and their relationship with antibiotics in the
 948 Huangpu River and the drinking water sources, Shanghai, China. *Sci. Total.*
 949 *Environ.* 458, 267-272.
 950 Jones, J.G., Gardener, S., Simon, B.M., Pickup, R.W., 1986. Factors affecting the

951 measurement of antibiotic resistance in bacteria isolated from lake water. J.
 952 Appl. Bacteriol. 60, 455-462.

953 Khan, G.A., Berglund, B., Khan, K.M., Lindgren, P.E., Fick, J., 2013. Occurrence and
 954 abundance of antibiotics and resistance genes in rivers, canal and near drug
 955 formulation facilities - a study in Pakistan. Plos One 8, e62712.

956 Knapp, C.W., Engemann, C.A., Hanson, M.L., Keen, P.L., Hall, K.J., Graham, D.W.,
 957 2008. Indirect evidence of transposon-mediated selection of antibiotic
 958 resistance genes in aquatic systems at low-level oxytetracycline exposures.
 959 Environ. Sci. Technol. 42, 5348-5353.

960 Knapp, C.W., Lima, L., Olivares-Rieumont, S., Bowen, E., Werner, D., Graham,
 961 D.W., 2012. Seasonal variations in antibiotic resistance gene transport in the
 962 Almendares River, Havana, Cuba. Front. Microbiol. 3, e396.

963 Koczura, R., Mokracka, J., Taraszewska, A., Lopacinska, N., 2016. Abundance of
 964 class 1 integron-integrase and sulfonamide resistance genes in river water and
 965 sediment is affected by anthropogenic pressure and environmental factors.
 966 Microb. Ecol. 72, 909-916.

967 Kondrashov, F.A., 2012. Gene duplication as a mechanism of genomic adaptation to a
 968 changing environment. Proc. Biol. Sci. 279, 5048-5057.

969 Kümmerer, K., 2009a. Antibiotics in the aquatic environment—A review—part I.
 970 Chemosphere 75, 417-434.

971 Kümmerer, K., 2009b. Antibiotics in the aquatic environment – A review – Part II.
 972 Chemosphere 75, 435-441.

973 Lamas, A., Franco, C.M., Regal, P., Miranda, J.M., Vázquez, B., Cepeda, A.
 974 High-Throughput platforms in real-time PCR and applications. in:
 975 Samadikuchaksaraei A., ed. Polymerase Chain Reaction for Biomedical
 976 Applications. Rijeka: InTech; 2016

977 LaPara, T.M., Burch, T.R., McNamara, P.J., Tan, D.T., Yan, M., Eichmiller, J.J., 2011.
 978 Tertiary-treated municipal wastewater is a significant point source of antibiotic
 979 resistance genes into Duluth-Superior Harbor. Environ. Sci. Technol. 45,
 980 9543-9549.

981 Lavilla Lerma, L., Benomar, N., Casado Muñoz, M.d.C., Gálvez, A., Abriouel, H.,
 982 2014. Antibiotic multiresistance analysis of mesophilic and psychrotrophic
 983 *Pseudomonas* spp. isolated from goat and lamb slaughterhouse surfaces
 984 throughout the meat production process. Appl. Environ. Microb. 80,
 985 6792-6806.

986 Lei, X.N., Lu, J.J., Liu, Z.L., Tong, Y.B., Li, S.M., 2015. Concentration and
 987 distribution of antibiotics in water-sediment system of Bosten Lake, Xinjiang.
 988 Environ. Sci. Pollut. Res. 22, 1670-1678.

989 Lekunberri, I., Balcázar, J.L., Borrego, C.M., 2018. Metagenomic exploration reveals
 990 a marked change in the river resistome and mobilome after treated wastewater
 991 discharges. Environ. Pollut. 234, 538-542.

992 Li, B., Yang, Y., Ma, L., Ju, F., Guo, F., Tiedje, J.M., Zhang, T., 2015. Metagenomic
 993 and network analysis reveal wide distribution and co-occurrence of
 994 environmental antibiotic resistance genes. ISME J. 9, 2490-2502.

- 995 Li, H., Helm, P.A., Metcalfe, C.D., 2010. Sampling in the Great Lakes for
996 pharmaceuticals, personal care products, and endocrine- disrupting substances
997 using the passive polar organic chemical integrative sampler. *Environ. Toxicol.*
998 *Chem.* 29, 751-762.
- 999 Li, S., Shi, W., Liu, W., Li, H., Zhang, W., Hu, J., Ke, Y., Sun, W., Ni, J., 2018. A
1000 duodecennial national synthesis of antibiotics in China's major rivers and seas
1001 (2005–2016). *Sci. Total. Environ.* 615, 906-917.
- 1002 Li, W.H., Shi, Y.L., Gao, L.H., Liu, J.M., Cai, Y.Q., 2012. Occurrence of antibiotics in
1003 water, sediments, aquatic plants, and animals from Baiyangdian Lake in North
1004 China. *Chemosphere* 89, 1307-1315.
- 1005 Lindim, C., van Gils, J., Georgieva, D., Mekenyan, O., Cousins, I.T., 2016. Evaluation
1006 of human pharmaceutical emissions and concentrations in Swedish river
1007 basins. *Sci. Total. Environ.* 572, 508-519.
- 1008 Liu, J.L., Wong, M.H., 2013. Pharmaceuticals and personal care products (PPCPs): A
1009 review on environmental contamination in China. *Environ. Int.* 59, 208-224.
- 1010 Liu, M., Zhang, Y., Yang, M., Tian, Z., Ren, L., Zhang, S., 2012. Abundance and
1011 distribution of tetracycline resistance genes and mobile elements in an
1012 oxytetracycline production wastewater treatment system. *Environ. Sci.*
1013 *Technol.* 46, 7551-7557.
- 1014 Liu, Y., Zhang, J., Gao, B., Feng, S., 2014. Combined effects of two antibiotic
1015 contaminants on *Microcystis aeruginosa*. *J. Hazard. Mater.* 279, 148-155.
- 1016 Luo, Y., Mao, D.Q., Rysz, M., Zhou, D.X., Zhang, H.J., Xu, L., Alvarez, P.J.J., 2010.
1017 Trends in Antibiotic Resistance Genes Occurrence in the Haihe River, China.
1018 *Environ. Sci. Technol.* 44, 7220-7225.
- 1019 Luo, Y., Xu, L., Rysz, M., Wang, Y., Zhang, H., Alvarez, P.J.J., 2011. Occurrence and
1020 transport of tetracycline, sulfonamide, quinolone, and macrolide antibiotics in
1021 the Haihe River Basin, China. *Environ. Sci. Technol.* 45, 1827-1833.
- 1022 Lyandres, O., 2012. Keeping great lakes water safe: Priorities for protecting against
1023 emerging chemical pollutants. Alliance for the Great Lakes,
1024 [https://www.csuohio.edu/urban/sites/csuohio.edu/urban/files/Drinking_Water_](https://www.csuohio.edu/urban/sites/csuohio.edu/urban/files/Drinking_Water_Great_Lakes.pdf)
1025 [Great_Lakes.pdf](https://www.csuohio.edu/urban/sites/csuohio.edu/urban/files/Drinking_Water_Great_Lakes.pdf).
- 1026 Magdaleno, A., Saenz, M.E., Juárez, A.B., Moretton, J., 2015. Effects of six
1027 antibiotics and their binary mixtures on growth of *Pseudokirchneriella*
1028 *subcapitata*. *Ecotox. Environ. Safe.* 113, 72-78.
- 1029 Manaia, C.M., 2017. Assessing the risk of antibiotic resistance transmission from the
1030 environment to humans: Non-direct proportionality between abundance and
1031 risk. *Trends Microbiol.* 25, 173-181.
- 1032 Martinez, J.L., Coque, T.M., Baquero, F., 2015. What is a resistance gene? Ranking
1033 risk in resistomes. *Nat. Rev. Microbiol.* 13, 116-123.
- 1034 Matyar, F., Kaya, A., Dinçer, S., 2008. Antibacterial agents and heavy metal resistance
1035 in Gram-negative bacteria isolated from seawater, shrimp and sediment in
1036 Iskenderun Bay, Turkey. *Sci. Total Environ.* 407, 279-285.
- 1037 McConnell, R.L., Abel, D.C. *Environmental Geology Today* eds. Sudbury, United
1038 States: Jones and Bartlett Publishers; 2013

- 1039 Meng, X.Z., Venkatesan, A.K., Ni, Y.L., Steele, J.C., Wu, L.L., Bignert, A., Bergman,
1040 Å., Halden, R.U., 2016. Organic contaminants in Chinese sewage sludge: A
1041 meta-analysis of the literature of the past 30 years. *Environ. Sci. Technol.* 50,
1042 5454-5466.
- 1043 Metch, J.W., Burrows, N.D., Murphy, C.J., Pruden, A., Vikesland, P.J., 2018.
1044 Metagenomic analysis of microbial communities yields insight into impacts of
1045 nanoparticle design. *Nat. Nanotechnol.* doi: 10.1038/s41565-017-0029-3.
- 1046 Muziasari, W.I., Pitkänen, L.K., Sørum, H., Stedtfeld, R.D., Tiedje, J.M., Virta, M.,
1047 2017. The resistome of farmed fish feces contributes to the enrichment of
1048 antibiotic resistance genes in sediments below Baltic Sea fish farms. *Front.*
1049 *Microbiol.*, 1491.doi: 1410.3389/fmicb.2017.01491. eCollection 02017.
- 1050 Na, G., Gu, J., Ge, L., Zhang, P., Wang, Z., Liu, C., Zhang, L., 2011. Detection of 36
1051 antibiotics in coastal waters using high performance liquid
1052 chromatography-tandem mass spectrometry. *Chinese J. Oceanol. Limnol.* 29,
1053 1093-1102.
- 1054 Nakata, H., Kannan, K., Jones, P.D., Giesy, J.P., 2005. Determination of
1055 fluoroquinolone antibiotics in wastewater effluents by liquid
1056 chromatography-mass spectrometry and fluorescence detection. *Chemosphere*
1057 58, 759-766.
- 1058 Nakayama, T., Hoa, T.T.T., Harada, K., Warisaya, M., Asayama, M., Hinenoya, A.,
1059 Lee, J.W., Phu, T.M., Ueda, S., Sumimura, Y., Hirata, K., Phuong, V.T.,
1060 Yamamoto, Y., 2017. Water metagenomic analysis reveals low bacterial
1061 diversity and the presence of antimicrobial residues and resistance genes in a
1062 river containing wastewater from backyard aquacultures in the Mekong Delta,
1063 Vietnam. *Environ. Pollut.* 222, 294-306.
- 1064 Narciso-da-Rocha, C., Manaia, C.M., 2016. Multidrug resistance phenotypes are
1065 widespread over different bacterial taxonomic groups thriving in surface
1066 water. *Sci. Total. Environ.* 563, 1-9.
- 1067 Ngumba, E., Gachanja, A., Tuhkanen, T., 2016. Occurrence of selected antibiotics and
1068 antiretroviral drugs in Nairobi River Basin, Kenya. *Sci. Total. Environ.* 539,
1069 206-213.
- 1070 Oh, S., Tandukar, M., Pavlostathis, S.G., Chain, P.S.G., Konstantinidis, K.T., 2013.
1071 Microbial community adaptation to quaternary ammonium biocides as
1072 revealed by metagenomics. *Environ. Microbiol.* 15, 2850-2864.
- 1073 Ouyang, W.Y., Huang, F.Y., Zhao, Y., Li, H., Su, J.Q., 2015. Increased levels of
1074 antibiotic resistance in urban stream of Jiulongjiang River, China. *Appl.*
1075 *Microbiol. Biotechnol.* 99, 5697-5707.
- 1076 Ozaktas, T., Taskin, B., Gozen, A.G., 2012. High level multiple antibiotic resistance
1077 among fish surface associated bacterial populations in non-aquaculture
1078 freshwater environment. *Water Res.* 46, 6382-6390.
- 1079 Park, M., Reckhow, D., Lavine, M., Rosenfeldt, E., Stanford, B., Park, M.H., 2014.
1080 Multivariate analyses for monitoring EDCs and PPCPs in a lake water. *Water*
1081 *Environ. Res.* 86, 2233-2241.
- 1082 Peng, X., Ou, W., Wang, C., Wang, Z., Huang, Q., Jin, J., Tan, J., 2014. Occurrence

1083 and ecological potential of pharmaceuticals and personal care products in
 1084 groundwater and reservoirs in the vicinity of municipal landfills in China. *Sci.*
 1085 *Total. Environ.* 490, 889-898.

1086 Peng, X., Zhang, K., Tang, C., Huang, Q., Yu, Y., Cui, J., 2011. Distribution pattern,
 1087 behavior, and fate of antibacterials in urban aquatic environments in South
 1088 China. *J. Environ. Monitor.* 13, 446-454.

1089 Pruden, A., Arabi, M., Storteboom, H.N., 2012. Correlation between upstream human
 1090 activities and riverine antibiotic resistance genes. *Environ. Sci. Technol.* 46,
 1091 11541-11549.

1092 Pruden, A., Pei, R., Storteboom, H., Carlson, K.H., 2006. Antibiotic resistance genes
 1093 as emerging contaminants: studies in northern Colorado. *Environ. Sci.*
 1094 *Technol.* 40, 7445-7450.

1095 Reuther, R., 2009. Lake and river sediment monitoring. *Environmental monitoring*
 1096 *Encyclopedia of Life Support Systems (EOLSS)* vol 2, 120-147.

1097 Rizzo, L., Manaia, C., Merlin, C., Schwartz, T., Dagot, C., Ploy, M.C., Michael, I.,
 1098 Fatta-Kassinos, D., 2013. Urban wastewater treatment plants as hotspots for
 1099 antibiotic resistant bacteria and genes spread into the environment: A review.
 1100 *Sci. Total. Environ.* 447, 345-360.

1101 Rodriguez-Mozaz, S., Chamorro, S., Marti, E., Huerta, B., Gros, M., Sanchez-Melsio,
 1102 A., Borrego, C.M., Barcelo, D., Balcazar, J.L., 2015. Occurrence of antibiotics
 1103 and antibiotic resistance genes in hospital and urban wastewaters and their
 1104 impact on the receiving river. *Water Res.* 69, 234-242.

1105 Romero, J.L., Grande Burgos, M.J., Pérez-Pulido, R., Gálvez, A., Lucas, R., 2017.
 1106 Resistance to antibiotics, biocides, preservatives and metals in bacteria
 1107 isolated from seafoods: Co-selection of strains resistant or tolerant to different
 1108 classes of compounds. *Front. Microbiol.* 8, e1650.

1109 Roose-Amsaleg, C., Laverman, A.M., 2016. Do antibiotics have environmental
 1110 side-effects? Impact of synthetic antibiotics on biogeochemical processes.
 1111 *Environ. Sci. Pollut. Res.* 23, 4000-4012.

1112 Seiler, C., Berendonk, T., 2012. Heavy metal driven co-selection of antibiotic
 1113 resistance in soil and water bodies impacted by agriculture and aquaculture.
 1114 *Front. Microbiol.* 3, e399.

1115 Selvam, A., Kwok, K., Chen, Y., Cheung, A., Leung, K.S.Y., Wong, J.W.C., 2017.
 1116 Influence of livestock activities on residue antibiotic levels of rivers in Hong
 1117 Kong. *Environ. Sci. Pollut. Res.* 24, 9058-9066.

1118 Staples, C.A., Woodburn, K.B., Klecka, G.M., Mihaich, E.M., Hall, A.T., Ortego, L.,
 1119 Caspers, N., Hentges, S.G., 2008. Comparison of four species sensitivity
 1120 distribution methods to calculate predicted no effect concentrations for
 1121 bisphenol A. *Hum. Ecol. Risk Assess.* 14, 455-478.

1122 Stepanauskas, R., Glenn, T.C., Jagoe, C.H., Tuckfield, R.C., Lindell, A.H., King, C.J.,
 1123 McArthur, J.V., 2006. Coselection for microbial resistance to metals and
 1124 antibiotics in freshwater microcosms. *Environ. Microbiol.* 8, 1510-1514.

1125 Stokes, H.W., Gillings, M.R., 2011. Gene flow, mobile genetic elements and the
 1126 recruitment of antibiotic resistance genes into Gram-negative pathogens.

1127 FEMs Microbiol. Rev. 35, 790-819.

1128 Storteboom, H., Arabi, M., Davis, J.G., Crimi, B., Pruden, A., 2010. Tracking
 1129 antibiotic resistance genes in the south Platte River Basin using molecular
 1130 signatures of urban, agricultural, and pristine sources. Environ. Sci. Technol.
 1131 44, 7397-7404.

1132 Su, H.C., Liu, Y.S., Pan, C.G., Chen, J., He, L.Y., Ying, G.G., 2018. Persistence of
 1133 antibiotic resistance genes and bacterial community changes in drinking water
 1134 treatment system: From drinking water source to tap water. Sci. Total.
 1135 Environ. 616-617, 453-461.

1136 Suzuki, S., Ogo, M., Koike, T., Takada, H., Newman, B., 2015. Sulfonamide and
 1137 tetracycline resistance genes in total- and culturable-bacterial assemblages in
 1138 South African aquatic environments. Front. Microbiol. 6, e796.

1139 Tamtam, F., Mercier, F., Le Bot, B., Eurin, J., Tuc Dinh, Q., Clément, M., Chevreuil,
 1140 M., Occurrence and fate of antibiotics in the Seine River in various
 1141 hydrological conditions. Sci. Total Environ. 393, 84-95.

1142 Thevenon, F., Adatte, T., Wildi, W., Pote, J., 2012. Antibiotic resistant bacteria/genes
 1143 dissemination in lacustrine sediments highly increased following cultural
 1144 eutrophication of Lake Geneva (Switzerland). Chemosphere 86, 468-476.

1145 Thomas, C.M., Nielsen, K.M., 2005. Mechanisms of, and barriers to, horizontal gene
 1146 transfer between bacteria. Nat. Rev. Microbiol. 3, 711.

1147 Vålitalo, P., Kruglova, A., Mikola, A., Vahala, R., 2017. Toxicological impacts of
 1148 antibiotics on aquatic micro-organisms: A mini-review. Int. J. Hyg. Envir. Heal
 1149 220, 558-569.

1150 Van Boeckel, T.P., Brower, C., Gilbert, M., Grenfell, B.T., Levin, S.A., Robinson, T.P.,
 1151 Teillant, A., Laxminarayan, R., 2015. Global trends in antimicrobial use in
 1152 food animals. Proc. Natl. Acad. Sci. USA 112, 5649-5654.

1153 Vila-Costa, M., Gioia, R., Acena, J., Perez, S., Casamayor, E.O., Dachs, J., 2017.
 1154 Degradation of sulfonamides as a microbial resistance mechanism. Water Res.
 1155 115, 309-317.

1156 Wang, C., Gu, X.C., Zhang, S.H., Wang, P.F., Guo, C., Gu, J., Hou, J., 2013.
 1157 Characterization of antibiotic-resistance genes in antibiotic resistance
 1158 *Escherichia coli* isolates from a lake. Arch. Environ. Con. Tox 65, 635-641.

1159 Wang, J., Wang, J., Zhao, Z., Chen, J., Lu, H., Liu, G., Zhou, J., Guan, X., 2017a.
 1160 PAHs accelerate the propagation of antibiotic resistance genes in coastal water
 1161 microbial community. Environ. Pollut. 231, 1145-1152.

1162 Wang, W., Wang, H., Zhang, W., Liang, H., Gao, D., 2017b. Occurrence, distribution,
 1163 and risk assessment of antibiotics in the Songhua River in China. Environ. Sci.
 1164 Pollut. Res. 24, 19282-19292.

1165 Wang, Z., Du, Y., Yang, C., Liu, X., Zhang, J., Li, E., Zhang, Q., Wang, X., 2017c.
 1166 Occurrence and ecological hazard assessment of selected antibiotics in the
 1167 surface waters in and around Lake Honghu, China. Sci. Total. Environ. 609,
 1168 1423-1432.

1169 Wright, G.D., Poinar, H., 2012. Antibiotic resistance is ancient: implications for drug
 1170 discovery. Trends Microbiol. 20, 157-159.

- 1171 Wu, C., Huang, X., Witter, J.D., Spongberg, A.L., Wang, K., Wang, D., Liu, J., 2014.
1172 Occurrence of pharmaceuticals and personal care products and associated
1173 environmental risks in the central and lower Yangtze river, China. *Ecotox.*
1174 *Environ. Safe.* 106, 19-26.
- 1175 Wu, Y., Yu, C.P., Yue, M., Liu, S.P., Yang, X.Y., 2016. Occurrence of selected PPCPs
1176 and sulfonamide resistance genes associated with heavy metals pollution in
1177 surface sediments from Chao Lake, China. *Environ. Earth Sci.* 75, 43.
1178 <https://doi.org/10.1007/s12665-015-4838-0> 75,
- 1179 Xie, Z.X., Lu, G.H., Liu, J.C., Yan, Z.H., Ma, B.N., Zhang, Z.H., Chen, W., 2015.
1180 Occurrence, bioaccumulation, and trophic magnification of pharmaceutically
1181 active compounds in Taihu Lake, China. *Chemosphere* 138, 140-147.
- 1182 Xie, Z.X., Lu, G.H., Yan, Z.H., Liu, J.C., Wang, P.F., Wang, Y.H., 2017.
1183 Bioaccumulation and trophic transfer of pharmaceuticals in food webs from a
1184 large freshwater lake. *Environ. Pollut.* 222, 356-366.
- 1185 Xiong, W., Sun, Y., Ding, X., Zhang, Y., Zeng, Z., 2014. Antibiotic resistance genes
1186 occurrence and bacterial community composition in the Liuxi River. *Front.*
1187 *Environ. Sci.* 2, 61. doi: 10.3389/fenvs.2014.00061.
- 1188 Xu, J., Zhang, Y., Zhou, C., Guo, C., Wang, D., Du, P., Luo, Y., Wan, J., Meng, W.,
1189 2014. Distribution, sources and composition of antibiotics in sediment,
1190 overlying water and pore water from Taihu Lake, China. *Sci. Total. Environ.*
1191 497, 267-273.
- 1192 Xu, L.K., Ouyang, W.Y., Qian, Y.Y., Su, C., Su, J.Q., Chen, H., 2016.
1193 High-throughput profiling of antibiotic resistance genes in drinking water
1194 treatment plants and distribution systems. *Environ. Pollut.* 213, 119-126.
- 1195 Xu, Y., Xu, J., Mao, D.Q., Luo, Y., 2017. Effect of the selective pressure of sub-lethal
1196 level of heavy metals on the fate and distribution of ARGs in the catchment
1197 scale. *Environ. Pollut.* 220, 900-908.
- 1198 Yan, C., Yang, Y., Zhou, J., Liu, M., Nie, M., Shi, H., Gu, L., 2013. Antibiotics in the
1199 surface water of the Yangtze Estuary: Occurrence, distribution and risk
1200 assessment. *Environ. Pollut.* 175, 22-29.
- 1201 Yan, Z.H., Yang, X.F., Lu, G.H., Liu, J.C., Xie, Z.X., Wu, D.H., 2014. Potential
1202 environmental implications of emerging organic contaminants in Taihu Lake,
1203 China: Comparison of two ecotoxicological assessment approaches. *Sci. Total.*
1204 *Environ.* 470, 171-179.
- 1205 Yang, J.F., Ying, G.G., Zhao, J.L., Tao, R., Su, H.C., Liu, Y-S., 2011. Spatial and
1206 seasonal distribution of selected antibiotics in surface waters of the Pearl
1207 Rivers, China. *J. Environ. Sci. Health Part B* 46, 272-280.
- 1208 Yang, Y., Cao, X., Lin, H., Wang, J., 2016. Antibiotics and antibiotic resistance genes
1209 in sediment of Honghu Lake and East Dongting Lake, China. *Microb. Ecol.*
1210 72, 791-801.
- 1211 Yang, Y., Liu, W., Xu, C., Wei, B., Wang, J., 2017a. Antibiotic resistance genes in
1212 lakes from middle and lower reaches of the Yangtze River, China: Effect of
1213 land use and sediment characteristics. *Chemosphere* 178, 19-25.
- 1214 Yang, Y., Wang, W., 2018. Benzyltrimethylammonium chloride shifts the

proliferation of functional genes and microbial community in natural water from eutrophic lake. *Environ. Pollut.* 236, 355-365.

Yang, Y., Xu, C., Cao, X., Lin, H., Wang, J., 2017b. Antibiotic resistance genes in surface water of eutrophic urban lakes are related to heavy metals, antibiotics, lake morphology and anthropic impact. *Ecotoxicology* 26, 831-840.

Yin, Q., Yue, D.M., Peng, Y.K., Liu, Y., Xiao, L., 2013. Occurrence and distribution of antibiotic-resistant bacteria and transfer of resistance genes in Lake Taihu. *Microbes Environ.* 28, 479-486.

Zhang, C.H., Tang, J.W., Wang, L.L., Gao, X.Y., He, X.D., 2015a. Occurrence of antibiotics in water and sediment from Zizhuyuan Lake. *Pol. J. Environ. Stud.* 24, 1831-1836.

Zhang, P., Zhou, H., Li, K., Zhao, X., Liu, Q., Li, D., Zhao, G., 2018. Occurrence of pharmaceuticals and personal care products, and their associated environmental risks in a large shallow lake in north China. *Environ. Geochem. Health.* <https://doi.org/10.1007/s10653-018-0069-0>

Zhang, Q.Q., Ying, G.G., Pan, C.G., Liu, Y.S., Zhao, J.L., 2015b. Comprehensive evaluation of antibiotics emission and fate in the river basins of China: Source analysis, multimedia modeling, and linkage to bacterial resistance. *Environ. Sci. Technol.* 49, 6772-6782.

Zhang, S.H., Pang, S., Wang, P.F., Wang, C., Han, N.N., Liu, B., Han, B., Li, Y., Anim-Larbi, K., 2016. Antibiotic concentration and antibiotic-resistant bacteria in two shallow urban lakes after stormwater event. *Environ. Sci. Pollut. Res.* 23, 9984-9992.

Zhang, W., Sturm, B.S.M., Knapp, C.W., Graham, D.W., 2009a. Accumulation of Tetracycline Resistance Genes in Aquatic Biofilms Due to Periodic Waste Loadings from Swine Lagoons. *Environ. Sci. Technol.* 43, 7643-7650.

Zhang, X.X., Zhang, T., Fang, H.P., 2009b. Antibiotic resistance genes in water environment. *Appl. Microbiol. Biotechnol.* 82, 397-414.

Zhang, X.X., Wu, B., Zhang, Y., Zhang, T., Yang, L.Y., Fang, H.H.P., Ford, T., Cheng, S.P., 2009c. Class 1 integronase gene and tetracycline resistance genes *tetA* and *tetC* in different water environments of Jiangsu Province, China. *Ecotoxicology* 18, 652-660.

Zhang, Y., Xu, J., Zhong, Z.X., Guo, C.S., Li, L., He, Y., Fan, W.H., Chen, Y.C., 2013. Degradation of sulfonamides antibiotics in lake water and sediment. *Environ. Sci. Pollut. Res.* 20, 2372-2380.

Zheng, J., Gao, R., Wei, Y., Chen, T., Fan, J., Zhou, Z., Makimilua, T.B., Jiao, Y., Chen, H., 2017. High-throughput profiling and analysis of antibiotic resistance genes in East Tiaoxi River, China. *Environ. Pollut.* 230, 648-654.

Zheng, S., Qiu, X., Chen, B., Yu, X., Liu, Z., Zhong, G., Li, H., Chen, M., Sun, G., Huang, H., Yu, W., Freestone, D., 2011. Antibiotics pollution in Jiulong River estuary: Source, distribution and bacterial resistance. *Chemosphere* 84, 1677-1685.

Zhong, Z.X., Xu, J., Zhang, Y., Li, L., Guo, C.S., He, Y., Fan, W.H., Zhang, B.P., 2013. Adsorption of sulfonamides on lake sediments. *Front. Environ. Sci. Eng.*

1259 7, 518-525.

1260 Zhou, L.J., Ying, G.G., Zhao, J.L., Yang, J.F., Wang, L., Yang, B., Liu, S., 2011.

1261 Trends in the occurrence of human and veterinary antibiotics in the sediments

1262 of the Yellow River, Hai River and Liao River in northern China. *Environ.*

1263 *Pollut.* 159, 1877-1885.

1264 Zhou, L.J., Wu, Q.L.L., Zhang, B.B., Zhao, Y.G., Zhao, B.Y., 2016. Occurrence,

1265 spatiotemporal distribution, mass balance and ecological risks of antibiotics in

1266 subtropical shallow Lake Taihu, China. *Environ. Sci.-Process Impacts* 18,

1267 500-513.

1268 Zhou, T., Lu, J.J., Tong, Y.B., Li, S.M., Wang, X.L., 2014. Distribution of antibiotic

1269 resistance genes in Bosten Lake, Xinjiang, China. *Water Sci. Technol.* 70,

1270 925-931.

1271 Zhou, Z.C., Zheng, J., Wei, Y.Y., Chen, T., Dahlgren, R.A., Shang, X., Chen, H., 2017.

1272 Antibiotic resistance genes in an urban river as impacted by bacterial

1273 community and physicochemical parameters. *Environ. Sci. Pollut. Res.* 24,

1274 23753-23762.

1275 Zhu, F., Jiao, S.J., Shan, Z.J., Kong, D.Y., Ge, F., Wang, N., 2015. Sorption and

1276 desorption behavior of doxycycline on a Wushantu soil in Taihu Lake region,

1277 China. *Fresen. Environ. Bull.* 24, 3295-3299.

1278 Zhu, Y.G., Johnson, T.A., Su, J.Q., Qiao, M., Guo, G.X., Stedtfeld, R.D., Hashsham,

1279 S.A., Tiedje, J.M., 2013. Diverse and abundant antibiotic resistance genes in

1280 Chinese swine farms. *Proc. Nat. Acad. Sci.* 110, 3435-3440.

1281 Zhu, Y.G., Zhao, Y., Li, B., Huang, C.L., Zhang, S.Y., Yu, S., Chen, Y.S., Zhang, T.,

1282 Gillings, M.R., Su, J.Q., 2017. Continental-scale pollution of estuaries with

1283 antibiotic resistance genes. *Nat. Microbiol.* 2, e16270.

1284 Zou, S.C., Zhu, C.J., He, Z.M., Luan, T.G., Xu, W.H., Zhang, G., 2009. Preliminary

1285 studies on the pollution levels of antibiotic resistance genes in the water of

1286 Beijiang River, South China. *Asian J. Ecotoxicol.* 4, 655-660.

1287 Zou, S., Xu, W., Zhang, R., Tang, J., Chen, Y., Zhang, G., 2011. Occurrence and

1288 distribution of antibiotics in coastal water of the Bohai Bay, China: Impacts of

1289 river discharge and aquaculture activities. *Environ. Pollut.* 159, 2913-2920.

1290

1291 Table 1 The number of studies and detected number of each class of antibiotics in
 1292 global lakes

Class	China		Other Countries	
Water	N ^a	n ^b	N	n
Sulphonamides and Trimethoprim	22	16	20	14
Tetracyclines	14	4	1	3
Quinolones	18	12	10	10
Macrolides and Lincosamides	9	10	8	7
Others	3	4	3	3
Sediment	N	n	N	n
Sulphonamides and Trimethoprim	10	15	0	0
Tetracyclines	7	4	0	0
Quinolones	9	11	1	4
Macrolides and Lincosamides	7	9	1	5
Others	3	5	1	1

1293 ^a: number of studies, ^b: number of studied compounds

1294

1295 Table 2 Comparison of bioaccumulation factors of antibiotics in freshwater system (Unit: L kg⁻¹).

Aquatic biota	SMZ	SMX	SDM	SCM	TC	OTC	ROX	ETM	OFL	NOR	ENR	FLE	CIP
Aquatic plants and algae													
Aquatic plants										554-4400 ^a	653-2590 ^a		10200-170000 ^a
Algae		25-183 ^b			30-977 ^b		28-2352 ^b ; 100-254 ^c	NA ^b ; 8.7-162 ^c	60-375 ^b	27-639 ^b			38-1202 ^b
Invertebrates													
Daphnia magna													
Snail													
Mussel, snail and bivalve		39-94 ^b			42-577 ^b		811-6352 ^b	447-3511 ^b	11-327 ^b	42-2780 ^b			99-4067 ^b
Mussel							406 ^c						
Crab		27-65 ^e				157 ^e	195 ^e			9 ^e	11050-34508 ^e		
Bivalve							510 ^c	32 ^c					
Shrimp		7.7-131 ^b			4.6-159 ^b		43-1704 ^b ; 294 ^c	NA ^b		84-3125 ^b	16600 ^a	4050 ^a	45-933 ^b

Fish										
Carp	1672 ^f				7410 ^a ; 200-376 ^c	32 ^c ; 4492 ^f		284 ^f	4490 ^a	3262 ^f
Blunt snout bream										4130 ^a
Topmouth gudgeon	3870 ^a									
Fish	NA ^b ; 13 ^e	339 ^e		3.1-362 ^b	17-3668 ^b ; 139 ^c	35-2029 ^b ; 44 ^c ; 52 ^e	12-970 ^b	35-2038 ^b	12059 ^e	23-13552 ^b
Lake anchovy					665 ^c	3.8 ^c				
Xingkai bitterling		527 ^e	232 ^e		5290 ^e	164 ^e			45407 ^e	
Crucian	13 ^e	355 ^e							31241 ^e	

1296 ^a(Li et al. 2012); ^b(Xie et al. 2017) ;^c(Xie et al. 2015) ; ^d(Ding et al. 2016); ^e(Bai et al. 2014); ^f(Gao et al. 2012)

1297 Sulfamethazine: SMZ; Sulfamethoxazole: SMX; Sulfadimethoxine: SDM; Sulfacetamide: SCM; Tetracycline: TC; Oxytetracycline: OTC;

1298 Roxithromycin: ROX; Erythromycin: ETM; Ofloxacin: OFL; Norfloxacin: NOR; Ciprofloxacin: CIP; Enrofloxacin: ENR; Fleroxacin: FLE.

Figure Captions

Fig. 1 The median values and 95% confidence intervals of average concentrations of antibiotics in bulk water of global lakes (ng L⁻¹, the number in the parentheses indicate the counts of data; invalid confidence intervals were not shown in the figure).

Fig. 2 The median values of average concentrations of antibiotics in sediment of global lakes (ng g⁻¹ dry weight, the number in the parentheses indicate the counts of data; invalid confidence intervals were not shown in the figure).

Fig. 3 The comparison of antibiotics in bulk water in lakes between China and other countries worldwide (n, the data number). Different letters over the bars indicate statistically significant differences at $p < 0.05$ level.

Fig. 4 The temporal trend of antibiotics in bulk water from Taihu Lake (a: (Xu et al. 2014); b: (Yan et al. 2014); c: (Wu et al. 2014) ; d: (Zhou et al. 2016); e: (Hu et al. 2017a), The number was the counts of sampling sites; sulfamethoxazole concentrations in 2013 was the mean values of the two references).

Fig. 5 The hazardous concentration for five percent of the species (HC₅) and their 2.5th and 97.5th percentile values for antibiotics on aquatic system (a) (Unit: µg L⁻¹; the number in the parentheses indicate the counts of species) and hazard quotient of antibiotics in freshwater lakes (b) (Sulfamethoxazole: SMX; Sulfadiazine: SDZ; Sulfamethazine: SMZ; Trimethoprim: TMP; Tetracycline: TC; Oxytetracycline: OTC; Chlortetracycline: CTC; Ofloxacin: OFL; Norfloxacin: NOR; Ciprofloxacin: CIP; Lomefloxacin: LOM; Enrofloxacin: ENR; Erythromycin: ERY; Clarithromycin: CLA; Lincomycin: LIN).

Fig. 6 The comparison of *sul* genes in bulk water and sediment between lakes and rivers (n, the number of studies).

Fig. 7 The transport of antibiotics and ARGs in different media and their possible exposure ways to human health (A: chemical pollutants and ARGs from sewage treatment plants and hospitals; B: chemical pollutants and ARGs from agriculture, aquaculture and animal husbandry; C: drinking water, D: recreation and swimming; E: ingestion of aquatic food; ① the sedimentation of pollutants from water; ② the

1328 resuspension of pollutants from sediment; ③ the microbial community and
1329 resistome changed due to the selection pressure of chemical pollution; ④ chemical
1330 pollutants accumulated by aquatic biota; ⑤ chemical pollutants and ARGs released
1331 from feces or dead body of aquatic animals; ⑥ ARGs ingested or adsorbed by
1332 aquatic biota; ⑦the transport of chemical pollutants and ARGs between aquatic
1333 biota).
1334

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

1349

1350

1351 **Fig. 7**

1352