University of Dundee

Reply to Gohar on 'Lungs, methotrexate and psoriasis', a comment on 'Fatal, incidental, idiopathic pulmonary fibrosis in a patient receiving long-term low-dose methotrexate for psoriasis'

Affleck, A.; Goudie, A.; Smith, R.

Published in:
Clinical and Experimental Dermatology

DOI:
[10.1111/ced.13994](https://doi.org/10.1111/ced.13994)

Publication date:
2019

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Affleck, A., Goudie, A., & Smith, R. (2019). Reply to Gohar on 'Lungs, methotrexate and psoriasis', a comment on 'Fatal, incidental, idiopathic pulmonary fibrosis in a patient receiving long-term low-dose methotrexate for psoriasis'. *Clinical and Experimental Dermatology*, 44(5), 595. <https://doi.org/10.1111/ced.13994>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Article Type: Correspondence

Lungs, methotrexate and psoriasis: reply from author

A. Affleck,¹ A. Goudie² and R. Smith²

Departments of ¹Dermatology and ²Respiratory Medicine, Ninewells Hospital,
Dundee, UK

Corresponding author: Andrew Affleck

E-mail: andrew.affleck@nhs.net

ORCID: 0000-0002-0385-3186

Funding: None

Conflicts of interest: None to declare

Thank you for your interest in our case report and discussion.¹ Pulmonary fibrosis arising in association with methotrexate therapy was first reported in the 1970s in the context of cancer chemotherapy with very high doses given (up to 3.5g/m²). Such patients accumulated a very high dose e.g. 41g over a few weeks.² In this specific

This is the peer reviewed version of the following article: Affleck, A., et al. "Lungs, methotrexate and psoriasis: reply from author", *Clinical and Experimental Dermatology* (2019), which has been published in final form at <https://doi.org/10.1111/ced.13994>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving. This article is protected by copyright. All rights reserved.

clinical scenario it is possible that methotrexate was causative although interestingly, the same author comments a year later that there are often many confounding factors e.g. use of concomitant agents which may also have pulmonary toxicity. They state, in practice, cytotoxic drug-induced lung disease may be difficult to establish with certainty because other potential causes of pulmonary disease are frequently present.³ Experimental mouse models have shown that administration of very high daily doses of methotrexate (3mg/kg) for 5 weeks causes pulmonary fibrosis.⁴ However, again, this is quite different from the low weekly doses of methotrexate given for psoriasis in clinical practice. Direct extrapolation from both these examples is not possible. Also, it should be emphasised that many apparent case reports of “pulmonary fibrosis” arising in the context of methotrexate treatment for psoriasis are in fact spurious including your reference 6 in which Karadag *et al*/ describe a case of resolving interstitial pneumonitis NOT fibrosis. With regards to the Japanese populations alluded to with psoriasis and interstitial pneumonia (IP) it is with interest that the prevalence of IPs is greater than the background prevalence of the most common form of chronic IP which is idiopathic pulmonary fibrosis (IPF).⁵ Regarding our patient, we suggest that there were 2 possibilities –

1. There was no connection between his psoriasis and interstitial lung disease; the apparent correlation was a coincidence.
2. There was a correlation between his psoriasis and interstitial lung disease in that both disorders shared underlying immunological pathways.

The same argument applies to the role of methotrexate in interstitial lung disease (ILD); active role or innocent bystander? We do not believe that methotrexate had any negative effect on the natural history of the ILD. Correlation does not prove

causation. We agree that there may be a group of patients with psoriasis who are at increased risk of developing ILD and identifying these individuals is desirable. However, at the present time we do not believe that routine chest CT scanning is indicated nor any blind preventative medical treatments both of which may have side-effects. Further discussion and debate is desirable.

References

1. Gohar A. Comment on "Fatal, incidental, idiopathic pulmonary fibrosis in a patient receiving long-term low-dose methotrexate for psoriasis
2. Sostman HD, Mattay RA, Putman CE *et al.* Methotrexate-induced pneumonitis. *Medicine* 1976; **55**: 371-385.
3. Sostman HD, Matthay RA, Putman CE. Cytotoxic drug-induced lung disease. *Am J Med* 1977; **62**: 608015.
4. Ohbayashi M, Susuki M, Yashiro Y, Fukuwaka *et al.* Induction of pulmonary fibrosis by methotrexate treatment in mice lung in vivo and in vitro. *J Toxicol Sci* 2010; **35**: 653-661.
5. Kawamoto, Hironori et al. "Interstitial Pneumonia in Psoriasis." *Mayo Clinic proceedings. Innovations, quality & outcomes* vol. 2,4 370-377. 20 Sep. 2018, doi:10.1016/j.mayocpiqo.2018.07.006