University of Dundee

Thiel Embalming

McDougall, Seaneen; Soames, Roger; Felts, Paul

Published in:
Clinical Anatomy

DOI:
[10.1002/ca.23491](https://doi.org/10.1002/ca.23491)

Publication date:
2020

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

McDougall, S., Soames, R., & Felts, P. (2020). Thiel Embalming: Quantifying histological changes in skeletal muscle and tendon and investigating the role of boric acid. *Clinical Anatomy*, 33(5), 696-704.
<https://doi.org/10.1002/ca.23491>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Thiel Embalming: Quantifying histological changes in skeletal muscle and tendon and investigating the role of boric acid

Journal:	<i>Clinical Anatomy</i>
Manuscript ID	CA-18-0309.R2
Wiley - Manuscript type:	Original Communications
Keywords:	anatomy
Additional Keywords:	Thiel embalming, anatomical teaching, histology, skeletal muscle, tendon

SCHOLARONE™
Manuscripts

Author Accepted Manuscript version of McDougall, S, Soames, R & Felts, P 2019, 'Thiel Embalming: Quantifying histological changes in skeletal muscle and tendon and investigating the role of boric acid', *Clinical Anatomy*. <https://doi.org/10.1002/ca.23491>
©2019 Wiley

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

1) Title and Abstract

Thiel Embalming: Quantifying histological changes in skeletal muscle and tendon and investigating the role of boric acid

Introduction: Cadaver preservation methods impact their utilisation in anatomical research and teaching. Thiel-embalmed cadavers show flexibility, however the cause remains poorly understood. This study aimed to i) describe qualitative and quantitative histological differences between Thiel-embalmed and formalin-fixed skeletal muscle and tendon tissue; ii) investigate whether boric acid in Thiel solution is solely responsible for modification of tissues, and iii) explore whether the modifications observed could potentially explain the mechanisms underpinning the flexibility of Thiel cadavers. **Materials and Methods:** Skeletal muscle and tendon samples were harvested from mice preserved using formalin, Thiel solution, or modified-Thiel solution (without Boric Acid). Using standard H&E and Gomori's trichrome histological methods, tissues were examined to determine whether differences were apparent between the preservative treatments. **Results:** Differences were present between the Thiel and formalin-fixed tissues; formalin-fixed samples remained substantially more intact while Thiel-embalmed samples showed fibre fragmentation and lack of nuclei. The mean cell diameter of Thiel-embalmed muscle (24.4 μm) was significantly smaller ($P<0.005$) than formalin-fixed muscle (40.7 μm). There was significantly greater ($P<0.005$) fragmentation in Thiel-embalmed muscle (631.5 per 1mm^2) compared to formalin-fixed muscle (75.4 per 1mm^2). Samples embalmed using modified-Thiel showed a severe lack of integrity within internal tissue structure. **Conclusions:** This suggests that Thiel solution significantly alters tissue structure at cellular level, with quantitative data demonstrating

measurable differences between Thiel and formalin-fixed specimens. While the precise mechanism for these alterations remains unknown, it is shown that boric acid is not the only component of Thiel responsible for degradation of internal tissue structure.

Keywords: Thiel embalming; anatomical teaching; histology; skeletal muscle; tendon

2) Text

Introduction

Teaching, research and understanding of anatomy is facilitated by the preservation and subsequent use of cadavers. The method of cadaver preservation impacts on how it can be utilised within the context of anatomical teaching and research. Traditionally, anatomy departments have used formalin embalming solutions. However, the resulting unrealistic texture, inflexibility and colour, combined with the deleterious effects of formalin on human health, have encouraged the search for alternative methods of preservation (Coleman and Kogan, 1998; Messmer et al., 2010; Hammer et al., 2011; Hammer et al., 2012). One alternative method is Thiel embalming (Thiel, 1992; Thiel, 2002). While Thiel embalming solution still contains a number of potentially hazardous ingredients including formalin, boric acid, chlorocresol and morpholine, these are found in very low concentrations, making the use of Thiel cadavers during dissection safer than traditional formalin cadavers (Eisma et al., 2010).

Cadavers preserved using the Thiel method demonstrate texture, flexibility and colour that is more similar to *in vivo* conditions for many of the body organs and tissues while articulated joints are freely movable (Thiel, 1992; Jaung et al., 2011). However, preservation of some organs, e.g. the brain, is problematic often resulting in additional fixation being required. Despite Thiel preserved cadavers being used in an increasing number of surgical and anatomical investigations (Benkhadra et al., 2009; Boryor et al., 2010; Eisma et al., 2010; McLeod et al., 2010; Fessel et al., 2011; Prasad Rai et al., 2012; Ernesto Ottone, 2016; Verstraete et al., 2016; Lone et al., 2017; Zwirner et al. 2019) a distinct paucity of information remains pertaining to how and why it produces cadavers with more life-like qualities.

Few studies have specifically investigated the microscopic effects of Thiel embalming on tissue, however Benkhadra et al. (2011), Hammer et al. (2015) and Martynuik et al. (2014) have reported qualitative signs of histological degradation in tissue samples embalmed using Thiel solution. Benkhadra et al. (2011) examined skeletal muscle and tendon samples from Thiel preserved human cadavers, Hammer et al. (2015) examined skeletal muscle from Thiel cadavers and Martynuik et al. (2014) examined porcine Thiel embalmed skeletal muscle and tendon. All three reported modifications to the tissue samples including fragmentation, acellularisation and loss of integrity, with the skeletal muscle being affected more than tendon. To date, no studies have attempted to establish whether these histological differences between Thiel-embalmed and formalin-fixed tissues are quantifiably significant.

Additionally, while the cross-linking effect of formalin-fixation on tissue is widely accepted, the components responsible for the modification of Thiel-embalmed tissue

1
2
3 remain poorly understood. Benkhadra et al. (2011) attempted to explain this
4
5 modification by highlighting the corrosive effect of acid on proteins. They suggested that
6
7 the boric acid component of Thiel solution may be denaturing the muscle proteins,
8
9 resulting in the breakdown of muscle fibre structure. However, while the degradative
10
11 effects of acids on proteins are well documented, boric acid may not hold the sole
12
13 responsibility for modifying the Thiel embalmed tissue - there are additional
14
15 components of the Thiel solution which are likely to have an effect on tissue structure
16
17 and these require further investigation. Both the Thiel perfusion and submersion
18
19 solutions contain high concentrations of chemical salts. Salts are known to alter tissue
20
21 structure, a phenomenon encouraged and harnessed in the meat technology industry in
22
23 order to influence 'sensory' characteristics of meat such as appearance and texture
24
25 (Desmond, 2006; Henney et al., 2010). It is therefore prudent to consider whether these
26
27 chemical salts are playing an important role in modifying the structure of Thiel-
28
29 embalmed tissues. Comparing a modified Thiel solution, with the Boric Acid removed, to
30
31 the standard Thiel solution and using formalin as a control will help to determine
32
33 whether it is the only component causing any modifications observed, or whether
34
35 further consideration needs to be given to the role other components, such as the
36
37 chemical salts, are playing in modifying Thiel-embalmed tissue. By better understanding
38
39 which components of Thiel are responsible for the modifications observed in skeletal
40
41 muscle and tendon, it may be possible to begin to better understand the mechanisms
42
43 underpinning the modifications and their potential role in creating the flexibility found in
44
45 Thiel cadavers.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The ability to understand the flexibility of Thiel-embalmed cadavers is important in the context of anatomical research and surgical training as this may impact on the types of procedures that can be reliably undertaken using the cadavers. Changes in the soft tissues surrounding bones, including skeletal muscle and tendon, are largely responsible for the flexibility, or inflexibility, of cadaver joints (Jaung et al., 2011), therefore the use of histology as a starting point to describe the gross structural changes occurring in skeletal muscle and tendon tissue may give an indication as to why this flexibility occurs in Thiel-embalmed cadavers.

The aims of this study were to use histology as a starting point to: i) describe differences between Thiel-embalmed and formalin-fixed skeletal muscle and tendon tissue both qualitatively and quantitatively; ii) investigate whether boric acid is solely responsible for the modifications seen in the tissue structure of Thiel-embalmed skeletal muscle and tendon tissue; and iii) ascertain whether the modifications observed could potentially indicate the mechanisms underpinning the flexibility of Thiel cadavers.

Materials and Methods

Due to the fact that part of the study involved the removal of the boric acid component of Thiel embalming solution and it could not be guaranteed that this modified solution would embalm effectively, mice were used as an animal model for human skeletal muscle and tendon tissue. Given that mice are routinely used as animal models for human skeletal muscle and tendon, and a pilot study showed there were no observable

1
2
3 differences between the effects of Thiel embalming on mouse tissue compared to
4
5 human tissue, it was deemed acceptable to use them as an animal model in this context.
6
7

8
9 All mice used in the study were sacrificed by CO₂ inhalation and immediately preserved.
10
11 Mice were embalmed in one of three ways; intra-cardiac perfusion of formalin, intra-
12
13 cardiac perfusion of Thiel solution (Table 1) followed by 2 weeks in Thiel submersion
14
15 solution; or intra-cardiac perfusion of modified Thiel (Table 2) followed by 2 weeks in
16
17 modified Thiel submersion solution. Given the small size of the animals used, we feel
18
19 that perfusion followed by two weeks in submersion fluid was ample to achieve full
20
21 fixation. After this time the mice were then processed through tissue harvesting and
22
23 histological examination. The number of mice used in each embalming treatment group,
24
25 as well as the tissues harvested and histological processing carried out is listed in Table
26
27 3. Tissue samples were discarded if they suffered structural damage during harvesting or
28
29 processing. Samples were selected to allow comparisons to be made within individual
30
31 animals, as well as between different animals.
32
33
34
35
36
37
38

39 The tissues were processed for histological examination using standard techniques.
40
41 Briefly, tissues were dehydrated through increasing concentrations of ethanol, passed
42
43 through xylene and embedded in paraffin wax. A Leica microtome was used to section
44
45 the samples at 6 µm before they were stained using haematoxylin and eosin (H & E),
46
47 examined and photographed using a light microscope (Leitz Orthoplan).
48
49
50

51
52 Further transverse muscle sections were taken at 4 µm from the formalin and Thiel-
53
54 embalmed samples and stained using Gomori's trichrome to visualise changes between
55
56 muscle, collagen and nuclei which stain red, green/blue and black respectively.
57
58
59
60

The qualitative parameters used to determine differences between the preservative treatments on the basis of visible histomorphology included overall cell structure, cell organisation and integrity, the presence or absence of nuclei, the appearance of skeletal muscle striations in the muscle samples, visibility of connective tissue within the skeletal muscle samples and collagen fibre alignment in the tendon samples. Similar parameters were assessed in a total of eight additional control Thiel-embalmed samples (4 skeletal muscle and 4 tendon) that were either 'washed' overnight in ethanol/water prior to staining or were post-fixed in formalin. This was a posthoc investigation to ensure that any fragmentation or lack of staining occurring in samples was not the result of salts embedded in the sample, or the tissue being inherently too fragile for processing.

Quantitative data were gathered by measuring the diameter of muscle fibres on transverse sections using ImageJ software (Schneider et al., 2012) which calculates the minimum Feret's diameter of each fibre. The minimum Feret's diameter is the 'minimum distance of parallel tangents at opposing borders of the muscle fibre' (Briguet et al., 2004) and is used to minimise the impact of issues such as orientation of the cutting angle during sample sectioning when measuring the diameter of skeletal muscle fibres.

In addition, the number of visible breakages within longitudinally sectioned muscle fibres was recorded, using the counter tool in ImageJ, in order to quantify fragmentation of the muscle tissue. Finally, the presence of any areas of disruption within longitudinally sectioned tendon fibres was recorded using Image J to quantify disruption of the tendon tissue.

Statistical Analysis

The statistical analyses of the comparison between the formalin and Thiel-embalmed sample groups, for the mean diameter of skeletal muscle fibres, the mean occurrence of muscle fibre fragmentation, and the mean disruption of the tendon fibres was carried out using a Mann-Whitney test for two independent samples for non-parametric data. All analyses were carried out using IBM SPSS Statistics 21 software package.

The study was approved by the relevant institutional review board.

Results:

Post-hoc Thiel samples 'washed' overnight in ethanol/water, or post-fixed in formalin, prior to staining, did not show any observable difference in staining or fragmentation compared to those samples included in the main study. This suggested that any changes observed were a result of the Thiel embalming process itself rather than histological processing.

Qualitative changes: Formalin vs. Thiel vs. Modified Thiel

Skeletal Muscle:

On visual inspection all muscle samples maintained their gross structure, however when examined histologically all Thiel-embalmed and modified-Thiel-embalmed muscle samples showed considerable differences compared to the formalin-fixed samples: Figure 1 shows typical attributes of the different tissue treatments in longitudinal

sections. Changes in tissue structure were substantial enough to be obvious at low magnifications. All Thiel-embalmed muscle samples, including the modified-Thiel samples, were affected in a number of ways, including a loss of nuclear staining and a loss of definitive striations in some cells, considerable fibre fragmentation and a loss of cell structure integrity. These changes appeared to be more severe in the modified-Thiel-embalmed muscle samples compared to the standard Thiel samples. These histological changes were observed in each of the three muscles sampled (vastus lateralis, pectoralis major and extensor digitorum longus), demonstrating that the changes were not limited to any specific muscle or to one individual animal. These observations were not present in the formalin-fixed muscle tissues, where the fibres and their nuclei retained their histological integrity.

Gomori-trichrome staining of the Thiel-embalmed transverse muscle sections, showed relatively intact collagen present within the endomysium, perimysium and epimysium of the muscle samples (Fig. 2), suggesting that perhaps collagen is stabilising the gross structure of the tissues.

Tendon

On visual inspection all tendon samples appeared to maintain integrity of gross structure. However, on histological examination, the Thiel-embalmed and modified-Thiel-embalmed tendon samples showed minor differences compared to the formalin-fixed tendon samples (Fig. 3), with the changes becoming more apparent at higher magnifications. The Thiel-embalmed and modified-Thiel-embalmed samples, while very

similar to each other in appearance, showed less uniformity in the collagen bundles, with a slightly disrupted appearance and more breakages when compared to the formalin samples. Despite these modifications to the microscopic structure the fibres remained aligned. The histological modifications were observed in both tendons examined (calcaneal and extensor digitorum longus tendons), in each of the animals, demonstrating that they were not specific to individual tendons or individual animals.

These observations were not present in the formalin-fixed tendon tissues, where the fibres retained their histological integrity.

Quantitative changes: Formalin vs. Standard Thiel

Skeletal Muscle

The mean diameter of Thiel-embalmed muscle fibre samples was significantly smaller ($P < 0.005$) at $24.4 \mu\text{m}$ ($n=10$, $SD=\pm 6.5$) than that of formalin-fixed samples at $40.7 \mu\text{m}$ ($n=10$, $SD = \pm 4.4$), using a Mann-Whitney test.

The mean occurrence of fragmentation within Thiel-embalmed samples was significantly greater ($P < 0.005$) at 631.5 per 1 mm^2 ($n=10$, $SD = \pm 245.36$) compared to 75.4 per 1 mm^2 ($n=10$, $SD = \pm 49.07$) in formalin-fixed samples, using a Mann-Whitney test.

Tendon

There was a significant increase ($P < 0.05$) in the average amount of disruption in Thiel embalmed tendon sample images at 60 per 1 mm^2 ($n=10$, $SD = \pm 30$) compared to the

formalin embalmed tendon sample images at 27.5 per 1mm² (n=10, SD = ±15), using a Mann-Whitney test.

Despite significant differences in the amount of disruption observed in Thiel and formalin tendon samples, the total disruption in Thiel embalmed tendon is very low compared to the disruptive fragmentation seen in Thiel embalmed muscle (631.5 per 1mm² in muscle compared to 60 per 1mm² in tendon).

Discussion:

It is clear that the two methods of preservation used in the current study are operating via different process: traditional formalin fixation produces crosslinking which preserves the tissue by preventing degradation, while Professor Thiel himself described his method as a process similar to ‘pickling’ (Thiel, 1992), a process which uses high concentrations of salts to preserve foods via anaerobic fermentation (Barrett, 2003). It is important to compare and contrast the effects of the two different preservation processes for a number of reasons. Comparing Thiel and formalin specimens serves as a control; there are fewer changes in structural integrity noted in formalin-fixed specimens, demonstrating that histological processing is not the likely cause of the degradation seen in Thiel-embalmed tissue, but rather that the degradation is inherently linked to the Thiel process itself. Perhaps even more importantly, if anatomy departments are to use soft-fix embalming methods such as Thiel, then they need to understand the effect this has on tissue structure, how it differs from the traditional formalin fixation used and what effect this may have on future anatomical and clinical research projects.

The present study has shown, both qualitatively and quantitatively, that the internal structure of skeletal muscle and tendon preserved using the Thiel method show significant changes. This supports the observations of both Benkhadra et al. (2011) and Martyniuk et al. (2014) who demonstrated similar degradative changes in skeletal muscle and tendon, as well as Hammer et al. (2015), who described Thiel-embalmed skeletal muscle cells as being 'dissolved' and having a 'washed-out' appearance, with a lack of nuclei and degraded structural borders. The quantitative data of the current study demonstrate that these tissue structure changes result in significant measurable difference between the Thiel and formalin-fixed specimens.

While changes in tissue characteristics of Thiel-embalmed samples have previously been described in a number of investigations (Benkhadra et al., 2011; Fessel et al., 2011; Wilke et al., 2011; Martyniuk et al., 2014; Hammer et al., 2015; Verstraete et al., 2015; Venne et al., 2018; Zwirner et al., 2019), only Benkhadra et al. (2011) have attempted to specifically explain which of the components of Thiel solution are responsible for the changes observed. They suggest that, given the denaturing effects of acids on proteins, boric acid must be responsible for the degradation of proteins involved in maintaining the integrity of skeletal muscle and tendon structure. This study sought to determine whether boric acid was the sole component responsible for tissue structure degradation by comparing tissue changes in a modified Thiel solution which contained no boric acid to those observed in the standard Thiel solution, with formalin-fixed samples serving as a control. The results of this study showed that the internal structure of skeletal muscle samples, from the modified-Thiel solution, underwent a severe change in internal

structure even though boric acid was excluded from the embalming solution, suggesting that other components also play a role in driving these changes.

As previously discussed, Thiel solution contains high concentrations of chemical salts in both the perfusion and submersion solutions (Thiel, 1992). The chemical salts in Thiel solution are considered to be responsible for fixation (Kerckaert et al. 2008) and indeed various salts have been used as part of the embalming process for thousands of years (Coleman and Kogan, 1998; Bremner, 2014). However, these salts may also affect tissue structure by solubilisation, a phenomenon encouraged and harnessed in the meat technology industry in order to influence the 'sensory' characteristics of meat such as appearance, texture and smell, (Desmond, 2006; Henney et al., 2010), although the process by which this occurs is complex and not completely understood. Links have previously been made between salt-based embalming processes and the similar process occurring during salting or curing of meat (Coleman and Kogan, 1998). It is, therefore, proposed that the chemical salts found in Thiel solution also play a major role, alongside boric acid, in denaturing the skeletal muscle myofibrillar proteins, leading to changes in fibre structure.

While the internal structure of Thiel and modified-Thiel samples appears to be affected, the external gross structure remains essentially intact. This could be explained by a lack of degradation of collagen in the structural membrane of the tissue samples. As shown using the Gomori trichrome stain, the collagen in the Thiel-embalmed muscle samples remains relatively stable and intact, despite the degradation and fragmentation of the muscle fibres. Preservation of collagen would also explain why tendon shows less disruption than skeletal muscle. It may also be the case that the proteins degraded

1
2
3 within the muscle samples play a role in helping to maintain the overall gross structure
4
5 of the muscle samples. Indeed, Wolff et al. (2008) suggest that protein denaturation
6
7 occurs due to the presence of high salt levels in Thiel solution, leading to precipitation
8
9 and homogenisation of the tissue which, together with additional precipitation and
10
11 linkage caused by the Thiel solution, helps to maintain the texture of the tissues. This is
12
13 similar to a process often termed the 'salting out effect' in the meat technology industry
14
15 where salts help to solubilise meat proteins which then form an exudate which acts as a
16
17 sticky 'cement' between meat pieces (Smith, 2001, Hutton, 2002). It is perhaps likely
18
19 that a combination of these two processes is helping to maintain the external tissue
20
21 structure while the internal structure is degraded.
22
23
24
25
26
27

28 The flexibility of cadavers preserved using Thiel solution compared to formalin-fixed
29
30 cadavers is one of the main advantages of this method as it allows more movement and
31
32 facilitates training in surgical procedures (Giger et al., 2008; Kerckaert et al., 2008; Eisma
33
34 et al., 2010; Prasad Rai et al., 2012; Yiasemidou et al., 2017); however very little is
35
36 known about how this flexibility arises. This becomes important in understanding the
37
38 role Thiel cadavers can play in education, research and surgical training. As the flexibility
39
40 of cadavers is largely related to the composition of soft tissues surrounding joints (Jaung
41
42 et al., 2011) it is highly likely that Thiel cadaver flexibility is, at least partially, a result of
43
44 the change in tissue structure integrity, particularly that of skeletal muscle tissue. The
45
46 widespread fragmentation observed in the Thiel fibres results in a lack of longitudinal
47
48 integrity, meaning that there will be very little resistance offered to any stretch applied
49
50 to the tissue. It is hypothesised that the mechanism underpinning the flexibility is likely a
51
52 complex interplay between the degradation of the internal structure of skeletal muscle,
53
54
55
56
57
58
59
60

relative preservation of collagen integrity, as well as perhaps a role for protein exudate 'stickiness'.

Study Limitations and Suggestions for Future Work

There are a number of limitations with this study, which should be addressed in future work. Firstly, as previously discussed, due to modification of the Thiel solution as part of the study, it was deemed inappropriate at this stage to use human donor material, however the study would ideally be carried out using human cadaveric material.

This histological study alone cannot provide a definitive answer as to what is causing the flexibility observed in Thiel cadavers. However it is part of a wider study including both biomechanical and protein analysis investigations, which may provide a more robust hypothesis as to the mechanisms underpinning flexibility of Thiel cadavers. The results of these studies highlight the potential role of collagen in maintaining overall gross structure while the internal muscle structure is degraded. It is recommended that future work would involve a more detailed analysis of collagen structure to determine whether the collagen is indeed remaining relatively intact and providing the overall support for the gross tissue structure. This may be aided by examining the histological structure of additional body tissues embalmed using Thiel solution; those with relatively higher collagen content such as skin and ligaments, compared to those with relatively lower collagen content such as kidney and spleen.

A further recommendation would be to specifically investigate the role of the other components in the Thiel solution, in order to understand whether they are playing a key

role in the degradation of the skeletal muscle structure, as seems to be indicated by the results of this study. While it is hypothesised that it is the chemical salts playing a major role in the degradation of internal tissue structure, it would be prudent to investigate this further, while perhaps also investigating the effects of other Thiel solution components such as the ethylene-glycol which is believed to be responsible for the plasticity or 'soft haptics' of Thiel cadavers (Hammer et al, 2015). While it is likely that there is a complex interplay between the components of the Thiel solution and their effects on tissue, it would be useful to better understand the effects of each component on tissue structure.

Understanding how Thiel differs from formalin in its effect on cadaveric tissue, which components are responsible for the changes seen and what the overall effects of these changes are, is very important in determining which anatomical and surgical training procedures cadavers and their tissues can be used for. Not only will this allow more robust and reliable data to be gathered from any associated studies, but it will also permit departments to plan their use of cadavers accordingly – ensuring that cadavers are used to their full potential within an anatomical and surgical teaching and research setting.

Concluding remarks

From the results of this study, it is concluded that there are observable and measureable differences between tissues embalmed using Thiel solution and formalin. These differences are not solely due to the effects of boric acid on the tissue proteins and it is suggested that perhaps the chemical salts present in the Thiel solution are playing a role

in the degradation of the internal tissue structure. The changes observed, with degradation and fragmentation of the muscle fibre cells, while the surrounding collagen remains relatively intact, are very likely to be influencing the flexibility of Thiel cadavers as there is not the same impediment to flexibility that occurs when the tissues are ‘hardened’ during formalin fixation via crosslinking.

1) References

Barrett DM. 2003. Pickling. In Caballero B, Trugo L, Finglas P, Editors. Encyclopaedia of food sciences and nutrition: Volumes 1-10 (2nd Ed). Elsevier Science BV. P 4563-4566.

Benkhadra M, Bouchot A, Gerard J, Genelot D, Trouilloud P, Martin L, Girard C, Danin A, Anderhuber F, Feigl G. 2011. Flexibility of Thiel’s embalmed cadavers: the explanation is probably in the muscles. Surg Radiol Anat, 33:365-368.

Benkhadra M, Savoldelli G, Fournier R, Gamulin Z, Francois C, Trouilloud P, Feigl G, Fasel J. 2009. A new anatomical technique to investigate nerves by imagery. Surg Radiol Anat, 31:221-224.

Boryor A, Hohmann A, Wunderlich A, Geiger M, Kilic F, Sander M, Sander C, Bockers T, Sander F. 2010. In-vitro results of rapid maxillary expansion on adults compared with finite element simulations. J Biomech, 43:1237-1242.

Briguet A, Courdier-Fruh I, Foster M, Meier T, Magyar J. 2004. Histological parameters for the quantitative assessment of muscular dystrophy in the mdx-mouse. Neuromuscul Disord. 14:675-682.

Coleman R. & Kogan I. 1998. An improved low-formaldehyde embalming fluid to preserve cadavers for anatomy teaching. J Anat, 192:443-446.

Desmond E. 2006. Reducing salt: A challenge for the meat industry. Meat Sci, 74:188-196.

- 1
2
3 Eisma R, Mahendran S, Majumdar S, Smith D, Soames R. 2010. A comparison of Thiel and formalin-
4 fixed cadavers for thyroid surgery training. *Surg-J R Coll Surg E*, 9:142-146.
5
6
7
8 Ernesto Ottone N, Vargas CA, Fuentes R, del Sol M. 2016. Walter Thiel's Embalming Method. Review
9 of Solutions and Applications in Different Fields of Biomedical Research. *Int J of Morph*. Dec 1;34(4).
10
11
12
13 Fessel G, Frey K, Schweizer A, Calcagni M, Ullrich O, Snedeker J. 2011. Suitability of Thiel-embalmed
14 tendons for biomechanical investigation. *Anat Anz*, 193:237-241.
15
16
17
18
19 Giger U, Frésard I, Häfliger A, Bergmann M, Krähenbühl L. 2008. Laparoscopic training on Thiel
20 human cadavers: a model to teach advanced laparoscopic procedures. *Surg Endo*. Apr 1;22(4):901-6.
21
22
23
24 Hammer N, Löffler S, Feja C, Bechmann I, Steinke H. 2011. Substitution of Formaldehyde in Cross
25 Anatomy Is Possible. *J Natl Cancer Inst*, 103:610-611.
26
27
28
29
30 Hammer N, Löffler S, Feja C, Sandrock M, Schmidt W, Bechmann I, Steinke H. 2012. Ethanol-glycerin
31 fixation with thymol conservation: A potential alternative to formaldehyde and phenol embalming.
32
33
34
35 Anat Sci Educ, 5:225-233.
36
37
38 Hammer N, Löffler S, Bechmann I, Steinke H, Hädrich C, Feja C. 2015. Comparison of modified Thiel
39 embalming and ethanol-glycerin fixation in an anatomy environment: potentials and limitations of
40 two complementary techniques. *Anat Sci Educ*. 8(1):74-85.
41
42
43
44
45 Henney J, Taylor C, Boon C. 2010. Strategies to reduce sodium intake in the United States.
46 Washington D.C. National Academy Press. Pg. 67
47
48
49
50 Hutton T. 2002. Sodium technological functions of salt in the manufacturing of food and drink
51 products. *Brit Food Journ.*, 104, 126-152.
52
53
54
55 Jaung R, Cook P, Blyth P. 2011. A comparison of embalming fluids for use in surgical workshops. *Clin*
56
57
58 Anat, 24:155-161.
59
60

- 1
2
3 Kerckaert I, Van Hoof T, Pattyn P, D'Herde K. 2008. Endogent: centre for anatomy and invasive
4 techniques. *Anatomy*, 2:28-33.
5
6
7
8 Lone M, McKenna JP, Balta JY, O'Mahony SM, Cryan JF, Downer EJ, Toulouse A. 2017. Assessment of
9 Thiel-Embalmed Cadavers as a Teaching Tool for Oral Anatomy and Local Anesthesia. *J of Den Educ*.
10 81(4):420-6.
11
12
13
14
15 Martyniuk A, Easteal R, Ellis R. 2014. Histological characterization of soft-embalmed porcine tendon
16 and muscle. *FASEB J*, 28:914-1.
17
18
19
20
21 McLeod G, Eisma R, Schwab A, Corner G, Soames R, Cochran S. 2010. An evaluation of Thiel-
22 embalmed cadavers for ultrasound-based regional anaesthesia training and research. *BMUS J*,
23 18:125-129.
24
25
26
27
28 Messmer C, Kellogg R, Zhang Y, Baiak A, Leiweke C, Marcus J, Levin S, Zenn M, Erdmann D. 2010. A
29 technique to perfuse cadavers that extends the useful life of fresh tissues: The Duke experience.
30 *Anat Sci Educ*, 3:191-194.
31
32
33
34
35 Prasad Rai B, Tang B, Eisma R, Soames R, Wen H, Nabi G. 2012. A qualitative assessment of human
36 cadavers embalmed by Thiel's method used in laparoscopic training for renal resection. *Anat Sci*
37 *Educ*, 5:182-186.
38
39
40
41
42
43 Schneider C, Rasband W, Eliceiri K. 2012. "NIH Image to ImageJ: 25 years of image analysis". *Nature*
44 *Methods*, 9:671-675.
45
46
47
48
49 Smith D. 2001. Functional properties of muscle proteins in processed poultry products. In: Sams, A.
50 (ed.) *Poultry Meat Processing*. Boca Raton: CRC Press.
51
52
53
54 Thiel W. 1992. The preservation of whole bodies in natural dyes - Die konservierung ganzer leichen
55 in natuerlichen farben. *Anat Anz*, 174:185-195.
56
57
58
59
60

Thiel W. 2002. Ergänzung für die Konservierung ganzer Leichen nach W. Thiel. *Anat Anz*, 184: 267-269.

Venne G., Welte L., Noel G. 2018. Quantitative and Qualitative Comparison of Thiel and Phenol-Based Soft-Embalmed skin for Surgical Training. *The FASEB Journal* 2018 32: (1_supplement), pp.632-6.

Verstraete M, Van Der Straeten C, Opsomer GJ, De Lepeleere B, Victor J. 2016. Evaluation of embalming and freezing on biomechanical properties of tendons. In *Orthopaedic Proceedings* 2016 Jan (Vol. 98, No. SUPP_4, pp. 98-98). Brit Ed Soc of Bone & Joint Surg.

Wilke H, Werner K, Häussler K, Reinehr M, Böckers T. 2011. Thiel-fixation preserves the nonlinear load–deformation characteristic of spinal motion segments, but increases their flexibility. *J Mech Behav Biomed Mater*, 4:2133-2137.

Wolff KD, Kesting M, Mücke T, Rau A, Hölzle F. 2008. Thiel embalming technique: a valuable method for microvascular exercise and teaching of flap raising. *Microsurgery: Off. J Intl Micros Soc and Euro Fed. Soc Micros.*;28(4):273-8.

Yiasemidou M, Roberts D, Glassman D, Tomlinson J, Biyani S, Miskovic D. 2017. A multispecialty evaluation of Thiel cadavers for surgical training. *World J Surg*. May 1;41(5):1201-7.

Zwirner J, Scholze M, Ondruschka B and Hammer N. 2019. Tissue biomechanics of the human head are altered by Thiel embalming, restricting its use for biomechanical validation. *Clin Anat*, doi:[10.1002/ca.23409](https://doi.org/10.1002/ca.23409)

2) Footnotes

There is no conflict of interest to declare.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

3) Figure Legends

Figure 1: Comparison of H and E stained longitudinal mouse skeletal muscle samples preserved using Thiel (A, B, C and D), modified-Thiel (E and F) and formalin (G and H) shown at increasing magnification (A, C, E and G x4, B, D, F and H x40). A, B, E, F, G and H represent samples of vastus lateralis muscle, C and D represent pectoralis major muscle.

Figure 2: Comparison of Gomori trichrome stained transverse mouse vastus lateralis skeletal muscle samples fixed with Thiel’s solution (A and B) or formalin (C and D), shown at increasing magnification (A and C x4, B and D x10).

Figure 3: Comparison of H and E stained mouse calcaneal tendon samples preserved using Thiel (A, B, C and D), modified-Thiel (E and F) and formalin (G and H) shown at increasing magnification (A, C, E and G x10, B, D, F and H x40). A, B, E, F, G and H represent samples of calcaneal tendon, C and D represent samples taken from extensor digitorum longus muscle/tendon complex.

4) Tables

- Table 1 – Thiel solution recipe as used at Institution**
- Table 2 – Modified Thiel solution recipe – without boric acid**
- Table 3 – Sample Data**

***Please see attached files**

Figure 1: Comparison of H and E stained longitudinal mouse skeletal muscle samples preserved using Thiel (A, B, C and D), modified-Thiel (E and F) and formalin (G and H) shown at increasing magnification (A, C, E and G x4, B, D, F and H x40). A, B, E, F, G and H represent samples of vastus lateralis muscle, C and D represent pectoralis major muscle.

160x81mm (300 x 300 DPI)

Figure 2: Comparison of Gomori trichrome stained transverse mouse vastus lateralis skeletal muscle samples fixed with Thiel's solution (A and B) or formalin (C and D), shown at increasing magnification (A and C x4, B and D x10).

80x61mm (300 x 300 DPI)

Figure 3: Comparison of H and E stained mouse calcaneal tendon samples preserved using Thiel (A, B, C and D), modified-Thiel (E and F) and formalin (G and H) shown at increasing magnification (A, C, E and G x10, B, D, F and H x40). A, B, E, F, G and H represent samples of calcaneal tendon, C and D represent samples taken from extensor digitorum longus muscle/tendon complex.

160x80mm (300 x 300 DPI)

Table 1 – Thiel solution recipe as used at Institution

Thiel Arterial Solution: 1 litre	Thiel Submersion Solution: 1.5 litre
(As used at Institution)	(As used at Institution)
Water 590ml	Water 1200ml
Boric acid 18g	Boric acid 41.25g
Propylene glycol 175ml	Propylene glycol 135ml
Ammonium nitrate 118g	Ammonium nitrate 135g
Potassium nitrate 30g	Potassium nitrate 67.5g
Stock II 35ml	Stock chlorocresol solution 28ml
Sodium sulphite 50g	Sodium sulphite 94g
Formalin (37%) 35ml	Formalin (37%) 28ml
Morpholine 10ml	
Alcohol 70ml	

Table 2 – Modified Thiel solution recipe – without boric acid

Thiel Arterial Solution: 1 litre (As used at Institution)	Thiel Submersion Solution: 1.5 litre (As used at Institution)
Water 608ml	Water 1241ml
Propylene glycol 175ml	Propylene glycol 135ml
Ammonium nitrate 118g	Ammonium nitrate 135g
Potassium nitrate 30g	Potassium nitrate 67.5g
Stock II 35ml	Stock chlorocresol solution 28ml
Sodium sulphite 50g	Sodium sulphite 94g
Formalin (37%) 35ml	Formalin (37%) 28ml
Morpholine 10ml	
Alcohol 70ml	

Table 3 – Tissue sample data

Treatment	No. of mice used	Samples Harvested (number in brackets)	Histological Processing & Analysis
Formalin Control - (8.9% formalin solution with phenol)	12	<ul style="list-style-type: none">- vastus lateralis (23)- pectoralis major (6)- extensor digitorum longus muscle/tendon complex (22)- calcaneal tendon (24)	<ul style="list-style-type: none">- H&E- Gomori's trichrome- Qualitative Analysis- Quantitative Analysis
Thiel	12	<ul style="list-style-type: none">- vastus lateralis (24)- pectoralis major (5)- extensor digitorum longus muscle/tendon complex (20)- calcaneal tendon (22)	<ul style="list-style-type: none">- H&E- Gomori's trichrome- Qualitative Analysis- Quantitative Analysis
Modified Thiel without boric acid	3	<ul style="list-style-type: none">- vastus lateralis (4)- pectoralis major (2)- extensor digitorum longus muscle/tendon complex (3)- calcaneal tendon (4)	<ul style="list-style-type: none">- H&E- Qualitative Analysis