University of Dundee

Cerebrofaciothoracic dysplasia

Deciphering Developmental Disorders (DDD) Study; Michael Yates, Thabo; Ng, Oon Hui; Offiah, Amaka C.; Willoughby, Josh; Berg, Jonathan N.

Published in:
American Journal of Medical Genetics Part A

DOI:
[10.1002/ajmg.a.60678](https://doi.org/10.1002/ajmg.a.60678)

Publication date:
2019

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Deciphering Developmental Disorders (DDD) Study, Michael Yates, T., Ng, O. H., Offiah, A. C., Willoughby, J., Berg, J. N., & Johnson, D. S. (2019). Cerebrofaciothoracic dysplasia: Four new patients with a recurrent TMCO1 pathogenic variant. *American Journal of Medical Genetics Part A*, 179(1), 43-49.
<https://doi.org/10.1002/ajmg.a.60678>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

This is a repository copy of *Cerebrofaciothoracic dysplasia: four new patients with a recurrent TMCO1 pathogenic variant.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/140464/>

Version: Accepted Version

Article:

Yates, T.M., Ng, O.-H., Offiah, A.C. orcid.org/0000-0001-8991-5036 et al. (4 more authors) (2018) Cerebrofaciothoracic dysplasia: four new patients with a recurrent TMCO1 pathogenic variant. American Journal of Medical Genetics Part A. ISSN 1552-4825

<https://doi.org/10.1002/ajmg.a.60678>

This is the peer reviewed version of the following article: Cerebrofaciothoracic dysplasia: four new patients with a recurrent TMCO1 pathogenic variant, which has been published in final form at <https://doi.org/10.1002/ajmg.a.60678> . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Cerebrofaciothoracic Dysplasia: four new patients with a recurrent TMCO1 pathogenic variant

Running title: TMCO1-related Cerebrofaciothoracic Dysplasia

T. Michael Yates¹, Oon-Hui Ng², Amaka C. Offiah³, Josh Willoughby⁴, DDD Study⁵, Jonathan N. Berg⁶, Diana S. Johnson¹

¹Sheffield Clinical Genetics Service, Sheffield Children's NHS Foundation Trust, Sheffield, UK

²Ninewells Hospital and Medical School, University of Dundee, Dundee, UK

³Academic Unit of Child Health, University of Sheffield, Sheffield, UK; Radiology Department, Sheffield Children's NHS Foundation Trust, Sheffield, UK

⁴Sheffield Diagnostic Genetics Service, Sheffield Children's NHS Foundation Trust, Sheffield, UK

⁵ DDD Study, Wellcome Trust Sanger Institute, Hinxton, Cambridge, UK

⁶Dundee Clinical Genetics Service, Ninewells Hospital and Medical School, Dundee, UK

Corresponding Author

Dr Diana Johnson

Sheffield Clinical Genetics Service

Sheffield Children's NHS Foundation Trust

Sheffield, UK

Ph: +44 114 2717025

Fax: +44 114 2737467

E-mail: diana.johnson@sch.nhs.uk

Conflict of interest

The authors declare no conflict of interest.

Abstract

Biallelic loss of function variants in the TMCO1 gene have been previously demonstrated to result in Cerebrofaciothoracic Dysplasia (CFTD; MIM #213980). The phenotype of this condition includes severe intellectual disability, as well as distinctive craniofacial features, including brachycephaly, synophrys, arched eyebrows, 'cupid's bow' upper lip, and microdontia. In addition, non-specific skeletal anomalies are common, including bifid ribs, scoliosis and spinal fusion. Only 19 molecularly confirmed patients have been previously described. Here, we present four patients with CFTD, including three brothers from a Pakistani background and an additional unrelated white Scottish patient. All share the characteristic craniofacial appearance, with severe intellectual disability and skeletal abnormalities. We further define the phenotype with comparison to the published literature, and present images to define the dysmorphic features in a previously unreported ethnic group. All of our patient series are homozygous for the same c.292_293del (p.Ser98*) TMCO1 pathogenic variant, which has been previously reported only in an isolated Amish population. Thus we provide evidence that CFTD may be more common than previously thought. The patients presented here further delineate the phenotypic spectrum of CFTD and provide evidence for a recurrent pathogenic variant in TMCO1.

Keywords: TMCO1, Cerebrofaciothoracic Dysplasia, Intellectual Disability

Introduction

Cerebrofaciothoracic dysplasia (CFTD; MIM #213980) is a rare, autosomal recessive, developmental disorder. It is characterized by distinctive craniofacial features (including brachycephaly, synophrys and microdontia), skeletal abnormalities, mostly affecting the ribs and vertebrae, and intellectual disability. CFTD is caused by biallelic (all reported are homozygous) loss of function variants in the TMCO1 gene (Alanay et al., 2013). Of note, Xin et al. (2010) identified a condition described as TMCO1 defect syndrome prior to the molecular basis of CFTD being identified (Alanay et al., 2013), although they did not make the link to CFTD clinically. It is now thought that TMCO1 defect syndrome is part of the CFTD spectrum (Alanay et al., 2013; Pehlivan et al., 2014).

Only 19 molecularly confirmed patients with CFTD have been reported to date (Alanay et al., 2013; Caglayan et al., 2013; Pehlivan et al., 2014; Xin et al., 2010). Here, we present three further familial patients together with an unrelated additional patient. All share a recurrent TMCO1 pathogenic variant. Description of these four new patients and review of those previously described allows further delineation of the genotypic and phenotypic spectrum of this condition.

Subjects and Methods

Editorial policies and Ethical Considerations

No ethical approval has been sought as this is a retrospective study with informed parental consent to publish.

Methods

Patients 1 and 4 were identified through the Wellcome Trust Deciphering Developmental Disorders (DDD) study. Recruitment to this study was via UK regional Clinical Genetics centers after routine referral. The affected person and their parents had microarray and trio-based exome sequencing and analysis as per Wright et al. (2015). Variants identified as pathogenic were confirmed via Sanger sequencing in a clinical laboratory setting. Patients 2 & 3 were tested by Sanger sequencing for the variant found in their sibling.

Results

Clinical Reports

Patient 1 (DECIPHER ID 259339)

This male patient was born to consanguineous parents (second cousins) of North-West Pakistani origin. The pregnancy was uncomplicated. He was delivered at 35 weeks by elective cesarean section and birth weight was 3.40 kg (98th centile). He spent one day following birth in Special Care due to hypoglycemia. He first sat upright at one year of age, walked at two years, and first words were at 18 months of age.

He was initially referred to his local Genetics service at the age of 13 years and 8 months, with severe learning disability and a diagnosis of Autism Spectrum Disorder. His verbal communication was limited to sentences of a few words.

On examination, he had blue irides, synophrys, downslanting palpebral fissures, macroglossia, high arched palate, everted lower lip, low set ears, narrow chest, generalized joint laxity and adducted halluces (Fig. 1), and an intention tremor. Head circumference was on the 91st centile, height on the 98th centile and weight between the 50th-75th centiles. MRI brain and spine at the age of ten years demonstrated low lying cerebellar tonsils.

His parents advised they had two other similarly affected male children, therefore individual genetic evaluations were initiated.

Patient 2 (DECIPHER ID 358411)

Patient 2 is the brother of Patient 1. The pregnancy was unremarkable. He was delivered at 42 weeks by emergency cesarean section due to cephalopelvic disproportion, with a birth weight of 4.89 kg (98th-99.6th centiles). He had bilateral talipes equinovarus. He first smiled at the age of six months, walked at seven years and first words were at four years of age.

On assessment at the age of twenty-nine years, he had synophrys, arched eyebrows, mild hypertelorism, 'cupid's bow' upper lip, everted lower lip, high narrow palate, grooved tip of the tongue and low-set ears. There was hypermobility of the distal joints of the hands, and he had a left-sided Sprengel deformity clinically (Fig.1). His head circumference was on the 99.6th centile. His behavior was generally affable.

A chest radiograph (Fig. 2) demonstrated a dorsal scoliosis concave to the right, fused left 4th and 5th ribs posteriorly, and an un-ossified posterior end of the left 6th rib. There was also a bony bridge leading to posterior fusion of the left 7th and 8th ribs. CT head demonstrated agenesis of the corpus callosum.

Patient 3 (DECIPHER ID 358412)

Patient 3 is the brother of Patients 1 and 2. Antenatal ultrasound had demonstrated an abnormal bladder, polyhydramnios, kinking of the thoracic spine and an enlarged 3rd cerebral ventricle. He was delivered at 38 weeks by elective cesarean section, with a birth weight of 4.02 kg (91st-98th centiles). At birth he was noted to have scoliosis, prominent left-sided ribs, left talipes equinovarus, a systolic heart murmur, and congenital macrocephaly (above 99.6th centile). His length at birth was on the 97th centile.

He also had a left-sided inguinal hernia, undescended left testis, Fallot's tetralogy and scoliosis, all of which required surgical treatment. He had severe congenital bilateral sensorineural hearing loss and severe developmental delay . He first sat upright at the age of six years and has never walked, but is able to bottom shuffle. He is non-verbal, and cannot feed himself.

On assessment at the age of twenty-six years, he had synophrys, mild hypertelorism, 'cupid's bow' upper lip, everted lower lip, low set and posteriorly rotated ears, gingival hypertrophy, a short neck and an intention tremor (Fig.1) . He

still has his primary dentition. His head circumference was between the 75th and 91st centiles.

Chest and spinal radiographs showed abnormalities of the spine with multiple block vertebrae in the midthoracic region associated with scoliosis. Posterior fusion defects were seen in the cervical spine, and there were multiple bifid ribs. CT head demonstrated enlarged ventricles and subarachnoid space.

Patient 4 (DECIPHER ID 267516)

Patient 4 is a white Scottish male from a non-consanguineous family. Bilateral talipes equinovarus was noted on antenatal ultrasound. He was delivered at 41 weeks gestation by elective cesarean section, with a birth weight of 3.77 kg (50th-75th centile). He had severe developmental delay. He first walked at five years of age, first words were at two years and six months of age, and he is not toilet trained.

On assessment at the age of eight years, he had brachycephaly, a flat face, frontal hair upsweep, highly arched eyebrows, hypertelorism, epicanthic folds, long eyelashes, wide nasal bridge, thin upper lip and everted lower lip. (Fig.1). His ears were low set and posteriorly rotated, and his tongue had a grooved tip. His head circumference was between the 25th and 50th centiles and height and weight both between the 50th and 75th centiles. He had three café au lait patches, and the anus was anteriorly placed. He had scoliosis and bilateral talipes equinovarus, and also primary generalized epilepsy. Radiographs demonstrated abnormal modelling and fusion of his ribs, predominantly on the right.

Molecular testing

Each patient was found to be homozygous for a c.292_293del p.(Ser98*) TMCO1 pathogenic variant (NM_019026.4 transcript). This variant has previously been reported in an Amish population (Xin et al., 2010). Microarray analysis for patients 1 & 4 was normal, and no additional variants were identified on exome sequencing.

Discussion

TMCO1 plays an important role in the maintenance of intracellular calcium homeostasis through dynamic interaction with the endoplasmic reticulum (ER). The ER is the principle reserve of intracellular calcium. TMCO1 forms a transmembrane channel to allow calcium release in the presence of excess ER stores. Tmco1 knockout mice demonstrate several of the features of CFTD patients, including intellectual disability, craniofacial dysmorphic features and abnormal bone formation (Wang et al., 2016).

Our patient series demonstrates the characteristic features of CFTD (Fig. 1), including distinctive craniofacial dysmorphic features (4/4; 100%), rib (3/4; 75%) and vertebral (2/4; 50%) abnormalities, and severe intellectual disability (4/4; 100%). The feature of an anteriorly placed anus is novel (although this was unconfirmed by measurement or specialty consultation in this case). Further reports are needed to determine if this is part of the phenotype. The phenotype of these patients has some overlap with Kleefstra syndrome, particularly in relation to the brachycephaly, synophrys, 'cupid's bow' upper lip and low set ears. Synophrys and 'cupid's bow' upper lip are also features of Coffin-Siris syndrome. Additionally, the dysmorphic

features, especially including synophrys, and neurodevelopmental phenotype have some similarity to Cornelia de Lange syndrome. These conditions should be considered in the differential, especially in the context of normal microarray.

Xin et al. (2010) reported a series of Amish patients, sharing the same p.(Ser98*) TMCO1 pathogenic variant as seen here. They have similar craniofacial dysmorphic features and a high prevalence of rib and/or vertebral anomalies. However, there are some notable differences. For example, all of the Amish patients had hypotonia and feeding problems; these were not seen in our series. In addition, Xin et al. (2010) reported gingival hyperplasia was a prominent feature whereas only 1 of 4 patients presented here have this. Overall, however, the two series share a strikingly similar phenotype.

Combining our series of four patients with those previously reported (Alanay et al., 2013; Caglayan et al., 2013; Pehlivan et al., 2014; Xin et al., 2010) allows further delineation of the phenotypic spectrum of this rare condition (Table 1). The majority of patients (22/23; 96%) were from consanguineous families. Neonatal hypotonia and poor feeding were seen in 16/21 (76%) patients. All patients have global developmental delay, with 4/18 (22%) non-verbal and 9/20 (45%) not able to walk. All patients have severe intellectual disability.

Common dysmorphic features include macrocephaly (12/21; 57%), brachycephaly (19/23; 83%), flat face (19/23; 83%), arched eyebrows (20/23; 87%), synophrys (22/23; 96%), hypertelorism (21/23; 91%), long eyelashes (16/21; 76%), wide nasal bridge (18/23; 78%), low set ears (23/23; 100%), microdontism (13/18; 72%) and

gingival hyperplasia (11/15; 73%). Affable behavior is often present (4/8; 50%). Rib (17/22; 77%) and vertebral anomalies (16/23; 70%) are common radiographic findings. Bilateral bifid and fused ribs are frequently seen, as well as scoliosis and spinal fusion.

As discussed, the p.(Ser98*) TMCO1 pathogenic variant has previously been found only in a multiply consanguineous Amish community (Xin et al., 2010). This is therefore the first report of the variant outside this population. The variant is predicted to result in nonsense-mediated decay. Interestingly, the brothers reported here and the additional unrelated patient share the same variant, indicating this is a recurrent variant in TMCO1 and may represent a mutational hotspot. This variant has a small but appreciable allele frequency of 0.0001974 in Non-Finnish Europeans, according to gnomAD data (Lek et al., 2016).

In conclusion, we present four patients with CFTD due to a recurrent TMCO1 pathogenic variant, expanding the genotypic spectrum and further delineating the associated phenotype.

Acknowledgements

We would like to thank the families involved for their consent to publish this work.

The DDD study presents independent research commissioned by the Health Innovation Challenge Fund [grant number HICF-1009-003], a parallel funding partnership between the Wellcome Trust and the Department of Health, and the Wellcome Trust Sanger Institute [grant number WT098051]. The views expressed in this publication are those of the author(s) and not necessarily those of the Wellcome Trust or the Department of Health. The study has UK Research Ethics Committee approval (10/H0305/83, granted by the Cambridge South REC, and GEN/284/12 granted by the Republic of Ireland REC). The research team acknowledges the support of the National Institute for Health Research, through the Comprehensive Clinical Research Network.

Conflict of Interest

The authors do not have any conflict of interest to disclose.

Consent

Informed consent was obtained for all subjects for inclusion in this study.

References

- Alanay, Y., Ergüner, B., Utine, E., Haçariz, O., Kiper, P.O., Taşkıran, E.Z., ...
Akarsu, N.A. (2013). TMCO1 Deficiency Causes Autosomal Recessive Cerebrofaciothoracic Dysplasia. *American Journal of Medical Genetics Part A*, 164A, 291-304. doi: 10.1002/ajmg.a.36248
- Caglayan, A.O., Per, H., Akgumus, G., Gumus, H., Baranoski, J., Canpolat, M., ...
Gunel, M. (2013). Whole-exome sequencing identified a patient with TMCO1 defect syndrome and expands the phenotic spectrum. *Clinical Genetics*, 84, 394-395. doi: 10.1111/cge.12088
- Lek, M., Karczewski, K.J., Minikel, E.V., Samocha, K.E., Banks, E., Fennell, T., ...
Exome Aggregation Consortium. (2016). Analysis of protein-coding genetic variation in 60,706 humans. *Nature*, 536, 285-291. doi: 10.1038/nature19057
- Pehlivan, D., Karaca, E., Aydin, H., Beck, C.R., Gambin, T., Muzny, D.M., ... Lupski, J.R. (2014). Whole-exome sequencing links TMCO1 defect syndrome with cerebro-facio-thoracic dysplasia. *European Journal of Human Genetics*, 22, 1145-1148. doi: 10.1038/ejhg.2013.291
- Wang, Q-C., Zheng, Q., Tan, H., Zhang, B., Li, X., Yang, Y., ... Tang, T.S. (2016). TMCO1 is an ER Ca²⁺ Load-Activated Ca²⁺ Channel. *Cell*, 165, 1454-1466. doi: 10.1016/j.cell.2016.04.051
- Wright, C.F., Fitzgerald, T.W., Jones, W.D., Clayton, S., McRae, J.F., van Kogelenberg, M., ... DDD study. (2015). Genetic diagnosis of developmental disorders in the DDD study: a scalable analysis of genome-wide research data. *Lancet*, 385, 1305-14. doi: 10.1016/S0140-6736(14)61705-0

Xin, B., Puffenberger, E.G., Turben, S., Tan, H., Zhou, A., Wang, H. (2010).

Homozygous frameshift mutation in TMCO1 causes a syndrome with
craniofacial dysmorphism, skeletal anomalies, and mental retardation.

Proceedings of the National Academy of Sciences of the United States of
America, 107, 258-263. doi: 10.1073/pnas.0908457107

Titles and legends to figures

Figure 1. Patient photographs (age in years). Patient 1 a-c (13), Patient 2 d-g (29), Patient 3 h-i (26), Patient 4 j-k (1), l-o (8). Note characteristic craniofacial appearance including synophrys and arched eyebrows. Patients 1 and 2 have hypermobility of the fingers. Patient 2 also has a Sprengel deformity.

Figure 2.

PA chest radiograph patient 2, age 25 years. Note likely dorsal scoliosis concave to the right. The left 4th and 5th ribs are fused posteriorly, giving the appearance of a single broad rib (arrows). The posterior end of the left 6th rib is unossified (triangle). There is a bony bridge leading to posterior fusion of the left 7th and 8th ribs (*)