University of Dundee

Preoperative prediction of margin requirement following a core biopsy result suggestive of a phyllodes tumour

Mberu, V.; Macaskill, E. J.; Purdie, C.; Evans, A.

Published in:
Clinical Radiology

DOI:
[10.1016/j.crad.2019.11.006](https://doi.org/10.1016/j.crad.2019.11.006)

Publication date:
2020

Licence:
CC BY-NC-ND

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Mberu, V., Macaskill, E. J., Purdie, C., & Evans, A. (2020). Preoperative prediction of margin requirement following a core biopsy result suggestive of a phyllodes tumour. *Clinical Radiology*, 75(4), 319.e21-319.e27. <https://doi.org/10.1016/j.crad.2019.11.006>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

© 2019. This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

1 **Keywords**

2 Phyllodes tumour, fibroadenoma, surgical excision, margin

3 **Introduction**

4 Phyllodes tumour is a rare fibroepithelial breast tumour which arises from the periductal
5 stroma of the breast, it accounts for less than 1% of all primary breast tumours. (1,2) They
6 can occur at any age but are most frequent in the fourth decade with earlier onset in those of
7 Asian origin.(3,4) Phyllodes tumours can be divided into benign, borderline and malignant
8 subtypes, depending on histological parameters such as nature of tumour borders, degree of
9 stromal cellularity and atypia, mitotic count and stromal overgrowth. (5,6) Fibroadenoma is a
10 very common fibroepithelial tumour which shares some characteristics with phyllodes
11 tumour, but has a younger age distribution.(1) Differentiating between phyllodes and
12 fibroadenoma tumours is difficult as clinical, radiological and histopathologic appearance
13 may mimic each other with a definitive diagnosis only being made after microscopic analysis
14 of the entire excised lesion.(1) Despite their similarities, fibroadenomas are benign and are
15 managed either conservatively or by enucleation without surgical margins while phyllodes
16 tumours require excision with clear margins to avoid recurrence particularly in the borderline
17 and malignant subtypes. (7–11) Accurate identification pre-operatively is critical for
18 appropriate surgical planning to avoid complications from overtreatment or inadequate
19 excision.(10)

20 Imaging modalities used in the assessment of breast disease include mammography,
21 ultrasound and Magnetic Resonance Imaging (MRI). Ultrasound elastography is the most
22 recent imaging modality employed in differentiating between malignant and benign lesions
23 by assessing the stiffness within lesions which has proved an accessory tool in assessing
24 breast masses.(1) Studies evaluating the use of MRI suggests that both phyllodes and
25 fibroadenoma show similar findings.(12) Immunohistochemical markers have been

suggested to increase accuracy in distinguishing between fibroepithelial tumours but histology remains the gold standard.(8,13)

Recent guidelines suggest benign phyllodes tumours do not need excision with margins but borderline and malignant phyllodes require excision with a clear margin to reduce the risk of local recurrence and potential distant metastases.(14) However, core biopsy cannot reliably differentiate between fibroadenomas and phyllodes tumour subtypes, which can be explained pathologically due their heterogeneity.(15) As such, it can be difficult for the surgeon to predict whether a margin is required or not from the preoperative information. If a more accurate prediction of the likely nature of the lesion as fibroadenoma or benign phyllodes versus borderline or malignant phyllodes were available, this could reduce unnecessary margin excision and cosmetic compromise, or avoid second operations to achieve clear margins.

Previous studies have compared the imaging features of fibroadenomas and phyllodes tumours (1,16,17) but none has compared the imaging features of those lesions requiring excision with a margin and those which do not.(18) In this study, we aimed to identify ultrasound and mammographic features associated with borderline and malignant phyllodes versus benign phyllodes and fibroadenomas to aid surgical planning.

Methodology

A prospective database of consecutive ultrasound visible masses was used to identify lesions with a core biopsy suggestive of a phyllodes tumour in a single unit between May 2010 and January 2019. Masses with an ultrasound core biopsy result of B3, raising the possibility of a phyllodes tumour were included.

The breast ultrasound features assessed were: mass shape and size; orientation; margin definition; presence of macro-lobulations and micro-lobulations; echogenicity; cystic spaces; echogenic clefts; skin involvement; surrounding oedema; Breast Imaging Reporting And Data System (BIRADS) score(19); posterior features and vascularity. BIRADS score allows

for unambiguous reporting of breast imaging and is graded from incomplete to proven malignancy, with scores of 0 to 6 respectively. For this study, only lesions with a BIRADS score of at least 3 were included.

Mammographic features assessed included presence of a mass; calcification; margin definition and BIRADS score. Breast density was assessed using the BIRADS classification(19) and a Visual Analogue Scale (VAS). (20) Breast density or composition, was assessed in terms of the proportion of fibroglandular tissue with the least dense breast graded as BIRADS density 'a' and the most dense graded as BIRADS density 'd'. In addition, a 10cm visual analogue scale presented on paper sheets was used, with the approximate percentage of dense breast tissue marked with the right side of line marking 100% density and the left marking 0% density. Inter-rater agreement was achieved via discussion.

Excision surgical pathology was recorded, and lesions were classified, using definitions described earlier (5,6) as a benign phyllodes/ fibroadenoma or as a borderline/malignant phyllodes if a margin was required. Ultrasound and mammographic features in these 2 groups were compared. Statistical analysis used Chi-square, Fisher's exact test and receiver-operating curve (ROC).

Attention was paid to features previously described to be suggestive of a phyllodes tumour. Imaging features were carried out by an experienced consultant radiologist who was blinded to the final pathology outcomes. For our study, benign phyllodes and fibroadenoma were grouped as not needing a margin whereas borderline and malignant phyllodes were grouped as needing a margin.(8,14)

Results

31 patients with 31 lesions meeting the inclusion criteria were identified. Of these, 6 lesions were screen-detected, all of which were benign, and 25 were symptomatic. There were 13 lesions requiring a margin (6 malignant, 7 borderline) and 18 benign lesions not requiring a

margin (13 benign phyllodes, 4 fibroadenoma and 1 other). The average age of study patients was 53 years old, (range 22- 84 years). 25 patients had mammograms performed, with the lesion being mammographically occult in 3, leaving 22 to be included in analysis. All 31 patients had a breast ultrasound.

Table 1. Categorical ultrasound features assessed in borderline/malignant phyllodes and fibroadenoma/benign phyllodes.

Table 2. Categorical mammographic features assessed in borderline/malignant phyllodes and fibroadenoma/benign phyllodes.

The following ultrasound features were found significantly more frequently in those lesions that were borderline/malignant phyllodes as shown in table 1; an irregular margin [8/13 (62%) vs 3/18 (17%) $p=0.01$], presence of micro-lobulations [7/13 (54%) vs 3/18 (17%) $p=0.028$], mixed echogenicity [9/13 (69%) vs 1/18 (6%) $p=0.0002$], echogenic clefts [6/13 (46%) vs 1/18 (6%) $p=0.007$], BIRADS score of more than 3 [11/13 (85%) vs 9/18 (50%) $p=0.047$], posterior enhancement (9/11 (82%) vs 6/18 (33%) $p=0.01$].

Large ultrasound size was significantly associated with borderline and malignant phyllodes tumours with an area under the curve (AUC) of 0.76, $p=0.003$ as shown in figure 1.

Stiffness at shock wave elastography (SWE) was also associated with borderline and malignant phyllodes, AUC 0.71, $p=0.026$ as shown in figure 2.

On mammography, fibroadenoma and benign phyllodes tumours had well-defined margins compared to the borderline/ malignant Phyllodes tumours [7/9 (78%) vs 4/13 (31%), $p=0.04$] as shown in table 2. No other mammographic features were statistically significant.

Mode of presentation was a significant factor with symptomatic lesions being more likely to require a margin than screen-detected lesions [13/13 (100%) vs 6/18 (33%) $p=0.005$], respectively.

Discussion

In this study, we have identified multiple pre-operative features that are significantly different between lesions requiring a margin and those that do not. In this study we grouped benign phyllodes and fibroadenoma as lesions not needing a margin and borderline and malignant phyllodes as lesions needing a margin. A previous study found that although differentiating between fibroadenomas and phyllodes tumours is difficult even for pathologists specialised in breast pathology, there is an inter-rater agreement when fibroadenomas and benign phyllodes are distinguished from malignant and borderline subtypes. (8)

This study shows that there are a number of pre-operative differentiating features between fibroadenoma and benign phyllodes versus borderline and malignant phyllodes found on ultrasound. The features suggesting borderline and malignant phyllodes, and thus the need for a surgical excision with margin include an irregular shape, micro-lobulations, high echogenicity, BIRADS score > 3, distal enhancement, large size and stiffness on SWE. In addition, a mammographic poorly-defined margin and symptomatic presentation suggest the need for a margin at excision.

We found increasing size to be a significant factor in predicting the need for a surgical margin. This is in agreement with a previous study which found that mean lesion size increased when comparing benign, borderline and malignant phyllodes tumours. (18) However, this study did not include any fibroadenomas.

As the number of differentiating factors shown is high, a multivariate analysis to find those with independent significance would be helpful. Unfortunately, our dataset is too small for multivariate analysis to be performed.

The lack of a radiological, well-defined margin in borderline and malignant phyllodes mirrors the pathological findings of infiltrative margins and stromal overgrowth in such tumours. (21)

The posterior enhancement may reflect the desmoplastic growth of tumour stromal cells in

borderline and malignant lesions as it is known that posterior enhancement is found more frequently in high grade invasive cancers than low grade. (22)

Micro-lobulations are probably a reflection of an infiltrative margin as they are common features of an invasive cancer but uncommon in fibroadenomas. When found in fibroadenomas, they are thought to be due to hyalinisation or infarction leading to fibrous changes that increase the stromal component.(23) The high ultrasound BIRADS score in the borderline and malignant phyllodes is likely to reflect both infiltrative margins and the increased heterogeneity seen in these lesions.

Increased stiffness in invasive breast cancer has been shown to reflect active tumour stromal interaction.(24) Although tumour stromal interaction in malignant phyllodes is not well understood, it is possible that increased stiffness at shear wave elastography reflects activated tumour associated fibroblasts and the production of stiff collagen.(25,26) Stiffness on shear wave elastography may also reflect the increased cellularity of the stroma in phyllodes tumour compared to fibroadenoma.(1)

Although the number of screening-detected lesions in the study is small, it is striking that they were all benign phyllodes or fibroadenomas. The reason for this is not clear but may reflect their small size and impalpability.

Although peri-lesional oedema was not found to be a significant feature in our study, some studies(16,27) have found that rapidly growing tumours have surrounding interstitial oedema due to the compressed lymphatics or mammary ducts best seen on T2-weighted MRI images. Patients in this study did not routinely undergo MRI and, as such, this could not be studied in this cohort.

The small sample size from a single centre is the largest limitation in this study. Further studies should aim to include more cases from multiple centres. A larger sample size would enable multivariate analysis to identify independent factors. In addition, there were not enough MRI scans for this modality to be evaluated.

Conclusion

We have identified multiple ultrasound and mammographic features that may be used to guide surgeons' decisions regarding the use of a margin when excising lesions suggestive of a phyllodes tumour. Due to our small sample size, further studies involving larger numbers are required to validate our results.

References

1. Li L-J, Zeng H, Ou B, Luo B-M, Xiao X-Y, Zhong W-J, et al. Ultrasonic Elastography Features of Phyllodes Tumors of the Breast: A Clinical Research. Coleman WB, editor. PLoS One [Internet]. 2014 Jan 15 [cited 2019 Jun 10];9(1):e85257. Available from: <http://dx.plos.org/10.1371/journal.pone.0085257>
2. Rowell MD, Perry RR, Hsiu J, Virginia N, Barranco SC, Beach V. Phyllodes Tumors. 1993;165(March):376–9.
3. Venter AC, Roşca E, Daina LG, Muţiu G, Pirte AN, Rahotă D. Phyllodes tumor: Diagnostic imaging and histopathology findings. Rom J Morphol Embryol. 2015;
4. McCarthy E, Kavanagh J, O'Donoghue Y, McCormack E, D'Arcy C, O'Keeffe SA. Phyllodes tumours of the breast: Radiological presentation, management and followup. Br J Radiol. 2014;
5. Tan BY, Acs G, Apple SK, Badve S, Bleiweiss IJ, Brogi E, et al. Phyllodes tumours of the breast: A consensus review. Histopathology. 2016.
6. Guidelines Working Group of the UK National Coordinating Committee for Breast Pathology. Pathology reporting of breast disease in surgical excision specimens incorporating the dataset for histological reporting of breast cancer. R Coll Pathol. 2016;v1(6):160.
7. Jacobs TW, Chen YY, Guinee DG, Holden JA, Cha I, Bauermeister DE, et al. Fibroepithelial lesions with cellular stroma on breast core needle biopsy: Are there

178 predictors of outcome on surgical excision? Am J Clin Pathol. 2005;

179 8. Lawton TJ, Acs G, Argani P, Farshid G, Gilcrease M, Goldstein N, et al. Interobserver
180 variability by pathologists in the distinction between cellular fibroadenomas and
181 phyllodes tumors. Int J Surg Pathol. 2014;

182 9. Tan EY, Hoon TP, Yong WS, Wong HB, Hui HG, Yeo AWY, et al. Recurrent phyllodes
183 tumours of the breast: Pathological features and clinical implications. ANZ J Surg.
184 2006;

185 10. Stoffel E, Becker AS, Wurnig MC, Marcon M, Ghafoor S, Berger N, et al. Distinction
186 between phyllodes tumor and fibroadenoma in breast ultrasound using deep learning
187 image analysis. Eur J Radiol Open. 2018;

188 11. Lu Y, Chen Y, Zhu L, Cartwright P, Song E, Jacobs L, et al. Local Recurrence of
189 Benign, Borderline, and Malignant Phyllodes Tumors of the Breast: A Systematic
190 Review and Meta-analysis. Annals of Surgical Oncology. 2019.

191 12. Kamitani T, Matsuo Y, Yabuuchi H, Fujita N, Nagao M, Kawanami S, et al.
192 Differentiation between benign phyllodes tumors and fibroadenomas of the breast on
193 MR imaging. Eur J Radiol. 2014;

194 13. Yang X, Kandil D, Cosar EF, Khan A. Fibroepithelial tumors of the breast: Pathologic
195 and immunohistochemical features and molecular mechanisms. Arch Pathol Lab Med.
196 2014;

197 14. Rageth CJ, O'Flynn EAM, Pinker K, Kubik-Huch RA, Mundinger A, Decker T, et al.
198 Second International Consensus Conference on lesions of uncertain malignant
199 potential in the breast (B3 lesions). Breast Cancer Research and Treatment. 2019.

200 15. Youn I, Choi SH, Moon HJ, Kim MJ, Kim EK. Phyllodes Tumors of the Breast:
201 Ultrasonographic Findings and Diagnostic Performance of Ultrasound-Guided Core
202 Needle Biopsy. Ultrasound Med Biol. 2013;

- 203 16. Duman L, Gezer NS, Balci P, Altay C, Başara I, Durak MG, et al. Differentiation
204 between Phyllodes Tumors and Fibroadenomas Based on Mammographic
205 Sonographic and MRI Features. Breast Care. 2016;
- 206 17. Kim GR, Choi JS, Han BK, Ko EY, Ko ES, Hahn SY. Combination of shear-wave
207 elastography and color Doppler: Feasible method to avoid unnecessary breast
208 excision of fibroepithelial lesions diagnosed by core needle biopsy. PLoS One. 2017;
- 209 18. Yii N, Read T, Tan CC, Ng SL, Bennett I. Diagnosing phyllodes tumours of the breast:
210 how successful are our current preoperative assessment modalities? ANZ J Surg.
211 2018;
- 212 19. D'Orsi CJ, Sickles EA, Mendelson EB, Morris EA et al. ACR BI-RADS® Atlas, Breast
213 Imaging Reporting and Data System. 5th Editio. Reston, VA, American College of
214 Radiology; 2013.
- 215 20. Morrish OWE, Tucker L, Black R, Willsher P, Duffy SW, Gilbert FJ. Mammographic
216 Breast Density: Comparison of Methods for Quantitative Evaluation. Radiology. 2015;
- 217 21. Taira N, Takabatake D, Aogi K, Ohsumi S, Takashima S, Nishimura R, et al.
218 Phyllodes tumor of the breast: Stromal overgrowth and histological classification are
219 useful prognosis-predictive factors for local recurrence in patients with a positive
220 surgical margin. Jpn J Clin Oncol. 2007;
- 221 22. Irshad A, Leddy R, Pisano E, Baker N, Lewis M, Ackerman S, et al. Assessing the
222 role of ultrasound in predicting the biological behavior of breast cancer. American
223 Journal of Roentgenology. 2013.
- 224 23. Thomassin-Piana J. Solid masses: what are the underlying histopathological lesions?
225 Diagn Interv Imaging. 2014;
- 226 24. Mei Rosa N, Joan SB. A stiff blow from the stroma: collagen crosslinking drives tumor
227 progression. Cancer Cell. 2009;

25. Shi XQ, Li J, Qian L, Xue X, Li J, Wan W. Correlation between elastic parameters and collagen fibre features in breast lesions. Clin Radiol. 2018;
26. Nie Y, Chen J, Huang D, Yao Y, Chen J, Ding L, et al. Tumor-associated macrophages promote malignant progression of breast phyllodes tumors by inducing myofibroblast differentiation. Cancer Res. 2017;
27. Wurdinger S, Herzog AB, Fischer DR, Marx C, Raabe G, Schneider A, et al. Differentiation of phyllodes breast tumors from fibroadenomas on MRI. Am J Roentgenol. 2005;

Figure 1: Receiver Operating Curve showing the association between ultrasound size of the lesion and borderline and malignant phyllodes tumour subtypes.

Figure 2: Receiver operating curve (ROC) showing the association between stiffness of the lesion and borderline and malignant phyllodes tumour subtypes.

Table 1: Categorical ultrasound features assessed in borderline/malignant phyllodes and fibroadenoma/benign phyllodes.

Lesion characteristics	Borderline/malignant phyllodes (%)	Fibroadenoma/benign phyllodes (%)	p-value
Mass shape			
Oval and round	5/13 (38)	15/18 (83)	0.01
Irregular	8/13 (62)	3/18 (17)	

Orientation ¹			
<i>Taller than wide</i>	0	0	>0.05
<i>Not taller than wide</i>	12/12 (100)	18/18 (100)	
Margin			
<i>Well-defined</i>	6/13 (46)	12/18 (67)	>0.05
<i>Poorly defined, mixed</i>	7/13 (54)	6/18 (33)	
Micro-lobulations			
<i>Yes</i>	7/13 (54)	3/18 (17)	0.028
<i>No</i>	6/13 (46)	15/18 (83)	
Macro-lobulations			
<i>Yes</i>	9/13 (69)	7/18 (39)	>0.05
<i>No</i>	4/13 (31)	11/18 (61)	
Echogenicity			
<i>Hypodense</i>	4/13 (31)	17/18 (94)	
<i>Mixed</i>	9/13 (69)	1/18 (6)	0.002
Cystic spaces			
<i>Yes</i>	5/13 (38)	2/18 (11)	>0.05
<i>No</i>	8/13 (62)	16/18 (89)	
Echogenic clefts			

¹ One lesion could not be assessed as it was too large to accurately determine the orientation.

Yes	6/13 (46)	1/18 (6)	0.007
No	7/13 (54)	17/18 (94)	
Skin changes			
Yes	2/13 (15)	1/18 (6)	> 0.05
No	11/13 (85)	17/18 (94)	
Bright surrounding fat/oedema ²			
Yes	1/12 (8)	0	>0.05
No	11/12 (92)	18/18 (100)	
BIRADS score			
3	2/13 (15)	9/18 (50)	0.047
4a, 4b, 4c and 5	11/13 (85)	9/18 (50)	
Posterior features			
None, shadowing	2/11 (18)	12/18(67)	0.01
Enhancement	9/11 (82)	6/18 (33)	
Vascularity			
Penetrating vessels ³			
Yes	9/11 (82)	6/9 (67)	>0.05
No	2/11 (18)	3/9 (33)	

² One lesion was unassessable.

³ Two lesions were unassessable.

Surrounding vessels ⁴			>0.05
Yes	7/9 (78)	7/9 (78)	
No	2/9 (22)	2/9 (22)	
Laterality			
Left	7/13 (54)	7/18 (39)	>0.05
Right	6/13 (46)	11/18 (61)	

245

246 Table 2: *Categorical mammographic features assessed in borderline/malignant phyllodes*
247 *and fibroadenoma/benign phyllodes.*

Lesion characteristics	Borderline/malignant phyllodes (%)	Fibroadenoma/benign phyllodes (%)	p-value
Presence of mass			
Yes	9/10 (90)	13/15 (87)	>0.05
No	1/10 (10)	2/15 (13)	
Calcification			
Yes	3/10 (30)	3/15 (20)	>0.05
No	7/10 (70)	12/15 (80)	
Margin ⁵			
Well defined	2/9 (22)	9/13 (69)	0.04
	7/9 (78)	4/13 (31)	

⁴ Two lesions were unassessable.

⁵ 3 lesions were mammographically occult.

<i>Poorly defined, mixed</i>			
BIRADS density			
<i>a-b</i>	3/10 (30)	7/15 (47)	>0.05
<i>c-d</i>	7/10 (70)	8/15 (53)	
BIRADS score			
<i>1-3</i>	5/10 (50)	12/15 (80)	>0.05
<i>4</i>	5/10 (50)	3/15 (20)	

¹ 3 lesions were mammographically occult.

248
249