University of Dundee

Feasibility of a trial to evaluate nicotinamide for chemoprevention of skin cancers in organ transplant recipients in the UK

Gollins, C. E.; Shah, A.; Sinha, K.; Khan, S.; Paul, N.; Meeajun, B.

Published in:
British Journal of Dermatology

DOI:
[10.1111/bjd.18982](https://doi.org/10.1111/bjd.18982)

Publication date:
2020

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Gollins, C. E., Shah, A., Sinha, K., Khan, S., Paul, N., Meeajun, B., Abbott, R. A., Blasdale, C., Cooper, H., Harwood, C. A., Ismail, F., Lear, J. T., Mackintosh, L., McCormack, S., Perrett, C. M., Proby, C. M., Durack, A., Patalay, R., & Matin, R. N. (2020). Feasibility of a trial to evaluate nicotinamide for chemoprevention of skin cancers in organ transplant recipients in the UK. *British Journal of Dermatology*, 183(2), 394-396. <https://doi.org/10.1111/bjd.18982>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

DR CHARLOTTE ELIZABETH GOLLINS (Orcid ID : 0000-0002-7539-4265)

DR AADARSH SHAH (Orcid ID : 0000-0003-3203-6975)

DR KHUSHBOO SINHA (Orcid ID : 0000-0003-2591-2260)

DR CATHERINE ANNE HARWOOD (Orcid ID : 0000-0002-1375-0965)

DR CHARLOTTE M PROBY (Orcid ID : 0000-0002-3292-4836)

Article type : Research Letter

Feasibility of a trial to evaluate nicotinamide for chemoprevention of skin cancers in organ transplant recipients in the UK

C. Gollins,¹ A. Shah,² K. Sinha,² S. Khan,³ N. Paul,⁴ B. Meeajun,⁵ R. A. Abbott,⁶ C. Blasdale,⁷ H. Cooper,⁸ C. A. Harwood,⁹ F. Ismail,¹⁰ J. T. Lear,¹¹ L. Mackintosh,¹² S. McCormack,¹³ C. M. Perrett,¹⁴ C. M. Proby,¹⁵ A. Durack,¹⁶ R. Patalay¹⁷ and R. N. Martin³

Affiliations:

1. Dermatology Department, St George's Hospital, St George's University Hospitals NHS Foundation Trust, UK
2. Dermatology Department, Basildon University Hospital, Basildon and Thurrock Hospitals NHS Foundation Trust, UK
3. Department of Dermatology, Churchill Hospital, Oxford University Hospitals NHS Foundation Trust, UK
4. Department of Dermatology, St Mary's Hospital, Imperial College Healthcare NHS Trust, UK
5. Department of Dermatology, Whittington Hospital, Whittington Health NHS Trust, UK
6. Department of Dermatology, University Hospital of Wales, Cardiff and Vale University Health Board, UK
7. Department of Dermatology, Royal Victoria Infirmary, The Newcastle upon Tyne Hospitals NHS Foundation Trust, UK
8. Dermatology Department, Queen Alexandra Hospital, Portsmouth Hospitals NHS Trust, UK

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/BJD.18982](#). This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

This article is protected by copyright. All rights reserved

9. Centre for Cell Biology and Cutaneous Research, Blizard Institute, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, UK
10. Department of Dermatology, Royal Free London NHS Foundation Trust, UK
11. Department of Dermatology, Manchester Royal Infirmary, Manchester University NHS Foundation Trust, UK
12. Department of Dermatology, NHS Greater Glasgow and Clyde, UK
13. Department of Dermatology, Queen Margaret Hospital, NHS Fife, UK
14. Department of Dermatology, University College Hospital, University College London Hospitals NHS Foundation Trust, UK
15. Department of Dermatology, Ninewells Hospital and Medical School, University of Dundee, UK
16. Department of Dermatology, Cambridge University Hospitals NHS Foundation Trust, UK
17. St John's Institute of Dermatology, Guy's and St Thomas' NHS Foundation Trust, UK

Corresponding author: Dr Charlotte Gollins

Email: charlotteelizabeth.gollins@nhs.net

Word count: 746

Figure Count: 0

There are no declarations of financial support or relationships that may pose a conflict of interest.

Dear Editor,

Keratinocyte cancers (KC) are the commonest malignancies in the UK population, with rates increasing annually.¹ Nicotinamide (vitamin B3) has recently been reported to reduce the incidence of KC in high-risk patients.² It is a precursor of nicotinamide adenine dinucleotide (NAD⁺) which replenishes cellular energy levels and enhances efficient repair of ultraviolet light (UV) induced DNA damage within keratinocytes.³ Solid organ transplant recipients (OTRs) are at particularly high risk of KC, with 150 times increased risk of cutaneous squamous cell carcinoma (cSCC) compared with the immunocompetent general population.⁴ Recent randomised controlled trial (RCT) data from Australia suggest that oral nicotinamide reduces cSCC by approximately 30% over a 12-month period in immunocompetent individuals with a past history of KC.²

Although trials investigating the role of nicotinamide in OTRs are planned or underway in Australia and Canada (<https://www.anzctr.org.au/Trial/Registration/TrialReview.aspx?id=372709>; <https://clinicaltrials.gov/ct2/show/NCT03769285>), to date, there have been no trials evaluating the chemopreventative role of nicotinamide in a UK OTR population.

We propose a randomised controlled trial (RCT) comparing nicotinamide versus placebo in risk reduction of cSCC in high-risk OTRs in the UK. To explore feasibility of undertaking this RCT we assessed willingness of UK clinicians and OTRs to participate and their potential concerns. An electronic Survey Monkey™ questionnaire was sent to Dermatologists and Transplant Units (August 2018 – April 2019). Anonymised paper-based surveys were self-completed by OTRs when attending dedicated Transplant Dermatology Clinics (January 2019 – May 2019). Data were entered and analysed using Microsoft Excel.

Clinician survey: 99 clinicians (1 male:1.4 female) from 40 UK National Health Service (NHS) Trusts (in England, Scotland, Wales and Northern Ireland) responded (79% consultants: 87% dermatologists, 12% transplant physicians; response rate approximately 10%). 38% of respondents reviewed an average of 1-2 OTR/week, with 50% reviewing > 2 OTRs/week. 67 of 99 replied to questions on systemic chemoprevention. 49 of 67 (73%) respondents had initiated systemic chemoprevention, with acitretin the most frequent first-line choice (40/67, 60%); 24% (16/67) had used nicotinamide previously. 72 respondents replied to the question of whether they would be willing to recruit patients to this study; of these 59 (82%) would be willing. Specific concerns included optimising the definition of high-risk OTRs in order to maximise event rate.

OTR survey: 271 of 301 OTRs from 11 NHS Trusts (England, Scotland and Wales) responded (response rate 90%), 1 male:1.6 female; mean age 57 years (y), range 18 – 83y; mean time from first transplant 14.4 y (range 0 – 51y). The majority were renal transplant recipients (65%) with 14% lung, 12% heart, 4% liver, 1% pancreas and 0.4% bowel transplant recipients. 66 (24%) reported a prior cSCC history, with 50% of this cohort (n=33) reporting > 5 KC. 6% of respondents had previously received systemic chemoprevention: 4% (10/271) with acitretin and 3% (7/271) nicotinamide. Topical chemoprevention used at least once included 5% 5-fluorouracil cream (15%), 5% imiquimod cream (2%) and diclofenac in hyaluronic acid gel (1%). Overall, 71% (193/271) were willing to participate in the proposed trial. For patients with a past history of cSCC, and therefore eligible for recruitment to the proposed RCT, this increased to 85% (56/66). Specific concerns raised included the need for additional hospital visits and blood tests and potential interactions between nicotinamide and regular medications.

Skin cancers in OTRs are more frequent and aggressive, with a greater risk of tumour recurrence, local invasion and metastasis than in the immunocompetent population.^{4, 5} This feasibility survey has confirmed that the systemic chemoprevention strategy used most frequently in the UK is oral acitretin, a drug with significant adverse effects, and that current use of nicotinamide is limited. It is important that chemoprevention is safe, inexpensive and has few side effects. Nicotinamide fulfils these criteria and, if effective in OTRs, has the potential to be a more appropriate first line chemopreventative agent than

acitretin, but could also be used in conjunction with acitretin in the highest-risk patients. These feasibility surveys demonstrated support for a UK-based trial amongst both clinicians and in OTRs. Given the low-cost and low toxicity of oral nicotinamide, confirmation of its efficacy in a UK OTR population has the potential to transform chemopreventative practice in this growing high-risk patient population.

Acknowledgements: We are grateful to Margaret McPhee (UK Dermatology Clinical Trials Network) for administrative support and Dr Alexa Shipman for collection of patient surveys. This project was developed with support from the UK Dermatology Clinical Trials Network (UK DCTN). The UK DCTN is grateful to the British Association of Dermatologists and the University of Nottingham for financial support of the Network.

References:

1. Venables ZC, Nijsten T, Wong KF et al. Epidemiology of basal and cutaneous squamous cell carcinoma in the UK. 2013-2015: a cohort study. *Br J Dermatol*. 2019; 181: 474-482
2. Chen AC, Martin AJ, Choy B et al. A Phase 3 Randomized Trial of Nicotinamide for Skin-Cancer Chemoprevention. *N Engl J Med*. 2015; 373(17): 1618-1626
3. Surjana D, Halliday GM, Damian DL. Nicotinamide enhances repair of ultraviolet radiation-induced DNA damage in human keratinocytes and ex vivo skin. *Carcinogenesis*. 2013; 34 (5): 1144-1149
4. Harwood CA, Mesher D, McGregor JM et al. A Surveillance Model for Skin Cancer in Organ Transplant Recipients: A 22-Year Prospective Study in an Ethnically Diverse Population. *Am J Transplant*. 2013; 13: 119-129
5. Ponticelli C, Cucchiari D, Bencini P. Skin cancer in kidney transplant recipients 2014. *J Nephrol*; 27: 385-394