University of Dundee

Albrecht Nagel on vision with two eyes

Wade, Nicholas J.

Published in:
Strabismus

DOI:
[10.1080/09273972.2020.1802181](https://doi.org/10.1080/09273972.2020.1802181)

Publication date:
2020

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Wade, N. J. (2020). Albrecht Nagel on vision with two eyes. *Strabismus*, 28(3), 115-118.
<https://doi.org/10.1080/09273972.2020.1802181>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Perspective

Albrecht Nagel on vision with two eyes

Nicholas J. Wade

Psychology, University of Dundee, Dundee DD1 4HN, UK

Email: n.j.wade@dundee.ac.uk

Abstract

Nagel's book on vision with two eyes was published in 1861, during a period in which German visual scientists were struggling to rescue the doctrine of identical retinal points from the evidence of stereoscopic depth. The long observational history of binocular vision has been dominated by the appearance of a single world with two eyes and its breakdown when the eyes are distorted abnormally. Early in the nineteenth century the flat horopter of Aguilonius (proposed two centuries earlier) assumed curvature in the form of the Vieth-Müller circle which was linked to identical retinal points: there were only two possible states of binocular perception – singleness with images on the Vieth-Müller circle and doubleness otherwise. This elegant edifice was undermined when Wheatstone demonstrated singleness and depth from images with slight retinal disparities. Nagel responded by providing observations on combining simple line stimuli in the two eyes. In the last part of chapter 3 of his book, Nagel describes experiments with lines varying in orientation or curvature with respect to the two eyes; it is in this section that Nagel draws attention to cyclofusion and the involvement of the extraocular muscles in it. Ocular torsion was an issue of considerable contention in nineteenth century visual science. The possibility of torsion in opposite directions seemed fanciful and yet this is what Nagel proposed in order to maintain cyclofusion for lines inclined in opposite directions relative to the horizontal. Similar rotations about the vertical resulted in a depth effect with no cyclovergence. The involvement of cyclovergence remained hotly debated until photographic recording of eye movements verified it.

Albrecht Nagel on vision with two eyes

Albrecht Nagel (1833-1895) published his book on *Das Sehen mit zwei Augen*¹ (Vision with two eyes) in 1861 (Figure 1). Three articles in the current volume of *Strabismus* present translations by H. J. Simonsz into English from Nagel's examination of stereoscopic vision. Nagel's book is dedicated to Helmholtz and von Gräfe under whom he had worked before taking up his position at Bonn; he was called to Tübingen in 1864 and remained there for the rest of his life. Part 1² of the translation (pp. 2-4 and 19-24 from Nagel's book) provides the Introduction and a section of Chapter 3 (on stereoscopic phenomena) concerned with binocular combinations of two rods. Part 2³ (pp. 24-38) addresses the question of fusion with disparate images like dots or lines. Part 3⁴ (pp. 38-51) presents experiments with lines varying in orientation or curvature with respect to the two eyes. It is in this section that Nagel draws attention to cyclofusion and the involvement of the extraocular muscles in it. In general, Nagel's book reflects the struggles within German visual science to rescue the doctrine of identical retinal points from the evidence of stereoscopic depth perception.

DAS
SEHEN MIT ZWEI AUGEN

UND
DIE LEHRE
VON DEN
IDENTISCHEN NETZHAUTSTELLEN

VON
D^r. ALBRECHT NAGEL,
Privatdocenten der Universität in Bonn.

Mit 4 lithographirten Tafeln und 37 in den Text gedruckten Holzschnitten.

LEIPZIG UND HEIDELBERG.
C. F. WINTERSCHE VERLAGSHANDLUNG.
1861.

Figure 1. Detail of a portrait of Albrecht Nagel and the title page of his book on vision with two eyes.

The long observational history of binocular vision has been dominated by the appearance of a single world with two eyes and its breakdown when the eyes are distorted abnormally, as in strabismus or with external movement of one eye alone. From the beginning of the seventeenth century, with a growing understanding of the anatomy and dioptrics of eyes, stimulation of the retina could be linked to singleness of vision. At the beginning of the nineteenth century the flat horopter of Aguilonius⁵ assumed curvature. First Bell⁶ then Prevost⁷ proposed that corresponding points fall on a circle passing through the point of bifixation and the centers of the eyes. That is, the horopter is a circle rather than a plane. This was formalized by Vieth⁸ and later by

Müller⁹, and it has become known as the Vieth-Müller circle. Müller¹⁰ augmented his geometrical description of the circle of single vision by linking it with identical retinal points: “Whenever an object lies out of the ‘horopter’, its image falls on non-identical points of the retinae, and it is seen double”^{11(p.1201)}. In this way, there were only two possible states of binocular perception – singleness when objects fell on the circumference of the Vieth-Müller circle and doubleness otherwise, and singleness was served by a fixed organic relation between nerve fibers. It is this elegant edifice that Wheatstone¹² undermined with his demonstration of singleness and depth from images with slight retinal disparities: he remarked that “objects whose pictures do not fall on corresponding points of the two retinae may still appear single”^{12(p.384)}. Thus, in the year that saw publication of Wheatstone’s article on stereoscopic depth perception we find a statement by Müller denying its possibility. Wheatstone was well aware of the originality of both his observations and his interpretation of them, hence the meticulousness of his experiments in their support. Not only did he argue that singleness and depth could be observed with stimulation of non-identical retinal points but also that the stimulation of identical points could result in double vision. Moreover, the stereoscope, perhaps more than any other instrument, ushered in the era of experimentation for spatial vision. It fulfilled the scientific desire to examine binocular vision by observation and experiment. As Towne¹³ put it: “The introduction of the stereoscope inaugurated a new epoch in the physiology of vision, opened a wide field for further inquiry, and suggested additional methods of investigation, while the theory of binocular vision has been greatly modified by results which have been obtained through the medium of the instrument”^{13(p.70)}.

The impact of Wheatstone’s experiments was felt acutely by sensory physiologists in Germany. It was fully appreciated by Nagel who commented that

Wheatstone had shaken the dogma of identical points for the first time and that hardly any physiologists agreed with him; Nagel was not among them. As he noted in Part 1, Volkmann voiced his views that stereoscopic vision threatened the received wisdom and Panum¹⁴ stated similar sentiments. Panum sought to salvage the dogma with his concept of fusional areas so that single vision was expanded to a region around the geometrically constrained circumference of the Vieth-Müller circle. Nagel tried to finesse the problem by proposing separate circles for each eye (see Figs. 7 and 8 in Plates I and II which are reproduced in Parts 1 and 2). Despite praising the stereoscope for facilitating experiments on binocular vision, it was not enlisted for the observations Nagel made with simple line stimuli – he adopted a free-fusion method: “The stereoscope itself is the most insignificant in the experiments that belong here”. Much of the material in Part 1 is concerned with describing and justifying free-fusion using the example of viewing two coloured, vertical rods each in the parallel visual axes of the eyes. The interpretation of the ensuing perception is psychological rather than physiological. Using this procedure, Part 2 is concerned with combining pairs of dots or lines with different separations in each eye. The observations with dots are essentially like those made by Wheatstone¹². Many more manipulations are made with paired vertical or near vertical lines in each eye. The separations and orientations of one or both members of a pair varied and the pairings are illustrated. Somewhat frustratingly, Nagel refers to them in the text but he does not specify the figure numbers to which the observations apply. The final image presented in Part 2 is of a single vertical line in one eye and an inclined one in the other – rather like the first pair of stereoscopic figures in Wheatstone¹² in which the half images were both inclined to the vertical. Wheatstone’s stimuli were surrounded by circles to facilitate binocular alignment whereas such assistance was not employed in Nagel’s figures. It is Nagel’s difficulty with defining the

apparent location in space of the single line combination that leads to Part 3.

Combinations of inclined lines, curves and crosses are described as well as rivalling vertical and horizontal lines like those examined by Panum¹⁴. The most novel combination was that involving lines inclined to the horizontal – fusion of the lines was not accompanied by depth but by strong feelings of muscular activity. Nagel argued that the fusion is attained by rotations of the eyes in opposite directions – cyclovergence.

Ocular torsion was an issue of considerable contention in nineteenth century visual science. Speculations that the oblique muscles could rotate the eyes around the z-axis were frequently stated. For example, Bell¹⁵ wrote: “By dissection and experiment it can be proved, that the oblique muscles are antagonists to each other, and that they roll the eye in opposite directions, the superior oblique directing the pupil downwards and outwards, and the inferior oblique directing it upwards and inwards”^{15(p.174)}. However, evidence supporting it in humans was hard to find and when it was presented it was usually contended¹⁶. The possibility of torsion in opposite directions seemed fanciful and yet this is what Nagel proposed in order to maintain cyclofusion for lines inclined in opposite directions relative to the horizontal. Similar rotations about the vertical resulted in a depth effect with no cyclovergence. As Nagel noted, achieving cyclofusion with single lines required a lot of practice and patience. He returned to the issue later¹⁷⁻¹⁹ and showed that the cyclofusion could be produced more readily with arrays of lines. It was largely as a consequence of this demonstration that Hering^{20,21} became convinced that cyclovergence occurred. The involvement of cyclovergence remained hotly debated until photographic recording of eye movements verified it and Crone and Everhard-Halm²² recommended that large stimuli should be used in order to measure it. Readers can observe this for themselves with Figure 2; cyclofusion can be experienced using a simple stereoscopic display with the aid of red/cyan glasses or they

can experience it with free-fusion in the manner of Nagel's observations. The arrays of lines inclined relative to horizontal in the monocular views will appear horizontal with two eyes.

Figure 2. *Nagel's cyclofusion* by Nicholas Wade. Upper, an anaglyph of Nagel which can be viewed with red/cyan glasses with the combination red/left eye and cyan/right eye. The two portraits and the embedded lines are inclined at 2° clockwise and counterclockwise of horizontal. Lower, the two components are presented in Universal View with the left eye, right eye, and left eye stimuli; the left pair is for uncrossed and the right pair for crossed cyclofusion. Nagel would have viewed the equivalent of the left pair.

References

1. Nagel A. *Das Sehen mit zwei Augen, die Lehre von den identischen Netzhautstellen*, Leipzig and Heidelberg, Germany: Winter; 1861
2. Nagel A. Vision with two eyes, the doctrine of the identical retina Points. Part 1: The stereoscopic phenomena, *Strabismus*, 28:55-59, 2020
3. Nagel A. Vision with two eyes, the doctrine of the identical retina Points. Part 2: Fusion of disparate images, *Strabismus*, 28:???-???, 2020
4. Nagel A. Vision with two eyes, the doctrine of the identical retina Points. Part 3: Stereoscopic experiment: Fusion of disparate images including cyclofusion, *Strabismus*, 28:???-???, 2020
5. Aguilonius F. *Opticorum Libri Sex. Philosophis juxta ac Mathematicis utiles*. Antwerp, Belgium: Moreti: 1613
6. Bell C. *The Anatomy of the Human Body*, Vol. 3. London, UK: Longman, Rees, Cadell and Davies; 1803
7. Prevost P. *Essais de Philosophie ou Étude de l'Esprit Humain*. Geneva, Switzerland: Paschoud
8. Vieth GUA. Ueber die Richtung der Augen. *Annalen Physik* 28:233-253, 1818
9. Müller J. *Zur vergleichenden Physiologie des Gesichtssinnes des Menschen und der Thiere, nebst einen Versuch über die Bewegung der Augen und über den menschlichen Blick*. Leipzig, Germany: Cnobloch; 1826
10. Müller J. *Handbuch der Physiologie des Menschen*, Vol. 2. Coblenz, Germany: Hölscher, 1838
11. Müller J. *Elements of Physiology*, Vol. 2. W. Baly, trans. London, UK: Taylor and Walton; 1843

12. Wheatstone C. Contributions to the physiology of vision.—part the first. On some remarkable, and hitherto unobserved, phenomena of binocular vision. *Philos Trans R Soc London*. 128:371–394, 1838
13. Towne J. Remarks on the stereoscopic theory of vision, with observations on the experiments of Professor Wheatstone—Section I. *Guy's Hosp Rep*, 8:70–80, 1862
14. Panum PL. *Physiologische Untersuchungen über das Sehen mit zwei Augen*. Kiel, Germany: Schwerssche Buhhandlung, 1858
15. Bell C. On the motions of the eye, in illustration of the uses of the muscles and of the orbit. *Philos Trans R Soc London*, 113:166-186, 1823
16. Simonsz HJ, Wade NJ. (2018) Why did Donders, after describing pseudotorsion, deny the existence of ocular counterrolling together with Ruete, Volkmann, von Graefe and von Helmholtz, until Javal reconfirmed its existence? *Strabismus*, 26:211-222, 2018
17. Nagel A. Ueber das Vorkommen von wahren Rollungen des Auges um die Gesichtslinie. *Alb v Graefes Arch f Ophth* 14:228–246, 1868
18. Nagel A. Ueber das Vorkommen von wahren Rollungen des Auges um die Gesichtslinie. Zweiter Artikel. *Alb v Graefes Arch f Ophth* 17:237-264, 1871
19. Nagel A. On the occurrence of true rolling of the eye about the line of sight. Second article. *Strabismus*, 8:33-38, 2000
20. Hering E. Ueber die Rollung des Auges um die Gesichtslinie *Alb v Graefes Arch f Ophth*. 15:1–16, 1869
21. Hering E. About the rolling of the eye around the line of sight. *Strabismus*, 27:230-234, 2019

22. Crone RA, Everhard-Halm Y. Optically induced eye torsion. I. Fusional cyclovergence. *Alb v Graefes Arch f Ophth.* 195:231-239, 1975