University of Dundee

Biomarker counseling, disclosure of diagnosis and follow-up in patients with mild cognitive impairment

Frederiksen, Kristian S.; Nielsen, Thomas R.; Appollonio, Ildebrando; Andersen, Birgitte Bo; Riverol, Mario; Boada, Mercè

Published in:
International Journal of Geriatric Psychiatry

DOI:
[10.1002/gps.5427](https://doi.org/10.1002/gps.5427)

Publication date:
2021

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Frederiksen, K. S., Nielsen, T. R., Appollonio, I., Andersen, B. B., Riverol, M., Boada, M., Ceccaldi, M., Dubois, B., Engelborghs, S., Frölich, L., Hausner, L., Gabelle, A., Gabryelewicz, T., Grimmer, T., Hanseeuw, B., Hort, J., Hugon, J., Jelic, V., Koivisto, A., ... Waldemar, G. (2021). Biomarker counseling, disclosure of diagnosis and follow-up in patients with mild cognitive impairment: A European Alzheimer's Disease Consortium survey. *International Journal of Geriatric Psychiatry*, 36(2), 324-333. <https://doi.org/10.1002/gps.5427>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Biomarker counselling, disclosure of diagnosis and follow-up in patients with mild cognitive impairment: A European Alzheimer's Disease Consortium survey

Kristian Steen Frederiksen¹, T Rune Nielsen¹, Ildebrando Appollonio², Birgitte Bo Andersen¹, Mario Riverol³, Mercè Boada^{4, 5}, Mathieu Ceccaldi⁶, Bruno Dubois⁷, Sebastiaan Engelborghs^{8,9}, Rüdiger Fölsch¹⁰, Lucrezia Hausner¹⁰, Audrey Gabelle¹¹, Tomasz Gabryelewicz¹², Timo Grimmer¹³, Bernard Hanseeuw¹⁴, Jakub Hort¹⁵, Jacques Hugon¹⁶, Vesna Jelic¹⁷, Anne Koivisto^{18,19}, Milica G. Kramberger²⁰, Thibaud Lebouvier²¹, Alberto Lleó²², Alexandre de Mendonça²³, Flavio Nobili^{24, 25}, Pierre-Jean Ousset²⁶, Robert Perneczky^{27,28,29,30}, Marcel Olde Rikkert³¹, David Robinson³², Olivier Rouaud³³, Elisabet Sánchez³⁴, Isabel Santana^{35, 36}, Nikolaos Scarmeas^{37, 38}, Katerina Sheardova³⁹, Stephanie Sloan⁴⁰, Luiza Spiru^{41, 42}, Elka Stefanova⁴³, Latchezar Traykov⁴⁴, Görsev Yener⁴⁵, Gunhild Waldemar¹

Corresponding author:

Kristian Steen Frederiksen

Danish Dementia Research Center,

Regionshospitalet

Copenhagen University Hospital

Blegdamsvej

DK-2100 Copenhagen

Denmark

Email: kristian.steen.frederiksen@regionh.dk

Telephone: +45 35456911

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/gps.5427](https://doi.org/10.1002/gps.5427). This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

This article is protected by copyright. All rights reserved.

¹ Danish Dementia Research Centre

Department of Neurology

Rigshospitalet

University of Copenhagen

Copenhagen

Denmark

² School of Medicine and Surgery and Milan Center for Neuroscience (NeuroMI)

University of Milano – Bicocca

Milan

Italy

³ Department of Neurology

Clinica Universidad de Navarra. University of Navarra

Pamplona and Madrid

Spain

⁴ Research Center and Memory Clinic, Fundació ACE, Institut Català de Neurociències Aplicades,

Universitat Internacional de Catalunya

Barcelona

Spain

⁵ Networking Research Center on Neurodegenerative Diseases (CIBERNED), Instituto de Salud

Carlos III

Madrid

Spain

⁶ Department of Neurology and Neuropsychology, CHU Timone, APHM
and Aix Marseille University, Inserm, Institut de Neurosciences des Systèmes
Marseille

France

⁷ Alzheimer Research Center (IM2A) and Department of Neurology

Salpêtrière University Hospital, AP-HP

Sorbonne University

Paris

France

⁸ Reference Center of Biological Markers of Dementia (BIODEM)

Institute Born-Bunge and University of Antwerp

Antwerp

Belgium

⁹ Department of Neurology and Center for Neurosciences

UZ Brussel and Vrije Universiteit Brussel (VUB)

Brussels

Belgium

¹⁰ Central Institute for Mental Health

Department of Geriatric Psychiatry

University of Heidelberg

Mannheim

Germany

¹¹ Memory Resources and Research Center

Department of Neurology

Gui de Chauliac Hospital

Montpellier University Hospital

University of Montpellier

Montpellier

France

¹² Department of Neurodegenerative Disorders

Mossakowski Medical Research Centre PAN

Warsaw

Poland

¹³ Department of Psychiatry and Psychotherapy,

Technical University of Munich, School of Medicine, Klinikum rechts der Isar

Munich

Germany

¹⁴ Cliniques Universitaires Saint-Luc, Department of Neurology and Université Catholique de Louvain, Institute of Neuroscience

Brussels

Belgium

¹⁵ Memory Clinic

Department of Neurology

Charles University

Second Faculty of Medicine

Motol University Hospital

Prague

Czech Republic

¹⁶ Center of Cognitive Neurology

Lariboisière Hospital Paris

University of Paris

Paris

France

¹⁷ Clinic for Cognitive Disorders, Theme Aging

Karolinska University Hospital – Huddinge

Stockholm

Sweden

¹⁸ Department of Neurology

University of Eastern Finland

Kuopio University Hospital

Kuopio

Finland

¹⁹ Department of Neurosciences

University of Helsinki

Department of Geriatrics

Helsinki University Hospital

Helsinki

Finland

²⁰ Center for Cognitive Impairments

University Medical Center Ljubljana

Ljubljana

Slovenia

²¹ Lille 2 University of Health and Law

Pôle de Neurologie

Lille

France

²² Neurology Department

Hospital de la Santa Creu i Sant Pau

Barcelona

Spain

²³ Faculdade de Medicina da Universidade de Lisboa

Lisbon

Portugal

²⁴ Department of Neuroscience (DINOEMI)

University of Genoa

Genoa

Italy

²⁵ IRCCS Ospedale Policlinico San Martino

Genova

Italy

²⁶ Memory Clinic

Clinical Research Center

Toulouse University Hospital

Toulouse

France

²⁷ Department of Psychiatry and Psychotherapy

University Hospital

Ludwig-Maximilians-University Munich

Munich

Germany

²⁸ German Center for Neurodegenerative Disorders (DZNE) Munich,

Munich

Germany

²⁹ Munich Cluster for Systems Neurology (SyNergy)

Munich

Germany

³⁰ Ageing Epidemiology (AGE) Research Unit

School of Public Health

Imperial College London

London

UK

³¹ Radboudumc Alzheimer Center

Dept Geriatrics

Radboud University Medical Center

Nijmegen

The Netherlands

³² Memory Clinic

St. James's Hospital

Dublin

Ireland

³³ Vaud University Hospital

Leenaards Memory Centre

Department of Clinical Neuroscience

Lausanne

Switzerland

³⁴ Servicio de geriatría.

Hospital Universitario Ramon y Cajal.

Spain

³⁵ Neurology Department

Centro Hospitalar e Universitário de Coimbra

Coimbra

Portugal

³⁶ Faculty of Medicine

University of Coimbra

Coimbra

Portugal

³⁷1st Department of Neurology

Aiginitio University Hospital

National and Kapodistrian University of Athens

Athens

Greece

³⁸ Department of Neurology

Columbia University Medical Center

New York

USA

³⁹ Memory Center ICRC

International Clinical Research Center

St. Anne's University Hospital

Brno

Czech Republic

⁴⁰ Neuroprogressive Disorders and Dementia Network

Ninewells Hospital

Dundee

Scotland

⁴¹ Carol Davila University of Medicine and Pharmacy

Geriatrics-Gerontology and Old Age Psychiatry (Alzheimer Unit) Clinical Department

Bucharest

Romania

⁴² Excellence Memory Clinic and Longevity Medicine

Ana Aslan International Foundation

Bucharest

Romania

⁴³ University of Belgrade, Faculty of Medicine

Neurology clinic, Clinical Center of Serbia

Belgrade

Serbia

⁴⁴ Department of Neurology

UH “Alexandrovska”

Sofia

Bulgaria

⁴⁵ Department of Neurosciences

Department of Neurology

Dokuz Eylül University Medical School

Izmir

Turkey

Abstract

Objectives

Mild cognitive impairment (MCI) is associated with an increased risk of further cognitive decline, partly depending on demographics and biomarker status. The aim of the present study was to survey the clinical practices of physicians in terms of biomarker counselling, management, and follow-up in European expert centers diagnosing patients with MCI.

Methods

An online email survey was distributed to physicians affiliated with European Alzheimer's Disease Consortium centers (Northern Europe: 10 centers; Eastern and Central Europe: 9 centers; Southern Europe: 15 centers) with questions on attitudes towards biomarkers and biomarker counselling in MCI and dementia. This included post-biomarker counselling and the process of diagnostic disclosure of MCI, as well as treatment and follow-up in MCI.

Results

The response rate for the survey was 80.9% (34 of 42 centers) across 20 countries. A large majority of physicians had access to biomarkers and found them useful. Pre- and post-biomarker counselling varied across centers, as did practices for referral to support groups and advice on preventive strategies. Less than half reported discussing driving and advance care planning with patients with MCI.

Conclusions

The variability in clinical practices across centers calls for better biomarker counselling and better training to improve communication skills. Future initiatives should address the importance of communicating preventive strategies and advance planning.

Keywords: mild cognitive impairment, dementia, diagnostic disclosure, biomarker counselling, biomarkers, Alzheimer's disease, survey, diagnosis

Key points

- Physicians' practice regarding biomarker counselling, disclosure of diagnosis and follow-up in patients with mild cognitive impairment is not known;
- Practices varied across European centers with regards to a number of issues including biomarker counselling and preventive strategies
- Communications training and development of guidelines on these issues may help to improve practices and realize less variability

Word count of the body text: 3084

1 INTRODUCTION

A growing number of patients are referred for diagnostic evaluation of possible cognitive impairment. This is presumably due to, in part, an increasing prevalence of dementia worldwide^{1,2} but may also be driven by an increased awareness of dementia in the population at large and among physicians³. Stakeholders also highlight the need for early diagnosis prior to the stage of dementia to enable adequate support and possibly, in the future, the ability to offer early disease-modifying therapy⁴. This is likely to increase the number of patients diagnosed with more subtle cognitive impairment, which include patients without dementia but with an underlying neurodegenerative disease and other possibly non-progressive conditions.

The term mild cognitive impairment (MCI) was created to capture a group of those patients with objectively measurable cognitive impairment not fulfilling the criteria for dementia (e.g., no impairment in activities of daily living)⁵⁻⁷ and not necessarily related to dementia disorders.

Although initially developed as a research tool, MCI has since been adopted into clinical practice at many centers. Over the years, the concept has further evolved, especially with the introduction of Alzheimer's disease (AD) biomarkers and the subsequent addition of MCI due to AD and prodromal AD to the diagnostic criteria⁸⁻¹⁰. Although patients with MCI have a higher risk of progression to dementia¹¹, this risk varies greatly depending on a variety of factors¹². For example, in one study, having MCI and an abnormal biomarker of amyloid and neurodegeneration was found to increase the lifetime risk of a 60-year-old from 78.1% to 95.6% versus only abnormal markers of neurodegeneration. Moreover, the risk decreased approximately 10 percentage points for a 75 year old compared to a 60 year old due to shorter life expectancy¹³.

It should be kept in mind that these estimates do not convey the individual patient's risk of progression but are estimates at group level. Another issue is that an age-dependent proportion of older people will have asymptomatic amyloidosis in the brain ¹⁴ and a relatively high incidence of other age-related conditions i.e. cerebrovascular disorders which may also affect biomarkers. This is also reflected in the fact that 'incidental' amyloidosis can be found in an equivalent proportion of patients with dementia not usually associated with amyloid pathology ¹⁵. This highlights the important issue of biomarker counselling prior to and following sampling in patients with MCI and the ethical dilemmas inherent to early biomarker-based diagnosis ¹⁶. Other issues may make the term MCI difficult to administer in a clinical setting. Whereas a substantial number of patients and caregivers are familiar with the term dementia ³, MCI is likely to be less familiar. Moreover, conveying the concept of MCI, i.e., cognitive deficit but no impairment in activities of daily living, may be challenging.

Studies have been conducted on the attitudes of physicians toward the concept of MCI ¹⁷, the perception of patients and caregivers concerning disclosure of dementia ¹⁸, the possible benefits of a timely diagnosis ¹⁹, and on disclosing a positive biomarker status to patients with MCI or no cognitive impairment ^{20–22}. Moreover, previous surveys have examined physician practices for diagnosing MCI, including how the diagnosis was disclosed, the terms that were used, and follow-up ^{23–25}. However, little is known about how physicians who manage patients with MCI carry out biomarker counselling or how the results and consequences of biomarker sampling are communicated to patients. Additionally, there also is no clinical standard established for biomarker use in MCI-patients.

Thus, the primary objective of the present study was to survey the clinical practices of European physicians in terms of biomarkers and biomarker counselling in MCI. We also assessed how the

concept of MCI and biomarker results were conveyed to patients, in addition to how the physician's characteristics may influence how this is done. Finally, we assessed the guidance and management, including follow-up, available to patients with MCI.

2 METHODS

2.1 Study design

The present study was designed as a survey of physicians working in European Alzheimer's Disease Consortium (EADC) centers. EADC is a European network of centers of excellence working in the field of AD and was established in 2001. The centers conduct research and carry out diagnosis and treatment of patients suspected of having MCI or dementia.

For the present study, we developed two online questionnaires. One was sent to a coordinating doctor (usually a senior specialist) at participating centers and the other to individual center physicians regularly diagnosing and doing follow-up with patients with MCI. To identify centers who were interested in participating, an email was sent to the contact person at each center. Each center was asked to identify at least five physicians who were interested in participating.

The online survey was conducted from February 1, 2019 to April 31, 2019. Participants received an email with a link to the survey, and four rounds of reminders were sent. The questionnaire for coordinating physicians asked about issues on a more organizational level, while the one for individual physicians was divided into three sections addressing attitudes towards biomarkers and biomarker counselling in MCI and dementia; post-biomarker counselling and the process of diagnostic disclosure of MCI; and treatment and follow-up in MCI. The latter questionnaire also

included sections on demographics, training, and experience. Physicians were explicitly asked to complete their questionnaire according to their present practice.

To facilitate the statistical analysis, the survey presented answers using a five-point Likert scale of “Always/almost always” to “Never/almost never”; “Very well” to “Not at all”; and “To a great extent” to “Not at all”. Where relevant, “Don’t know” was also an option, just as space was available to make comments. For ease of reporting, some categories were collapsed into one. The data that support the findings of this study are available from the corresponding author upon reasonable request.

2.2 Statistical analysis

To explore the factors associated with practice and attitudes in disclosure of diagnosis, we carried out statistical analyses to assess the impact of age, years of experience, whether respondents actively recruited patients with MCI for research, and whether respondents had received training in the process of disclosing a diagnosis of dementia or similar devastating conditions. This was done using the Mann-Whitney U test for independent samples. Where relevant, we also compared differences in respondent practices between patients with MCI and patients with dementia using the Wilcoxon signed-rank test for dependent samples. Statistical analyses were carried out using Intercooled Stata 9.2 for Macintosh (StataCorp LLC, College Town, Texas, USA). Level of significance was set at $p < 0.05$ (two-tailed test).

2.3 Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

3 RESULTS

All 69 EADC member centers were emailed and 42 centers agreed to participate. The response rate was 80.9 % (35 out of 42) for center coordinating physicians (i.e. number of questionnaires received from the coordinating physician on organizational matters) and 50.6 % (110 out of 213) of individual physicians responded to the survey. Some coordinating physicians also completed the individual physician questionnaire. There was a median of three respondents per center (Range: 1–7). Twenty-three centers (67.7 %) were based in neurology, 7 (20.6 %) in geriatrics and 4 (11.8 %) in psychiatry. Centers had a median of 600 (range 200–5000) new visits per year. Among newly referred patients, a median of 31.5 % (range 15–60%) were diagnosed with MCI per month. Every center conducted research, and 32 conducted research that recruited patients with MCI. Three European regions were represented: Northern Europe: 11 centers (Belgium 2, Denmark 1, Finland 1, Germany 3, Ireland 1, the Netherlands 1, Sweden 1, UK 1); Eastern and Central Europe (based on Organisation for Economic Cooperation and Development definition) and Turkey: 9 centers (Czech Republic 2, Poland 1, Romania 1, Serbia 1, Slovenia 1, Switzerland 2, Turkey 1); and Southern Europe: 15 centers (France 5, Greece 1, Italy 3, Portugal 2, and Spain 4). All individual respondents reported diagnosing and following patients with MCI. The mean age of individual physicians was 42.1 years (standard deviation 10.1). Respondents reported having a mean of 95.6 (range 4–160) consultations per month. Fifteen respondents (14%) reported that they were not involved in research, and 33 (30%) reported spending 50% or more of their time on research. Table 1 reports additional baseline characteristics for individual respondents.

3.1 Attitudes towards biomarkers and biomarker counselling in MCI

Almost all respondents had access to magnetic resonance imaging (98.2%; n=108) and cerebrospinal fluid (CSF) sampling (91.8%; n=101), whereas fewer had access to ^{18}F -FDG-PET (74.5%; n=82) and amyloid PET (50.9%; n=56). A majority reported always or usually ordering a magnetic resonance imaging scan (MCI: 81.8%; n=90, dementia: 76.4%; n=84), whereas less than half always or usually ordered cerebrospinal fluid sampling or ^{18}F -FDG-PET (Supplementary Figure 1).

Respondents were also asked about the value of biomarkers. Biomarkers reflecting amyloid pathology was found to be the most valuable to predict progression and rate of progression in MCI patient. Very few found that the biomarkers had no value in this respect (Figure 1, supplementary Figure 2).

Most, but not all respondents, always or usually discussed the decision to order biomarkers with patients with MCI (85.7%; n=90) and dementia (81.1%; n=86). A large majority said that they discussed this more in-depth with patients with MCI. Individual physicians recruiting patients with MCI for research were more likely to do so (Table 2). Most, but not all of the respondents always or usually discussed the ability to diagnose the underlying etiology in patients suspected of having MCI (actively recruiting p=0.002) (Table 2). Fewer respondents always or usually discussed the ability to predict progression (MCI: 61.0%; n=61, dementia: 68.1%; n=64) and the uncertainties of biomarker interpretation with patients prior to sampling (MCI: 60.6%; n=60, dementia: 53.3%; n=56).

3.2 Post biomarker counselling and diagnostic disclosure of MCI

The diagnosis of dementia was found to be more meaningful to more respondents than MCI ($p=0.0002$) (Table 2). Most respondents (79.1%; $n=87$) never or seldom found that the diagnosis of MCI was unethical. For MCI disclosure guidelines, 28.3% ($n=30$) reported having access to guidelines (dementia: 46.2%; $n=49$).

Almost all respondents disclosed the MCI diagnosis when it was suspected. Risk of progression and the probable underlying etiology but not the probable rate was often discussed with patients with MCI (Table 2). About half disclosed the risk of progression and underlying cause regardless of whether the patient asked. A substantial minority only did so if asked by the patient (Table 3).

Linguistically the term reported used most often was MCI and rarely or never “a form of mild dementia” and “predementia stage” (Figure 2). Regarding tools used when disclosing the diagnosis of MCI, a little over half always or usually showed brain imaging scans, whereas about a quarter seldom or never showed brain imaging scans (communications training: $z=.04$; $p=0.04$) (Table 2).

3.3 Management of patients with MCI

Almost all respondents reported following up on MCI (95.2%; $n=100$) and patients with dementia (90.48%; $n=95$). Half (50.5%; $n=53$) reported following patients with MCI for ≥ 5 years and 45.3% ($n=48$) for dementia. Regarding frequency of visits, 37.7% ($n=40$) reported seeing patients with MCI twice a year, while 47.6% ($n=50$) did so for patients with dementia. A total of 67.3% ($n=70$) respondents reported that local support groups were available. Treatment with cholinesterase inhibitors in patients with MCI was offered always or usually by 23.6% ($n=25$) and seldom or never by 50.0% ($n=53$). Data on the prevalence of testing for the Butyrylcholinesterase K variant in

patients started on cholinesterase inhibitors was not collected. A majority also addressed non-pharmacological treatment (Supplementary Figure 3).

A little less than half always or usually reported discussing driving when giving the MCI diagnosis, whereas most discussed this with patients with dementia (Table 2). A similar pattern emerged for legal matters (Table 2).

4 DISCUSSION

Our study presents the results of a survey of 34 EADC centers of excellence working in the field of AD and 110 individual physicians affiliated to the centers on various aspects of the diagnostic disclosure and management of MCI, including biomarker counselling. Our most important finding is that there is a high degree of heterogeneity across centers, particularly regarding counselling (e.g., pre-biomarker counselling). In addition, a relatively high number of physicians did not discuss preventative measures with patients or planning for the future for instance by mentioning advance directives.

One of the arguments for early diagnosis is to offer support and possibly treatment, including the possibility of participating in trials with potentially disease-modifying therapy, to patients with MCI²⁶. For this reason, all patients should ideally have the opportunity to participate in support groups, and all patients should be offered counselling on how to mitigate the risk of progression. About two-thirds of respondents reported that it was possible to refer patients to local support groups. About three-fourths mentioned physical exercise as an intervention for MCI, and fewer than three-fourths discussed other possible strategies to reduce the risk of progression. There was also a clear difference in how often respondents discussed driving and advance planning with patients

diagnosed with MCI versus dementia. Although it would be logical to assume that advance planning is best handled at an early stage of cognitive impairment, reluctance to engage in possibly difficult issues may be related to the attitudes of the physician but may also be due to patient preference to avoid dealing with emotionally difficult issues. For many patients, the issue of whether the cognitive impairment may affect driving, is a sensitive one. Regardless it may be relevant even in patients with MCI to discuss driving, and possibly especially so in certain cognitive subtypes of MCI such as patients with dys-executive syndrome or prominent visuo-cognitive impairment.

A large majority of respondents found that biomarkers were helpful in predicting progression to dementia in patients with MCI. Respondents saw AD biomarkers (tau and beta-amyloid) as the most valuable, which could indicate that MCI is often seen within a clinico-biological AD framework. However, CSF sampling and amyloid PET were nevertheless reported as performed in a minority of patients suspected of MCI in our study, which is in line with previous findings²⁷. This may be due to funding issues, reimbursement and access to PET facilities and tracers. Although 85.1% of respondents reported always or usually discussing biomarker sampling with patients, it follows that 14.9% do not. This distribution was the same for patients with dementia. In patients with dementia, refraining from discussing biomarker sampling may reflect the perception that conveying this type of information is difficult due to impaired capacity to consent. However, even if the patients would be unable to give informed consent, a legal representative could substitute. Providing inadequate information to the patient prior to biomarker sampling is problematic for several reasons. For example, the patient has the right to both know and not know what their prognosis is¹⁶. Thus, inadequate biomarker counselling may compromise non-maleficence or the ethical principle of autonomy.

Another issue is that biomarkers may be perceived as potentially more harmful in MCI due to the uncertainty related to individual patient prognosis. Although the probability of progression from MCI to dementia on a group level is highly increased depending on the biomarker status²⁸, it is difficult to determine at the individual patient level, with some patients progressing after variable time periods, some remaining stable, and some reverting back to normal cognition^{6,29–31}. Modelling of the risk of progression at the individual patient level is underway³² and likely to improve the ability of physicians to counsel patients about the individual risks of progression. The present study found that slightly more than half of respondents always or usually discussed the uncertainty of the biomarker results with their patients. In another study, in patients suspected of having dementia, physicians did not discuss the uncertainty related to the diagnosis in about a third of consultations

³³.

Physicians in routine clinical settings may fail to undertake a discussion of the uncertainty regarding biomarkers for several different reasons. For example, they may be a lack of knowledge and unfamiliarity with (CSF) biomarker sampling, or generally a disbelief that biomarkers are accurate, or variability in the distribution of the types of patients individual physicians are faced with. It may also be that the probability of conflicting biomarker results is high implying that the interpretation of AD biomarkers is complicated by multiple biomarker constellations³⁴. Or there may be a reluctance to introduce uncertainty into the diagnosis, or a belief that uncertainty may weaken the patients' trust in the physician. In our study, around 60% always or usually included information on how biomarkers may help estimate the risk of progression. Physicians may also avoid prognostication due to various perceptions or feelings, such as a sense of discomfort in terms of uncertainty, delivering bad news, or taking away hope. However, the right to know, which

derives from the moral value of respect for autonomy, is a central argument in favor of biomarker testing³⁵. Furthermore, withholding information and dishonesty may have consequences for the patient-doctor relationship and thus, ultimately, for the patient³⁶. Moreover, not being open about the risk of progression may deprive the patient of the chance to plan for this eventuality. Lastly, evidence suggests that disclosing amyloid biomarker status is safe³⁷, which means that, with the right support and information, it is unharmed to be forthcoming about biomarker results. As always, an individualized approach is advisable as there also is a wish not to know their prognosis^{16,18,38}. Indeed, comments from respondent in the present study indicate that they try to tailor information and diagnostic disclosure to avoid, e.g., not overwhelming the patient, which is a risk³⁹.

One way to ensure adequate pre and post biomarker counselling is to have guidelines available. A total of 28.3% of respondents reported that national or local guidelines were available on diagnostic disclosure of MCI, while 46.8% reported the same for dementia. This is in line with previous findings^{23,40}. To our knowledge, no international guidelines have been published on this topic, although some recommendations exist^{41,42}. Such guidelines would be relevant for centers with relatively easy access to biomarkers, but less so in areas where access is limited.

Our study has limitations. Since we exclusively surveyed EADC expert centers, our findings may only be generalizable to tertiary centers with a high degree of specialization and access to biomarkers. In less specialized centers, using biomarkers may play a lesser role in diagnostic disclosure and MCI as a diagnosis may instead primarily be used to describe the functional level of patients rather than their clinico-biological trajectory. Nevertheless, our survey sample may reflect other parameters in memory clinics, for instance distributions between medical specialties.

Moreover, although we explicitly asked respondents to answer according to their actual practice, it is not possible to distinguish attitudes from actions.

In conclusion, we found that biomarkers are widely used in patients with MCI, but that not all patients receive adequate pre- and post-biomarker counselling. Clinical dementia practice varied greatly across centers, which may indicate that physicians lack guidance on issues related to diagnostic disclosure, including biomarker counselling. Training that enhances communication skills may represent one way of improving diagnostic disclosure. At present, because disease-modifying therapies are not available for patients with prodromal AD, additional emphasis must be put on preventive strategies, such as encouraging exercise and smoking cessation, but also on discussing advance planning and continued participation in clinical trials of emerging new treatments.

REFERENCES

1. Prince M, Wimo A, Guerchet M, Ali G, Wu Y, Prina M. The Global Impact of Dementia: An Analysis of Prevalence, Incidence, Cost and Trends. London: Alzheimer's Disease International. World Alzheimer Report 2015. 2015;
2. Taudorf L, Nørgaard A, Islamoska S, Jørgensen K, Laursen TM, Waldemar G. Declining incidence of dementia: A national registry-based study over 20 years. *Alzheimer's Dement.* 2019;15(11):1383–1391.
3. McParland P, Devine P, Innes A, Gayle V. Dementia knowledge and attitudes of the general public in Northern Ireland: an analysis of national survey data. *Int Psychogeriatrics.*

2012;24(10):1600–1613.

4. Prince M, Bryce R, Ferri C. World Alzheimer Report 2011 The benefits of early diagnosis and intervention Alzheimer's Disease International World Alzheimer Report 2011 The benefits of early diagnosis and intervention. London Alzheimer's Dis Int. 2011;
5. Petersen RC, Smith GE, Waring SC, Ivnik RJ, Tangalos EG, Kokmen E. Mild Cognitive Impairment: Clinical Characterization and Outcome. Arch Neurol. 1999;56(3):303–308.
6. Petersen RC, Roberts RO, Knopman DS, et al. Mild cognitive impairment: ten years later. Arch Neurol. 2009;66(12):1447–1455.
7. Winblad B, Palmer K, Kivipelto M, et al. Mild cognitive impairment - Beyond controversies, towards a consensus: Report of the International Working Group on Mild Cognitive Impairment. J Intern Med. 2004;256(3):240–246.
8. Albert MS, DeKosky ST, Dickson D, et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. Alzheimers Dement. 2011;7(3):270–279.
9. Cummings JL, Dubois B, Molinuevo JL, Scheltens P. International Work Group criteria for the diagnosis of Alzheimer disease. Med Clin North Am. 2013;97(3):363–368.
10. Dubois B, Feldman HH, Jacova C, et al. Advancing research diagnostic criteria for Alzheimer's disease: the IWG-2 criteria. Lancet Neurol. 2014 Jun;13(6):614–629.
11. Hu C, Yu D, Sun X, Zhang M, Wang L, Qin H. The prevalence and progression of mild cognitive impairment among clinic and community populations: a systematic review and meta-analysis. Int psychogeriatrics. 2017;29(10):1595–1608.
12. Song Y-N, Wang P, Xu W, et al. Risk Factors of Rapid Cognitive Decline in Alzheimer's Disease and Mild Cognitive Impairment: A Systematic Review and Meta-Analysis. J

Alzheimers Dis. 2018;66(2):497–515.

13. Brookmeyer R, Abdalla N. Estimation of lifetime risks of Alzheimer's disease dementia using biomarkers for preclinical disease. *Alzheimer's Dement.* Elsevier Inc.; 2018;14(8):981–988.
14. Jansen WJ, Ossenkoppele R, Knol DL, et al. Prevalence of Cerebral Amyloid Pathology in Persons Without Dementia. *JAMA.* 2015;313(19):1924.
15. Bergeron D, Gorno-Tempini ML, Rabinovici GD, et al. Prevalence of amyloid- β pathology in distinct variants of primary progressive aphasia. *Ann Neurol.* 2018 Nov 21;84(5):729–740.
16. Vanderschaeghe G, Dierickx K, Vandenberghe R. Review of the Ethical Issues of a Biomarker-Based Diagnoses in the Early Stage of Alzheimer's Disease. *J Bioeth Inq.* 2018;15(2):219–230.
17. Mitchell T, Woodward M, Hirose Y. A survey of attitudes of clinicians towards the diagnosis and treatment of mild cognitive impairment in Australia and New Zealand. *Int Psychogeriatrics.* 2008;20(1):77–85.
18. Pinner G, Bouman WP. Attitudes of patients with mild dementia and their carers towards disclosure of the diagnosis. *Int Psychogeriatrics.* 2003;15(3):279–288.
19. Dubois B, Padovani A, Scheltens P, Rossi A, Agnello GD. Timely diagnosis for alzheimer's disease: A literature review on benefits and challenges. *J Alzheimer's Dis.* 2016;49(3):617–631.
20. Visser PJ, Wolf H, Frisoni G, Gertz H-J. Disclosure of Alzheimer's disease biomarker status in subjects with mild cognitive impairment. *Biomark Med.* 2012;6(4):365–368.
21. Lingler JH, Klunk WE. Disclosure of amyloid imaging results to research participants: Has the time come? *Alzheimer's Dement.* 2013 Nov;9(6):741-744.e2.

22. Lim YY, Maruff P, Getter C, Snyder PJ. Disclosure of positron emission tomography amyloid imaging results: A preliminary study of safety and tolerability. *Alzheimer's Dement*. 2016;12(4):454–458.
23. Nielsen TR, Svensson BH, Rohr G, et al. The process of disclosing a diagnosis of dementia and mild cognitive impairment: A national survey of specialist physicians in Denmark. *Dementia*. 2018;147130121877744.
24. Bertens D, Vos S, Kehoe P, et al. Use of mild cognitive impairment and prodromal AD/MCI due to AD in clinical care: A European survey. *Alzheimer's Res Ther*. 2019;11(1):1–12.
25. Roberts JS, Karlawish JH, Uhlmann WR, Petersen RC, Green RC. Mild cognitive impairment in clinical care: a survey of American Academy of Neurology members. *Neurology*. 2010 Aug 3;75(5):425–431.
26. Brooker D, La Fontaine J, Evans S, Bray J, Saad K. Public health guidance to facilitate timely diagnosis of dementia: ALzheimer's COoperative Valuation in Europe recommendations. *Int J Geriatr Psychiatry*. 2014 Jul;29(7):682–693.
27. Bocchetta M, Galluzzi S, Kehoe PG, et al. The use of biomarkers for the etiologic diagnosis of MCI in Europe: An EADC survey. *Alzheimer's Dement [Internet]*. 2015 Feb;11(2):195-206.e1. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1552526014024650>
28. Vos SJB, Verhey F, Frölich L, et al. Prevalence and prognosis of Alzheimer's disease at the mild cognitive impairment stage. *Brain*. 2015;138(5):1327–1338.
29. Visser PJ, Kester A, Jolles J, Verhey F. Ten-year risk of dementia in subjects with mild cognitive impairment. *Neurology*. 2006;67(7):1201–1207.
30. Xue H, Hou P, Li Y, Mao X, Wu L, Liu Y. Factors for predicting reversion from mild cognitive impairment to normal cognition: A meta-analysis. *Int J Geriatr Psychiatry*. 2019;34(10):1361–1368.

31. Malek-Ahmadi M. Reversion From Mild Cognitive Impairment to Normal Cognition: A Meta-Analysis. *Alzheimer Dis Assoc Disord*. 30(4):324–330.
32. IS van M, hanie J Vos, Isabelle Bos, Femke H Bouwman, Charlotte E Teunissen, Philip Scheltens, Frederik Barkhof, Lutz Frolich, Johannes Kornhuber, Jens Wiltfang, Wolfgang Maier, Oliver Peters, Eckart Rüther, Flavio Nobili, Giovanni WM van der F. Biomarker-based prognosis for people with mild cognitive impairment (ABIDE): a modelling study. 2019;4422(19):1–11.
33. Visser LNC, Pelt SAR, Kunneman M, et al. Communicating uncertainties when disclosing diagnostic test results for (Alzheimer's) dementia in the memory clinic: The ABIDE project. *Health Expect*. 2019;
34. Weise D, Tiepolt S, Awissus C, et al. Critical Comparison of Different Biomarkers for Alzheimer's Disease in a Clinical Setting. Buchert R, editor. *J Alzheimer's Dis* [Internet]. 2015 Sep 9;48(2):425–432. Available from: <https://www.medra.org/servlet/aliasResolver?alias=iospress&doi=10.3233/JAD-150229>
35. Smedinga M, Tromp K, Schermer MHN, Richard E. Ethical Arguments Concerning the Use of Alzheimer's Disease Biomarkers in Individuals with No or Mild Cognitive Impairment: A Systematic Review and Framework for Discussion. *J Alzheimer's Dis*. 2018;66(4):1309–1322.
36. A controlled trial to improve care for seriously ill hospitalized patients. The study to understand prognoses and preferences for outcomes and risks of treatments (SUPPORT). The SUPPORT Principal Investigators. *JAMA*. 274(20):1591–1598.
37. Wake T, Tabuchi H, Funaki K, et al. The psychological impact of disclosing amyloid status to Japanese elderly: a preliminary study on asymptomatic patients with subjective cognitive decline. *Int Psychogeriatrics*. 2017;1–5.

38. Ahalt C, Walter LC, Yourman L, Eng C, Pérez-Stable EJ, Smith AK. Knowing is better": Preferences of diverse older adults for discussing prognosis. *J Gen Intern Med*. 2012;27(5):568–575.
39. Mastwyk M, Ames D, Ellis K a., Chiu E, Dow B. Disclosing a dementia diagnosis: What do patients and family consider important? *Int Psychogeriatrics*. 2014;26(8):1263–1272.
40. Gilliard J, Gwilliam C. Sharing the diagnosis: A survey of memory disorders clinics, their policies on informing people with dementia and their families, and the support they offer. *Int J Geriatr Psychiatry*. 1996;11:1001–1003.
41. Grill JD, Apostolova LG, Bullain S, et al. Communicating mild cognitive impairment diagnoses with and without amyloid imaging. *Alzheimer's Res Ther*. *Alzheimer's Research & Therapy*; 2017;9(1):1–8.
42. Herukka SK, Simonsen AH, Andreasen N, et al. Recommendations for cerebrospinal fluid Alzheimer's disease biomarkers in the diagnostic evaluation of mild cognitive impairment. *Alzheimer's Dement*. 2017;13(3):285–295.

CONFLICT OF INTEREST STATEMENT

Kristian Steen Frederiksen, T Rune Nielsen, Ildebrando Appollonio, Birgitte Bo Andersen, Mario Riverol, Mercè Boada, Mathieu Ceccaldi, Bruno Dubois, Sebastiaan Engelborghs, Lutz Frölich, Lucrezia Hausner, Audrey Gabelle, Tomasz Gabryelewicz, Bernard Hanseeuw, Jakub Hort, Jacques Hugon, Vesna Jelic Anne Koivisto, Milica G. Kramberger, Thibaud Lebouvier, Alberto Lleó, Alexandre de Mendonça, Flavio Nobili, Pierre-Jean Ousset, Robert Perneczky, Marcel Olde Rikkert, David Robinson, Olivier Rouaud, Elisabet Sánchez, Isabel Santana, Katerina Sheardova,

Stephanie Sloan, Luiza Spiru, Elka Stefanova, Latchezar Traykov, Görsev Yener, Gunhild

Waldemar have no conflicts of interest regarding this manuscript.

Outside the submitted work Dr. Grimmer reported having received consulting fees from Actelion, Biogen, Eli Lilly, Iqvia/ Quintiles; MSD; Novartis, Quintiles, Roche Pharma, lecture fees from Biogen, Lilly, Parexel, Roche Pharma, and grants to his institution from Actelion and PreDemTech. Outside the submitted work Dr. Scarmeas reports grants from Alzheimer's Association, grants from European Social Fund , grants from Ministry for Health Greece, during the conduct of the study; personal fees from Merck Consumer Health , personal fees from NIH, grants from EISAI, personal fees from EISAI Korea, grants from EPAD,

Table 1. Individual physician characteristics

Age, n[†] (%)	
<i>n ≤40 years</i>	50 (45.5)
<i>n >40 years</i>	60 (54.5)
Sex (female), n (%) (n=109)	64 (58.7)
Specialists, n (%)	
<i>Neurologist</i>	60 (54.6)
<i>Geriatrician</i>	10 (9.1)
<i>Psychiatrist</i>	11 (10.0)
<i>Old age psychiatrist</i>	8 (7.3)
<i>Other specialty</i>	4 (3.6)
<i>No</i>	17 (15.5)
Clinical experience with dementia patients, n (%)	
<i>≤5 years</i>	32 (29.1)
<i>>5 years</i>	78 (70.9)

Communications training	
<i>Has received formal communications training</i>	33 (32.4)
<i>Has not received formal communications training</i>	77 (67.6)

† n=110 unless otherwise stated

Table 2. Results from questions on attitudes towards biomarkers

	Always/ almost always	Usually	About half the time	Seldom	Never/ almost never	P-value
Tend to discuss more in-depth with MCI [†] versus dementia [‡]						
Actively recruiting MCI	43.8% (n=32)	30.1% (n=22)	8.2% (n=6)	12.3% (n=9)	5.5% (n=4)	0.008*
Not actively recruiting MCI	13.3% (n=4)	43.3% (n=13)	10.0% (n=3)	13.3% (n=4)	20.0% (n=6)	
Discuss ability to diagnose underlying cause of MCI						
Actively recruiting MCI	80.0% (n=56)	14.3% (n=10)	1.4% (n=1)	4.3% (n=3)	0	0.002*
Not actively recruiting MCI	45.8% (n=11)	41.7% (n=10)	12.5% (n=3)	0	4.2% (n=1)	

Meaningfulness of diagnosis (physician)						
<i>MCI</i>	49.5% (n=54)	44.0% (n=48)	2.8% (n=3)	1.8% (n=2)	1.8% (n=2)	p=0.0002**
<i>Dementia</i>	75.5% (n=83)	17.3% (n=19)	0.9% (n=1)	6.4% (n=7)	0	
Meaningfulness of diagnosis (patient)						
<i>MCI</i>	29.1% (n=32)	52.7% (n=58)	5.5% (n=6)	12.7% (14)	0	p<0.00001**
<i>Dementia</i>	64.6% (n=71)	20.9% (n=23)	6.4% (n=7)	6.4% (n=7)	1.8% (n=2)	
Meaningfulness of diagnosis (caregiver)						
<i>MCI</i>	30.6% (n=33)	48.2% (n=52)	9.3% (n=10)	11.1% (n=12)	0.9% (n=1)	p<0.00001**
<i>Dementia</i>	80.0% (n=88)	12.7% (n=14)	2.7% (n=3)	3.6% (n=4)	0.9% (n=1)	
Discuss risk of progression						
<i>MCI</i>	39.8% (n=41)	36.9% (n=38)	11.7% (n=12)	6.8% (n=7)	4.9% (n=5)	p<0.00001**
<i>Dementia</i>	48.1% (n=50)	30.8% (n=32)	9.6% (n=10)	9.6% (n=10)	1.9% (n=2)	
Discuss probable underlying cause						
<i>MCI</i>	30.7% (n=27)	45.5% (n=40)	13.6% (n=12)	5.7% (n=5)	4.6% (n=4)	p<0.00001**
<i>Dementia</i>	59.1% (n=62)	27.6% (n=29)	6.7% (n=7)	3.8% (n=4)	2.9% (n=3)	
Shows brain imaging when disclosing MCI diagnosis €						
<i>Communications training</i>	45.5% (n=15)	24.2% (n=8)	12.1% (n=4)	3% (n=1)	15.2% (n=5)	p=0.04**
<i>No communications training</i>	22.9% (n=16)	30.0% (n=21)	14.3% (n=10)	17.1% (n=12)	33.3% (n=11)	
Use other aids when disclosing MCI diagnosis §						
<i>Communications training</i>	35.7% (n=25)	50.0% (n=35)	4.3% (n=3)	7.1% (n=5)	2.9% (n=2)	p=0.002**
<i>No communications training</i>	14.7% (n=5)	29.4% (n=10)	5.9% (n=2)	26.5% (n=9)	23.5% (n=8)	
Discuss driving						
<i>MCI</i>	21.0% (n=22)	27.6% (n=29)	17.1% (n=18)	27.6% (n=29)	6.7% (n=7)	p<0.00001
<i>Dementia</i>	62.7% (n=66)	27.6% (n=29)	7.6% (n=8)	1.9% (n=2)	0	
Discuss other legal matters						
<i>MCI</i>	14.3% (n=15)	12.4% (n=13)	18.1% (n=19)	35.2% (n=37)	20.0% (n=21)	p<0.00001
<i>Dementia</i>	29.8% (n=31)	33.7% (n=35)	12.5% (n=13)	16.4% (n=17)	7.7% (n=8)	

Actively recruiting refers to whether individual physicians at the time of the survey were actively recruiting patients with MCI to research trials

† MCI: Mild cognitive impairment

‡ Likert scale items for this question were: Very well, Well, Fairly well, Poorly, and Very poorly

* P-values show results from Man-Whitney U test for independent samples comparing respondents actively recruiting patients with MCI. Non-significant results were found for age and years of experience for all questions

** P-values are for results from Wilcoxon signed-rank test for dependent samples comparing MCI versus dementia

Table 3. Disclosure of diagnosis and prognosis in MCI and dementia

	Yes, regardless of whether the patient asks about it	Only if the patient asks about it	Never, even if the patient asks about it	It depends on whether I think the patient may benefit from it
Probability of progression				
<i>MCI</i>	54.3%	35.2%	0	10.5%
<i>Dementia</i>	55.9%	34.3%	0	9.8%
Possible/probable rate of progression				
<i>MCI</i>	30.5%	50.5%	8.6%	10.5%
<i>Dementia</i>	30.4%	52.9%	2.9%	11.8%
Possible future symptoms				

<i>MCI</i>	22.9%	57.1%	5.7%	14.3%
<i>Dementia</i>	33.3%	50.0%	1.0%	15.7%
Possible underlying pathology				
<i>MCI</i>	51.9%	30.2%	2.8%	15.1%
<i>Dementia</i>	71.8%	22.3%	0	5.8%

MCI: mild cognitive impairment

Figure 1. Biomarkers for predicting progression

The figure displays individual physicians' evaluation of the value of biomarkers for predicting progression from MCI. "Don't know" replies are not displayed for ease of interpretation

Figure 2. Terms used when disclosing a diagnosis of mild cognitive impairment

The figure displays individual physicians' response with regards to questions on language used when disclosing a diagnosis of MCI

AD: Alzheimer's disease

MCI: Mild cognitive impairment

Supporting information

Supplementary figure 1: How often, when diagnosing patients with MCI do you order the following biomarkers

Supplementary figure 2: Do you find that the following biomarkers are valuable in predicting the rate at which individual MCI patients will progress

Supplementary figure 3: When disclosing a diagnosis of dementia how often do you discuss the following non-pharmacological treatment/prevention