University of Dundee

A new filaggrin gene mutation in a Korean patient with ichthyosis vulgaris

Ohguchi, Yuka; Nomura, Toshifumi; Suzuki, Shotaro; Mizuno, Osamu; Nomura, Yukiko; Nemoto-Hasebe, Ikue

Published in:
European Journal of Dermatology

DOI:
[10.1684/ejd.2014.2410](https://doi.org/10.1684/ejd.2014.2410)

Publication date:
2014

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Ohguchi, Y., Nomura, T., Suzuki, S., Mizuno, O., Nomura, Y., Nemoto-Hasebe, I., Okamoto, H., Sandilands, A., Akiyama, M., Irwin McLean, W. H., & Shimizu, H. (2014). A new filaggrin gene mutation in a Korean patient with ichthyosis vulgaris. *European Journal of Dermatology*, 24(4), 491-493. <https://doi.org/10.1684/ejd.2014.2410>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

A new filaggrin gene mutation in a Korean patient with ichthyosis vulgaris

Short title: *FLG* mutation in the Korean population

Yuka Ohguchi¹, Toshifumi Nomura¹, Shotaro Suzuki¹, Osamu Mizuno¹, Yukiko Nomura¹,
Ikue Nemoto-Hasebe¹, Hiroyuki Okamoto², Aileen Sandilands³, Masashi Akiyama⁴, W.
H. Irwin McLean³, Hiroshi Shimizu¹

¹ Department of Dermatology, Hokkaido University Graduate School of Medicine,
Sapporo, Japan

² Department of Dermatology, Kansai Medical University, Hirakata, Japan

³ Dermatology and Genetic Medicine, College of Life Sciences and College of Medicine,
Dentistry and Nursing, University of Dundee, Dundee, UK

⁴ Department of Dermatology, Nagoya University Graduate School of Medicine, Nagoya,
Japan

Correspondence: Toshifumi Nomura, M.D., Ph.D.

Department of Dermatology, Hokkaido University Graduate School of Medicine, North
15 West 7, Kita-ku, Sapporo 060-8638, Japan

Tel: +81-11-706-7387, Fax: +81-11-706-7820, E-mail: nomura@huhp.hokudai.ac.jp

Number of Words (main text): 734

Number of Figures: 1

Number of References: 10

Ichthyosis vulgaris (IV; OMIM 146700) is the most common genetic disorder of keratinization inherited in an autosomal semidominant fashion with incomplete penetrance [1]. IV is clinically characterized by dry skin and scaling, especially on the flexor limbs and the trunk [1]. Palmoplantar hyperlinearity and keratosis pilaris are also associated in most cases. Loss-of-function mutations in the gene encoding filaggrin (*FLG*), a crucial protein for epidermal barrier function, have been identified as a cause of IV [1]. These mutations also have been demonstrated as an important predisposing factor for atopic eczema (AE) and other allergic disorders, such as asthma, rhinitis and food allergies [2]. *FLG* mutations have been well characterized, especially in the European, Chinese, Singaporean-Chinese and Japanese populations and, to date, 23, 28, 22 and 10 *FLG*-null variants have been reported, respectively [2, 3, 4, 5, 6]. Interestingly, these studies have also highlighted the presence of specific *FLG*-null mutations distinct to each population, suggesting that each population is likely to have a unique set of *FLG* mutations [5, 7]. To date, only three loss-of-function mutations in *FLG*—p.Arg501Ter, c.3321delA and p.Tyr1767Ter—have been identified in Korean patients with IV and/or AE [8, 9]. Given that more than 10 *FLG*-null variants were identified in each of the Chinese, Singaporean-Chinese and Japanese populations, we speculate that there may be

more *FLG* mutations to be discovered in the Korean population. To clarify more precisely the genetic architecture of *FLG* mutations in the Korean population, we recruited a Korean patient with IV and performed mutation analysis of *FLG* in this study.

A 48-year-old female of South Korean origin living in Japan visited our hospital presenting with dry skin that she had suffered from since early childhood. Physical examination revealed marked dry skin and scaling on the extensor limbs and the trunk (*Figure 1A*). She also showed palmar hyperlinearity. She had a past history of allergic rhinitis, but had never suffered from other atopic diseases such as AE or asthma. Thus, a clinical diagnosis of IV was made. Genomic DNA from the patient was obtained from peripheral blood using QIAamp DNA Blood Maxi Kit (Qiagen, Maryland, USA) and was subjected to *FLG* mutation analysis. This study was approved by the medical ethics committee of Hokkaido University Graduate School of Medicine and conducted according to the Declaration of Helsinki Principles. The patient gave written informed consent. The entire coding regions of *FLG* were amplified and sequenced as described previously [10], which led to identification of a heterozygous nonsense mutation, p.Gln1790Ter, in repeat 5 of exon 3 of *FLG* (*Figure 1B*). Since this mutation creates a *BsaBI* restriction site, the presence of the mutation was further confirmed by restriction

digest, as described previously [6]. Briefly, a 449-bp PCR fragment was amplified with forward primer 5' GTAGTCGGAGACAGTGGAA 3' and reverse primer 5' ACATCAGACCTTTCCTGGGAC 3'. After *Bsa*BI restriction digestion, the mutant allele was resolved as 258-bp and 191-bp fragments, whereas the wild-type allele yielded a 449-bp fragment (*Figure 1C*). Her brother was also affected, but DNA from him and the other family members was not available for *FLG* mutation analysis.

Mutation p.Gln1790Ter was first reported in the Chinese population and shown to be carried by 1.3% and 4.8% of Han Chinese patients with AE and IV, respectively [3]. Recently, we identified the same mutation in a Japanese patient with IV who developed food-dependent exercise-induced anaphylaxis [6]. In this study, we further screened 224 Japanese patients with AE and 113 Japanese control individuals for the mutation p.Gln1790Ter. Notably, the mutation was also carried by one patient with AE, but not by any of the control individuals in the Japanese population. Furthermore, our study has revealed that this *FLG*-null variant is also present in the Korean population. Thus, the p.Gln1790Ter allele is shared by the Korean, Japanese and Chinese populations, but not by other populations. We speculate that the p.Gln1790Ter allele might originate from a single founder in East Asian countries. Nevertheless, the limited scope of this study

(single case report) has not allowed for haplotype analysis. To date, only three *FLG*-null variants have been reported in the Korean population. This study brings the total number of *FLG*-null mutations identified in the Korean population to four. Since we had no access to other Korean DNA samples in this study, further studies are warranted to clarify the prevalence of this mutation and its impact on the development of IV or atopic diseases in the Korean population.

Acknowledgments

We are grateful to the patient for her participation in this study. We also thank Ms. Masae Takeda for her fine technical assistance in *FLG* mutation analysis. This work was supported by Grants-in-Aid for Young Scientists (B) 25860925 (to T. Nomura) and (B) 24791129 (to I. Nemoto-Hasebe), both from the Japan Society for the Promotion of Science. The Centre for Dermatology and Genetic Medicine, University of Dundee is supported by a Strategic Award from The Wellcome Trust (098439/Z/12/Z). There are no conflicts of interest to declare.

References

1. Smith FJ, Irvine AD, Terron-Kwiatkowski A, *et al.* Loss-of-function mutations in the gene encoding filaggrin cause ichthyosis vulgaris. *Nat Genet* 2006; 38: 337-42.
2. Irvine AD, McLean WH, Leung DY. Filaggrin mutations associated with skin and allergic diseases. *N Engl J Med* 2011; 365: 1315-27.
3. Zhang H, Guo Y, Wang W, Shi M, Chen X, Yao Z. Mutations in the filaggrin gene in Han Chinese patients with atopic dermatitis. *Allergy* 2011; 66: 420-7.
4. Li M, Liu Q, Liu J, *et al.* Mutations analysis in filaggrin gene in northern China patients with atopic dermatitis. *J Eur Acad Dermatol Venereol* 2013; 27: 169-74.
5. Chen H, Common JEA, Haines RL, *et al.* Wide spectrum of filaggrin-null mutations in atopic dermatitis highlights differences between Singaporean Chinese and European populations. *Br J Dermatol* 2011; 165: 106-14.
6. Mizuno O, Nomura T, Ohguchi Y, *et al.* Loss-of-function mutations in the gene encoding filaggrin underlie a Japanese family with food-dependent exercise-induced anaphylaxis. *J Eur Acad Dermatol Venereol*, *in press*.
7. Akiyama M. *FLG* mutations in ichthyosis vulgaris and atopic eczema: spectrum of mutations and population genetics. *Br J Dermatol* 2010; 162: 472-7.

8. Yu HS, Kang MJ, Jung YH, *et al.* Mutations in the filaggrin are predisposing factor in Korean children with atopic dermatitis. *Allergy Asthma Immunol Res* 2013; 5: 211-5.
9. Kim EJ, Jeong MS, Li K, *et al.* Genetic polymorphism of FLG in Korean ichthyosis vulgaris patients. *Ann Dermatol* 2011; 23: 170-6.
10. Sandilands A, Terron-Kwiatkowski A, Hull PR, *et al.* Comprehensive analysis of the gene encoding filaggrin uncovers prevalent and rare mutations in ichthyosis vulgaris and atopic dermatitis. *Nat Genet* 2007; 39: 650-4.

Figure Legends

Figure 1.

(A) Marked dry skin and dark brownish scales on the right shin.

(B) A heterozygous transition mutation c. 5368C>T was identified in the patient, resulting in p.Gln1790Ter.

(C) A 449-bp PCR fragment was amplified with forward primer 5'

GTAGTCGGAGACAGTGGAA 3' and reverse primer 5'

ACATCAGACCTTTCCTGGGAC 3'. After *Bsa*BI restriction digestion, the mutant

allele was resolved as 258-bp and 191-bp fragments whereas the wild-type allele

yielded a 449-bp fragment. Mutant-specific bands (258-bp and 191-bp, arrowheads)

were identified in the patient but not in the control individual.