University of Dundee

Pairs of surface wave packets with zero-sum energy in the Hawking radiation analog

Guha, Anirban; Heifetz, Eyal; Gupta, Akanksha

Published in:
Physical Review D

DOI:
[10.1103/PhysRevD.102.104061](https://doi.org/10.1103/PhysRevD.102.104061)

Publication date:
2020

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Guha, A., Heifetz, E., & Gupta, A. (2020). Pairs of surface wave packets with zero-sum energy in the Hawking radiation analog. *Physical Review D*, 102(10), [104061]. <https://doi.org/10.1103/PhysRevD.102.104061>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Pairs of surface wave packets with zero-sum energy in the Hawking radiation analog

Anirban Guha^{*}

School of Science and Engineering, University of Dundee, Dundee DD1 4HN, United Kingdom

Eyal Heifetz[†]

Department of Geophysics, Porter school of the Environment and Earth Sciences, Tel Aviv University, Tel Aviv 69978, Israel

Akanksha Gupta[‡]

Department of Mechanical Engineering, Indian Institute of Technology, Kanpur, U.P. 208016, India

(Received 21 August 2020; accepted 27 October 2020)

Here we propose a minimal analog gravity setup and suggest how to select two surface gravity wave packets in order to mimic some key aspects of Hawking radiation from the horizon of nonrotating black holes. Our proposed setup, unlike the scattering problem conventionally studied, constitutes of a constant mean flow over a flat bathymetry, in which the two wave packets possess the same amount of wave action but equal and opposite (sign) amount of energy, thereby mimicking virtual particles created out of near horizon vacuum fluctuations. Attention is given to the physical mechanism relating to the signs of the wave action and energy norm with the wave's intrinsic and total phase speeds. We construct narrow wave packets of equal wave action, the one with positive energy and group speed propagates against the mean flow and escapes from the black hole as Hawking radiation, while the other with negative energy and group speed is drifted by the mean flow and falls into it. Hawking's prediction of low frequency mode amplification is satisfied in our minimal model by construction. We find that the centroid wave numbers and surface elevation amplitudes of the wave packets are related by simple analytical expressions.

DOI:

I. INTRODUCTION

Direct probing of Hawking radiation in gravitational black holes (BHs) seems to be unlikely in the near future. Hence, laboratory studies of the phenomena in analogous physical systems, obeying similar equations of motion as the fields around BHs, provide tools to examine and demonstrate different features of Hawking radiation. In the pursuit of finding laboratory analogs of BH radiation (c.f. Barceló [1] for an updated review), Schutzhold and Unruh [2] theoretically demonstrated how surface gravity waves, in the presence of a countercurrent flow in a shallow basin, can be used to simulate phenomena around BHs in the laboratory. Rousseaux *et al.* [3] reported the first successful analog gravity experiment mimicking white hole (WH) horizons by surface gravity waves. Weinfurter *et al.* [4] used localized obstacle to block the upstream propagation of a long wave, converting it into a pair of short waves with opposite-signed energy, one with positive and the other with negative energy. This experiment successfully demonstrated the thermal nature of the

stimulated Hawking process at an analog WH horizon. Hawking radiation in analog wave-current systems have been further established experimentally and numerically in recent years, see Refs. [5–7]. Specifically, Euvé *et al.* [5] established analog quantum Hawking radiation using correlation of the randomly fluctuating free surface downstream of the obstacle.

The objective in this paper is more modest. It aims to propose a minimal water wave analog of pairs of virtual particles with equal and opposite energy, created out of near horizon vacuum fluctuations, where the particle with the positive energy escapes to infinity, and the one with negative energy falls into the BH, leading to BH evaporation [8,9]. As this phenomena by itself is not necessarily related to wave scattering, it is enough to assume here a flow system with a constant mean countercurrent over a flat bathymetry (i.e., constant water depth, see Fig. 1).

II. PSEUDOENERGY AND PSEUDOMOMENTUM

Consider for simplicity a rectangular quasi-2D domain (x, z) of the size $(0, L) \times (-H, \eta')$, filled with water (assumed here to be inviscid and incompressible), where L is the horizontal length, H is the mean fluid depth, and

^{*}anirbanguha.ubc@gmail.com

F1:1 FIG. 1. Schematic diagram of the black hole analog setup. For
F1:2 details about the various symbols, see text.

67 $\eta'(x, t)$ denotes the free surface elevation about the mean
68 depth (e.g., Fig. 1). For this setup the continuity and Euler's
69 momentum equations read:

$$\nabla \cdot \mathbf{u} = 0; \quad \frac{D\mathbf{u}}{Dt} \equiv \left(\frac{\partial}{\partial t} + \mathbf{u} \cdot \nabla \right) \mathbf{u} = -\frac{\nabla p}{\rho} + \mathbf{g}. \quad (1a, b)$$

Here $\nabla \equiv (\partial/\partial x, \partial/\partial z)$ is the 2D gradient operator, $\mathbf{u} = (u, w)$ denotes velocity, p denotes pressure, ρ is the density of water (assumed constant), and $\mathbf{g} = -g\hat{\mathbf{z}}$ is the gravity vector pointing downwards.

Assuming periodic boundary conditions at $x = 0$ and L , it is straightforward to show that both the domain-integrated momentum in the x direction (P) and the total fluid energy (E):

$$P = \rho \int_{x=0}^L \int_{z=-H}^{\eta'} u dx dz, \quad (2a)$$

$$E = \frac{\rho}{2} \int_{x=0}^L \left[\left(\int_{z=-H}^{\eta'} |\mathbf{u}|^2 dz \right) + g(\eta'^2 - H^2) \right] dx, \quad (2b)$$

are conserved [10]. The two terms in the rhs of Eq. (2b) are, respectively, the fluid kinetic and potential energy. Consider a steady mean current in the negative x direction: $\mathbf{u} = (-\bar{U}, 0)$ with $\bar{U} > 0$, and a constant mean height H satisfying hydrostatic balance. This flow is a solution of Eq. (1a,b) and posses the domain integrated momentum and energy

$$\bar{P} = -\rho L H \bar{U}, \quad \bar{E} = \frac{\rho L H}{2} (\bar{U}^2 - gH). \quad (3a, b)$$

Now suppose that on top of this steady base state we add a perturbation that is composed of surface gravity waves of the form $\eta'(x, t) = ae^{i(kx - \omega t)} + \text{c.c.}$, where a and k , respectively, denote amplitude and wave number (defined positive here), $\omega = kc_p$ denotes frequency, c_p is the phase speed, and c.c. denotes complex conjugate. Then

$$\omega = \hat{\omega} - k\bar{U} = k(\hat{c}_p - \bar{U}) = kc_p, \quad (4)$$

where the intrinsic surface gravity wave frequency and phase speeds (denoted by hat) are given by the familiar dispersion relation:

$$\hat{\omega} = k\hat{c}_p = \pm\sqrt{gk \tanh kH}. \quad (5)$$

Denoting the wave fields by prime so that $\mathbf{u} = (-\bar{U} + u', w')$, we obtain

$$P = \bar{P} + \delta P, \quad \delta P = \rho \int_{x=0}^L \int_{z=0}^{\eta'} u' dx dz, \quad (6a)$$

$$\begin{aligned} E &= \bar{E} + \delta E, & \delta E &= E' - \bar{U} \delta P, \\ E' &= \frac{\rho}{2} \int_{x=0}^L \left(\int_{z=-H}^{\eta'} |\mathbf{u}'|^2 dz + g \eta'^2 \right) dx. \end{aligned} \quad (6b)$$

The quantities δP and δE are, respectively, known by (the somewhat confusing terms) pseudomomentum and pseudoenergy. As is evident from Eqs. (6a) and (6b), they are simply the momentum and energy contribution of the waves to the system. Since $\bar{P}$ and $\bar{E}$ are constant, δP and δE are also conserved (in the Appendix we explicitly show that δE in the shallow water limit is equivalent to the energy density integral in Schützhold and Unruh [2] [Eqs. (67) and (68)]). Note that E' —the positive definite wave eddy energy—is only one of the contributions by the surface waves to the total change in the energy (as will be clarified further in the next section). Hence, neither the pseudomomentum nor the pseudoenergy are sign definite; negative pseudoenergy implies that the addition of linear waves to the base flow reduces the energy of the system below its mean value $\bar{E}$, whereas positive pseudoenergy increases the energy above its mean value.

III. PAIRS OF ZERO-SUM PSEUDOENERGY WAVE PACKETS

The essential idea in this analogy is that confined surface gravity wave packets represent virtual particles. Therefore we aim to choose superposition pairs of wave packets with equal and opposite values of pseudoenergy δE in a way that the sign of their group velocity (in the frame of rest) will be equal to the sign of their pseudoenergy. When this is achieved, the wave packet with the positive pseudoenergy manages to overcome the leftward countercurrent $-\bar{U}$ and escapes rightward (from the BH horizon into the outer space), whereas the negative pseudoenergy wave packet is drifted leftward with the base flow (into the BH). Consequently, the energy in the left region (inside the BH) is reduced on average and become $\bar{E} - |\delta E|$. Eventually when the leftward wave packet dissipates, it is expected to reduce the mean energy of BH, so that the new mean energy $\bar{E}_{\text{new}} \approx \bar{E} - |\delta E|$.

(a)

(b)

F2:1 FIG. 2. Schematic description of the fact that (a) rightward propagating surface waves have a positive pseudomomentum, while
 F2:2 (b) leftward propagating surface waves have a negative pseudomomentum.

137 Next we wish to suggest how to choose excited pairs of
 138 oppositely signed pseudoenergy wave packets based on
 139 their physical properties. We first note that for surface
 140 waves it can be shown, after some algebra, that the wave
 141 eddy energy satisfies

$$E' = \frac{1}{2} \rho g L a^2 = \hat{c}_p \delta P, \quad (7)$$

143 implying that $\hat{c}_p$ and δP are of the same sign. This sign
 144 agreement can be understood from Fig. 2. The mechanism
 145 of surface wave propagation is such that the horizontal
 146 convergence (divergence) results in upward (downward)
 147 motion that translates the vertical height anomaly η' . Hence
 148 for rightward or positive propagation, $\hat{c}_p > 0$ [Fig. 2(a)],
 149 and u' is in phase with η' . Therefore the vertical integration
 150 of positive u' from the bottom to the wave crests exceeds
 151 the vertical integration of negative u' from the bottom to the
 152 wave troughs and consequently δP is positive, in agreement
 153 with Eq. (6a). By the same argument it follows that δP is
 154 negative when $\hat{c}_p$ is negative [Fig. 2(b)]. Equations (4),
 155 (6b), and (7) then imply the following relations:

$$\delta E = (\hat{c}_p - \bar{U}) \delta P = c_p \delta P = \left(1 - \frac{\bar{U}}{\hat{c}_p}\right) E'. \quad (8)$$

156 Consider then two waves with different wave numbers
 158 k^+ and k^- (both defined positive), where both waves have a
 159 positive $\hat{c}_p$ (and hence a positive δP). Thus both waves
 160 are “trying” to propagate to the right (in the positive x
 161 direction) against the mean current $-\bar{U}$, see Fig. 1. If we
 162 assume a situation such that
 163

$$\hat{c}_p^- < \bar{U} < \hat{c}_p^+,$$

164 then Eq. (8) implies that $\delta E^+ > 0$ while $\delta E^- < 0$. In other
 166 words, the wave that manages to counterpropagate against
 167 the current with a positive phase speed in the rest frame
 168 ($c_p^+ > 0$) carries a positive pseudoenergy, whereas the wave
 169 whose intrinsic phase speed is not large enough to match
 170 the opposed current ($c_p^- < 0$) carries a negative pseudoenergy
 171 and consequently propagates to the left in the rest
 172 frame (despite that the pseudomomentum of both waves

being positive), as shown in Fig. 1. This statement can be
 173 written in terms of frequency and wave action. Defining the
 174 wave action as $\delta A \equiv \delta P/k$, we obtain from Eq. (8) that
 175 $\delta E = \omega \delta A$. Consider δA as an analog for $\hbar$, then for
 176 positive δA the sign of the pseudoenergy is determined
 177 by the sign of its frequency ω . This suggests that we can set
 178 a perturbation of *zero* pseudoenergy composed of two
 179 waves ($\delta E = \delta E^+ + \delta E^- = 0$) with the same positive
 180 value of wave action $\delta A^+ = \delta A^- > 0$. These in combina-
 181 tion yield
 182

$$\Omega^+ = -\Omega^- > 0 \Rightarrow \hat{\Omega}^+ + \hat{\Omega}^- = \alpha^+ + \alpha^-, \quad (9a)$$

$$\left(\frac{\alpha^-}{\alpha^+}\right)^2 = \frac{\hat{\Omega}^-}{\hat{\Omega}^+} = \sqrt{\frac{\alpha^- \tanh \alpha^-}{\alpha^+ \tanh \alpha^+}}. \quad (9b)$$

183 Here we have used the following nondimensionaliza-
 184 tions: $\alpha^{+(-)} \equiv k^{+(-)} H$, $\hat{\Omega}^{+(-)} \equiv \hat{\omega}^{+(-)} H / \bar{U}$ and $\Omega^{+(-)} \equiv$
 185 $\omega^{+(-)} H / \bar{U}$. Additionally Eq. (4) has also been used, from
 186 which we obtain $\Omega^{+(-)} = \hat{\Omega}^{+(-)} - \alpha^{+(-)}$, where $\hat{\Omega}^{+(-)} =$
 187 $Fr^{-1} \sqrt{\alpha^{+(-)} \tanh \alpha^{+(-)}}$, in which the Froude number
 188 $Fr \equiv \bar{U} / \sqrt{gH}$. According to Eq. (9a), the waves have
 189 equal and opposite frequencies. Hence in the rest frame, the
 190 “+” wave will propagate to the right against the mean
 191 current whereas the “-” wave will be drifted to the left,
 192 following the scenario depicted in Fig. 1. Furthermore,
 193 Eq. (9b) provides a direct relation of the amplitude ratio of
 194 the “+” and “-” waves. An interesting point to notice from
 195 Eq. (9b) is that the condition of zero pseudoenergy super-
 196 position does *not* imply that the free surface should be
 197 initially flat.
 198

199 While the pseudomomentum of a monochromatic sinus-
 200 oidal wave is perfectly well defined, its position is
 201 obviously not. Therefore, in order to generate an initial
 202 zero pseudoenergy perturbation whose position and
 203 momentum are both reasonably well defined, we should
 204 construct pairs of narrow wave packets rather than pairs of
 205 monochromatic waves. Hence, the positive (negative)
 206 pseudoenergy wave packet should propagate with a posi-
 207 tive (negative) group speed c_g (or in nondimensional
 208 terms, $C_g^{+(-)} \equiv c_g^{+(-)} / \bar{U}$), satisfying:

F3:1 FIG. 3. Dispersion curves: (a) Ω versus α , and (b) C_p versus α . The blue, yellow and green curves, respectively, denote
 F3:2 $Fr = 0.4, 0.6$ and 0.8 . The short red lines in (a) denotes the slope of the blue curve, which equals to the group speed. The “*”s of same
 F3:3 color denote a pair wave; the one above the zero line has $\delta A > 0$ and $\delta E > 0$, while that below the zero line has $\delta A > 0$ and $\delta E < 0$.

$$C_g^{+(-)} \equiv \frac{\partial \Omega^{+(-)}}{\partial \alpha^{+(-)}} = -1 + \frac{1}{2Fr} \sqrt{\frac{1}{\alpha^{+(-)} \tanh \alpha^{+(-)}}} \left[1 + \frac{2\alpha^{+(-)}}{\sinh 2\alpha^{+(-)}} \right]. \quad (10)$$

209 Furthermore, the centroid group and phase speeds of each
 210 wave packet should possess the same sign. This is because
 211 the sign of c_p (or in nondimensional terms, $C_p^{+(-)} \equiv$
 212 $c_p^{+(-)}/\bar{U}$) determines the sign of δE whereas the sign of
 213 c_g determines the wave packet’s direction of propagation.
 214 Consider the positive branch of Ω and address only
 215 subcritical flows, i.e., $Fr < 1$, in order to enable wave’s
 216 counterpropagation. The variations of Ω and C_p with α for
 217 different Fr values are respectively plotted in Figs. 3(a)
 218 and 3(b). Two wave packets with equal wave action, and
 219 equal and opposite pseudoenergy, consist of a “pair wave”
 220 (denoted by the same colored “*”s), and therefore satisfies
 221 Eqs. (9a) and (9b). The “+” (“-”) wave packet’s frequency,
 222 phase and group speeds are all positive (negative), and
 223 hence escapes into space (falls into the BH), in analogy
 224 with Hawking radiation. Notice that for subcritical flows,
 225 this condition fails in the shallow-water limit (since the
 226 pseudoenergy is always positive); see the Appendix.

227 Figure 4 shows a pair of wave packets (both having
 228 positive wave action but equal and opposite pseudoenergy)
 229 in a countercurrent flow over a flat bathymetry. This
 230 configuration is numerically simulated using an in-house
 231

high-order spectral code, detailed in Raj and Guha [11].
 As already mentioned, a zero-sum pseudoenergy does not
 necessarily imply that the superposition of the wave packet
 would render the free surface flat, as clearly shown in
 Fig. 4(a), which is the configuration at $t = 0$. The back-
 ground flow is subcritical with $Fr = 0.7$. The “+” wave
 packet (centroid wave number $\alpha^+ = 0.8$) emits as Hawking
 radiation while the “-” wave packet (centroid wave
 number $\alpha^- = 2.47$) falls inside the BH; the wave pair
 has the same magnitude of centroid frequency as per
 Eq. (9a). Here the definition of the event horizon is
 arbitrary; however it must be located to the left of the
 superposed wave packets at $t = 0$. The fact that $\alpha^- > \alpha^+$
 is evident from the dispersion curve in Fig. 3(a). A conse-
 quence of $\alpha^- > \alpha^+$ is that $a^- > a^+$ as per Eq. (9b), which
 is also clear from Fig. 4(b).

IV. PARALLELS WITH THE RATIO OF BOGOLIUBOV COEFFICIENTS AND LOW-FREQUENCY MODE AMPLIFICATION

The study of classical and quantum fields around BHs
 reveals that a pair wave created with a temporal frequency
 Ω satisfies [2,8]:

$$\left(\frac{\beta^-}{\beta^+} \right)^2 = \exp \left(-\frac{\Omega}{T} \right), \quad (11)$$

where $\beta^{+(-)}$ are referred to as the positive (negative) norm
 amplitudes (also known as the Bogoliubov coefficients),

F4:1 FIG. 4. Simulation of zero-sum pseudoenergy wave packet pair for $Fr = 0.7$. (a) Configuration at $t = 0$, and (b) configuration at a
 F4:2 later time when the “+” wave packet escapes the BH while the “-” wave packet falls inside it.

and T denotes an effective temperature proportional to the surface gravity of a BH. According the Hawking's prediction $(\beta^-)^2 = [\exp(\Omega/T) - 1]^{-1}$, which implies divergence as $\Omega \rightarrow 0$ since for this limit, $(\beta^-)^2 \approx T/\Omega$.

In analog gravity experiments with surface waves in a countercurrent flow over a localized obstacle, parallels between Eq. (11) and the scattering coefficients were first established in Weinfurter *et al.* [4], and then in subsequent studies, e.g., see Refs. [5,6]. The scattering coefficients in the analog-gravity experiments correspond to the wave action of the “+” and “-” waves [4]. We emphasize that here we have *not* solved a scattering problem, therefore its relevancy to Eq. (11) is somewhat limited. Yet, it is interesting to see that in the current analysis $\delta A^+ = \delta A^-$, hence the $\Omega \rightarrow 0$ limit of Eq. (11) is always satisfied. Furthermore, noting that

$$\delta A^{+(-)} = \frac{\rho g L}{2} \frac{\{a^{+(-)}\}^2}{\omega^{+(-)} + k^{+(-)} \bar{U}}, \quad (12)$$

we readily find that $\delta A^+ \rightarrow \infty$ when $\hat{\omega}^+ \rightarrow 0$, leading to both $k^+ \rightarrow 0$ and $\omega^+ \rightarrow 0$ [c.f. Fig. 3(a)]. Hence by construction $\delta A^- \rightarrow \infty$, however the denominator in Eq. (12) for this case does not vanish, rather $a^- \rightarrow \infty$. This fact can also be clearly observed from Eq. (9b). In summary, the aspect of low-frequency mode amplification in Hawking's prediction is satisfied by this minimal model.

V. DISCUSSION

The aim of this paper is to characterize the properties of zero-sum energy pair wave packets in the hydrodynamic analogy of Hawking radiation. First we wished to clarify the somewhat non-intuitive physical meaning of positive and negative energy norms (pseudoenergy), how those are related to the wave propagation mechanism, and how the general energy norm converges to the one suggested by Schützhold and Unruh [2] in the shallow water limit.

Next we considered a simple setup consisting of a constant subcritical countercurrent flow over a flat bathymetry; this setup was enough to demonstrate the analog phenomena where positive (negative) energy wave packets escape from (drifted into) the black hole. The combined requirements of a wave packet pair with equal (and positive in our case) wave action, and equal and opposite signed pseudoenergy, determine their centroid wave numbers as well as their surface elevation amplitude.

While forming such pairs of wave packets in the laboratory might not be a simple task, it is straight forward to numerically simulate stochastic generation of such zero-sum energy pairs, mimicking near-horizon vacuum fluctuations. The nonlinear effects of wave dissipation and wave-mean flow interaction, which feedback into the countercurrent and shift the horizon position, are under ongoing numerical investigation and will be published in a follow-up paper.

ACKNOWLEDGMENTS

A. G. thanks Alexander von Humboldt foundation for supporting the research visit to Tel-Aviv University, Israel.

APPENDIX: PSEUDOENERGY OF SHALLOW WATER GRAVITY WAVE

Writing the pseudoenergy explicitly, using Eqs. (6a) and (6b) we obtain

$$\delta E = \frac{\rho}{2} \int_{x=0}^L \left[\int_{z=-H}^{\eta'} (|\mathbf{u}'|^2 - 2\bar{U}u') dz + g\eta'^2 \right] dx. \quad (A1)$$

In the shallow water limit, $|\mathbf{u}'|^2 \Rightarrow u'^2$, and u' is not a function of z . Consequently the pseudo-energy expression for shallow water gravity waves for this setup becomes

$$\delta E_{\text{SW}} = \frac{\rho}{2} \int_{x=0}^L (Hu'^2 + g\eta'^2 - 2\bar{U}u'\eta') dx. \quad (A2)$$

Let us define the perturbation velocity potential ϕ' to satisfy $\mathbf{u}' = \nabla\phi'$, then for the shallow water the linearized, time-dependent Bernoulli's potential equation (or equivalently, the linearized momentum in the x direction) implies

$$\left(\frac{\partial}{\partial t} - \bar{U} \frac{\partial}{\partial x} \right) \phi' = -g\eta'. \quad (A3)$$

This relation allows writing the integrand of Eq. (A2) solely in terms of ϕ'

$$\delta E_{\text{SW}} = \frac{\rho}{2g} \int_{x=0}^L \left[gH \left(\frac{\partial \phi'}{\partial x} \right)^2 + \left(\frac{\partial \phi'}{\partial t} \right)^2 - \left(\bar{U} \frac{\partial \phi'}{\partial x} \right)^2 \right] dx, \quad (A4)$$

which is equivalent to the energy norm defined in Eqs. (67) and (68) in Schützhold and Unruh [2]. Furthermore, for the shallow water surface gravity wave, the amplitudes of the vertical displacement a , and the velocity potential amplitude $|\phi|$, are related by [12]

$$a = \frac{\alpha |\phi|}{\sqrt{gH}}.$$

Using Eq. (8) and $\hat{c}_p = \pm \sqrt{gH}$, we can express the pseudoenergy in terms of $|\phi|$ as

$$\delta E_{\text{SW}} = \frac{\rho L}{2H} \alpha^2 |\phi|^2 (1 \mp Fr). \quad (A5)$$

Hence pseudoenergy for shallow-water waves is *always* positive for subcritical flows ($Fr < 1$). Therefore pairs of opposite pseudoenergy wave packets in subcritical flows require nonshallow water dynamics.

- 341 [1] C. Barceló, Analogue black-hole horizons, *Nat. Phys.* **15**, 358
342 210 (2019). 359
- 343 [2] R. Schützhold and W.G. Unruh, Gravity wave 360
344 analogues of black holes, *Phys. Rev. D* **66**, 044019 361
345 (2002). 362
- 346 [3] G. Rousseaux, C. Mathis, P. Maïssa, T.G. Philbin, and U. 363
347 Leonhardt, Observation of negative-frequency waves in a 364
348 water tank: A classical analogue to the Hawking effect?, 365
349 *New J. Phys.* **10**, 053015 (2008). 366
- 350 [4] S. Weinfurter, E. W. Tedford, M. C. Penrice, W. G. Unruh, 367
351 and G. A. Lawrence, Measurement of Stimulated Hawking 368
352 Emission in an Analogue System, *Phys. Rev. Lett.* **106**, 369
353 021302 (2011). 370
- 354 [5] L.-P. Euvé, F. Michel, R. Parentani, T. G. Philbin, and G. 371
355 Rousseaux, Observation of Noise Correlated by the Hawking 372
356 Effect in a Water Tank, *Phys. Rev. Lett.* **117**, 121301 373
357 (2016). 374
- [6] S. Robertson, F. Michel, and R. Parentani, Scattering of 375
gravity waves in subcritical flows over an obstacle, *Phys.*
Rev. D **93**, 124060 (2016). 376
- [7] L.-P. Euvé, S. Robertson, N. James, A. Fabbri, and G. 377
Rousseaux, Scattering of Co-Current Surface Waves on an 378
Analogue Black Hole, *Phys. Rev. Lett.* **124**, 141101 (2020). 379
- [8] S. W. Hawking, Black hole explosions?, *Nature (London)* 380
248, 30 (1974). 381
- [9] S. W. Hawking, Particle creation by black holes, *Commun.*
Math. Phys. **43**, 199 (1975). 382
- [10] O. Bühler, *Waves and Mean Flows* (Cambridge University 383
Press, Cambridge, England, 2009). 384
- [11] R. Raj and A. Guha, On Bragg resonances and wave triad 385
interactions in two-layered shear flows, *J. Fluid Mech.* **867**, 386
482 (2019). 387
- [12] P. Kundu and I. Cohen, *Fluid Mechanics* (Elsevier, 388
New York, 2004). 389