University of Dundee

Zygomaticus major muscle bony attachment site

Sarilita, E.; Rynn, C.; Mossey, P. A.; Black, S.

Published in:
Morphologie

DOI:
[10.1016/j.morpho.2020.06.009](https://doi.org/10.1016/j.morpho.2020.06.009)

Publication date:
2021

Licence:
CC BY-NC-ND

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Sarilita, E., Rynn, C., Mossey, P. A., & Black, S. (2021). *Zygomaticus major* muscle bony attachment site: a Thiel-embalmed cadaver study. *Morphologie*, 105(348), 24-28. <https://doi.org/10.1016/j.morpho.2020.06.009>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Morphologie

Zygomaticus major muscle bony attachment site: a Thiel embalmed cadaver dissection study

--Manuscript Draft--

Manuscript Number:	MORPHO-D-20-00041R1
Article Type:	Original Article
Section/Category:	Anatomy
Keywords:	zygomaticus major; Thiel embalmed; cadaver dissection
Corresponding Author:	Erli Sarilita Universitas Padjadjaran INDONESIA
First Author:	Erli Sarilita
Order of Authors:	Erli Sarilita Christopher Rynn Peter A Mossey Sue Black
Abstract:	<p>Summary</p> <p>Objective</p> <p>The zygomaticus major is a principal muscle of facial expression which is engaged when smiling. The zygomaticus major origin on the zygomatic bone is often discussed relevant to its importance in the field of plastic surgery. In addition, the zygomaticus major attachment site is also significant for forensic craniofacial reconstruction, separating the cheek into frontal and lateral surfaces. However, there are discrepancies amongst published articles regarding the precise origin of the zygomaticus major muscle. The aim of this study is to investigate more distinctive and palpable landmarks as the bony attachment of the zygomaticus major.</p> <p>Methods</p> <p>This project is the first zygomaticus major dissection study utilising Thiel embalmed cadavers. Fifty-two facial dissections were investigated in 26 Thiel embalmed bodies, bequeathed to the Centre for Anatomy and Human Identification at The University of Dundee between 2013 and 2015.</p> <p>Results</p> <p>This study found that the origin of zygomaticus major muscle was located at the superior margin of the temporal process on the lateral surface of zygomatic bone. Moreover, the zygomaticus major muscle overlapped the anterosuperior border of the masseter muscle. One out of 52 zygomaticus major muscles presented bifurcation.</p> <p>Conclusion</p> <p>The origin site of zygomaticus major is considered important to increase resemblance in forensic craniofacial reconstruction. Furthermore, since zygomaticus major is a salient muscle involved in facial expression, the potential effects for cosmetic/surgical procedures are also relevant to the medical field and successful surgical outcomes. The current study provided easily palpable landmarks of zygomaticus major origin site which is beneficial for both surgeons and forensic craniofacial reconstruction practitioners.</p>

Highlights

Zygomaticus major muscle bony attachment site: a Thiel embalmed cadaver dissection study

- This article observed the bony origin site of zygomaticus major muscle on Thiel-embalmed cadavers.
- Previous studies reported variations in the location of bony attachment.
- This study found that the origin of zygomaticus major muscle was located at the superior margin of the temporal process on the lateral surface of zygomatic bone.
- The zygomaticus major muscle overlapped the anterosuperior border of the masseter muscle.
- The current study provided easily palpable landmarks of zygomaticus major origin site which is beneficial for both surgeons and forensic craniofacial reconstruction practitioners.

Original article

Zygomaticus major muscle bony attachment site: a Thiel embalmed cadaver dissection study

Erli Sarilita^{1,2,3}, Christopher Rynn², Peter A Mossey³, Sue Black⁴

¹ Faculty of Dentistry, Universitas Padjadjaran, Indonesia

² Centre for Anatomy and Human Identification, University of Dundee, Scotland, UK

³ School of Dentistry, University of Dundee, Scotland, UK

⁴ Vice Chancellor's Office, Lancaster University, UK

Corresponding author:

Erli Sarilita, Jl. Raya Bandung Sumedang KM 21, Jatinangor 45363, West Java, Indonesia
(erli.sarilita@fkg.unpad.ac.id)

Tel: +62 22 7794120

Summary

Objective. The zygomaticus major is a principal muscle of facial expression which is engaged when smiling. The zygomaticus major origin on the zygomatic bone is often discussed relevant to its importance in the field of plastic surgery. In addition, the zygomaticus major attachment site is also significant for forensic craniofacial reconstruction, separating the cheek into frontal and lateral surfaces. However, there are discrepancies amongst published articles regarding the precise origin of the zygomaticus major muscle. The aim of this study is to investigate more distinctive and palpable landmarks as the bony attachment of the zygomaticus major.

Methods. This project is the first zygomaticus major dissection study utilising Thiel embalmed cadavers. Fifty-two facial dissections were investigated in 26 Thiel embalmed bodies, bequeathed to the Centre for Anatomy and Human Identification at The University of Dundee between 2013 and 2015.

Results. This study found that the origin of zygomaticus major muscle was located at the superior margin of the temporal process on the lateral surface of zygomatic bone. Moreover, the zygomaticus major muscle overlapped the anterosuperior border of the masseter muscle. One out of 52 zygomaticus major muscles presented bifurcation.

Conclusion. The origin site of zygomaticus major is considered important to increase resemblance in forensic craniofacial reconstruction. Furthermore, since zygomaticus major is a salient muscle involved in facial expression, the potential effects for cosmetic/surgical procedures are also relevant to the medical field and successful surgical outcomes. The current study provided easily palpable landmarks of zygomaticus major origin site which is beneficial for both surgeons and forensic craniofacial reconstruction practitioners.

Keywords: zygomaticus major, Thiel embalmed, cadaver dissection.

Site d'attachement osseux musculaire majeur de Zygomaticus: une étude de dissection de cadavres embaumés par Thiel

Résumé

Objectif. Le **zygomatique majeur** est un muscle principal de l'expression faciale qui est engagé en souriant. **L'origine du muscle zygomatique majeur** sur l'os zygomatique est souvent discutée en fonction de son importance dans le domaine de la chirurgie plastique. De plus, le site d'attachement majeur du zygomaticus est également important pour la reconstruction craniofaciale médico-légale, séparant la joue en surfaces frontales et latérales. Cependant, il existe des divergences entre les articles publiés concernant l'origine précise **du muscle zygomatique majeur**. Le but de cette étude est d'étudier des repères plus distinctifs et plus palpables comme l'attachement osseux du zygomaticus major.

Méthode. Ce projet est la première étude de **dissection sur le muscle zygomatique majeur** utilisant des cadavres embaumés Thiel. Cinquante-deux dissections faciales ont été étudiées dans 26 corps embaumés Thiel, légués au Centre d'anatomie et d'identification humaine de l'Université de Dundee entre 2013 et 2015.

Résultats. Cette étude a révélé que l'origine de zygomaticus major **était située au bord supérieur du processus temporal** sur la surface latérale de l'os zygomatique. De plus, le muscle majeur zygomatique chevauchait le bord antéro-supérieur du muscle masséter. Un des 52 principaux muscles zygomatiques présentait une bifurcation.

Conclusion. Le site d'origine de zygomaticus major est considéré comme important pour augmenter la ressemblance dans la reconstruction craniofaciale médico-légale. De plus, comme le zygomaticus major est un muscle saillant impliqué dans l'expression faciale, les effets potentiels pour les procédures cosmétiques / chirurgicales sont également pertinents pour le domaine médical et les résultats chirurgicaux réussis. L'étude actuelle a fourni des repères facilement palpables du site d'origine majeur de zygomaticus, ce qui est bénéfique pour les chirurgiens et les praticiens de la reconstruction craniofaciale médico-légale.

Introduction

The zygomaticus major muscle is the main muscle of facial expression engaged in smiling. It is one of the five upper lip elevator muscles, along with zygomaticus minor, levator anguli oris, levator labii superioris, levator labii superioris alaeque nasi. Zygomaticus major extends from zygomatic bone and blends with orbicularis oris at the lateral corner forming modiolus. The width of the superior origin of the zygomaticus major is reported to be 8 to 10 mm [1].

Previous articles regarding zygomaticus major muscle were published in the facial reconstructive surgery journals which indicate the importance of zygomaticus major in that field [1-7]. The caudal insertion of zygomaticus major has been clearly defined at the modiolus, slightly lateral to the corner of the mouth, together with the orbicularis oris, risorius, buccinator, depressor anguli oris and levator anguli oris muscles. However, it is the origin of this muscle that is less well explained [4].

In 2019, Ryu and Kahng performed facelift on a 66-year-old female patient and found anatomical variation of the zygomatic branches of the facial nerve which entered the deep fascia approximately 10 mm to the zygomaticus major origin site [20]. Paralytic injection intended for orbicularis oculi muscle may unintentionally expose zygomaticus major to temporary paralysis resembling the Bell's Palsy appearance which may extend to several months. Although temporary, this outcome is unfavorable for both clinicians and patients [5]. Therefore, knowledge of zygomaticus major muscle attachment location as part of spatial anatomy of facial layers plays important role in safe surgery.

In the field of forensic craniofacial anthropology, the zygomaticus major also plays a significant role. One of the branches of forensic craniofacial anthropology is craniofacial reconstruction (CFR) otherwise known as forensic facial approximation. CFR is a process of scientifically estimating and depicting the shape of the face from the underlying skull. Whilst CFR methodology is also used to depict faces from historical/archaeological skulls, the importance lies in its forensic application which may serve to assist the recognition of unidentified victims following either severe facial trauma or putrefaction [8-9]. CFR techniques are varied and include 2D, 3D manual (sculptural, whether physical or virtual) and 3D computerised (automated), but most involve both scientific and artistic input. The accuracy and reliability of CFR is still a controversy among practitioners and researchers [9].

The University of Dundee employs a combinatorial method of three aspects: (i) anatomical muscle sculpting; (ii) facial feature morphology estimation, and (iii) overlaying a skin layer that is guided by average soft tissue depth (ASTD) pegs attached to the craniofacial landmarks [9]. The anatomical modelling stage requires the sculpting of facial musculature onto the 3D skull (on skull replica by manual method or on skull model by using computer software) before adding the skin layer [9]. One of the sculpted muscles is the zygomaticus major.

The literature describes several variations for the attachment of zygomaticus major [2-7]. Freilinger et al. in 1987 dissected 12 formaldehyde-phenol cadavers and located the zygomaticus major origins anterior to the zygomaticotemporal suture [2]. In 1994, Tremolada et al. described that the zygomaticus major attached to the inferior border of the zygomatic bone where it was intersected by a line between the lateral canthus to the angle of the mandible based on the dissection of 8 fresh and 3 embalmed cadavers [3]. Mowlavi and Wilhelmi performed dissection on 13 cadavers in 2004 and discovered that the upper attachment of the zygomaticus major was at the most anterior-inferior part of the temporal fossa at the intersection of the frontal process and the temporal process of the zygomatic bone [4]. Facial dissection of 24 cadavers by Spiegel and DeRosa in 2005 found that zygomaticus major origins were 1.4 and 1.5 cm below the Frankfort horizontal line at 1 cm lateral to and at the lateral canthus [5]. Miller et al. confirmed the reliability of the subzygomatic fossa as the zygomaticus major origin through hemifacial dissection of 23 fresh cadavers in 2007 [6]. In 2018, Ryu et al. reported that mean distances between the centre of the zygomaticus major origin and the posterior border of tragus 61 mm through facelift procedure on 20 patients [7].

Due to the variation of the zygomaticus major origin in the literature, this study aimed to identify a simple and palpable bony landmark of the zygomaticus major muscle attachment site to aid the successful outcome of both CFR and facial surgery. Palpability of the bony landmark is advantageous for clinicians, whereas bony landmark is particularly beneficial for CFR practitioners in which cases only skeletal structures available.

Materials and methods

Fifty-two facial dissections were conducted on 26 cadavers bequeathed to the Centre for Anatomy and Human Identification, University of Dundee and comprised 15 males and 11 females. The age at death for the cadavers ranged from 61 to 101 years with a mean age of 80.5 ± 11.49 years. All cadavers were preserved using the Thiel soft fix embalming method [10].

Routine dissection tools such as scalpel blade, scalpel handle, haemostat forceps, tweezer forceps, and scissors, were used in the dissection procedure. The first author performed the facial dissection. Zygomaticus major muscle was identified by firstly removing the skin and the subcutaneous tissue carefully. The dissected zygomaticus major was photographed using a COOLPIX S33 digital camera (Nikon, Japan) from an oblique angle (approximately 35 to 45°) to capture the whole cheek region. The first author observed and investigated the chosen landmark which represents the precise location of the origin of the zygomaticus major muscle whilst the second author (CR) verified the dissection findings. A third observer, with craniofacial anatomy and morphology experience, evaluated the bilateral digital images of 3 cadaver dissections for inter-observer study.

Results

All cadaver face dissections showed that the zygomaticus major consistently originated from the superior margin of the temporal process on the lateral surface of zygomatic bone (Fig. 1). The zygomaticus major overlapped the superoanterior border of the masseter muscle. The superior attachment of the masseter muscle was located along the inferior margin of the zygomatic bone in all cadavers (Fig. 1). Zygomaticus major was represented by a single band in all cadavers except one (male, 77 years), where a single band bifurcated caudally towards the modiolus, unilaterally on the left side. An inter-observer study to evaluate agreement was conducted, whether the zygomaticus major attachment was located at the superior margin of the temporal process of zygomatic bone which yielded 100% agreement between the 2 observers.

Discussion

The Thiel embalmed preservation technique is used in the Centre for Anatomy and Human Identification University of Dundee for undergraduate and postgraduate teaching and

research. Thiel preservation shows superiority compared to formalin in the flexibility of the soft tissue including the muscles, fascia and connective tissue [10-13]. Dissection of the Thiel bodies is considered easier and faster, compared to the formalin cadavers, due to the easy separation of the superficial tissues and structures which can be performed by precise blunt dissection. The 'life like' feel and response of the cadaveric tissue is rated highly by surgeons [10].

The zygomaticus major is commonly bilateral. In this study, two types of zygomaticus major were identified, single and bifid. The present study found one male cadaver with a bifid zygomaticus major, which was comparable to the results from the previous studies [14-18, 22]. The muscle fibres commenced bifurcation in the middle region of the muscle and ran inferiorly and inserted at the modiolus at two different, but closely located, regions. This variation may present as cheek dimples at skin level [14,16]. The finding of this study suggested an easily distinguishable and palpable bony landmark for the origin of the zygomaticus major muscle. This is beneficial for both fields of clinical reconstructive surgery and forensic CFR.

Earlier studies have attempted to identify the exact location of the zygomaticus major muscle origin [2-7]. The samples of five previous studies and the present study were derived from European ancestral group [2-6]. However, the findings varied among European studies. The results of the present study showed closest similarity to the study performed by Freilinger *et al.* as well as Mowlavi and Wilhelmi [2,4]. In contrast, studies by Tremolada *et al.* and Miller *et al.* reported that the zygomaticus major originated from the inferior margin of the zygoma, whereas the present study showed that the lowest border of the zygomatic bone including the subzygomatic fossa, was in fact the attachment site for the masseter muscle [3,6]. The zygomaticus major muscle passed over the area of the subzygomatic fossa but did not attach there: instead it attached to the superior margin of the temporal process of zygomatic bone.

A Korean cadaveric dissection study by Hu *et al.* was conducted to discover the topography of the masseter for botulinum toxin ("botox") injection application [19]. Some photographs of dissection showed the position of the zygomaticus major as it is located near to the masseter. The result of the present study corresponds well with the photographs of Hu *et al.* and Elvan *et al.* dissection studies, that the zygomaticus major overlaps the superior anterior attachment of masseter [19, 22].

Furthermore, the current study was not a quantitative linear measurements study in which distortion can occur after angle alteration. The differences in the zygomaticus major bony origin between the current study and the earlier cadaveric dissection studies may be due to tissue differences resulting from the different embalming methods and dissection techniques. Having the experience of dissecting both types of the embalmed body, the author felt that the formalin-fixed dissection was more challenging and required a more aggressive dissection approach whereas the Thiel cadavers were more amenable to gentler blunt dissection.

The cheek surface can be divided into 2 areas by the diagonal line formed by the underlying zygomaticus major (frontal cheek and lateral cheek). The attachment site of the zygomaticus major on the zygomatic bone is relevant to CFR due to its influence in depicting facial angle (frontal face and lateral face) and cheek shape. Changes in facial angle may influence the CFR result. Positioning this muscle attachment correctly might therefore enhance the accuracy of CFR, and the results of this study shows an alteration in the resulting facial depiction between the two approaches.

Periorbital rhytids treatment requires a clear definition of zygomaticus major origin landmark. Botulinum toxin injection targeting orbicularis oculi rhytids should avoid the the zygomaticus major fibers. The upper part of zygomaticus major is either overlapped by or adjacent to orbicularis oculi [5, 7, 22]. This study provided information for the clinicians that the zygomaticus major begins from the superior margin of the temporal process on the lateral surface of the zygomatic bone.

For the most part, the location of facial musculature in previous zygomaticus major dissection studies corroborated well with the findings of the current study. However, the consensus from this dissection study supported the idea that the results of the current study can be applied to the revised CFR guideline. The observation of bony attachment of the zygomaticus major on Thiel soft-fix cadavers is relevant to clinical facial reconstructive surgery as well as the forensic CFR field.

Conclusion

The zygomaticus major muscle attachment site is important to surgeons and CFR practitioners. This dissection study utilised Thiel embalmed cadavers which are more akin to

1 fresh tissue response than formalin-fixed remains. This study found that the zygomaticus
2 major bony attachment was located at the superior margin of the temporal process on the
3 lateral surface of zygomatic bone. The current study provided evidence of a variation from
4 this traditional standpoint and thus advocated an easier bony landmarks for zygomaticus
5 major attachment location compared to previous literature.
6
7
8
9

10 **Author contributions**

11
12 Concept/design: ES, CR, PM; data collection: ES; data analysis/interpretation: ES, CR;
13 craniofacial reconstruction: ES; craniofacial reconstruction supervision: CR; drafting the
14 manuscripts: ES; critical revision of the manuscript: ES, CR, PM, SB; approval of the
15 manuscript: ES, CR, PM, SB. All authors have read and approved the manuscript.
16
17
18
19
20
21
22
23
24

25 **Acknowledgements**

26
27 This study was a part of a doctoral thesis of the first author which is funded by the
28 Directorate General of Higher Education Ministry of Research, Technology, and Higher
29 Education of the Republic of Indonesia (666/E4.4/K/2014) and awarded a research grant by
30 the Indonesia Endowment Fund for Education (PRJ-1536/LPDP.3/2017). Preliminary result of
31 this study was presented as oral presentation at Anatomical Society Winter Meeting, 18th-
32 20th December 2017, Dundee, Scotland.
33
34
35
36
37
38
39
40

41 **Disclosure of interest**

42
43 The authors declare that there are no conflicts of interest.
44
45
46

47 **Ethical statement**

48
49 The bodies were procured from people who signed inform consent to donate their bodies
50 for teaching and research prior to death.
51
52
53
54

55 **References**

- 56
57 1. Mendelson BC, Muzaffar AR, Adams WP. Surgical anatomy of the midcheek and malar
58 mounds. *Plast Reconstr Surg* 2002;110:885-896.
59
60
61
62
63
64
65

2. Freilinger G, Gruber H, Happak W, Pechmann U. Surgical anatomy of the mimic muscle system and the facial nerve: importance for reconstructive and aesthetic surgery. *Plast Reconstr Surg* 1987;80:686-690.
3. Tremolada C, Fissette J, Candiani P. Anatomical basis for a safe and easier approach to composite rhytidectomy. *Aesthet Plast Surg* 1994;18:387-391.
4. Mowlavi A, Wilhelmi BJ. The extended SMAS facelift: Identifying the lateral zygomaticus major muscle border using bony anatomic landmarks. *Ann Plast Surg* 2004;52:353-357.
5. Spiegel JH, DeRosa J. The anatomical relationship between the orbicularis oculi muscle and the levator labii superioris and zygomaticus muscle complexes. *Plast Reconstr Surg* 2005;116:1937-1942.
6. Miller PJ, Sarah S, Shah A. The subzygomatic fossa: A practical landmark in identifying the zygomaticus major muscle. *Arch Facial Plast Surg* 2007;9:271-274.
7. Ryu MH, Moon VA, Yin W. The inclusion of orbicularis oculi muscle in the SMAS flap in Asian facelift: anatomical consideration of orbicularis muscle and zygomaticus major muscle. *Aesthetic Plast Surg* 2018;4: 471-478.
8. Prag J, Neave R. *Making Faces: Using Forensic and Archaeological Evidence*. London: British Museum Press; 1997.
9. Wilkinson C. Facial reconstruction—anatomical art or artistic anatomy? *J Anat* 2010;216:235-250.
10. Eisma R, Lamb C, Soames RW. From formalin to Thiel embalming: What changes? One anatomy department's experiences. *Clin Anat* 2013;26:564-571.
11. Brenner E. Human body preservation—old and new techniques. *J Anat* 2014;224:316-344.
12. Thiel W. Die Konservierung ganzer Leichen in natürlichen Farben. *Ann Anat* 1992;174:185–195.
13. Thiel W. Ergänzung für die Konservierung ganzer Leichen nach W. Thiel. *Ann Anat* 2002;184, 267–269.
14. Pessa JE, Zadoo VP, Garza PA, Adrian EK, Dewitt AI, Garza JR. Double or bifid zygomaticus major muscle: Anatomy, incidence, and clinical correlation. *Clin Anat* 1998;11:310-313.
15. Phan K, Onggo J. Prevalence of bifid zygomaticus major muscle. *J Craniofac Surg* 2019;30:758-760.

16. Lightoller GHS. Facial Muscles: The Modiolus and Muscles surrounding the Rima Oris with some remarks about the Panniculus Adiposus. *J Anat* 1925;60:1-85.
17. Hu KS, Jin GC, Youn KH, Kwak HH, Koh KS, Fontaine C, Kim HJ. An anatomic study of the bifid zygomaticus major muscle. *J Craniofac Surg* 2008;19:534-536.
18. Farahvash MR, Abianeh SH, Farahvash B, Farahvash Y, Yagoobi A, Nazparvar B. Anatomic variations of midfacial muscles and nasolabial crease: a survey on 52 hemifacial dissections in fresh Persian cadavers. *Aesthet Surg J* 2010;30:17-21.
19. Hu KS, Kim ST, Hur MS, Park JH, Song WC, Koh KS, Kim HJ. Topography of the masseter muscle in relation to treatment with botulinum toxin type A. *Oral Surg Oral Med O* 2010;110:167-171.
20. Ryu MH, Kahng D. Anatomical variation of zygomatic nerve branches around zygomaticus major muscle in facelift. *Plast Reconstr Surg Glob Open* 2017;5:2.
21. Lee WJ, Wilkinson CM, Hwang HS. An accuracy assessment of forensic computerized facial reconstruction employing cone-beam computed tomography from live subjects. *J Forensic Sci* 2012;57:318-327.
22. Elvan Ö, Örs AB, Tezer MS. Anatomical Evaluation of Zygomaticus Major Muscle With Relation to Orbicularis Oculi Muscle and Parotid Duct [published online ahead of print May 19, 2020]. *J Craniofac Surg*. doi: 10.1097/SCS.00000000000006396.

Fig. 1 Bony attachment of the zygomaticus major (ZMaj).

Figure 1