University of Dundee

The ability of maize roots to grow through compacted soil is not dependent on the amount of roots formed

Vanhees, Dorien J.; Loades, Kenneth W.; Bengough, A. Glyn; Mooney, Sacha J.; Lynch, Jonathan P.

Published in:
Field Crops Research

DOI:
[10.1016/j.fcr.2020.108013](https://doi.org/10.1016/j.fcr.2020.108013)

Publication date:
2021

Licence:
CC BY-NC-ND

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Vanhees, D. J., Loades, K. W., Bengough, A. G., Mooney, S. J., & Lynch, J. P. (2021). The ability of maize roots to grow through compacted soil is not dependent on the amount of roots formed. *Field Crops Research*, 264, [108013]. <https://doi.org/10.1016/j.fcr.2020.108013>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Revised manuscript with changes accepted:

Title: The ability of maize roots to grow through compacted soil is not dependent on the amount of roots formed

Authors:

Dorien J. Vanhees ^{a,b}

Kenneth W. Loades ^b

A. Glyn Bengough ^{b,c}

Sacha J. Mooney ^a

Jonathan P. Lynch ^{a (*)}

^a Division of Agricultural and Environment Sciences, School of Biosciences, University of Nottingham, Sutton Bonington Campus, Leicestershire, LE12 5RD, UK

^b The James Hutton Institute, Errol Road Invergowrie, DD2 5DA, UK

^c School of Science and Engineering, The University of Dundee, Dundee, DD1 4HN, UK

Author email addresses: dorien.vanhees@nottingham.ac.uk
Kenneth.Loads@hutton.ac.uk
a.bengough@dundee.ac.uk
sacha.mooney@nottingham.ac.uk
jpl4@psu.edu

(*) corresponding author. E-mail: jpl4@psu.edu Telephone: (+1) 814-863-2256

21 **Abstract**

22 **Highlights:**

- 23 • Soil compaction influenced the proportion of coarse roots (i.e. > 1 mm diameter)
- 24 of maize in a genotype-dependent manner
- 25 • Rooting depth was reduced and root distribution within the soil profile changed
- 26 when grown in compacted soil
- 27 • Rooting depth and total root length, total coarse root length and total fine root
- 28 length were not correlated with each other in compacted soil
- 29 • The ability of roots of different genotypes to reach a certain depth was not
- 30 related to the amount of roots formed

31 **Keywords**

32 Impedance; compaction; root length; root distribution; compensatory growth

33

Abstract

Mechanical impedance is a primary constraint to root growth and hence the capture of soil resources. To investigate whether rooting depth and root length under mechanical impedance caused by compaction are correlated we evaluated 12 maize lines at two field sites. To distinguish between lateral and nodal roots, roots were sorted into different diameter classes. Coarse roots had diameters >1 mm and represent nodal root axes. Greater proportions of coarse roots on compacted plots were found at both field sites however results were driven by genotypic variation. Soil compaction reduced total rooting depth (in all diameter classes) and coarse rooting depth at both sites compared to non-compacted plots. Root distribution was influenced by compaction with greater root length densities closer to the soil surface. Root length and root depth were not related to each other under impeded conditions. Coarse roots of some genotypes became obstructed on the compacted plots, while other genotypes were capable of growing through the impeding soil and reached deeper soil strata resulting in differential distribution of roots through the soil profile. On compacted plots we observed genotypes with similar root depths but with contrasting coarse root lengths. The ability of roots to grow through compacted soils is therefore not dependent solely on the coarse root length formed by the root system.

1. Introduction

The ability of plants to acquire nutrients and water is dependent on soil exploration. Mechanical impedance can lead to reduced total root length and/or a redistribution of root length within the soil profile (Pfeifer *et al.*, 2014a; Shierlaw and Alston, 1984), which could affect the acquisition of water and nutrients. As soils get denser and stronger with depth, due to overburden pressure (Gao *et al.*, 2012, 2016), mechanical impedance will often restrict deeper rooting root phenotypes more than topsoil foraging root phenotypes. Periodic droughts are common in many ecosystems and drier soils are generally harder (Gao *et al.*, 2012; To and Kay, 2005; Vaz *et al.*, 2011; Whalley *et al.*, 2005; Suralta *et al.*, 2018). However, plants with root systems that grow deeper are in general better adapted to drought (Chimungu *et al.*, 2014a; Lilley and Kirkegaard, 2016; Lynch, 2013; Zhan *et al.*, 2015). Certain soils offer very large mechanical impedance to roots, for example hard-setting soils in Australia (Mullins *et al.*, 1987) or rainfed lowland rice cultivation systems (Suralta *et al.*, 2018). Different agricultural management approaches can also introduce compaction and plough pans by wheeled traffic or trampling (Batey, 2009; Hamza and Anderson, 2005). Depending on the soil textural characteristics, suboptimal soil conditions during trafficking (such as high moisture contents) will exacerbate compaction (Horn *et al.*, 1995; Raper, 2005). Roots can become confined to surface soil strata when not capable of penetrating through a hard soil layer such as a plough pan (Barraclough and Weir, 1988; Ehlers *et al.*, 1983). Root systems are able to compensate root growth by exploiting the lesser impeded regions of the soil, as illustrated by split pot experiments (Bingham and Bengough, 2003; Pfeifer *et al.*, 2014a) or layered pot systems (Shierlaw and Alston, 1984). Roots of maize (Chimungu *et al.*, 2015), rice (Chandra Babu *et al.*,

2001; Clark *et al.*, 2000, 2002; Yu *et al.*, 1995), wheat (Botwright Acuña and Wade, 2005; Kubo *et al.*, 2006) and common bean (Rivera *et al.*, 2019) show substantial genotypic variability for penetrating strong wax layers simulating mechanical impedance.

Root systems consist of distinct root classes which vary by taxa, for example many dicot taxa have a dominant taproot, while monocots, such as cereals, form nodal roots from shoot nodes (Hochholdinger *et al.*, 2004; Lynch and Brown, 2012; Rich and Watt, 2013). Adult maize root systems consist of primary, seminal, crown (belowground nodal) and brace (aboveground nodal) roots, all these classes form lateral roots. For monocotyledons, nodal roots are the main parent axes of lateral roots present at depth as these laterals proliferate from nodal roots (Cairns *et al.*, 2004; Nagel *et al.*, 2012).

Genotypic variation for lateral root phenotypes has functional consequences in maize (Postma *et al.*, 2014; Zhan *et al.*, 2015; Zhan and Lynch, 2015; Jia *et al.*, 2018). Root classes have different elongation rates that vary greatly as a function of time. For maize, lateral roots have been found to elongate at 2.2 cm day⁻¹ for 2.5 days, while nodal roots elongated at a rate of 3 cm day⁻¹ over a 5 week period (Cahn *et al.*, 1989). Under non-impeded conditions primary roots of maize elongated at 4.8 cm day⁻¹, while seminals only elongated at 3.2 cm day⁻¹ (Veen and Boone, 1990). Differences in elongation rates between root types can lead to soils being differentially explored with time by each root type and could affect the volume and depth of bulk soil that can be explored within a certain time by different root types. Biomechanical properties also vary according to root class, with seminal roots being stronger than lateral roots (Loades *et al.*, 2013). Whether this translates to specific penetration ability under impeded soil conditions according to root class remains to be investigated. It has been hypothesised that the contrasting phenotypes of distinct root classes adds to a plants' plasticity and flexibility when interacting with different environments (Chochois *et al.*, 2015; Wu *et al.*, 2016) but the functional implications of the differential effects of mechanical impedance on distinct root classes are poorly understood.

Root system size differs among genotypes and different soil conditions (Gao and Lynch, 2016; Nakhforoosh *et al.*, 2014). Root system size, expressed as total root length or root length density, can be split between coarse and fine roots (Cahn *et al.*, 1989; Steinemann *et al.*, 2015; Varney *et al.*, 1991). Small grain cereals such as wheat or barley are characterised by fine axial roots, maize has thicker axial roots, while dicots and perennials have very coarse axial roots. But for all these species, a distinction between a main root axes and smaller diameter lateral roots can be made. Coarser roots are needed in order to deploy finer roots within the soil profile. Studies on wheat suggest that wheat genotypes with more root axes had greater penetration of wax layers (Whalley *et al.*, 2013).

Mechanical impedance not only affects root growth, it also has an impact on shoot growth. Root to shoot ratios can decrease under compaction (Andrade *et al.*, 1993; Hoffmann and Jungh, 1995; Pfeifer *et al.*, 2014a). Aboveground plant growth is impacted as leaf elongation rates can be reduced (Andrade *et al.*, 1993; Young *et al.*,

1997) and the rate of leaf appearance decreases (Beemster and Masle, 1996) when roots experience mechanical impedance. The reduction of shoot and root growth due to mechanical impedance can result in decreasing yield (Kirkegaard *et al.*, 1992; Tuzzin de Moraes *et al.*, 2020).

Better root growth under mechanical impedance can be attributed to different traits. For instance, the frictional component of mechanical impedance is reduced when roots produce mucilage or border cell sloughing (Iijima *et al.* 2000, 2004; Bengough and McKenzie, 1997). Smaller root tip radius to length ratios are linked to greater elongation rates under mechanical impedance (Colombi *et al.*, 2017b). Another beneficial trait is the presence of root hairs which can provide anchorage for roots to cross from loose to harder soil layers (Bengough *et al.*, 2011; Haling *et al.*, 2013). Root hairs also maintain water uptake when soils dry (Carminati *et al.*, 2017). Root anatomical traits such as greater cortical cell diameter have been linked to reduced energy costs under impeded conditions (Colombi *et al.*, 2019). It has been suggested that smaller outer cortical cells prevent buckling, which facilitate penetration of harder layers (Chimungu *et al.*, 2015).

Genotypes can adjust their root distribution with depth in response to compaction (Barracough and Weir, 1988) however few studies have compared different genotypes and their redistribution of roots under compaction. Little is known about root system size for those root systems that do manage to grow deeper in compacted soils. The hypothesis that rooting depth and root length are not related to each other on compacted plots was tested for deeper rooting genotypes.

2. Material and Methods

2.1. Plant material and growth conditions

Twelve maize (*Zea mays* L.) recombinant inbred lines from a study by Chimungu *et al.* (2015) were selected for different levels of root penetrability of a wax layer. These genotypes were planted in a split-plot design in order to study their root growth in compacted conditions at two field sites. Seeds were obtained from Dr. Shawn Kaeppeler (University of Wisconsin, Madison WI, USA – Genetics Cooperations Stock Center, Urbana, IL, USA). Genotypes were grown at the Apache Root Biology Centre (ARBC), Willcox Arizona, USA (32°01'N, 109°41'W), planted on June 16, 2016, and the Russell E. Larson Agricultural Research Center in Rock Springs (further referred to as PSU), Pennsylvania, USA (40°42'N, 77°57'W), planted on July 10, 2017. Field sites differed in soil texture, the ARBC site has a soil classified as a Grabe series (coarse-loamy, mixed, superactive, calcareous thermic Torrifuvent) and has a clay loam texture, while the PSU site is classified as a Hagerstown series (silt-loam, fine, mixed, semiactive, mesic Typic Hapludalf) and has a silt-loam texture. Compaction was introduced by passing over the treated plots with heavy machinery. At ARBC a 4 wheel tractor (4 tonnes with 8 passes) and at PSU a 3-axle truck (20 tonnes with 4 passes) were used. Penetration resistance (Figure 1) as well as an increase of dry bulk density were measured over the soil profile in order to verify increased soil

strength and soil density. A FieldScout SC900 Compaction Meter (Spectrum Technologies Inc., Aurora, IL, USA) fitted with a 1/2 inch cone was used to measure the penetration resistance. Dry bulk density was calculated as the mass of oven dried soil per unit soil sample volume (Blake and Hartge, 1986). Soil was oven dried at 105°C until a constant weight was achieved. Irrigation was managed on the basis of soil moisture content to avoid water deficit stress (PR2/6-tubes at ARBC (Delta-T Devices Ltd, Cambridge UK) and multiplexed TDR-100 probes at PSU (Campbell Scientific Inc., Logan, UT, USA). PR2/6-tubes were installed on both the compacted and non-compacted plots at ARBC, while TDR-100 probes were installed at a nearby field site on non-compacted conditions to determine if irrigation was needed (Figure S1: gravimetric water contents were calculated from volumetric water contents and soil dry bulk density). As the ARBC field site was based in the desert, the plots were heavily irrigated, while PSU plots did not require any supplemental irrigation. Nutrients and pesticides were applied based on standard agronomic practices (Table S1).

2.2. Root sampling

When all genotypes were tasselling (55 and 51 days after planting for ARBC (coarse-loam) and PSU (silt-loam) respectively) one soil core was taken from each subplot. Coring tubes (60 cm deep, 5.1 cm diameter) fitted with a plastic sleeve (4.5 cm diameter) were driven into the soil between 2 plants in a row (Trachsel *et al.*, 2013). Cores were stored at 4° C until root washing could be carried out, up to a maximum of 2 weeks. Cores were divided into six 10 cm increments and roots were washed out of the soil over a 850 µm sieve for each depth profile. Roots were temporarily stored in 75% ethanol in water (v/v). Root length per section was measured by scanning roots on a flatbed scanner (Epson Perfection V700 photo, Epson America, Inc., Long Beach, USA) and analysis was carried out with WinRHIZO Pro 2013e system (Regent Systems Inc., Quebec, Canada). Each core increment was captured by images taken at a resolution of 400 dpi (15.75 pxls/mm), speed priority setting off and dust removal on high. Axial roots (nodal, primary, seminal) and lateral roots have been identified as having a diameter >0.6 cm and <1.0 cm for maize respectively (Cahn *et al.*, 1989; Varney *et al.* 1991, Hund *et al.* 2009) and are therefore used instead of the Böhm classification (Böhm, 1979). Using root diameter classes 0 – 0.5, 0.5 – 1.0, 1 – 1.5, 1.5 – 2.0, 2.0 – 2.5, 2.5 – 3.0, 3.0 – 3.5, 3.5 – 4.0, >4 mm permitted discrimination of coarse (> 1 mm diameter) and fine (< 1 mm diameter) roots and attributed coarse roots to nodal root classes from the third node and upward. However individual nodal root classes could not be distinguished from cores as there is no reference to root crown position. Root length measurements (total, coarse and fine) and proportions (coarse and fine) were made for the entire soil core. Root distributions were compared on the basis of root length densities measurements within 10 cm increments. D₉₅ and D₇₅ are the rooting depth above which 95 and 75% of the total root length within a core were located. These rooting depth measurements were calculated by linear interpolation (Schenk and Jackson, 2002). When applied to the coarse root fraction in the core, in order to calculate the coarse rooting depth, these measurements are indicated as D_{95c},

D_{75c}. An overview of the different measurements directly measured or calculated from the winRHIZO scans and their definition can be found in Table 1.

2.3. Plant sampling

Two plants per subplot (4 replicate subplots per compaction treatment) were sampled at tasselling using the 'shovelomics' method (Trachsel *et al.*, 2011). Subsequent measurements per subplot were obtained by averaging between the two harvested plants per subplot. Root crowns were carefully washed and removed from the stem above the brace roots, brace roots not reaching the soil were clipped off at the base of the stem to expose the crown roots. Root crowns were then imaged to obtain information about the root angle in order to establish that root angle did not affect rooting depth (Figure S2). We used a Nikon D70s camera with a sigma DC 18-50mm lens which produced 4000 x 6000 pxl images. Illumination was achieved by white fluorescent indoor lighting with a white cloth cover for light diffusion. Root angle from the horizontal was manually measured in ImageJ (Schneider *et al.*, 2012). Above ground plant parts were dried at 60° C for 3 days and dry weight of the biomass recorded (Figure S3).

2.4. Statistical Analysis

Genotypes were planted in a completely randomised split-plot design with compaction treatment at the whole-plot level (167.26 m²) and twelve genotypes as subplots, replicated four times in each field site. Each subplot was 3.05 m x 4.57 m and ordering of genotypes (subplots) was randomised within each whole-plot. Every subplot was then planted with 4 rows of the appropriate genotype, with 23 cm within row spacing and 76 cm between row spacing reaching a planting density of approximately 57500 plants per hectare. This planting density is sparser than standard agronomic practices which reduced intra-plant competition and aided sampling. All root distribution variables derived from coring (1 soil core per subplot) were transformed using a Box-Cox transformation to achieve normality before analysing the data in a split plot ANOVA. Total root length per genotype was plotted against averaged penetration resistance across both field sites. Root proportions per genotype were plotted per field site and post-hoc comparisons between compacted and non-compacted treatments were carried out using a Tukey HSD test. The same was done for coarse and total rooting depth, where additionally a linear regression was tested between these measurements. A generalised linear model was applied to assess the effect of field site, compaction and genotype and coarse and total root lengths on rooting depth. Relationships between variables were first assessed by correlation plots on pooled data across all genotypes. For the relationship between D₇₅ and D_{75c} an analysis of covariance (ANCOVA) was performed, followed by linear regression. As genotypic effects were identified by the split plot analysis on certain variables, individual linear regressions were made on the averaged genotypic values. Rooting depth data (coarse and total) and total root length averaged per genotype were normally distributed within impedance level and field site datasets. Root proportional data was analysed by using a beta regression (Cribari-Neto and Zeileis, 2015). To further analyse the variable

relationships among each other, a principal component analysis (PCA) was conducted within field site – compaction treatment combinations. Principal components were retained based on eigenvalues greater than 1. In order to investigate different types of root distributions under compacted conditions rooting depth data was plotted against total root length data and genotypes with either similar root length and contrasting depth or with similar depth and contrasting root length were identified. An ANOVA was used to test the effect of genotype, compaction, total rooting depth (D_{75}), total coarse rooting depth (D_{75c}), total root length (TRL) and total fine root length (TRL_f) on aboveground biomass. Graphpad Prism (Graphpad Software, 2017) was used for visualising data and R version 3.5.0 (R Core Team, 2018) was used for data analysis.

3. Results

3.1. Decrease in root length on compacted soil depends on field site

Total root length (TRL) from ARBC (coarse-loam) soil cores was greater than the total root length in PSU (silt-loam) cores in both compacted and non-compacted plots (Figure 2, S4). On coarse-loam (ARBC) total root length was reduced by 47.4% on average across all genotypes when grown on the compacted plots and total root length was clearly reduced for each genotype (Figure 2, Table 2). As total coarse root length represents only a small part of the total root length (Figure 2), total root length reduction on coarse-loam (ARBC) is mainly due to reduced total fine root length (Figure 2, Table 2). In contrast, on silt-loam (PSU), compaction did not significantly alter total root length (Table 2), possibly due to greater penetration resistance on both compacted and non-compacted plots in comparison with the coarse-loam (ARBC) plots (Figure S4).

Total coarse root length (TRL_c) was differentially affected by the compaction treatment at both field sites (Figure 2, Table 2). A compaction x genotype interaction was present on coarse-loam (ARBC), but not on silt-loam (PSU) (Table 2). The overall average of total coarse root length decreased from $48.9 \text{ cm} \pm 3.4 \text{ (se)}$ to $39.0 \text{ cm} \pm 3.3 \text{ (se)}$ under compaction on coarse-loam (ARBC), while it increased from $16.3 \text{ cm} \pm 2.2 \text{ (se)}$ to $23.7 \text{ cm} \pm 3.3 \text{ (se)}$ under compaction on silt-loam (PSU). Total fine root length (TRL_f) was negatively affected by the compaction treatment on coarse-loam (ARBC) (decreasing from $1755.9 \text{ cm} \pm 77.9 \text{ (se)}$ to $809.1 \text{ cm} \pm 37.3 \text{ (se)}$), but was not affected on silt-loam (PSU) (Table 2). At both field sites no genotypic differences were present for total fine root length (Table 2). A positive effect of compaction was noted on overall coarse root proportion (P_c) at ARBC (coarse-loam) with an increase from 2.8% to 4.4% under compacted soil conditions (Table 2, Figure 3). At PSU (silt-loam) we observed an increase from 3.4% to 4.4% for P_c , (Figure 3) although no compaction treatment effect was noted (Table 2). Genotype had significant effect on the proportion of coarse roots (P_c) and fine roots (P_f) at both field sites and for coarse-loam (ARBC) there was an interaction between compaction treatment and genotype present (Table 2). Compaction increased the proportion of coarse roots for most genotypes (Figure 3). The only genotype that had greater P_c under impeded conditions at both field sites was IBM051. Other genotypes manifesting increased P_c under impeded conditions

were OhW122, OhW119, NyH126, IBM146 and IBM059 at ARBC (coarse-loam), but these did not show increased P_c in silt-loam (PSU). In silt-loam (PSU) other genotypes such as OhW128 and IBM284 increased their P_c , while they did not in coarse-loam (ARBC). A genotype that did not have greater coarse root proportions in response to compaction in either location was IBM086, this genotype had similar coarse root proportions in coarse-loam (ARBC), while in silt-loam (PSU) the coarse root proportions appeared smaller under compaction.

3.2. Total rooting depth versus coarse rooting depth

Absolute and relative measures of root length density per depth increment provided insight into how roots were growing within the soil profile and how root distributions change in response to compaction (Figure 4). Root distribution changes became clearer when relative root length density measures were considered (Figure 4B-D). Distributions of coarse root length density and root length density differed (Figure 4) illustrated by differences in values of D_{95} and D_{75} (rooting depth considering all roots diameter classes) versus D_{95c} and D_{75c} (rooting depth considering coarse roots) (D_{75} and D_{75c} shown in figure 4, 5). Total rooting depth and coarse rooting depth measurements were correlated (Figure 5F). D_{75} and D_{75c} were significantly reduced by compaction at both locations, while D_{95} and D_{95c} were only reduced at ARBC (coarse-loam) (Table 2). A genotypic effect on rooting depth was present in coarse-loam (ARBC), but absent in silt-loam (PSU) (Table 2). On coarse-loam (ARBC) some genotypes had significantly shallower total and coarse rooting depths under impeded conditions (IBM014, IBM059, IBM146, OhW119, OhW122) (Figure 5A-C). Other genotypes such as OHW128 had shallower total root length under impeded conditions, but coarse rooting depth was not significantly reduced (Figure 6C). In contrast, IBM323, IBM178, IBM284 and IBM086 had shallower coarse rooting depth under compaction, but total rooting depth was not reduced (Figure 5A-C). On silt-loam (PSU) an effect of compaction was present on D_{75c} and D_{75} (Table 1), however only genotype, IBM059, showed significantly shallower coarse root distributions (Figure 5B).

3.3. Relationships between root distribution variables

Relationships between the different variables can be further explored through the correlation plot across all genotypes (Figure S5, S6, S7) as well as the PCA plots per field site with treatment combination (Figure S8). Individual linear regressions between root distribution variables depicting the different genotypes can be found in Figures 6, S6 and S7. Across all field sites and levels of compaction rooting depth variables (D_{95} , D_{75} , D_{95c} and D_{75c}) positively correlated to each other (Figure S8). Likewise root length variables total root length (TRL), total fine root length (TRL_f) and total coarse root length (TRL_c) correlated strongly with each other (Figure S8). Relationships between rooting depth and the other root distribution variables are discussed below.

3.3.1. The relationship between total rooting depth and other root distribution variables

A negative relationship between total root length and total rooting depth was found under ARBC (coarse-loam) non-compacted conditions at a probability of 0.10 (Figure 6A, S5). General linear modelling indicated integrative effects of field site and compaction on the relationship between total root length and total rooting depth (Table 3). As total root length mainly consists of fine roots, a relationship persists between total fine root length and total rooting depth (Figures S6). No such relationship was seen on silt-loam (PSU) or under compaction (Figures 6B, S6). No relationship was found for total rooting depth and total coarse root length (Figure S6C-D) and coarse root proportion (Figure S6E-F) under any scenario.

3.3.2. The relationship between coarse rooting depth and other root distribution variables

A positive relationship between coarse root proportion and coarse rooting depth (D_{75c}) was present under non-compacted conditions at the coarse loam (ARBC) field site (Figure 6C). This correlation was not observed under compacted conditions, nor at the other field site (Figure 6C-D). Coarse rooting depth was also not correlated with total root length, total coarse root length or total fine root length under any of the field site with compaction treatment combinations (Figures 6E-F, S7). This could also be deduced from the general linear model (Table 3).

3.4. Root length density distributions show field-site dependent genotypic adjustments to compacted conditions

Genotype had an effect on coarse rooting depth on coarse-loam (ARBC) but on silt-loam (PSU) (Table 2). Coarse (Figure 7) and total (Figure S9) root length distributions over the soil profile at PSU (silt-loam) had smaller root length densities than on coarse-loam (ARBC). Distribution differences with depth between genotypes were less evident on silt-loam (PSU) (Figure 7), no significant statistical effect of genotype alone was noted on D_{75} or D_{75c} (Table 2), which could be attributed to larger standard errors at PSU (Figure 5D) and generally less roots found. The general linear model (Table 3) shows that genotype in combination with other factors did have a significant effect on D_{75c} . Different genotypes were found from each field site that showed similar coarse root length but contrasted in root depth or showed similar coarse root depth but contrasted in total root length. (Figure S10). For these measurements we observed genotypes with similar total coarse root length, but different rooting depths representing shallow and deeper root systems with similar root system sizes (IBM284 versus IBM323 for coarse loam (ARBC) and IBM051 and OhW122 for silt-loam (PSU)) (Figure 8A). A similar analysis was carried out based on total root length and depth (Figure S11, S12).

3.5. Relationship between root distribution variables and aboveground biomass

Compaction influenced aboveground biomass significantly (Figure 9, S2) but could also be related to other root measurements derived from coring. For both coarse-loam (ARBC) and silt-loam (PSU) soils, plant biomass was higher when coarse roots were

able to grow deeper due to being less impeded (Figure 9). In coarse-loam (ARBC) however TRL_f also played a role, while in silt-loam (PSU) greater plant biomass was reached on non-compacted plots without TRL_f having a role.

4. Discussion

In this study, on two different soils with compacted and non-compacted plots, we found total root length reduction due to compaction f was field site dependent (Figures 2, Table 2). Coarse root proportions were influenced by genotype at both field sites (Figure 4, Table 2). Rooting depth of coarse and total roots were strongly correlated (Figure 5). Root length and rooting depth variables were not correlated when plants were grown on compacted plots (Figures 6, S5, S6, S7, S8). Our results support the hypothesis that the ability of roots to grow to depth through compacted soils is not dependent on the amount of roots formed by the root system. Furthermore, we propose that root length density distributions are either characterised by avoidance or by adaptive strategies for different genotypes when grown in compaction.

4.1. Root phenotypes show high levels of plasticity

4.1.1. Field site effects on root systems

Total root lengths (TRL), total fine root lengths (TRL_f) and total coarse root lengths (TRL_c) were greater on coarse-loam (ARBC) than on silt-loam (PSU) (Figures 2, S6). The significant reduction of the fine root length due to compaction on coarse-loam (ARBC) could influence the proportions of fine and coarse roots. Greater changes in coarse root proportions were observed at ARBC (coarse-loam) versus PSU (silt-loam) (Figure 3), which could potentially be driven by a disproportionally greater reduction of total fine root length versus that of total coarse root length (causing a shift towards greater proportion of coarse roots). Rooting depths $D75$ and $D75_c$ were different at the two field sites (Figures 5, S6, S7, Table 3). Differences between field sites for observations considering root length, root proportions and root depth could be related to differences in soil parameters. Maize seedlings had significantly longer seminal roots in a sandy loam versus a sandy clay loam (Panayiotopoulos *et al.*, 1994) while rooting depths of grapevines were deeper in coarse textured soils than fine textured soils (Nagarajah, 1987). Greater root length was possibly attained on coarse-loam (ARBC) because of the greater sand fraction in the soil versus silt-loam (PSU). Another possible explanation for the root length differences between coarse- and silt-loam could be a difference in root-soil contact between the field sites. On the non-compacted plots of PSU (silt-loam), smaller dry bulk densities could mean reduced levels of root-soil contact, which in turn reduces water and nutrient uptake (Veen *et al.*, 1992). The coarse-loam field site (ARBC) consisted of a more uniform, less structured soil, while the silt-loam field site (PSU) had more pronounced soil structure in terms of aggregation observed in the field. Roots can take advantage of cracks or bio-pores from earthworms or old root channels present to bypass compacted layers (Atwell, 1993; Hatano *et al.*, 1988; Stirzaker *et al.*, 1996). Cracks and pores will impose lower axial pressures on roots than bulk soil (Jin *et al.*, 2013). It is likely that the

presence of such low-resistance channels in the soil structure at PSU (in silt-loam) could have permitted deeper rooting than at ARBC (coarse-loam).

4.1.2. Compaction influences root system distribution

Compaction influenced root growth at both field sites, but more significantly at ARBC (coarse-loam), where all rooting variables were significantly affected (Table 2). At both field sites the compaction treatment influenced the average total coarse root length across genotypes in different ways (Figure 3, Table 2). Total coarse root length decreased on coarse-loam (ARBC), which could be due to the effect compaction had on root system size in general. Total and fine root length were more significantly reduced than total coarse root length under compaction (Table 2, Figure 2). Reductions in root length in compacted soil has been reported for different species including maize (Grzesiak, 2009; Iijima and Kono, 1991). At PSU (silt-loam) total and fine root length were not significantly affected by compaction (Table 2) and total coarse root length increased (Figure 3, Table 2). Increased total coarse root length could potentially be caused by radial expansion as roots generally increase in diameter when experiencing mechanical impedance. Elongation is slowed compared to elongation rates at lower levels of mechanical impedance, which in turn decreases root length (Bengough *et al.*, 2006; Bengough and Mullins, 1991; Bengough and Young, 1993). While all root length measures decreased on coarse-loam (ARBC), these observations on root length were different on silt-loam (PSU) where fine and coarse roots were differentially affected by the compaction treatment. Coarser roots such as seminal or nodal root axes were more impeded than lateral roots possibly reflecting the fewer macropores present under compacted conditions. Such effects have been found in barley growing in glass ballotini of different sizes, with larger pores only restricting seminal growth and smaller pores restricting both laterals and seminal growth of barley (Goss, 1977). Laterals capable of growing in pores larger than their own diameters would encounter less impedance than those laterals forced to grow through bulk soil or smaller pores (Iijima and Kono, 1991).

Under compaction both rooting depth (D_{75}) and rooting depth of coarse roots (D_{75c}) decreased at both field sites (Tables 1, 2; Figures 5, 6, S6, S7). Reduction of D_{75c} due to compaction could be linked to reduced aboveground biomass (Figure 9). Shallower rooting depths probably reflect slower root elongation rates, so it will take longer for a root to reach deep soil strata. Smaller differences in rooting depth of compacted and non-compacted plots at PSU (silt-loam) (Figure 5) could be due to the smaller differences in penetration resistance with increasing depth versus ARBC (coarse-loam) (Figure 1). Roots at the PSU field site (silt-loam) would initially experience greater levels of mechanical impedance, but once they pass this zone should be able to elongate more normally. The reduction in rooting depth under compaction is in agreement with observations with wheat (Barracough and Weir, 1988; Chen *et al.*, 2014). Compaction altered root distribution, generally shifting root distribution to shallower strata (Figures 4, 5, 7, S9). Multiple studies have described similar redistributions of roots under impeded field conditions for various crops (Barracough

and Weir, 1988; Brereton *et al.*, 1986; Chen and Weil, 2011; Chen *et al.*, 2014). For maize specifically, roots of 2-3 week old plants were confined to surface layers under compaction (Veen and Boone, 1990). A similar observation was made during a 4 week growing period for maize grown in root boxes (Iijima *et al.*, 1991) and in the field up to tasselling (Laboski *et al.*, 1998). Soil compaction reduces soil porosity, hydraulic conductivity and air permeability and this increases the risk of hypoxia (Laboski *et al.*, 1998; Kuncuro *et al.*, 2014; Xiong *et al.*, 2020). In this set-up of compacting entire field sites, the stress caused by impedance and hypoxia cannot be mutually excluded. Therefore hypoxia might also have played a role in reducing growth of the root system. A recent studies by Xiong *et al.* (2020) illustrates that higher moisture contents negatively influenced root growth in compacted soils.

4.1.3. Compaction influenced genotypes differently

Under compaction at both field sites most genotypes had a greater proportion of coarse roots (Figure 3) and genotypes differed in this response (Table 3). An increased proportion of coarse roots could either be attributed to (1) the reduction of the fine root proportion, (2) the increase in diameter of roots grown under impeded conditions due to thickening or (3) a combination of the two. On coarse-loam (ARBC), total fine root length was significantly reduced (Table 3, Figure 2), which in turn would influence root proportions. However, as there was no such reduction of total fine root length on silt-loam (PSU), root thickening, which is the increase in radial diameter of the roots, is probably the main cause of a shift in root proportions on silt-loam.

Genotypic differences were found for total and coarse rooting depth variables (Figures 5, 7, 8, S6, S7, Tables 2, 3). No relationship between root length variables existed (with the exception of negative relationship between total root length and total rooting depth under non-compacted conditions at ARBC (coarse-loam)). Deeper rooting was not associated with root system size. Root phenes that have been found to contribute to overcoming impedance include anatomical traits such as reduced cell file number and increased levels of aerenchyma (Lynch and Wojciechowski, 2015; Vanhees *et al.*, 2020) and is thought to be linked to reduction of metabolic costs, reduced energy and/or facilitating O₂ diffusion (Hanbury and Atwell, 2005; Lynch, 2015; Colombi *et al.*, 2019). It has also been suggested that anatomical traits such as smaller outer cortical region cells will stabilize a root during the penetration of a harder soil layers (Chimungu *et al.* 2015). Other phenes are sharper root tip shape, the presence of root hairs, the production of mucilage, root cap sloughing and steeper growth angles (Bengough *et al.*, 2011; Colombi *et al.*, 2017b; Haling *et al.*, 2013; Iijima *et al.*, 2000, 2004; Jin *et al.*, 2013, Lynch and Wojciechowski, 2015). How all these phenes can synergistically work together is worthy of further investigation. Other studies have shown that plasticity could play a role, for instance for rice it was observed that upon rewetting after drought differences in nodal root elongation through a hardpan were present between genotypes (Suralta *et al.*, 2018).

4.2. The relationship between root length and root depth varies among genotypes

Root length and rooting depth are not related under impeded conditions (Figures 6, S6, S7, S8). Coarse rooting depth, representing nodal roots, responded differently to compaction among genotypes. Genotypes such as IBM059 or IBM323 at ARBC (coarse-loam) or IBM086 at PSU (silt-loam) grew deeper under non-compacted conditions but reduced their coarse root length under compaction (Figure 5A-B). IBM178 grew intermediately deep in coarse-loam (at ARBC), and deep in silt-loam (at PSU) but did not reduce its coarse rooting depth to the same extent as the aforementioned genotypes (Figure 5A-B). This suggests that coarse roots of some genotypes were obstructed by the compaction treatment while coarse roots of other genotypes were capable of growing through.

We found genotypes with similar root system size that reached different rooting depths (Figure 8A, S12) as well as genotypes with similar deeper coarse rooting depth but with different total coarse root length (Figure 8B, S12). Coarse rooting depths can thus be reached in different ways as the root system with smaller values for coarse root length densities were able to grow as deep as the root systems that have greater coarse root length density at depth. Shoots can be sustained by different root system sizes and rooting depths as long as water and nutrients are available. Therefore rooting depth under compaction is not simply related to the amount of roots formed. In the following section we discuss how each type of root distribution with depth could sustain plant growth.

4.2.1. Root systems with equal coarse root length reach different depths

Coarse roots of some genotypes were obstructed, while others managed to grow through impeded soil domains and reached deeper strata (Figure 8A). If nodal roots are sufficiently impeded, these and any laterals roots emerging from them will automatically be located within the upper soil strata. However laterals may grow downwards from a shallow starting point when they experience less impedance than nodal roots by, for instance, making use of smaller pores (Goss, 1977). Increased lateral branching has been observed in the non-impeded parts of the soil (Montagu *et al.*, 2001) and will enable a plant to extract water and nutrients when root length is maintained and sufficient soil resources are available in the unimpeded soil (Barracough and Weir, 1988). Compensatory root growth introduces more roots in the less impeded domains, often in the upper soil strata (Barracough and Weir, 1988; Materechera *et al.*, 1993; Nosalewicz and Lipiec, 2014; Pfeifer *et al.*, 2014a). A similar redistribution can be seen in the compacted plots (Figures 7, S9). Compensatory mechanisms may influence nutrient and water acquisition.

Rooting depth has been linked to water acquisition, especially under drought conditions where deeper rooting increases yield (Gao and Lynch, 2016; Hund *et al.*, 2009; Lynch, 2013, 2018; Chimungu *et al.*, 2014a, 2014b; Lynch *et al.* 2014; Zhan *et al.*, 2015). Our study did not employ a water deficit, but it has been shown that compaction can make water deficit stress more severe (Grzesiak *et al.*, 2014): even without the presence of water deficit stress, increased water uptake from the topsoil can be present on compacted soils. This in turn will increase the penetration resistance

within the topsoil and further limit access to the deeper soil layers (Colombi *et al.*, 2018). A root system that is limited to shallow soil strata will thus be more at risk for water deficit both in terms of reduced access, as well as increased water depletion in its local soil environment. It is likely that in the coarse soil (ARBC) compacted treatment the air-filled porosity was often <10% in the surface 20cm, but generally >10% in the other treatments. The root system might therefore have been influenced by poor oxygen availability (hypoxia) however this can be seen as common feature of compacted soils in the field where porosity is reduced. Clear shifts in root distribution occurred in our field sites, we observed changes in root proportion, changes in rooting depth, and changes in root distribution (Figures 3, 5, 7, S9). How these shifts influence resource acquisition under impeded field conditions merits further investigation.

Stresses such as waterlogging have been found to have a more severe impact in impeded soils (Grzesiak *et al.*, 2014). Environmental effects such as temperature fluctuations or soil drying by direct evaporation pose additional threats to more shallow root systems (Lynch, 2018). Overall compensatory root growth can be seen as a stress avoidance strategy as plants come less into direct contact with the impeded soil regions and grow where impedance is lower. This can be considered as an indirect adaptation or response to the impeding conditions. As soils are a typically heterogeneous, roots could take advantage of cracks or pores present to bypass compacted layers (Hatano *et al.*, 1988; Atwell, 1993; Stirzaker *et al.*, 1996) which would be another avoidance strategy. Rasse and Smucker (1998) showed that maize can make use of root-induced macropores from a previous alfalfa crop. Preferential growth towards artificial pores has been observed in compacted soils (Stirzaker *et al.*, 1996; Pfeifer *et al.*, 2014b; Colombi *et al.*, 2017a; Atkinson *et al.*, 2020) but whether this is due to oxytropism or locally reduced penetration rates is still under debate, although promising attempts have been made to model the process (Landl *et al.*, 2017). Roots adapted to impedance are characterised by traits that help them overcome impedance, enabling them to grow better in harder soils. Those genotypes capable of rooting deeper and of overcoming impedance stress are at less at risk of nutrient deficiencies, of lack of access to water and of other environmental stresses.

4.2.2. Equal depths can be reached by root systems of different sizes

We observed genotypes that contrasted in root system size (measured as total coarse root length) were able to reach similar depths on compacted plots (Figure 8B, S12). Greater amounts of coarse roots (measured as greater TRL_c) would be found when a maize plant forms more root axes per nodal position, additionally greater amounts of coarse roots may also be caused by root thickening. We found that no rooting depth measure correlated with TRL_c (Figures 6E-F, S6A-D, S7). The ability of a root system to grow deeper in compacted soils is therefore not dependent on the amount of roots formed as both large and parsimonious root systems can reach similar depths on compacted plots at both field sites. This is in contrast with observations on wheat, where penetrability of a harder wax layer was related to amount of root axis formed (Whalley *et al.*, 2013), or that denser root systems of lupin are deeper rooting (Chen

et al., 2014). On the other hand, comparisons between species show that species with a larger number of roots in the top layers of a layered medium did not automatically have greater penetration rates through the compacted layer (Materechera *et al.*, 1993). A field study with two rice varieties showed that varieties with a greater root density were able to root deeper under control conditions, but under greater penetrometer resistances became more strongly affected than others with lower rooting densities (Cairns *et al.*, 2004).

The formation of more roots can have benefits such as increased foraging for water and nutrients or reduced risk of root loss due to pests and diseases (Lynch, 2003, 2018, 2019). Increased root formation can however come at a substantial costs (York *et al.*, 2013, Lynch, 2003). Greater elongation rate, greater root diameter, increased branching or greater formation of axial roots increase the metabolic cost of the root system (York *et al.*, 2013, Lynch, 2018). Second, the formation of too many roots will introduce competition for internal and external resources (Lynch, 2018). Excessive root formation not only induces intraplant competition for resources, it also increases root maintenance and formation costs. Other traits, such as increased aerenchyma formation, large cortical cell size, reduced cortical cell file number or reduced crown root number bring costs down (Lynch, 2003, 2018) which would enable these plants to allocate resources elsewhere. For instance, it has been shown that maize with fewer crown roots are able to allocate roots deeper (Saengwilai *et al.*, 2014; Gao and Lynch, 2016, Lynch 2018). Recent experiments by Guo and York (2019) showed excising nodal roots stimulated greater shoot biomass and root length at depth under low N conditions as biomass was reallocated to lateral and early nodal roots. Under impeded conditions, metabolic cost reduction might be significant. A recent study by Colombi (2019) found energy costs were linked to cortical cell diameters, with greater cell diameters reducing the metabolic cost under impeded conditions. As both large as well as parsimonious root systems were able to reach similar coarse root depth (Figures 8B, S12B) we suggest that parsimonious phenotypes could potentially allocate more resources to shoot growth. This effect could be apparent under high input systems, where improved conversion of soil resources to yield would be greater for parsimonious phenotypes (Lynch and Brown, 2012; Lynch, 2018).

Conclusions

Rooting depth and root length were not correlated under impeded conditions. Different coarse rooting depths were reached by genotypes characterised by similar root system sizes. We suggest genotypes better adapted to impedance (and therefore rooting deeper) are less at risk of additional stresses such as nutrient deficiency, soil drying, lack of access to water and other environmental conditions. We hypothesise that excessive root formation will introduce greater competition for internal and external resources, furthermore larger root systems have greater metabolic costs associated with them. We also suggest that parsimonious phenotypes will be able to steer resource allocation to shoot growth and improved yields. We found that the

617 amount of roots formed by the root system does not determine the ability of those roots
618 to grow deeper under impeded conditions.

619 **Acknowledgements**

620 This research was supported by the Howard G. Buffett Foundation, the University of
621 Nottingham and the James Hutton Institute. The James Hutton Institute receives
622 funding from the Rural & Environment Science & Analytical Services Division of the
623 Scottish Government. We thank Hannah Schneider, Stephanie Klein, Chris Strock,
624 Kimo Jin and other members of the PSU roots lab for their help and support during the
625 execution of these field trials.

626

References

- Aggarwal, P., Choudhary, K.K., Singh, A.K., Chakraborty, D., 2006. Variation in soil strength and rooting characteristics of wheat in relation to soil management. *Geoderma* 136, 353–363. doi:10.1016/j.geoderma.2006.04.004
- Atkinson, J.A., Hawkesford, M.J., Whalley, W.R., Zhou, H., Mooney, S.J., 2020. Soil strength influences wheat root interactions with soil macropores. *Plant Cell Environ.* 43, 235–245. doi: 10.1111/pce.13659
- Atwell, B.J., 1993. Response of roots to mechanical impedance. *Environ. Exp. Bot.* 33, 27–40. doi:10.1016/0098-8472(93)90053-I
- Barracough, P.B., Weir, A.H., 1988. Effects of a compacted subsoil layer on root and shoot growth, water use and nutrient uptake of winter wheat. *J. Agric. Sci.* 110, 207–216. doi:10.1017/S0021859600081235
- Batey, T., 2009. Soil compaction and soil management - A review. *Soil Use Manag.* 25, 335–345. doi:10.1111/j.1475-2743.2009.00236.x
- Beemster, G.T.S., Masle, J. 1996. Effects of soil resistance to root penetration on leaf expansion in wheat (*Triticum aestivum* L.): composition, number and size of epidermal cells in mature blades. *J. Exp. Bot.* 47, 1651–1662. doi: 10.1093/jxb/47.11.1651
- Bengough, A.G., Bransby, M.F., Hans, J., McKenna, S.J., Roberts, T.J., Valentine, T.A., 2006. Root responses to soil physical conditions; growth dynamics from field to cell. *J. Exp. Bot.* 57, 437–447. doi:10.1093/jxb/erj003
- Bengough, A.G., McKenzie, B.M. 1997. Sloughing of root cap cells decreases the frictional resistance to maize (*Zea mays* L.) root growth. *J. Exp. Bot.* 48, 885–893. doi:10.1093/jxb/48.4.885
- Bengough, A.G., McKenzie, B.M., Hallett, P.D., Valentine, T.A., 2011. Root elongation, water stress, and mechanical impedance: a review of limiting stresses and beneficial root tip traits. *J. Exp. Bot.* 62, 59–68. doi:10.1093/jxb/erq350
- Bengough, A.G., Mullins, C.E., 1991. Penetrometer resistance, root penetration resistance and root elongation rate in two sandy loam soils. *Plant Soil* 131, 59–66. doi:10.1007/BF00010420
- Bengough, A.G., Young, I.M., 1993. Root elongation of seedling peas through layered soil of different penetration resistances. *Plant Soil* 149, 129–139. doi:10.1007/BF00010770
- Bennie, A.T.P., 1996. Growth and mechanical impedance, in: *Plant Roots: The Hidden Half*. pp. 453–470.
- Bingham, I.J., Bengough, A.G., 2003. Morphological plasticity of wheat and barley roots in response to spatial variation in soil strength. *Plant Soil* 250, 273–282. doi:10.1023/A:1022891519039
- Blake, G.R., Hartge, K.H. 1986. Bulk density in: Klute, A. (Eds.), *Methods of soil analysis: Part 1 Physical and mineralogical methods*. American Society of Agronomy, Inc; Soil Science Society of America, Inc., Madison, Wisconsin, pp.

669 363-376.

670 Botwright Acuña, T.L., Wade, L.J., 2005. Root penetration ability of wheat through
671 thin wax-layers under drought and well-watered conditions. *Aust. J. Agric. Res.*
672 56, 1235–1244. doi:10.1071/AR05067

673 Brereton, J.C., McGowan, M., Dawkins, T.C., 1986. The sensitivity of barley, field
674 beans and sugar beet to soil compaction. *F. Crop. Res.* 13, 223–237.

675 Böhm, W. (1979). *Methods of studying root systems*. Ecological studies. Springer,
676 Berlin.

677 Cahn, M.D., Zobel, R.W., Bouldin, D.R., 1989. Relationship between root elongation
678 rate and diameter and duration of growth of lateral roots of maize. *Plant Soil*
679 119, 271–279. doi:10.1007/BF02370419

680 Cairns, J.E., Audebert, A., Townend, J., Price, A.H., Mullins, C.E., 2004. Effect of soil
681 mechanical impedance on root growth of two rice varieties under field drought
682 stress. *Plant Soil* 267, 309–318. doi:10.1007/s11104-005-0134-1

683 Carminati, A., Passioura, J.B., Zarebanadkouki, M., Ahmed, M.A., Ryan, P.R., Watt,
684 M., Delhaize, E. 2017. Root hairs enable high transpiration rates in drying soils.
685 *New Phytol.* 216, 771-781. doi:10.1111/nph.14715

686 Chandra Babu, R., Shashidar, H.E., Lilley, J.M., Thanh, N.D., Ray, J.D., Sadasivam,
687 S., Sarkarung, S., O'Toole, J.C., Nguyen, H.T., 2001. Variation in root
688 penetration ability, osmotic adjustment and dehydration tolerance among
689 accessions of rice adapted to rainfed lowland and upland ecosystems. *Plant*
690 *Breed.* 120, 233–238. doi:10.1046/j.1439-0523.2001.00578.x

691 Chen, G., Weil, R.R., 2011. Root growth and yield of maize as affected by soil
692 compaction and cover crops. *Soil Tillage Res.* 117, 17–27.
693 doi:10.1016/j.still.2011.08.001

694 Chen, Y.L., Palta, J., Clements, J., Buirchell, B., Siddique, K.H.M., Rengel, Z., 2014.
695 Root architecture alteration of narrow-leaved lupin and wheat in response to soil
696 compaction. *F. Crop. Res.* 165, 61–70. doi:10.1016/j.fcr.2014.04.007

697 Chimungu, J.G., Brown, K.M., Lynch, J.P., 2014a. Large root cortical cell size
698 improves drought tolerance in maize. *Plant Physiol.* 166, 2166–2178.
699 doi:10.1104/pp.114.250449

700 Chimungu, J.G., Brown, K.M., Lynch, J.P., 2014b. Reduced root cortical cell file
701 number improves drought tolerance in maize. *Plant Physiol.* 166, 1943-1955.
702 doi:10.1104/pp.114.249037

703 Chimungu, J.G., Loades, K.W., Lynch, J.P., 2015. Root anatomical phenes predict
704 root penetration ability and biomechanical properties in maize (*Zea Mays*). *J.*
705 *Exp. Bot.* 66, 3151–3162. doi:10.1093/jxb/erv121

706 Chochois, V., Vogel, J.P., Rebetzke, G.J., Watt, M., 2015. Variation in Adult Plant
707 Phenotypes and partitioning among seed and stem-borne roots across
708 *Brachypodium distachyon* accessions to exploit in breeding cereals for well-
709 watered and drought environments . *Plant Physiol.* 168, 953–967.
710 doi:10.1104/pp.15.00095

711 Clark, L.J., Aphale, S.L., Barraclough, P.B., 2000. Screening the ability of rice roots
712 to overcome the mechanical impedance of wax layers: Importance of test
713 conditions and measurement criteria. *Plant Soil* 219, 187–196.
714 doi:10.1023/A:1004753900945

715 Clark, L.J., Cope, R.E., Whalley, W.R., Barraclough, P.B., Wade, L.J., 2002. Root
716 penetration of strong soil in rainfed lowland rice: Comparison of laboratory
717 screens with field performance. *F. Crop. Res.* 76, 189–198. doi:10.1016/S0378-
718 4290(02)00039-4

719 Colombi, T., Braun, S., Keller, T., Walter, A., 2017a. Artificial macropores attract crop
720 roots and enhance plant productivity on compacted soils. *Sci. Tot. Env.* 574,
721 1283-1293. doi: 10.1016/j.scitotenv.2016.07.194

722 Colombi, T., Herrmann A.M., Valenback P, Keller T. 2019. Cortical cell diameter is
723 key to energy costs of root growth in wheat. *Plant Physiol.* 180, 2049-2060.
724 doi:10.1104/pp.19.002262

725 Colombi, T., Kirchgessner, N., Walter, A., Keller, T., 2017b. Root tip shape governs
726 root elongation rate under increased soil strength. *Plant Physiol.* 174, 2289–
727 2301. doi:10.1104/pp.17.00357

728 Colombi, T., Torres, L.C., Walter, A., Keller, T., 2018. Feedbacks between soil
729 penetration resistance, root architecture and water uptake limit water
730 accessibility and crop growth – A vicious circle. *Sci.Total Environ.* 626, 1026–
731 1035. doi:10.1016/j.scitotenv.2018.01.129

732 Colombi, T., Walter, A., 2016. Root responses of triticale and soybean to soil
733 compaction in the field are reproducible under controlled conditions. *Funct. Plant*
734 *Biol.* 43, 114–128. doi:10.1071/FP15194

735 Cribari-Neto, F., Zeileis, A., 2015. Beta regression in R . *J. Stat.I Soft.* 34.
736 doi:10.18637/jss.v034.i02

737 Croser, C., Bengough, A.G., Pritchard, J., 2000. The effect of mechanical impedance
738 on root growth in pea (*Pisum sativum*). II. Cell expansion and wall rheology
739 during recovery. *Physiol. Plant.* 109, 150–159. doi:10.1034/j.1399-
740 3054.2000.100207.x

741 de Moreas, M.T., Debiasi, H., Franchini, J.C., Mastroberti, A.A., Levien, R., Leitner,
742 D., Schnepf, A., 2020. Soil compaction impacts soybean root growth in an
743 Oxisol from subtropical Brazil. *Soil Tillage Res.* 200, 104611. doi:
744 10.1016/j.still.2020.104611

745 Ehlers, W., Köpke, U., Hesse, F., Böhm, W., 1983. Penetration resistance and root
746 growth of oats in tilled and untilled loess soil. *Soil Tillage Res.* 3, 261–275.
747 doi:10.1016/0167-1987(83)90027-2

748 Gao, W., Hodgkinson, L., Jin, K., Watts, C.W., Ashton, R.W., Shen, J., Ren, T.,
749 Dodd, I.C., Binley, A., Phillips, A.L., Hedden, P., Hawkesford, M.J., Whalley,
750 W.R., 2016. Deep roots and soil structure. *Plant Cell Environ.* 39, 1662–1668.
751 doi:10.1111/pce.12684

752 Gao, W., Watts, C.W., Ren, T., Whalley, W.R., 2012. The effects of compaction and
753 soil drying on penetrometer resistance. *Soil Tillage Res.* 125, 14–22.

754 doi:10.1016/j.still.2012.07.006

755 Gao, Y., Lynch, J.P., 2016. Reduced crown root number improves water acquisition
756 under water deficit stress in maize (*Zea mays* L.). J. Exp. Bot. 67, 4545–4557.
757 doi:10.1093/jxb/erw243

758 Goss, M.J., 1977. Effects of mechanical impedance on root growth in barley
759 (*Hordeum vulgare* L.): I. Effects on the elongation and branching of seminal root
760 axes. J. Exp. Bot. 28, 96–111. doi:10.1093/jxb/28.1.96

761 Grzesiak, M.T., 2009. Impact of soil compaction on root architecture, leaf water
762 status, gas exchange and growth of maize and triticale seedlings. Plant Root 3,
763 10–16. doi:10.3117/plantroot.3.10

764 Grzesiak, M.T., Ostrowska, A., Hura, K., Rut, G., Janowiak, F., Rzepka, A., Hura, T.,
765 Grzesiak, S., 2014. Interspecific differences in root architecture among maize
766 and triticale genotypes grown under drought, waterlogging and soil compaction.
767 Acta Physiol. Plant. 36, 3249–3261. doi:10.1007/s11738-014-1691-9

768 Guo, H., York, L.M., 2019. Maize with fewer nodal roots allocates mass to more
769 lateral and deep roots that improve nitrogen uptake and shoot growth. J. Exp.
770 Bot. <https://doi.org/10.1093/jxb/erz258>

771 Haling, R.E., Brown, L.K., Bengough, A.G., Young, I.M., Hallett, P.D., White, P.J.,
772 George, T.S., 2013. Root hairs improve root penetration, root-soil contact, and
773 phosphorus acquisition in soils of different strength. J. Exp. Bot. 64, 3711–3721.
774 doi:10.1093/jxb/ert200

775 Hamza, M.A., Anderson, W.K., 2005. Soil compaction in cropping systems: A review
776 of the nature, causes and possible solutions. Soil Tillage Res. 82, 121–145.
777 doi:10.1016/j.still.2004.08.009

778 Hanbury, C.D., Atwell, B.J., 2005. Growth dynamics of mechanically impeded lupin
779 roots: does altered morphology induce hypoxia? Ann. Bot. 96, 913–924. doi:
780 10.1093/aob/mci243

781 Hatano, R., Iwanaga, K., Okajima, H., Sakuma, T., 1988. Relationship between the
782 distribution of soil macropores and root elongation. Soil Sci. Plant Nutr. 34, 535–
783 546. doi:10.1080/00380768.1988.10416469

784 Hochholdinger, F., Woll, K., Sauer, M., Dembinsky, D., 2004. Genetic dissection of
785 root formation in maize (*Zea mays*) reveals root-type specific developmental
786 programmes. Ann. Bot. 93, 359–368. doi:10.1093/aob/mch056

787 Hoffmann, C., Jungk, A. 1995. Growth and phosphorus supply of sugar beet as
788 affected by soil compaction and water retention. Plant Soil 176, 15–25. doi:
789 10.1007/BF00017671

790 Horn, R., Domzal, H., Slowińska-Jurkiewicz, A., van Ouwerkerk, C., 1995. Soil
791 compaction processes and their effects on the structure of arable soils and the
792 environment. Soil Tillage Res. 35, 23–36. doi.org/10.1016/0167-1987(95)00479-
793 C

794 Hund, A., Ruta, N., Liedgens, M., 2009. Rooting depth and water use efficiency of
795 tropical maize inbred lines, differing in drought tolerance. Plant Soil 318, 311–
796 325. doi:10.1007/s11104-008-9843-6

797 Iijima, M., Griffiths, B., Bengough, A.G., 2000. Sloughing of cap cells and carbon
798 exudation from maize seedling roots in compacted sand. *New Phytol.* 145, 477–
799 482. doi:10.1046/j.1469-8137.2000.00595.x

800 Iijima, M., Higuchi, T., Barlow, P.W., 2004. Contribution of root cap mucilage and
801 presence of an intact root cap in maize (*Zea mays*) to the reduction of soil
802 mechanical impedance. *Ann. Bot.* 94, 473–477. doi:10.1093/aob/mch166

803 Iijima, M., Kono, Y., 1991. Interspecific differences of the root system structures of
804 four cereal species as affected by soil compaction. *Japan. Jour. Crop Sci.* 60,
805 130–138. doi:10.1626/jcs.60.130

806 Iijima, M., Kono, Y., Yamauchi, A., Pardales, J.R., 1991. Effects of soil compaction
807 on the development of rice and maize root systems. *Environ. Exp. Bot.* 31, 333–
808 342. doi:10.1016/0098-8472(91)90058-V

809 Jia, X., Liu, P., Lynch, J.P., 2018. Greater lateral root branching density in maize
810 improves phosphorus acquisition from low phosphorus soil. *J. Exp. Bot.* 69,
811 4961–4790. doi:10.1093/jxb/ery252

812 Jin, K., Shen, J., Ashton, R.W., Dodd, I.C., Parry, M.A.J., Whalley, W.R., 2013. How
813 do roots elongate in a structured soil? *J. Exp. Bot.* 64, 4761–4777.
814 doi:10.1093/jxb/ert286

815 Kirkegaard, J.A., So, H.B., Troedson, R.J., Wallis, E.S. 1992. The effect of
816 compaction on the growth of pigeonpea on clay soils. I. Mechanisms of crop
817 response and seasonal effects on a vertisol in a sub-humid environment. *Soil*
818 *Tillage Res.* 24, 107–127. doi: 10.1016/0167-1987(92)90096-T

819 Konôpka, B., Pagès, L., Doussan, C., 2009. Soil compaction modifies morphological
820 characteristics of seminal maize roots. *Plant, Soil Environ.* 55, 1–10. doi:
821 10.17221/380-PSE

822 Kubo, K., Iwama, K., Yanagisawa, A., Watanabe, Y., Terauchi, T., Jitsuyama, Y.,
823 Mikuma, T., 2006. Genotypic variation of the ability of root to penetrate hard soil
824 layers among Japanese wheat cultivars. *Plant Prod. Sci.* 9, 47–55.
825 doi:10.1626/Pps.9.47

826 Kuchenbuch, R.O., Barber, S.A., 1988. Significance of temperature and precipitation
827 for maize root distribution in the field. *Plant Soil* 106, 9–14.
828 doi:10.1007/BF02371189

829 Kuncoro, P.H., Koga, K., Satta, N., Muto, Y., 2014. A study on the effect of
830 compaction on transport properties of soil gas and water. II: Soil pore structure
831 indices. *Soil Tillage Res.* 143, 180–187. doi:10.1016/j.still.2014.01.008

832 Laboski, C.A.M., Dowdy, R.H., Allmaras, R.R., Lamb, J.A., 1998. Soil strength and
833 water content influences on corn root distribution in a sandy soil. *Plant Soil* 203,
834 239–247. doi:10.1023/A:1004391104778

835 Landl, M., Huber, K., Schnepf, A., Vanderborght, J., Javaux, M., Bengough, A.G.,
836 and Vereecken, H. 2017. A new model for root growth in soil with macropores.
837 *Plant Soil* 415, 99–116. doi:10.1007/s11104-016-3144-2

838 Lilley, J.M., Kirkegaard, J.A., 2016. Farming system context drives the value of deep
839 wheat roots in semi-arid environments. *J. Exp. Bot.* 67, 3665–3681.

840 doi:10.1093/jxb/erw093

841 Lin, L.R., He, Y.B., Chen, J.Z., 2016. The influence of soil drying- and tillage-induced
842 penetration resistance on maize root growth in a clayey soil. *J. Integr. Agric.* 15,
843 1112–1120. doi:10.1016/S2095-3119(15)61204-7

844 Lipiec, J., Horn, R., Pietrusiewicz, J., Siczek, A., 2012. Effects of soil compaction on
845 root elongation and anatomy of different cereal plant species. *Soil Tillage Res.*
846 121, 74–81. doi:10.1016/j.still.2012.01.013

847 Loades, K.W., Bengough, A.G., Bransby, M.F., Hallett, P.D., 2013. Biomechanics of
848 nodal, seminal and lateral roots of barley: Effects of diameter, waterlogging and
849 mechanical impedance. *Plant Soil* 370, 407–418. doi:10.1007/s11104-013-
850 1643-y

851 Lynch, J.P., 2003. Rhizoeconomics: Carbon costs of phosphorus acquisition. *Plant*
852 *Soil* 269, 45-56 doi:10.1007/s11104-004-1096-4

853 Lynch, J.P., 2013. Steep, cheap and deep: An ideotype to optimize water and N
854 acquisition by maize root systems. *Ann. Bot.* 112, 347–357.
855 doi:10.1093/aob/mcs293

856 Lynch, J.P., 2015. Root phenes that reduce the metabolic costs of soil exploration:
857 Opportunities for 21st century agriculture. *Plant Cell Environ.* 38, 1775–1784.
858 doi:10.1111/pce.12451

859 Lynch, J.P., 2018. Rightsizing root phenotypes for drought resistance. *J. Exp. Bot.*
860 69, 327-3292. doi: 10.1093/jxb/ery048

861 Lynch, J.P., Brown, K.M., 2012. New roots for agriculture: Exploiting the root
862 phenome. *Philos. Trans. R. Soc. Biol. Sci.* 367, 1598–1604.
863 doi:10.1098/rstb.2011.0243

864 Lynch, J.P., Chimungu, J.G., Brown K.M., 2014. Root anatomical phenes associated
865 with water acquisition from drying soil: Targets for crop improvement. *J. Exp.*
866 *Bot.* 65, 6155-6166. doi: 10.1093/jxb/eru162

867 Lynch, J.P., Wojciechowski, T., 2015. Opportunities and challenges in the subsoil:
868 pathways to deeper rooted crops. *J. Exp. Bot.* 66, 2199-210.
869 doi:10.1093/jxb/eru508

870 Materechera, S.A., Alston, A.M., Kirby, J.M., Dexter, A.R., 1993. Field evaluation of
871 laboratory techniques for predicting the ability of roots to penetrate strong soil
872 and of the influence of roots on water sorptivity. *Plant Soil* 149, 149–158.
873 doi:10.1007/BF00016604

874 Materechera, S.A., Dexter, A.R., Alston, A.M., 1991. Penetration of very strong soils
875 by seedling roots of different plant species. *Plant Soil* 135, 31–41.
876 doi:10.1007/BF00014776

877 Merotto, A.J., Mundstock, C.M., 1999. Wheat root growth as affected by soil
878 strength. *Rev. Bras. Cienc. do Solo* 23, 197–202. doi:10.1590/S0100-
879 06831999000200002

880 Montagu, K.D., Conroy, J.P., Atwell, B.J., 2001. The position of localized soil
881 compaction determines root and subsequent shoot growth responses. *J. Exp.*

882 Bot. 52, 2127–2133. doi:10.1093/jexbot/52.364.2127

883 Mullins, C.E., Young, I.M., Bengough, A.G., Ley, G.J., 1987. Hard-setting soils. Soil
884 Use Manag. 3, 79–83. doi:10.1111/j.1475-2743.1987.tb00715.x

885 Nagarajah, S., 1987. Effects of soil texture on the rooting patterns of thompson
886 seedless vines on own roots and on ramsey rootstock in irrigated vineyards.
887 Am. J. Enol. Vitic. 38, 54–59.

888 Nagel, K.A., Putz, A., Gilmer, F., Heinz, K., Fischbach, A., Pfeifer, J., Faget, M.,
889 Blossfeld, S., Ernst, M., Dimaki, C., Kastenholz, B., Kleinert, A.-K., Galinski, A.,
890 Scharr, H., Fiorani, F., Schurr, U., 2012. GROWSCREEN-Rhizo is a novel
891 phenotyping robot enabling simultaneous measurements of root and shoot
892 growth for plants grown in soil-filled rhizotrons. Funct. Plant Biol. 39, 891.
893 doi:10.1071/fp12023

894 Nakhforoosh, A., Grausgruber, H., Kaul, H.P., Bodner, G., 2014. Wheat root diversity
895 and root functional characterization. Plant Soil 380, 211–229.
896 <https://doi.org/10.1007/s11104-014-2082-0>

897 Nosalewicz, A., Lipiec, J., 2014. The effect of compacted soil layers on vertical root
898 distribution and water uptake by wheat. Plant Soil 375, 229–240.
899 doi:10.1007/s11104-013-1961-0

900 Panayiotopoulos, K.P., Papadopoulou, C.P., Hatjioannidou, A., 1994. Compaction
901 and penetration resistance of an Alfisol and Entisol and their influence on root
902 growth of maize seedlings. Soil Tillage Res. 31, 323–337. doi:10.1016/0167-
903 1987(94)90039-6

904 Pfeifer, J., Faget, M., Walter, A., Blossfeld, S., Fiorani, F., Schurr, U., Nagel, K.A.,
905 2014a. Spring barley shows dynamic compensatory root and shoot growth
906 responses when exposed to localised soil compaction and fertilisation. Funct.
907 Plant Biol. 41, 581–597. doi:10.1071/FP13224

908 Pfeifer, J., Kirchgessner, N., Walter, A., 2014b. Artificial pores attract barley roots
909 and can reduce artifacts of pot experiments. J. Plant Nutr. Soil Sci. 177, 903-
910 913. doi:10.1002/jpln.201400142

911 Postma, J.A., Dathe, A., Lynch, J.P., 2014. The optimal lateral root branching density
912 for maize depends on nitrogen and phosphorus availability. Plant Physiol. 166,
913 590–602. doi:10.1104/pp.113.233916

914 Potocka, I., Szymanowska-Pulka, J., 2018. Morphological responses of plant roots to
915 mechanical stress. Ann. Bot. 122, 711–723. doi:10.1093/aob/mcy010

916 Potocka, I., Szymanowska-Pulka, J., Karczewski, J., Nakielski, J., 2011. Effect of
917 mechanical stress on Zea root apex. I. Mechanical stress leads to the switch
918 from closed to open meristem organization. J. Exp. Bot. 62, 4583–4593.
919 doi:10.1093/jxb/err169

920 R Core Team, 2018. R: A language and environment for statistical computing
921 (software).

922 Rasse, D.P., Smucker, A.J.M., 1998. Root recolonization of previous root channels
923 in corn and alfalfa rotations. Plant Soil 204, 203–212.
924 doi:10.1023/A:1004343122448

925 Raper, R., 2005. Agricultural traffic impacts on soil. J. Terramechanics 42, 259–280.
 926 doi.org/10.1016/j.jterra.2004.10.010

927 Rich, S.M., Watt, M., 2013. Soil conditions and cereal root system architecture:
 928 Review and considerations for linking Darwin and Weaver. J. Exp. Bot. 64,
 929 1193–1208. doi:10.1093/jxb/ert043

930 Rivera, M., Polanía, J., Ricaurte, J., Borrero, G., Beebe, S., Rao, I., 2019. Soil
 931 compaction induced changes in morpho-physiological characteristics of
 932 common bean. Journal of Soil Sci. Plant Nutr. 1–11. doi:10.1007/s42729-019-
 933 0007-y

934 Saengwilai, P., Tian, X., Lynch, J.P., 2014. Low crown root number enhances
 935 nitrogen acquisition from low-nitrogen soils in maize. Plant Physiol. 166, 581–
 936 589. doi:10.1104/pp.113.232603

937 Schenk, J.H., Jackson, R.B., 2002. The global biogeography of roots. Ecol. Monogr.
 938 72, 311–328. doi:10.1080/00223131.2002.10875464

939 Schneider, C.A., Rasband, W.S., Eliceiri, K.W., 2012. NIH Image to ImageJ: 25
 940 yearsof image analysis. Nature Meth. 9, 671–675. doi: 10.1038/nmeth.2089

941 Shierlaw, J., Alston, A.M., 1984. Effect of soil compaction on root growth and uptake
 942 of phosphorus. Plant Soil 77, 15–28. doi:10.1007/BF02182808

943 Simojoki, A., Jaakkola, A., Alakukku, L., 1991. Effect of compaction on soil air in a
 944 pot experiment and in the field. Soil Tillage Res. 19, 175–186.
 945 doi:10.1016/0167-1987(91)90085-C

946 Steinemann, S., Zeng, Z., McKay, A., Heuer, S., Langridge, P., Huang, C.Y., 2015.
 947 Dynamic root responses to drought and rewatering in two wheat (*Triticum*
 948 *aestivum*) genotypes. Plant Soil 391, 139–152. https://doi.org/10.1007/s11104-
 949 015-2413-9

950 Stirzaker, R.J., Passioura, J.B., Wilms, Y., 1996. Soil structure and plant growth:
 951 Impact of bulk density and biopores. Plant Soil 185, 151–162.
 952 doi:10.1007/BF02257571

953 Suralta, R.R., Niones, J.M., Kano-Nakata, M., Tran, T.T., Mitsuya, S., Yamauchi, A.
 954 2018. Plasticity in nodal root elongation through the hardpan triggered by
 955 rewatering during soil moisture fluctuation in rice. Scientific Reports 8:4341, 1-
 956 11.

957 To, J., Kay, B.D., 2005. Variation in penetrometer resistance with soil properties: The
 958 contribution of effective stress and implications for pedotransfer functions.
 959 Geoderma 126, 261–276. doi:10.1016/j.geoderma.2004.08.006

960 Trachsel, S., Kaeppler, S.M., Brown, K.M., Lynch, J.P., 2013. Maize root growth
 961 angles become steeper under low N conditions. F. Crop. Res. 140, 18–31.
 962 doi:10.1016/j.fcr.2012.09.010

963 Trachsel, S., Kaeppler, S.M., Brown, K.M., Lynch, J.P., 2011. Shovelomics: High
 964 throughput phenotyping of maize (*Zea mays* L.) root architecture in the field.
 965 Plant Soil 341, 75–87. doi:10.1007/s11104-010-0623-8

966 Vanhees D.J., Loades, K.W., Bengough, A.G., Mooney, S.J., Lynch, J.P., 2020. Root

967 anatomical traits contribute to deeper rooting of maize under compacted field
968 conditions. *J.Exp.Bot* 71, 4243-4257. doi: 10.1093/jxb/eraa165

969 Varney, G.T., Canny, M.J., Wang, X.L., McCully, M.E., 1991. The branch roots of
970 Zea. I. First order branches, their number, sizes and division into classes. *Ann.*
971 *Bot.* 67, 357–364. <https://doi.org/10.1093/oxfordjournals.aob.a088203>

972 Vaz, C.M.P., Manieri, J.M., de Maria, I.C., Tuller, M., 2011. Modeling and correction
973 of soil penetration resistance for varying soil water content. *Geoderma* 166, 92–
974 101. doi:10.1016/j.geoderma.2011.07.016

975 Veen, B.W., Boone, F.R., 1990. The influence of mechanical resistance and soil
976 water on the growth of seminal roots of maize. *Soil Tillage Res.* 16, 219–226.
977 doi:10.1016/0167-1987(90)90031-8

978 Veen, B.W., Van Noordwijk, M., De Willigen, P., Boone, F.R., Kooistra, M.J., 1992.
979 Root-soil contact of maize, as measured by thin-section technique. *Plant Soil*
980 139, 131-138. doi: 10.1007/BF00012848

981 Whalley, W.R., Dodd, I.C., Watts, C.W., Webster, C.P., Phillips, A.L., Andralojc, J.,
982 White, R.P., Davies, W.J., Parry, M.A.J., 2013. Genotypic variation in the ability
983 of wheat roots to penetrate wax layers. *Plant Soil* 364, 171–179.
984 doi:10.1007/s11104-012-1342-0

985 Whalley, W.R., Leeds-Harrison, P.B., Clark, L.J., Gowing, D.J.G., 2005. Use of
986 effective stress to predict the penetrometer resistance of unsaturated
987 agricultural soils. *Soil Tillage Res.* 84, 18–27. doi:10.1016/j.still.2004.08.003

988 White, R.G., Kirkegaard, J.A., 2010. The distribution and abundance of wheat roots
989 in a dense, structured subsoil - Implications for water uptake. *Plant, Cell and*
990 *Environment* 33, 133–148. doi:10.1111/j.1365-3040.2009.02059.x

991 Whitmore, A.P., Whalley, W.R., 2009. Physical effects of soil drying on roots and
992 crop growth. *J. Exp. Bot.* 60, 2845–2857. doi:10.1093/jxb/erp200

993 Wu, Q., Pagès, L., Wu, J., 2016. Relationships between root diameter, root length
994 and root branching along lateral roots in adult, field-grown maize. *Ann. Bot.* 117,
995 379–390. doi:10.1093/aob/mcv185

996 Xiong, P., Zhang, Z., Hallett, P.D., Peng, X., 2020. Variable responses of maize root
997 architecture in elite cultivars due to soil compaction and moisture. *Plant Soil*
998 455, 79-91. doi: 10.1007/s11104-020-04673-3

999 York, L.M., Nord, E.A., Lynch, J.P., 2013. Integration of root phenes for soil resource
1000 acquisition. *Front. Plant Sci.* 4, 1–15. doi:10.3389/fpls.2013.00355

1001 Young, I.M., Montagu, K., Conroy, J, Bengough, A.G., 1997. Mechanical impedance
1002 of root growth directly reduces leaf elongation rates of cereals. *New Phytol.* 135,
1003 613-619. doi:10.1046/j.1469-8137.1997.00693.xP

1004 Yu, L.-X., Ray, J.D., O'Toole, J.C., Nguyen, H.T., 1995. Use of wax-petrolatum
1005 layers for screening rice root penetration. *Crop Sci.* 35, 684–687.
1006 doi:10.2135/cropsci1995.0011183X003500030005x

1007 Zhan, A., Lynch, J.P., 2015. Reduced frequency of lateral branching improves N
1008 capture from low-N soils in maize. *J. Exp. Bot.* 66, 2055-2065.

1009 doi:10.1093/jxb/erv007
1010 Zhan, A., Schneider, H., Lynch, J.P., 2015. Reduced lateral root branching density
1011 improves drought tolerance in maize. *Plant Physiol.* 168, 1603–1615.
1012 doi:10.1104/pp.15.00187
1013
1014
1015
1016

1017 Artwork
1018 Figures

1019
1020 **Figure 1** – Average penetrometer resistances $\pm$ SE for compacted (red) and non-
1021 compacted (blue) treatments at (A) the ARBC field site (coarse-loam) and (B) the PSU
1022 field site (silt-loam) before planting. Mean soil moisture content (v/v) was measured
1023 for each 10 cm increment, together with soil dry bulk density, both written on graph
1024 within relevant depth ranges. Figure adopted from Vanhees *et al.* (2020).

1025

Figure 2 - Average total root length (cm) $\pm$ SE split into coarse (dark blue, dark red) and fine (light blue and light red) root length (cm) for maize genotypes tested at acoarse-loam (ARBC) versus silt-loam (PSU) field site. Coarse roots are defined as having diameters larger than 1 mm, while fine roots are those with diameter smaller than 1 mm. Compacted measurements in red, non-compacted measurement in blue. Error bars represent standard deviations. If differences between the field sites (***, $p \leq 0.001$), treatments (A/B, $p \leq 0.05$) and genotypes (a/ab/b, $p \leq 0.05$) were present.

Figure 3 – Proportions of coarse (>1.0 mm diameter) root length (%) ± SE found in cores of different genotypes in two field sites. Non-compacted data in blue, compacted data in red. IBM059 (coarse-loam at ARBC) and OHW128 (silt loam at PSU) have such small standard errors they could not be visualised. Post-hoc Tukey comparisons within field site indicate when treatment effect was significant for each genotype at significance level ° $P \leq 0.10$, * $P \leq 0.05-0.01$, ** $P \leq 0.01-0.001$, *** $P \leq 0.001$.

Figure 4 – Illustration of difference between absolute versus relative root length density distributions for genotype IBM014 considering total root length and coarse root length at the coarse-loam field site (ARBC). (A) + (C) Absolute distributions of root length densities, (B) + (D) Relative distributions of root length densities. Compacted data in red and non-compacted data in blue. Error bars represent standard errors. The rooting depth (cm) ± SE where 75% of the total root length (D_{75}) or coarse root length (D_{75c}) was visualised by the striped line, coloured region represents SE for the depth measurements. No error bars shown when standard error was too small to visualise.

Figure 5 – Coarse and total rooting depth and their correlation for both field sites under compaction (red) and non-compacted (blue) conditions.. (A) + (B) Average coarse rooting depth (D_{75c}), (C) + (D) Average total rooting depth, (E) + (F) Correlation between D_{75} and D_{75c} . Error bars represent standard errors. (A) + (C) + (E): ARBC field site (coarse-loam) and (B) + (D) + (F): PSU field site (silt-loam). Post hoc Tukey comparisons between compaction and noncompaction within each field site for each genotype were carried out on rooting depth data (panels A-D). Coarse and total rooting depth are correlated (E-F) Levels of significance + $P \leq 0.10$, * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$.

Figure 6 – Linear regressions between root distribution variables at the two different field sites. Field site ARBC (coarse-loam) visualised in A, C, E and field site PSU (silt-loam) visualised in B, D, F. Compacted data (red) and non-compacted data (blue). Each datapoint represents the averaged value across the replicates for each genotype tested. Normal linear regression was used for A–B and E–F, and a betaregression was used for C–D as data was proportional. When a significant relationship was found this was at a level of significance of [†] $P \leq 0.10$ or * $P \leq 0.05$.

Figure 7 - Genotypic variation in the coarse root length density (cm cm⁻³) per depth increment across two field sites and two compaction treatments. Non-compacted data in blue and compacted data in red. The ARBC field site (coarse-loam) and PSU field site (silt-loam) had different soil textures. The striped lines are the averages across all genotypes, lighter coloured lines are the average for individual genotypes tested. Similar plots for total root length density distributions can be found in Figure S9.

Figure 8 – Coarse root length densities (cm cm^{-3}) $\pm$ SE distributions with soil depth on compacted plots comparing (A) two genotypes per field site (coarse loam at ARBC and silt-loam at PSU) with similar total coarse root length but with different associated rooting depths and (B) two genotypes with similar rooting depths but with different total coarse root lengths. For (A) striped lines stands for the deeper rooting genotype and associated D_{75c} , while the solid line stands for the shallower rooting genotypes and associated D_{75c} . For (B), the solid line is used for the genotype that produces less roots but reaches equally deep as the genotype that produces more roots (striped lines). No error bars shown when standard error was too small to visualise. Selection comparison can be found in Figure S10. Similar plots for total root length density distributions can be found in Figure S12.

ARBC: Biomass ~ Genotype + D75 _c + D75 + TRL _f + TRL _c + Compaction		
	F-value	p-value
Genotype	1.144	ns
D75 _c	140.93	***
D75	0.231	ns
TRL _f	37.629	***
TRL _c	2.287	ns
Compaction	5.83	*
PSU: Biomass ~ Genotype + D75 _c + D75 + TRL _f + TRL _c + Compaction		
	F-value	p-value
Genotype	0.649	ns
D75 _c	14.491	**
D75	5.244	ns
TRL _f	0.083	ns
TRL _c	3.75	ns
Compaction	9.713	*

Figure 9 – Relationships between average biomass and root distribution values and are visualised in the accompanied plots where squares are data from the ARBC (coarse-loam) field site while circles are data from the PSU (silt-loam) field site. Compacted data in red, non-compacted data in blue. Level of significance in the ANOVA analysis are *** $p < 0.001$, ** $p < 0.01$ and * $p < 0.05$.

1098 **Tables**

1099 **Table 1** – Definitions of the different measurements obtained after WinRHIZO analysis of the soil cores.

Measurement	Abbreviation	Definition	Unit
Total root length	TRL	The summation of all individual root sections per 10 cm increment of the entire soil core	cm
Total coarse root length	TRL _c	The summation of all individual root sections per 10 cm increment with a diameter greater than 1 mm over the entire soil core	cm
Total fine root length	TRL _f	The summation of all individual root sections per 10 cm increment with a diameter smaller than 1 mm over the entire soil core	cm
Coarse root proportion	P _c	The ratio of total coarse root length versus total root length	%
Fine root proportion	P _f	The ratio of total fine root length versus total root length	%
Root length density	-	The root length found in the soil volume of a 10 cm increment of the soil core and this including all root diameter classes	cm cm ⁻³
Coarse root length density	-	The total coarse root length found in the soil volume of a 10 cm increment of the soil core	cm cm ⁻³
Relative root length density	-	The ratio of total root length density of a single 10 cm increment versus the sum of the total root length density found over the entire core	%
Relative coarse root length density	-	The ratio of coarse root length density of a single 10 cm increment versus the sum of the coarse root length density found over the entire core	%
Rooting depth	D ₉₅	The rooting depth above which 95% of the total root length is located	cm
	D ₇₅	The rooting depth above which 75% of the total root length is located	cm
Coarse rooting depth	D _{95c}	The rooting depth above which 95% of the total coarse root length is located	cm
	D _{75c}	The rooting depth above which 75% of the total coarse root length is located	cm

1100

1101

Table 2 F-values for split plot analysis results of the different root distribution variables at the two field sites. P-values tested at the following levels of significance: [†] $p \leq 0.10$, * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$. Subscript c stands for coarse and f stands for fine when measurements are made on a separate root class. TRL stands for total root length, P stands for proportion of coarse or fine roots. D75 and D95 stand for rooting depth at which 75 and 95 percent of the total root length can be found.

		ARBC (coarse-loam)	PSU (silt-loam)
TRL	Compaction	77.12 ***	1.37
	Genotype	0.67	0.54
	Compaction x Genotype	0.87	0.85
TRL _c	Compaction	4.59 [†]	3.61 [†]
	Genotype	1.58	1.12
	Compaction x Genotype	2.11 *	1.35
TRL _f	Compaction	78.81 ***	1.25
	Genotype	0.67	0.56
	Compaction x Genotype	0.81	0.83
P _c	Compaction	18.29 **	3.43
	Genotype	2.60 **	1.97 *
	Compaction x Genotype	2.12 *	1.34
P _f	Compaction	18.62 **	2.63
	Genotype	2.46 *	1.77 [†]
	Compaction x Genotype	1.93 [†]	1.30
D _{75c}	Compaction	76.53 ***	4.65 [†]
	Genotype	3.15 **	1.67
	Compaction x Genotype	0.71	0.55
D _{95c}	Compaction	42.29 ***	0.78
	Genotype	3.86 ***	0.65
	Compaction x Genotype	1.33	0.60
D ₇₅	Compaction	17.31 **	6.78 *
	Genotype	2.74 **	1.08
	Compaction x Genotype	0.87	0.36
D ₉₅	Compaction	25.02 ***	1.56
	Genotype	2.70 **	1.11
	Compaction x Genotype	1.33	0.33

1110 **Table 3** – Summary of general linear model results for the linear regression of total or
 1111 coarse rooting depth (D_{75} or D_{75c}) with total root length (TRL) or total coarse root length
 1112 (TRL_c). P-values tested at the following levels of significance: † $p \leq 0.10$, * $p \leq 0.05$, **
 1113 $p \leq 0.01$ and *** $p \leq 0.001$.

1114

$D_{75} \sim \text{TRL} + \text{Field site} + \text{Compaction} + \text{Genotype}$		
	F-value	p-value
Field site	57.36	***
Compaction	12.21	*
Genotype	1.22	
Total root length	3.09	†

$D_{75} \sim \text{TRL}_c + \text{Field site} + \text{Compaction} + \text{Genotype}$		
	F-value	p-value
Field site	106.37	***
Compaction	10.2	*
Genotype	1.17	
Total root length	0.34	

$D_{75c} \sim \text{TRL} + \text{Field site} + \text{Compaction} + \text{Genotype}$		
	F-value	p-value
Field site	35.83	***
Compaction	25.51	***
Genotype	2.12	*
Total root length	2.73	

$D_{75c} \sim \text{TRL}_c + \text{Field site} + \text{Compaction} + \text{Genotype}$		
	F-value	p-value
Field site	41.39	***
Compaction	34.77	***
Genotype	1.99	*
Total root length	1.39	

1115 **Appendices**
1116 **Supplementary data**

1117
1118 **Figure S1** – Volumetric water content was used to steer irrigation (ARBC, coarse-loam
1119 field site) or to decide if irrigation was needed (PSU, silt-loam field site). On coarse-
1120 loam PR2-tubes were installed while on silt-loam TDR-probes were used to measure
1121 volumetric water content; gravimetric water content was calculated from this and the
1122 dry bulk density. Compacted site (red), non-compacted (blue).

Figure S2 – Relationship between crown root angle and coarse rooting depth for ARBC (coarse-loam) and PSU (silt-loam) field sites. Figures adjusted from supplementary Figure 1 from Vanhees *et al.* (2020). Compacted site (red), non-compacted (blue).

Figure S3 – Biomass $\pm$ SE at both field sites under compacted (red) and non-compacted (blue) conditions for each genotype. The ARBC field site has a coarse-loam soil while the PSU field site has a silt loam soil texture. Figure adjusted from supplementary Figure 3 from Vanhees *et al.* (2020).

Figure S4 – Total root length of each genotype plotted at the averaged penetrometer resistance of the 2 field trial compaction treatment combinations.

Figure S5 – Correlation plots between tested variables averaged over all genotypes across field sites (ARBC (coarse-loam) or PSU (silt-loam)) and compacted (C) or non-compacted plots (NC) combinations. The correlation coefficient is visualised by the scale bar, negative correlations are orange and positive correlations are blue. A cross represents a non-significant correlation at significance $p \leq 0.05$.

1153 relationship was detected at significance level $p \leq 0.10$, other relationships were non-
 1154 significant (ns).

1155 **Figure S7** – Relationships between total rooting depth (D_{75c}) and other root distribution
 1156 variables across field sites and compaction treatments. Panels A and C represent field
 1157 site ARBC (coarse-loam) and panels B and D field site PSU (silt-loam). Non-
 1158 compacted data in blue, compacted data in red. No significant (ns) linear relationships
 1159 were detected.
 1160

1161

Figure S8 – Principle component analysis per field site (ARBC (coarse-loam) or PSU (silt-loam)) – compaction treatment (C – compacted; NC – non-compacted) combination illustrating relationships between root distribution variables within respective environmental conditions.

Figure S9 – Genotypic variation in total root length density (cm cm⁻³) per depth increment across two field sites and two compaction treatments. The ARBC field site has a coarse-loam soil texture and the PSU field site has a silt-loam soil texture. Non-compacted data in blue and compacted data in red. The striped line are the averages across all genotypes, lighter coloured lines are the average for individual genotypes tested.

Figure S10 – Selection of genotypes to be compared based on their coarse rooting depth and coarse total root length. Genotypes indicated with an arrow were selected on the bases of similar coarse root length but different coarse rooting depths (shallow versus deep) and genotypes indicated with a triangle were selected on the basis of similar coarse rooting depth but are different according to total coarse root length (few versus many roots for deeper rooting genotypes). . The ARBC field site has a coarse-loam soil texture and the PSU field site has a silt-loam soil texture.

Figure S11 – Selection of genotypes to be compared based on their total rooting depth and total root length. Genotypes indicated with an arrow were selected on the bases of similar coarse root length but different coarse rooting depths (shallow versus deep) and genotypes indicated with a triangle were selected on the basis of similar coarse rooting depth but are different according to total coarse root length (few versus many roots for deeper rooting genotypes). . The ARBC field site has a coarse-loam soil texture and the PSU field site has a silt-loam soil texture.

Figure S12 - Total root length densities (cm cm^{-3}) $\pm$ SE distributions with soil depth on compacted plots comparing (A) two genotypes per field sites with similar total coarse root length but with different associated rooting depths under compaction and (B) two genotypes with similar rooting depths but with different total coarse root lengths under compaction. For (A) solid lines stands for the deeper rooting genotype and associated D_{75} , while the striped line stands for the shallower rooting genotypes and associated D_{75} . For (B), the solid line is used for the genotype that produces less roots but reaches equally deep then the genotype that produces more roots (striped lines). The ARBC field site has a coarse-loam soil texture and the PSU field site has a silt-loam soil texture.

1205 **Table S1** – Field applications during the field season. Table adjusted from supplementary Table 1 from Vanhees *et al.* (2020).

Field applications						
Irrigation		Fertilizers		Pesticides		
coarse-loam	18/06/2016	0.94 mm	04/06/2016	ProSol (15 gallons/acre)	17/06/2016	Atrazine and S-metolachlor
	21/06/2016	0.60 mm	14/06/2016	ProSol (12.5 gallons/acre)	14/06/2016	Copper, Azoxystrobin and Chlorantraniliprole
	22/06/2016	1.20 mm	15/06/2016	UAN (38.3 lbs/acre)	23/06/2016	Chlorantraniliprole
	23/06/2016	0.32 mm	16/06/2016	ProSol (12.5 gallons/acre)		
	25/06/2016	1.60 mm				
	28/06/2016	0.30 mm				
	04/07/2016	0.24 mm				
	06/07/2016	0.20 mm				
	08/07/2016	0.50 mm				
	10/07/2016	0.72 mm				
	17/07/2016	0.75 mm				
	20/07/2016	0.50 mm				
	24/07/2016	0.50 mm				
	27/07/2016	0.50 mm				
	09/08/2016	0.50 mm				
silt-loam	No irrigation applied as moisture content remained high enough during growing season		urea Nitrogen (200 lbs/acre) applied prior to planting		No pesticides were applied	

1206

1207 **References in supplementary**

1208 Vanhees D.J., Loades, K.W., Bengough, A.G., Mooney, S.J., Lynch, J.P., 2020. Root
1209 anatomical traits contribute to deeper rooting of maize under compacted field conditions.
1210 J.Exp.Bot 71, 4243-4257. doi: 10.1093/jxb/eraa16

1