University of Dundee

Effect of unsteady flows on the development and magnitude of bed forms

Wang, Le; Cuthbertson, Alan; Pender, Gareth; Cao, Zhixian

Publication date:
2004

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Wang, L., Cuthbertson, A., Pender, G., & Cao, Z. (2004). *Effect of unsteady flows on the development and magnitude of bed forms*. Abstract from EGU General Assembly 2021.

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

EGU21-5854, updated on 04 Mar 2021

<https://doi.org/10.5194/egusphere-egu21-5854>

EGU General Assembly 2021

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.

Effect of unsteady flows on the development and magnitude of bed forms

Le Wang, Alan Cuthbertson, Gareth Pender, and Zhixian Cao

North China Electric Power University, School of Water Resources and Hydropower Engineering, China

(lewang@ncepu.edu.cn)

Sediment transport and associated morphological changes in alluvial rivers occur primarily under unsteady flow conditions that are manifested as well-defined flood hydrograph events. At present, typical bed forms generated by such unsteady flows is far less studied and, thus, more poorly understood, than equivalent bed forms generated under steady flow conditions. In view of this, the objective of this work is to investigate the development of morphological bed features, and specifically variability in the length, height and steepness of bed forms that develop in a mobile coarse-sand bed layer under unsteady flow hydrographs under zero sediment feed conditions. A series of laboratory flume experiments is conducted within which different flow hydrograph events are simulated physically by controlling their shape, unsteadiness and magnitude. Experimental results indicate that different categories of bed forms such as dunes, alternate bars or transitional dune-bar structures develop within the erodible bed layer when subject to varying hydrograph flow conditions. Examination of relative importance of three parameters used to describe the hydrograph characteristics (i.e. asymmetry, unsteadiness and total water work) on bed form dimensional descriptors (i.e. wavelength, height and steepness) reveals that hydrograph unsteadiness and total water work are the primary and second-order controls on bed deformations or corresponding bed form dimensions. By contrast, hydrograph asymmetry appears to have minimal or negligible influence on bed form development in terms of their type and magnitude. Based on these findings, a physical model was developed and tested to describe the effect of unsteady flow hydrographs with varying unsteadiness and total water work on the nature and size of resulting bed forms that are generated in sand-bed layers.