University of Dundee

The intertwined roles of specialized metabolites within the *Bacillus subtilis* biofilm

Kalamara, Margarita; Stanley-Wall, Nicola R.

Published in:
Journal of Bacteriology

DOI:
[10.1128/JB.00431-21](https://doi.org/10.1128/JB.00431-21)

Publication date:
2021

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Kalamara, M., & Stanley-Wall, N. R. (2021). The intertwined roles of specialized metabolites within the *Bacillus subtilis* biofilm. *Journal of Bacteriology*, 203(22), [e00431-21]. <https://doi.org/10.1128/JB.00431-21>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 Margarita Kalamara ¹ and Nicola R. Stanley-Wall ^{1*}

2 ¹ Division of Molecular Microbiology, School of Life Sciences, University of Dundee, Dundee, UK

3 **The intertwined roles of specialised metabolites within the *Bacillus subtilis* biofilm**

4 * Corresponding author: n.r.stanleywall@dundee.ac.uk

5

6 **ABSTRACT** (75 words limit)

7 Bacteria produce specialised metabolites with a range of functions. In this issue of the *Journal of*
8 *Bacteriology* Schoenborn *et al.* study the production and role of secondary metabolites during
9 biofilm development and sporulation in *Bacillus subtilis*. Most metabolites studied are produced
10 during differentiation and six are required for the development of biofilms and/or spores. The
11 authors propose a model for the timing of production and role in differentiation exerted by each
12 secondary metabolite.

13 **KEYWORDS**

14 *Bacillus subtilis*, biofilm, spore, differentiation, specialised metabolites

15

COMMENTARY

Social interactions in the microbial world are incredibly complex, but we are slowly starting to understand some of the intricate mechanisms that underpin them. Specialised metabolites, which are a diverse range of molecules with a broad range of functions, are major players in defining these social dynamics. These molecules can impact populations of microbes through killing or inhibiting growth, triggering differentiation between physiological states, or manipulating nutrient availability in the environment (1). The prevalence of these molecules, and their pervasive impact on many aspects of microbial life, means that they have crucial roles in defining the composition and emergent properties of inter-kingdom, inter-species, intra-species, and single strain communities of microbes.

Biofilms are highly heterogeneous structures where microenvironments and gradients in nutrient availability, oxygen levels, and cells types develop (2) (Figure 2A). *Bacillus subtilis* is a Gram-positive soil bacterium that has been extensively used for the study of social interactions in the context of biofilm formation. Examples of the cell types that make up a *B. subtilis* biofilm include motile cells, biofilm matrix producers, exoprotease producers, and endospores, facilitating the division of labor and the sharing of public goods between the community members (3). The regulatory processes leading to differentiation of cells into these physiological states is highly complex and relies on input from a variety of environmental signals, some of which are specialised metabolites produced by the cells in the biofilm themselves (3).

B. subtilis is well known for the plethora of specialised metabolites it produces. Most of these specialised metabolites are primarily linked with their antimicrobial properties and include

37 bacilysin (4), bacillaene (5), subtilosin A (6), plipastatin (7), surfactin (8), sporulation killing factor
38 (9) and sublancin 168 (10). Other specialised metabolites produced by *B. subtilis* include the iron-
39 chelating molecule pulcherriminic acid (11) and the siderophore bacillibactin (12) (Figure 2B). The
40 specialised metabolites with a known impact on cell state differentiation in *B. subtilis*
41 communities are surfactin and the pheromone ComX, which act as extracellular signals to induce
42 the differentiation of cells into biofilm matrix producers and “cannibals” (13). Cannibals are a
43 subpopulation of cells that produce the specialised metabolites sporulation killing factor (SKF)
44 and sporulation delaying protein (SDP), which function to lyse sister cells in the community to
45 use them as a nutrient source and delay sporulation (9, 13). The siderophore bacillibactin has
46 also been found to be involved in the development of *B. subtilis* biofilms (14). However, while we
47 know that some specialised metabolites are crucial for cell fate differentiation and biofilm
48 formation, there has not been a comprehensive systematic study of the interplay between
49 secondary metabolite production and differentiation until now.

50 In this paper, Schoenborn *et al.* tested the role of nine specialised metabolites during biofilm
51 formation and sporulation by examining the expression, production, and impact of deletion
52 mutants on differentiation (they deleted biosynthetic genes necessary to produce specialised
53 metabolites; these genes are often referred to as "clusters" based on their genomic structure).
54 The authors demonstrated that most clusters (those needed for surfactin, subtilosin A, ComX,
55 SDP, SKF, bacilysin, and bacillaene production) are expressed at a higher level under
56 differentiation-inducing conditions, except for the plipastatin and bacillaene clusters. Largely
57 mirrored by this analysis, all metabolites examined, including plipastatin and bacillaene, were
58 produced in significantly higher amounts under conditions that promote biofilm formation and

59 differentiation. The higher production of specialised metabolites during cell fate differentiation
60 points to these molecules having a role during these processes. Interestingly however, the ability
61 of cells to produce most of these molecules was not essential for biofilm formation, at least
62 individually, as deletion of genomic regions required for biosynthesis of the specialised
63 metabolites did not impact biofilm structure. The exception was surfactin, the absence of which
64 resulted in a biofilm deficient strain when analysed by the pellicle biofilm model. These findings
65 are in contrast to a recent study showing that lack of surfactin does not impact pellicle biofilm
66 development, but consistent with surfactin being required for architecturally complex colony
67 biofilms to form (15). In line with an impact to colony biofilm formation, in this study, the lack of
68 surfactin caused a decrease in the expression of the biofilm matrix protein encoding gene *tapA*.
69 The reduction in *tapA* expression was also found to be the case for the mutant lacking the ability
70 to produce ComX, which is consistent with both ComX and surfactin being important for
71 differentiation of cells into biofilm matrix producers. Another two molecules, subtilisin A and
72 bacillaene, impacted matrix gene expression, but this was at the later stages of the pellicle biofilm
73 formation, after around 16 hours of growth. At this point the increase in expression of the biofilm
74 matrix protein starts to level off in the wild type but continued to increase in the subtilisin A and
75 bacillaene mutants. Looking at sporulation dynamics, the authors showed that lack of surfactin,
76 plipastatin, bacilysin, subtilisin A, ComX, and bacillibactin all impacted sporulation. At 16 hours,
77 the number of spores was significantly lower for the strains incapable of producing the
78 specialised metabolites compared to the wild type, suggesting that these molecules are required
79 for triggering spore formation.

We are gaining more and more understanding about the multifaceted nature of specialised metabolites and some of the *B. subtilis*-produced specialised metabolites are now known to be multifunctional. For example, bacillaene protects *B. subtilis* from predation by other bacteria (16), can modulate production of secondary metabolites by competing bacteria (17), impacts the composition of mixed species bacterial communities (18), inhibits biofilm formation by other bacteria (19), and has been suggested to impact biofilm development of *B. subtilis* biofilms at subinhibitory concentrations (20). This paper by Schoenborn *et al.* reveals an additional role for some of the less widely explored specialised metabolites produced by *B. subtilis* in differentiation. Plipastatin (which has been studied for its ability to inhibit the growth of multiple plant pathogenic fungal species (21)), the bacteriocin subtilosin A (6), and bacillibactin (which is a siderophore), are now known to also function as signals that regulate sporulation in *B. subtilis* mixed communities. Therefore, it is clear that these molecules have a function in both competition against others, either through their antimicrobial functions or in limiting available nutrients in the environment by sequestering them, and in impacting cooperative dynamics in a single species biofilm. One can speculate about the multipurpose role for the molecules. Bacteria produce a relatively limited number of molecules with which they need to navigate an incredibly complex world. *B. subtilis* can be found in the gastrointestinal tract of animals, in association with plant roots, in bulk soil, and in marine environments and is likely to have to interact with its eukaryotic hosts, other species of microbes, and members of its own species. It therefore makes sense, from an evolutionary perspective, for bacteria to ensure that the limited molecules they produce have a variety of functions to help them thrive in an ever-changing environment.

102 **ACKNOWLEDGEMENTS**

103 Biofilm related work in the laboratory of N.R.S.-W. is funded by the Biotechnology and Biological
104 Science Research Council (BBSRC) [BB/P001335/1 and BB/R012415/1]. M.K. is supported by a
105 Biotechnology and Biological Sciences Research Council studentship [BB/M010996/1]

106

107 REFERENCES

- 108 1. Stubbendieck RM, Straight PD. 2016. Multifaceted Interfaces of Bacterial Competition. *J Bacteriol*
109 198:2145-55.
- 110 2. Flemming HC, Wingender J, Szewzyk U, Steinberg P, Rice SA, Kjelleberg S. 2016. Biofilms: an
111 emergent form of bacterial life. *Nat Rev Microbiol* 14:563-75.
- 112 3. Lopez D, Kolter R. 2010. Extracellular signals that define distinct and coexisting cell fates in *Bacillus*
113 *subtilis*. *FEMS Microbiol Rev* 34:134-49.
- 114 4. Kenig M, Abraham EP. 1976. Antimicrobial activities and antagonists of bacilysin and anticapsin. *J*
115 *Gen Microbiol* 94:37-45.
- 116 5. Patel PS, Huang S, Fisher S, Pirnik D, Aklonis C, Dean L, Meyers E, Fernandes P, Mayerl F. 1995.
117 Bacillaene, a novel inhibitor of procaryotic protein synthesis produced by *Bacillus subtilis*:
118 production, taxonomy, isolation, physico-chemical characterization and biological activity. *J*
119 *Antibiot (Tokyo)* 48:997-1003.
- 120 6. Babasaki K, Takao T, Shimonishi Y, Kurahashi K. 1985. Subtilosin A, a new antibiotic peptide
121 produced by *Bacillus subtilis* 168: isolation, structural analysis, and biogenesis. *J Biochem* 98:585-
122 603.
- 123 7. Umezawa H, Aoyagi T, Nishikiori T, Okuyama A, Yamagishi Y, Hamada M, Takeuchi T. 1986.
124 Plipastatins: new inhibitors of phospholipase A2, produced by *Bacillus cereus* BMG302-fF67. I.
125 Taxonomy, production, isolation and preliminary characterization. *J Antibiot (Tokyo)* 39:737-44.
- 126 8. Chen H, Wang L, Su CX, Gong GH, Wang P, Yu ZL. 2008. Isolation and characterization of
127 lipopeptide antibiotics produced by *Bacillus subtilis*. *Lett Appl Microbiol* 47:180-6.
- 128 9. Gonzalez-Pastor JE, Hobbs EC, Losick R. 2003. Cannibalism by sporulating bacteria. *Science*
129 301:510-3.
- 130 10. Paik SH, Chakicherla A, Hansen JN. 1998. Identification and characterization of the structural and
131 transporter genes for, and the chemical and biological properties of, sublancin 168, a novel
132 lantibiotic produced by *Bacillus subtilis* 168. *J Biol Chem* 273:23134-42.
- 133 11. Arnaouteli S, Matoz-Fernandez DA, Porter M, Kalamara M, Abbott J, MacPhee CE, Davidson FA,
134 Stanley-Wall NR. 2019. Pulcherrimin formation controls growth arrest of the *Bacillus subtilis*
135 biofilm. *Proc Natl Acad Sci U S A* 116:13553-13562.
- 136 12. May JJ, Wendrich TM, Marahiel MA. 2001. The *dhb* operon of *Bacillus subtilis* encodes the
137 biosynthetic template for the catecholic siderophore 2,3-dihydroxybenzoate-glycine-threonine
138 trimeric ester bacillibactin. *Journal of Biological Chemistry* 276:7209-7217.
- 139 13. Gonzalez-Pastor JE. 2011. Cannibalism: a social behavior in sporulating *Bacillus subtilis*. *FEMS*
140 *Microbiol Rev* 35:415-24.
- 141 14. Qin Y, He Y, She Q, Larese-Casanova P, Li P, Chai Y. 2019. Heterogeneity in respiratory electron
142 transfer and adaptive iron utilization in a bacterial biofilm. *Nat Commun* 10:3702.
- 143 15. Therien M, Kiesewlter HT, Auria E, Charron-Lamoureux V, Wibowo M, Maroti G, Kovacs AL,
144 Beauregard P. 2020. Surfactin production is not essential for pellicle and root-associated biofilm
145 development of *Bacillus subtilis*. *Biofilm* 2:100021.
- 146 16. Muller S, Strack SN, Hoefler BC, Straight PD, Kearns DB, Kirby JR. 2014. Bacillaene and sporulation
147 protect *Bacillus subtilis* from predation by *Myxococcus xanthus*. *Appl Environ Microbiol* 80:5603-
148 10.
- 149 17. Vargas-Bautista C, Rahlwes K, Straight P. 2014. Bacterial competition reveals differential
150 regulation of the *pks* genes by *Bacillus subtilis*. *J Bacteriol* 196:717-28.

- 151 18. Kieseewalter HT, Lozano-Andrade CN, Strube ML, Kovacs AT. 2020. Secondary metabolites of
152 *Bacillus subtilis* impact the assembly of soil-derived semisynthetic bacterial communities.
153 *Beilstein J Org Chem* 16:2983-2998.
- 154 19. Erega A, Stefanic P, Dogsa I, Danevcic T, Simunovic K, Klancnik A, Smole Mozina S, Mandic Mulec
155 I. 2021. Bacillaene Mediates the Inhibitory Effect of *Bacillus subtilis* on *Campylobacter jejuni*
156 Biofilms. *Appl Environ Microbiol* 87:e0295520.
- 157 20. Li H, Han X, Dong Y, Xu S, Chen C, Feng Y, Cui Q, Li W. 2021. Bacillaenes: Decomposition Trigger
158 Point and Biofilm Enhancement in *Bacillus*. *ACS Omega* 6:1093-1098.
- 159 21. Kieseewalter HT, Lozano-Andrade CN, Wibowo M, Strube ML, Maroti G, Snyder D, Jorgensen TS,
160 Larsen TO, Cooper VS, Weber T, Kovacs AT. 2021. Genomic and Chemical Diversity of *Bacillus*
161 *subtilis* Secondary Metabolites against Plant Pathogenic Fungi. *mSystems* 6.
- 162 22. Blin K, Shaw S, Kloosterman AM, Charlop-Powers Z, van Wezel GP, Medema MH, Weber T. 2021.
163 antiSMASH 6.0: improving cluster detection and comparison capabilities. *Nucleic Acids Res*
164 doi:10.1093/nar/gkab335.
- 165 23. Grant JR, Stothard P. 2008. The CGView Server: a comparative genomics tool for circular genomes.
166 *Nucleic Acids Res* 36:W181-4.

Figure 1: Secondary metabolites and *B. subtilis* biofilms. (A) Vertical cross-section of a colony biofilm formed by *B. subtilis* NCIB 3610 with a schematic representation of the concepts covered in this work. The cross-section was prepared and imaged by Dr. Sofia Arnaouteli while visiting the laboratory of Prof. Lars Dietrich. (B) Schematic of the genome of *B. subtilis* strain NCIB 3610 showing the locations on the chromosome of the secondary metabolite biosynthesis clusters and other explored molecules. The secondary metabolite biosynthesis clusters were predicted using AntiSMASH version 6.0 (22) and the genome map was constructed using GCView (23).