



University of Dundee

Regulation of mRNA capping in the cell cycle

Aregger, Michael; Cowling, Victoria H.

Published in:
RNA Biology

DOI:
[10.1080/15476286.2016.1251540](https://doi.org/10.1080/15476286.2016.1251540)

Publication date:
2017

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Aregger, M., & Cowling, V. H. (2017). Regulation of mRNA capping in the cell cycle. *RNA Biology*, 14(1), 11-14.
<https://doi.org/10.1080/15476286.2016.1251540>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Running title: CDK1 regulation of mRNA cap methylation

Point of View

Regulation of mRNA capping in the cell cycle

Michael Aregger^{1,2} and Victoria H. Cowling^{1*}

¹Centre for Gene Regulation and Expression, School of Life Sciences

University of Dundee, Dow Street, Dundee DD1 5EH, UK

²Current address: Donnelly Centre for Cellular and Biomolecular Research, University of Toronto, Toronto, ON M5S 3E1, Canada

*Corresponding author: v.h.cowling@dundee.ac.uk; Centre for Gene Expression, School of Life Sciences, University of Dundee, Dow Street, Dundee, United Kingdom, DD1 5EH

Key words

transcription, translation, gene expression, mRNA, mRNA cap, RNMT, phosphorylation, cell cycle

No potential conflicts of interest, financial or otherwise.

Abstract

The mRNA cap structure, which is added to nascent RNA pol II transcripts, recruits the protein complexes required for pre-mRNA transcript processing, mRNA export and translation initiation. The enzymes which catalyse mRNA cap synthesis are regulated by cellular signaling pathways which impact on their expression, localisation and activity. Here we discuss the recent observation that the mRNA cap methyltransferase, RNMT, is phosphorylated on Thr-77 by CDK1-cyclin B1, which regulates its activity and the proteins with which it interacts. RNMT Thr-77 phosphorylation provides a burst of

mRNA cap methyltransferase activity during early G1 phase at a time when transcription is reactivated following completion of the cell cycle. This co-ordination of transcription and mRNA capping makes an important contribution to gene expression in the cell; preventing RNMT Thr-77 phosphorylation inhibits cell proliferation. Here we discuss these findings and how mRNA cap synthesis may be regulated in other scenarios.

Introduction

Eukaryotic cells contain a variety of RNA molecules, the vast majority of which assist with mRNA processing and translation including ribosomal RNA and transfer RNA. Transcripts which mature to become protein-encoding messenger RNA (mRNA) only account for about 5% all cellular transcripts. The mRNA cap, a modification added to the first transcribed nucleotide of pre-mRNA and other RNA pol II transcripts, flags these RNAs for appropriate processing, nuclear export and translation into protein¹⁻³. The structure of the mRNA cap also protects transcripts from exonucleases. The importance of the mRNA cap for gene expression was recognised shortly after its discovery by a series of experiments *in vitro* and in yeast. More recently formation and degradation of the mRNA cap has been observed to be regulated by cellular signalling pathways in mammalian cells^{4,5}.

The mRNA cap consists of a 7-methylguanosine group linked via a 5' to 5' triphosphate bridge to the first transcribed nucleotide of RNA pol II transcripts (Figure 1)^{2,3}. The first few nucleotides can also be methylated, but the position and extent of methylation varies between species. Transcripts are synthesised with a 5' triphosphate on the first transcribed nucleotide to which the mRNA cap is added by the action of a series of enzymes^{1,4}. In mammals the basic mRNA cap structure, cap 0, is formed by the sequential action of CE/RNGTT and RNMT-RAM, two enzymes recruited to phosphorylated RNA pol II at the initiation of transcription. CE/RNGTT has triphosphatase and guanylyltransferase activities which cleave the transcript terminal

phosphate and catalyse addition of guanosine monophosphate, to form the cap structure, G(5')ppp(5')N (N is the first transcribed nucleotide). Subsequently, RNMT-RAM catalyses N-7 methylation of the guanosine cap to create, m7G(5')ppp(5')N. This cap structure is functional in recruiting the Cap-binding complex (CBC) and the eukaryotic initiation factor 4E, eIF4E, and other factors which promote transcript splicing, export and translation initiation^{5,6}. Furthermore, in mammals the first two nucleotides are methylated on the O-2 position by CMTR1 and CMTR2^{7,8}. O-2 methylation has roles in translation initiation and identifying transcripts as “self” in innate immunity⁴.

Regulation of mRNA cap methylation

The mammalian mRNA cap methyltransferase, RNMT-RAM, consists of the enzymic subunit, RNMT (RNA guanine-7 methyltransferase) and the activator subunit, RAM (RNMT-activating miniprotein)^{9,10}. RNMT has a catalytic core, which is homologous in structure and function to other eukaryotic cap methyltransferases¹¹. In addition, RNMT contains an N-terminal non-catalytic domain, which is conserved in mammals but not in other eukaryotes (Figure 2)⁹. *S.cerevisiae* null for the cap methyltransferase are inviable and can be rescued by the expression of murine RNMT catalytic domain¹². RAM activates RNMT by altering the position of key active site residues, increasing methyl donor binding¹¹. RAM also has a high-affinity RNA binding domain which may increase the rate of recruitment of transcripts in general or may have specificity for particular transcript motifs^{10,13}

In human cells, RNMT-RAM is recruited via the N-terminal domain of RNMT to RNA pol II proximal to transcription initiation sites^{14,15}. The RNMT N-terminal domain also impacts on cap methyltransferase activity. Its removal increases catalytic activity, suggesting that this domain may have some regulatory function. We questioned whether RNMT could be regulated by N-terminal domain post-translational modifications or interacting proteins, which could potentially change the conformation or action of the N-terminus. To address this question we performed proteomic analyses to map post-translational modifications of RNMT^{16,17}.

Our mass spectrometry analysis performed in HEK293 cells identified several phosphorylated, methylated and acetylated residues (Figure 2)¹⁶ (VHC/MA unpublished observations). All of the phosphorylation sites, along with others, were also indicated in the PhosphositePlus database, which contains data from high-throughput phosphoproteomic experiments from many research groups (Figure 2)¹⁷. Interestingly, the RNMT post-translational modifications were predominantly found in the N-terminus of RNMT and not in the catalytic domain. Sites of post-translational modification that arose in the catalytic domain may have been detrimental to activity and suppressed during evolution. The mammalian RNMT N-terminal domain may have evolved in order to allow regulation of cap methylation by signaling pathways. The phosphorylation sites found in the N-terminus cover a range of different amino acid motifs suggesting that several different kinases act on RNMT. Furthermore, the different methylation and acetylation events identified in our study demonstrate that several types of enzyme can modify RNMT. Therefore, the RNMT N-terminus is receiving a range of signaling

information from different pathways, some of which may be regulating RNMT and thus formation of the mRNA cap.

Regulation and function of RNMT T77 phosphorylation

In HeLa cells, Thr-77 is the predominant site of phosphorylation. We have also observed RNMT Thr-77 phosphorylation in embryonic stem cells, mammary epithelial cells, T lymphocytes and HEK293 cells ¹⁶ (VHC lab unpublished observations). RNMT Thr-77 is present in a CDK (cyclin-dependent kinase) recognition site adjacent to a cyclin binding site. The cell cycle is governed by the sequential activation of different CDK-cyclin complexes. RNMT Thr-77 phosphorylation levels oscillate throughout the cell cycle with highest phosphorylation levels during late S phase, through G2/M phase and into early G1 phase. The major kinase operational during these phases is CDK1-cyclin B1, although other CDK-cyclins are active at this time. We provided evidence in cells and *in vitro* that CDK1-cyclin B1 can phosphorylate RNMT directly. However, other CDK-cyclins may also phosphorylate RNMT Thr-77 and other proline-directed kinases

RNMT Thr-77 phosphorylation directly increases cap methyltransferase activity ¹⁶.

Presumably phosphorylation reverses some of the inhibitory effects of the N-terminal domain. The structure of the RNMT N-terminus has not been solved limiting molecular insight into Thr-77 function. Following reformation of the nuclear envelope after mitosis, factors involved in transcription are reassembled on chromatin and transcription initiates. Thus CDK1-dependent Thr-77 phosphorylation co-ordinates the peak of RNMT cap

methyltransferase activity with transcription initiation. Furthermore, high levels of transcripts carrying the mRNA cap may also contribute to the increase in cap-dependent translation that is observed during the initiation of G1 phase ¹⁸.

The mRNA cap binds to factors involved in transcription, processing and translation, and also protects the transcript from exonucleases ⁴. However, the precise role of the mRNA cap on these different gene expression processes on a genome-wide level is only beginning to be documented in mammalian cells. Mutation of RNMT Thr-77 to alanine (T77A) which blocks phosphorylation did not have a detectable effect on transcript level, despite significant depth of RNA sequencing analysis ¹⁶. However, the T77A mutation did inhibit the expression of a subset of proteins, and inhibited cell proliferation.

Although RNMT Thr-77 phosphorylation is cell cycle-dependent, the proteins whose expression is dependent on this modification were not found to be associated with synthesis at a specific stage of the cell cycle. It is important to highlight that this gene expression analysis was performed in unsynchronized cells and therefore a small effect on transcript levels during a specific phase of the cell cycle may have been undetected.

RNMT has a relatively long half-life in cells investigated to date (over 12 hours) (VHC lab unpublished observations). Therefore, the observation that RNMT Thr-77 phosphorylation is lost rapidly during G1 phase is likely to be the result of a rapidly acting phosphatase. The identity of the phosphatase or phosphatases involved is not known but a change in the activity or expression of these enzymes has the power to alter

Does RNMT Thr-77 have other functions?

Although RNMT Thr-77-dependent phosphorylation has a role in co-ordinating transcription and mRNA cap methylation during early G1, it is also highly phosphorylated at the end of S phase and through G2/M¹⁶. Although the majority of gene expression occurs during G1 phase of the cell cycle, translation has been described to occur at other phases including during mitosis^{19,20}. It may be that in certain circumstances, in particular cell lineages, the increase in mRNA cap methyltransferase activity driven by RNMT Thr-77 phosphorylation is needed to drive the expression of a subset of transcripts during late S or G2/M phase.

RNMT is imported into the nucleus following synthesis or after mitosis by interaction with importin-alpha^{21,22}. Three classical lysine-rich nuclear localisation signals lie in the RNMT N-terminus. Although we observed that phosphorylation of RNMT Thr-77 suppresses RNMT interaction with importin-alpha, in HeLa cells nuclear import of the RNMT T77A mutant or T77D mutant (phosphor mimic mutation) was not altered¹⁶. RAM carries a nuclear localization signal and thus in the presence of Thr-77 phosphorylation RNMT may rely on its interaction with RAM for nuclear import¹². However, it may be that in certain situations Thr-77 phosphorylation regulates RNMT

We observed that importin-alpha inhibits RNMT activity. If RNMT has a role in methylating mature cytoplasmic transcripts, T77 phosphorylation-dependent inhibition of importin-alpha binding would increase cytoplasmic cap methyltransferase activity, which

may have a positive effect on gene expression^{23,24}. This role could be particularly relevant when the nuclear envelope breaks down during mitosis and RNMT is dispersed into the cytoplasm. Increased RNMT activity and delocalisation from importin-alpha may be required for RNMT to function when it is dissociated from chromatin.

RNMT Thr-77 phosphorylation specificity

In HeLa cells, blocking RNMT Thr-77 phosphorylation did not affect the expression of all genes equally¹⁶. What is the nature of the specificity? Why are only certain genes dependent on RNMT Thr-77 phosphorylation? As with all enzymes, specificity comes from either the enzyme itself or from the availability of the substrate. RNMT Thr-77 is in the N-terminus, the domain that interacts with RNA pol II (probably indirectly)¹⁵. However, no differential recruitment of RNMT wild type and the T77A mutant was observed by chromatin immunoprecipitation performed on candidate genes. It remains a possibility that RNMT Thr-77 phosphorylation influences transcript recruitment in a gene-specific manner. Sequence motifs and structures in the 5' untranslated region of mRNAs may contribute to sensitivity to RNMT phosphorylation. Thus, analysis of the transcripts bound by RNMT/RAM using CLIP-Seq, or a more thorough analysis of the changes in translation rates of specific mRNAs using ribosome profiling/Ribo-Seq during specific stages of the cell cycle may provide further insights into the specificity of RNMT

Future perspectives

mRNA capping is now established as a gene expression mechanism which integrates cellular signaling pathways, resulting in reshaping of the gene expression landscape and alterations in cell physiology and fate. Transcription factors upregulate RNA pol II phosphorylation and methionine metabolism resulting in gene-specific regulation of mRNA cap methylation²⁵⁻²⁸. Furthermore, embryonic stem cell differentiation requires ERK1/2-dependent phosphorylation of RAM, which triggers its degradation and loss of pluripotency associated gene expression²⁹. It seems likely that many other signaling pathways will be regulating mRNA capping enzymes. The challenge going forward is to understand the role of mRNA capping regulation on a gene-specific level and how it impacts on mammalian development and disease.

Acknowledgements

Michael Aregger is funded by postdoctoral fellowship from the Swiss National Science Foundation. Victoria Cowling is funded by an Medical Research Council UK Senior Fellowship (MR/K024213/1) and Lister Institute Fellowship at the Centre for Gene Regulation and Expression, University of Dundee (Wellcome Trust Centre Award

Bibliography

- 1 Shuman, S. RNA capping: progress and prospects. *Rna* **21**, 735-737, doi:10.1261/rna.049973.115 (2015).
 - 2 Shatkin, A. J. Capping of eucaryotic mRNAs. *Cell* **9**, 645-653 (1976).
 - 3 Furuichi, Y. Discovery of m(7)G-cap in eukaryotic mRNAs. *Proc Jpn Acad Ser B Phys Biol Sci* **91**, 394-409, doi:10.2183/pjab.91.394 (2015).
 - 4 Ramanathan, A., Robb, G. B. & Chan, S. H. mRNA capping: biological functions and applications. *Nucleic Acids Res*, doi:10.1093/nar/gkw551 (2016).
 - 5 Topisirovic, I., Svitkin, Y. V., Sonenberg, N. & Shatkin, A. J. Cap and cap-binding proteins in the control of gene expression. *Wiley Interdiscip Rev RNA* **2**, 277-298, doi:10.1002/wrna.115 (2013).
- Gonatopoulos-Pournatzis, T. & Cowling, V. H. The Cap Binding Complex. *Biochem J* **457**, 231-242, doi:10.1042/BJ20131214 (2014).
- Belanger, F., Stepinski, J., Darzynkiewicz, E. & Pelletier, J. Characterization of hMTr1, a human Cap1 2'-O-ribose methyltransferase. *The Journal of biological chemistry* **285**, 33037-33044, doi:10.1074/jbc.M110.155283 (2010).
- Werner, M. *et al.* 2'-O-ribose methylation of cap2 in human: function and evolution in a horizontally mobile family. *Nucleic Acids Res* **39**, 4756-4768, doi:10.1093/nar/gkr038 (2011).
- Cowling, V. H. Regulation of mRNA cap methylation. *Biochem J* **425**, 295-302 (2009).
- Gonatopoulos-Pournatzis, T., Dunn, S., Bounds, R. & Cowling, V. H. RAM/Fam103a1 Is Required for mRNA Cap Methylation. *Mol Cell* **44**, 585-596, doi:10.1016/j.molcel.2011.11.017 (2012).

- 11 Varshney, D. *et al.* Molecular basis of RNA guanine-7 methyltransferase (RNMT) activation by RAM. *Nucleic Acids Res*, doi:10.1093/nar/gkw637 (2016).
- 12 Saha, N., Schwer, B. & Shuman, S. Characterization of human, *Schizosaccharomyces pombe*, and *Candida albicans* mRNA cap methyltransferases and complete replacement of the yeast capping apparatus by mammalian enzymes. *The Journal of biological chemistry* **274**, 16553-16562 (1999).
- 13 Gonatopoulos-Pournatzis, T. & Cowling, V. H. RAM function is dependent on Kapbeta2-mediated nuclear entry. *Biochem J* **457**, 473-484, doi:10.1042/BJ20131359 (2014).
- 14 Glover-Cutter, K., Kim, S., Espinosa, J. & Bentley, D. L. RNA polymerase II pauses and associates with pre-mRNA processing factors at both ends of genes. *Nat Struct Mol Biol* **15**, 71-78 (2008).
- 15 Aregger, M. & Cowling, V. H. Human cap methyltransferase (RNMT) N-terminal non-catalytic domain mediates recruitment to transcription initiation sites. *Biochem J* **455**, 67-73, doi:10.1042/BJ20130378 (2013).
- Aregger, M. *et al.* CDK1-Cyclin B1 Activates RNMT, Coordinating mRNA Cap Methylation with G1 Phase Transcription. *Mol Cell* **61**, 734-746, doi:10.1016/j.molcel.2016.02.008 (2016).
- Hornbeck, P. V. *et al.* PhosphoSitePlus: a comprehensive resource for investigating the structure and function of experimentally determined post-translational modifications in man and mouse. *Nucleic Acids Res* **40**, D261-270, doi:10.1093/nar/gkr1122 (2012).
- Pyronnet, S. & Sonenberg, N. Cell-cycle-dependent translational control. *Curr Opin*
- Stumpf, C. R., Moreno, M. V., Olshen, A. B., Taylor, B. S. & Ruggero, D. The translational landscape of the mammalian cell cycle. *Mol Cell* **52**, 574-582, doi:10.1016/j.molcel.2013.09.018 (2013).

- 20 Tanenbaum, M. E., Stern-Ginossar, N., Weissman, J. S. & Vale, R. D. Regulation of mRNA translation during mitosis. *Elife* **4**, doi:10.7554/eLife.07957 (2015).
- 21 Shafer, B., Chu, C. & Shatkin, A. J. Human mRNA cap methyltransferase: alternative nuclear localization signal motifs ensure nuclear localization required for viability. *Mol Cell Biol* **25**, 2644-2649 (2005).
- 22 Wen, Y. & Shatkin, A. J. Cap methyltransferase selective binding and methylation of GpppG-RNA are stimulated by importin-alpha. *Genes Dev* **14**, 2944-2949 (2000).
- 23 Kiss, D. L., Oman, K., Bundschuh, R. & Schoenberg, D. R. Uncapped 5' ends of mRNAs targeted by cytoplasmic capping map to the vicinity of downstream CAGE tags. *FEBS Lett* **589**, 279-284, doi:10.1016/j.febslet.2014.12.009 (2015).
- 24 Mukherjee, C. *et al.* Identification of cytoplasmic capping targets reveals a role for cap homeostasis in translation and mRNA stability. *Cell Rep* **2**, 674-684, doi:10.1016/j.celrep.2012.07.011 (2012).
- 25 Aregger, M. & Cowling, V. H. E2F1-dependent methyl cap formation requires RNA pol II phosphorylation. *Cell Cycle* **11**, 2146-2148, doi:10.4161/cc.20620 (2012).
- Fernandez-Sanchez, M. E., Gonatopoulos-Pournatzis, T., Preston, G., Lawlor, M. A. & Cowling, V. H. S-adenosyl homocysteine hydrolase is required for Myc-induced mRNA cap methylation, protein synthesis, and cell proliferation. *Mol Cell Biol* **29**, 6182-6191, doi:10.1128/MCB.01117-09 (2009).
- Cole, M. D. & Cowling, V. H. Specific regulation of mRNA cap methylation by the c-Myc and E2F1 transcription factors. *Oncogene* **28**, 1169-1175 (2009).
- Cowling, V. H. & Cole, M. D. The Myc transactivation domain promotes global phosphorylation of the RNA polymerase II carboxy-terminal domain independently of direct DNA binding. *Mol Cell Biol* **27**, 2059-2073 (2007).
- 29 Grasso, L. *et al.* mRNA Cap Methylation in Pluripotency and Differentiation. *Cell Rep* **16**, 1352-1365, doi:10.1016/j.celrep.2016.06.089 (2016).




Figure 1 mRNA cap structure and mRNA capping enzymes

Structure of mRNA cap and action of mRNA capping enzymes.




Figure 2 Post-translational modifications of RNMT

Human RNMT 1-120 is the N-terminal domain and RNMT 121-476 is the catalytic domain and interacts with RAM. Above depiction of RNMT are phosphorylation sites (P) obtained from PhosphositePlus (sites with only 1 mass spectrometry/high-throughput proteomics reference excluded). Below depiction of RNMT are phosphorylation, acetylation (Ac) and methylation (Me) sites identified from analysis in Aregger et al., 2016.