Interaction and Signalling Networks

Kirkpatrick, Clare L.; Lesouhaitier, Oliver; Malone, Jacob G.; An, Shi-Qi; Caly, Delphine L.

Published in:
Microbiology

DOI:
10.1099/mic.0.000401

Publication date:
2017

Document Version
Peer reviewed version

Microbiology

Interaction and Signalling Networks: a report from the fourth 'Young Microbiologists Symposium on Microbe Signalling, Organisation and Pathogenesis'

--Manuscript Draft--

Manuscript Number:	MIC-D-16-00311R1
Full Title:	Interaction and Signalling Networks: a report from the fourth 'Young Microbiologists Symposium on Microbe Signalling, Organisation and Pathogenesis'
Short Title:	A report from the fourth Young Microbiologists Symposium
Article Type:	Meeting Report
Section/Category:	Host-microbe interaction
Corresponding Author:	Delphine L Caly Institut Charles Viollette Villeneuve d'Ascq, FRANCE
First Author:	Clare L Kirkpatrick
Order of Authors:	Clare L Kirkpatrick
	Oliver Lesouhaitier
	Jacob G Malone
	Shi-Qi An
	Delphine L Caly

Interaction and Signalling Networks: a report from the fourth 'Young Microbiologists Symposium on Microbe Signalling, Organisation and Pathogenesis'

Clare L. Kirkpatrick[1], Olivier Lesouhaitier[2], Jacob G. Malone[3,4], Shi-Qi An[5] and Delphine L. Caly[6,*]

[1]Department of Microbiology & Molecular Medicine, Institute of Genetics & Genomics in Geneva (iGE3), Faculty of Medicine/CMU, University of Geneva, Switzerland

[2]Laboratory of Microbiology Signals and Microenvironnement LMSM, EA 4312, Normandie Université, Université de Rouen Evreux, France

[3]John Innes Centre, Norwich Research Park, Norwich, United Kingdom

[4]University of East Anglia, Norwich, United Kingdom

[5]Division of Molecular Microbiology, School of Life Sciences, University of Dundee

[6]Univ. Lille, EA 7394, ICV - Institut Charles Viollette, Lille, France.

* Corresponding author : Delphine L. Caly, Institut Charles Viollette, Cite Scientifique, 59655 Villeneuve d'Ascq, France. Email : Delphine.Caly@polytech-lille.fr.

Running title: A report from the fourth Young Microbiologists Symposium

Keywords: gene regulation, signalling, secretion, host-pathogen interactions, microbe-microbe interactions.

Abstract

At the end of June, over 120 microbiologists from 18 countries gathered in Dundee, Scotland for the fourth edition of the Young Microbiologists Symposium on "Microbe Signalling, Organisation and Pathogenesis". The aim of the symposium was to give early career microbiologists the opportunity to present their work in a convivial environment and to interact with senior world-renowned scientists in exciting fields of microbiology research. The meeting was supported by the Microbiology Society, the Society of Applied Microbiology, the American Society for Microbiology with further sponsorship from the European Molecular Biology Organisation and The Royal Society of Edinburgh. In this report, we highlight some themes that emerged from the many interesting talks and poster presentations, and some of the other activities that were on offer at this energetic meeting.

Introduction

The fourth Young Microbiologists Symposium (YMS2016) took place at the Apex City Quay Hotel in Dundee, Scotland on the 29th and 30th June 2016. The conference gathered 126 scientists coming from 18 countries and was organized by **Helge Dorfmueller** and **Robert Ryan**, from University of Dundee, and **Delphine Caly** from University of Lille in France. The main objective of the YMS2016 was to bring together early career microbiologists. The symposium programme covered several hot topics in microbiology and touched on current areas of interest to microbiologists including intracellular signalling, antibiotic resistance, bacterial secretion and host-microbe interactions. Renowned experts, who led sessions, and the many junior microbiologists who attended provided insight and new findings into these exciting areas. A novelty to this year's meeting was that participants were given the opportunity to attend a PLOS Pathogens writing and publishing workshop, chaired by **Neil Mabbott** from

the Roslin Institute and University of Edinburgh in Scotland, which provided valuable advice for PhD students and junior post-docs on how to write scientific papers and achieve successful publication.

Sensing, transduction and intracellular signalling

The YMS2016 kicked off with the FEMS keynote lecture from **Ute Römling** (Karolinska Institutet, Sweden), who described the identification of the *Pseudomonas aeruginosa* clone C strain cluster prevalent in patients, clinics and the environment worldwide. As part of this research, Ute discussed how her group identified the PACGI-1 genomic island in this cluster, and showed that it contributes to heat-shock resistance by encoding protein quality-control systems (Lee *et al.*, 2015). Next, Ute described her group's work on the ubiquitous bacterial second messenger signal cyclic-di-GMP in *Salmonella enterica* serovar Typhimurium, which controls rdar (red dry and rough) biofilm formation and virulence as part of a complex regulatory network involving the transcriptional regulator CsgD. Ute explained how her lab have identified and characterised several key players in this network, including the diguanylate cyclase AdrA, the cellulose synthase cyclic di-nucleotide-binding protein BcsE, and the degenerate phosphodiesterase STM1697, which controls flagellar gene transcription through binding to the master regulator FlhDC (Ahmad *et al.*, 2013; Le Guyon *et al.*, 2015) and gave perspectives on novel regulatory pathways.

These themes were built upon in the first session, which was opened by **Max Dow** (University College Cork, Ireland). Max discussed the structure-function relationship of HD-GYP domains which degrade the second messenger cyclic-di-GMP. Max began with a summary of his lab's work on the protein RpfG, which contains a HD-GYP domain, and controls virulence and motility in the plant pathogen *Xanthomonas campestris* (Ryan *et al.*, 2010). Recently, Max and

collaborators have determined the structures of PmGH, an enzymatically active HD-GYP protein from *Persephonella marina* (Bellini et al., 2014) and PA2572, an enzymatically-inactive YN-GYP variant from *P. aeruginosa* (Bellini *et al.*, unpublished). The work on PmGH suggested that active HD-GYP domains could be sub-divided into those with two or three metal-ion cofactors. In contrast, PA2572 carried no metals but was able to interact with other proteins via the GYP 'loop'.

Lisa Bowman (Imperial College London, UK) described a second, equally interesting dinucleotide second messenger; cyclic-di-AMP. Pioneering work from the Gründling lab has shown that cyclic-di-AMP regulates potassium and osmolyte uptake in *Staphylococcus aureus*, and is produced by the membrane bound cyclase DacA (Corrigan *et al.*, 2011). Lisa discussed her work to expand on the existing model for cyclic-di-AMP signalling by explaining her inventive use of a BioLog phenotypic microarray to determine the function of YbbR, an uncharacterised component of the DacA membrane protein complex. Based on this screen and suppressor mutagenesis, Lisa proposed that YbbR acts as a localisation determinant for DacA at the membrane, controlling local pools of c-di-AMP especially under stress conditions.

In the final talk in this session, **Francesca D'Angelo** (University Roma Tre, IT) attracted significant interest and many audience questions with her talk on the generation of synthetic cells. These synthetic cells consist of liposomes containing biological molecules, and represent an ambitious new approach to drug delivery (Stano *et al.*, 2012). After demonstrating that the HSL signal could be produced *in vitro*, Francesca built on this by encapsulating the functional HSL production system in her synthetic cells, protecting the HSL pathway from externally added inhibitors. The next step for this project will be to generate synthetic cells that can sense signals as well as produce an output.

Symbiosis, pathogenesis and mechanisms of host interaction

The ASM keynote lecture was presented by **Scott Hultgren** (Washington University, USA). Scott gave a fantastic and informative overview of his research into urinary tract infections (UTIs) by *E. coli*, which are mediated by the activities of type I pili. Building on structural models of pili, Scott first showed that high and low-affinity mannose-binding forms of the terminal FimH adhesin exist in equilibrium, with both states required for effective infection. He then moved on to a discussion of the clinical aspects of UTI, showing that bladder cells are remodelled by sensitisation to UTI, and thereafter are significantly more likely to become re-infected. Scott's talk finished with a description of several promising lines of research into UTI treatment, including an anti-pilus vaccines, and drugs targeting both pili and the FimH adhesin.

The host-microbe interactions session covered a large spectrum of topics introduced in the ASM lecture including polymicrobial infection, the use of new tools for studying host-microbe interactions in real time and the impact of both host communication signals and small metabolic compounds.

Marvin Whiteley (University of Texas, USA) showed that microbe-microbe interactions increase bacterial resistance to host defences (Ramsey & Whiteley, 2009) and allow synergistic effect for some pathogenic bacteria (Turner *et al.*, 2015), using various examples of interactions, such as *P. aeruginosa* and *S. aureus* in the cystic fibrosis lungs or *Aggregatibacter actinomycetemcomitans* and *Streptococcus gordonii* that form biofilms in the oral cavity. The highly organised wound communities and the precise spacing between bacteria during polymicrobial infection are required for infectious success (Stacy *et al.*, 2015), and Marvin explained why understanding this process could help in improving therapeutic strategies. The following talk was given by **Andrew Roe** (University of Glasgow, UK) who presented a new tool for studying protein interactions specifically dedicated to the host-pathogen interaction

research field. This tool, named LOV for light-oxygen-voltage sensing domain, enables the visualisation of bacterial cells attached to host cells. In parallel, Andrew showed how the LOV tool could be very suitable to study the direct translocation of bacterial type III effectors into host cells. Andrew's talk was illustrated by amazing images obtained by the fusion of a LOV-based reporter with the *Shigella flexneri* effector IpaB, demonstrating the interaction with the host cell actin network (Gawthorne *et al.*, 2016).

The use of mass spectrometry imaging in microbiology was discussed by **Heather Hulme** (University of Glasgow, UK), who showed that it could be a valuable tool for identifying biomarkers during an infection process. Using the example of mesenteric lymph node infection by *Salmonella*, Heather showed that palmitoylcarnitine (PalC), which is localised and accumulates in the damaged infected tissue, could be measured and used as a potential biomarker of infection.

The host environment encountered by bacteria plays a role in the success of infections. In this context, **Tuuli Ahlstrand** (University of Turku, Finland) showed that biofilms formed by the opportunistic pathogen *A. actinomycetemcomitans* could disrupt the host inflammation response by binding and internalising the proiflammatory cytokine interleukin-1β (Paino *et al.*, 2012), which is enhanced by a specific bacterial sensor named bacterial interleukin receptor I (BilRI) (Ahlstrand *et al.*, 2016; Paino *et al.*, 2013). In the same vein, **James Connolly** (University of Glasgow, UK) demonstrated how pathogenic *E. coli* integrates host signals in order to regulate its ability to colonize the urinary tract. More precisely, James demonstrated how D-serine influences both gene content and virulence factor expression in pathogenic *E. coli* (Connolly *et al.*, 2015) and how bacteria use a D-serine sensing system to adapt to their environment (Connolly *et al.*, 2016). Another way to prevent bacterial infection, using inhibitors of multivalent adhesion molecule 7 (MAM7), was described by **Daniel Stones** (University of Birmingham, UK) who described a bead-coupled recombinant MAM7 that not

only prevented bacterial adhesion and infection in rats, but also did not affect cytokines release and the wound healing process, suggesting a promising drug to counteract infection (Krachler *et al.*, 2011).

Bacterial shape, secretion and development

This session began and ended with a review of new developments in our understanding of the operation of the bacterial type VI secretion system (T6SS). This multi-protein complex is a delivery system for protein-based toxins targeted at other bacteria or at eukaryotic cells, while the bacteria that are the source of the toxins also express specific immunity proteins to protect themselves. **Alain Filloux** (Imperial College London, UK) presented a recently published structural study (Planamente *et al.*, 2016), focused on a previously uncharacterised component of the complex, the TssA baseplate. The Filloux group showed that TssA forms a circular baseplate-like structure that assembles onto the membrane-facing end of the TssBC sheath, sharing structural and functional homology with the gp6 baseplate of T4 bacteriophage, and is essential for T6SS activity.

Bacterial lifestyle changes often require remodelling of the cell envelope, whether to permit the entry of extracellular DNA during competence or to generate a spore that will be more resistant to the external environment than the mother cell from which it develops. **Emma Denham** (University of Warwick, UK) presented her group's ongoing work on the role of small RNAs in bacterial growth heterogeneity using *Bacillus subtilis* as their model system. This talk focused on one notable sRNA-controlled process, the AbrB-dependent transition from exponential to stationary phase (Mars *et al.*, 2015), where AbrB expression is regulated by the small RNA S1022. Modified AbrB levels lead to phenotypic heterogeneity, suggesting a novel sRNA-regulated bet-hedging strategy.

Tessa Quax (University of Freiburg, Germany) provided the conference's only talk on Archaea, specifically on archaellum-mediated motility in these organisms. Named "archaellum" due to its extreme structural difference to the bacterial flagellum, this substructure resembles the type IV pili seen in bacteria in terms of its components and assembly mechanism. Surprisingly, Tessa showed it can also interact with a CheY-like component of a chemotaxis system as the bacterial flagellum does, despite the extreme evolutionary divergence between these two kingdoms of life and the completely different composition of their respective motility organelles. Finally, **Francesca Cianfanelli** from the Coulthurst group (University of Dundee, UK) presented her work on the T6SS of *Serratia marcescens* and the specific interactions of VgrG and PAAR proteins at the tip of the T6SS "spike". This showed that PAAR proteins are essential for T6SS function and that particular VgrG-PAAR combinations are required for full T6SS-dependent antibacterial activity, including activity mediated by cargo adaptors that are not normally considered dependent on specific VgrG proteins (Cianfanelli *et al.*, 2016).

Bacterial inter-species and inter-kingdom interactions

The final session covered the topic of inter-species and inter-kingdom interactions, which included talks regarding interactions within complex communities, between microbes, and the various host signals/triggers that shape the interactions within these communities. A captivating example of the former was presented by **Christoph Tang** (University of Oxford, UK) who delivered the EMBO lecture. Christoph described that temperature is one of the most important environmental cues that act on regulatory networks of pathogenic microbes. His group discovered and characterised the RNA thermometer CssA from *Neisseria meningitidis*, an elegant mechanism that this microbe uses to adapt to different temperature changes.

Christoph explained how using NMR spectroscopy and SHAPE (Selective 2′-OH acylation analysed by primer extension) assays, the group discovered that at low temperature (30°C), all base pair regions of CssA are stably formed, and the ribosome cannot access the RBS which is fully occluded (Barnwal *et al.*, 2016). As the temperature is raised, the RNA structure starts to unfold and by 42°C, the thermometer structure is fully open, leading to efficient translation. Taken together, it suggests that CssA acts as a rheostat, whose stability is optimized to respond in a small temperature range such as occurs within the upper airways during infection.

Continuing with the theme of environmental cues altering the response of the microbial community during infection, **Vanessa Sperandio** (UT Southwestern Medical Center, USA) showed that enterohaemorrhagic *E. coli* (EHEC) senses fucose cleaved from the mucus layer in the colon by *Bacteroides thetaiotaomicron* through the histidine kinase FusK. It then rewires its transcription, repressing the expression of the LEE and fucose utilisation genes (Pacheco *et al.*, 2012). However, without mucus as a carbon source, *B. thetaiotaomicron* starts to secrete succinate, which upon being taken up by EHEC is sensed by the Cra transcription factor as a clue to a gluconeogenic environment. Cra binds to another transcription factor, KdpE, which is a response regulator (RR) phosphorylated by the QseC adrenergic sensor, to integrate adrenergic and sugar sensing to activate virulence gene expression at the interface with the intestinal epithelium. Through the interaction with another RR; QseB, QseC also represses the expression of the *fusKR* genes, further derepressing the virulence regulon. These data suggest a new layer of complexity in the inter kingdom signalling that underlies EHEC pathogenicity.

Given what is now known regarding the contribution of the host microbiota to health there is an urgent need for relevant animal models. **Beckie Ingram** (Queens College Belfast, UK) gave an inspiring talk about her group's work on developing appropriate murine models for understanding the pathophysiology of lung inflammation and the pathogenesis of lung disease in cystic fibrosis. These approaches will become crucial in improving our understanding of

microbial community interactions in the field of infectious diseases. Finally, **Clare Kirkpatrick** (University of Geneva, Switzerland) discussed the role of toxin-antitoxin (TA) systems in bacterial interactions and how they can shape the community. Clare discussed her recent work on the HigBA system from *Caulobacter crescentus* and revealed that this TA system acts as a switch to regulate bacterial growth and induce cell death upon antibiotic-induced DNA damage (Kirkpatrick *et al.*, 2016). This novel regulatory mechanism could potentially be used to develop new treatments to clear bacterial infections.

Conclusions

This symposium, like previous meetings (Caly *et al.*, 2012, 2014; Ryan *et al.*, 2009), covered many fascinating areas of microbiology. As always the forum allowed the attendees to gain many insights into up and coming areas and techniques in bacteriology, and provided junior microbiologists the opportunity to present and discuss their work. This was successfully achieved judging the numerous interactions between junior and senior scientists observed during and between scientific sessions.

After the final session, a number of awards were distributed. These included the Frontiers in Microbiology short talk prize that went to **Fang-Fang Wang** (Chinese Academy of Sciences Beijing, China) for her excellent presentation entitled, "Receptor histidine kinase directly binds plant chemical to promote bacterial adaptation in host plant". The Nature Reviews in Microbiology, Trends in Microbiology, Biochemical Journal and Molecular Microbiology poster prizes went to several PhD students working on outstanding projects. The meeting finished on relaxed note with a Ceilidh organised in the Apex hotel following the conference dinner.

Overall, the feedback from attendees was very positive; participants appreciated the quality of the scientific programme and the intimate atmosphere of the small conference. A post-meeting survey reported that 71% of the survey participants (n = 68) found the scientific programme 'very good' and 83% were interested in attending a future YMS conference (n = 65). One of the participants, who gave a talk as a junior post-doc at the YMS2012 and is now setting up her laboratory, used this opportunity to advertise for positions and made several promising contacts. This bodes well for further iterations of the meeting in the future.

Acknowledgements

We are deeply grateful to the participants who agreed for their work to be described in this report and we apologise to those whose work could not be mentioned due to space constraints. We also would like to thank all the speakers and participants for contributing to the success of this meeting. The organisers are extremely grateful to Erin Stanbridge, Kushal Rugjee, Birte Hollmann, Anne Six, the members of the Division of Molecular Microbiology in University of Dundee and Debbie Ree from the Dundee and Angus Convention Bureau for their help and support with the organisation of the meeting. We also thank the American Society for Microbiology, the European Molecular Biology Organization, the Federation of European Microbiological Societies, the Microbiology Society, the Society for Applied Microbiology and the Royal Society of Edinburgh and all our other sponsors for their financial support.

References

Ahlstrand, T., Tuominen, H., Beklen, A., Torittu, A., Oscarsson, J., Sormunen, R., Pöllänen, M. T., Permi, P. & Ihalin, R. (2016). A novel intrinsically disordered outer membrane lipoprotein of Aggregatibacter actinomycetemcomitans binds various cytokines and plays a role in biofilm response to interleukin-1β and interleukin-8.

Virulence 0.

Ahmad, I., Wigren, E., Le Guyon, S., Vekkeli, S., Blanka, A., el Mouali, Y., Anwar, N., Chuah, M. L., Lünsdorf, H. & other authors. **(2013).** The EAL-like protein STM1697 regulates virulence phenotypes, motility and biofilm formation in *Salmonella typhimurium*. *Mol Microbiol* **90**, 1216–1232.

Barnwal, R. P., Loh, E., Godin, K. S., Yip, J., Lavender, H., Tang, C. M. & Varani, G. (2016). Structure and mechanism of a molecular rheostat, an RNA thermometer that modulates immune evasion by *Neisseria meningitidis*. *Nucleic Acids Res* gkw584. Oxford University Press.

Bellini, D., Caly, D. L., Mccarthy, Y., Bumann, M., An, S. Q., Dow, J. M., Ryan, R. P. & Walsh, M. A. (2014). Crystal structure of an HD-GYP domain cyclic-di-GMP phosphodiesterase reveals an enzyme with a novel trinuclear catalytic iron centre. *Mol Microbiol* **91**, 26–38.

Caly, D. L., Coulthurst, S. J., An, S. -q., Helaine, S., Malone, J. G. & Ryan, R. P. (2014). Communication, Cooperation, and Social Interactions: a Report from the Third Young Microbiologists Symposium on Microbe Signalling, Organisation, and Pathogenesis. *J Bacteriol* **196**, 3527–3533.

Caly, D. L., Coulthurst, S. J., Geoghegan, J. A., Malone, J. G. & Ryan, R. P. (2012). Socializing, networking and development: a report from the second 'Young Microbiologists Symposium on Microbe Signalling, Organization and Pathogenesis'. *Mol Microbiol* **86**, 501–512.

Cianfanelli, F. R., Alcoforado Diniz, J., Guo, M., De Cesare, V., Trost, M., Coulthurst, S. J., Costa, T., Felisberto-Rodrigues, C., Meir, A. & other authors. **(2016).** VgrG and PAAR Proteins Define Distinct Versions of a Functional Type VI Secretion System.

PLOS Pathog **12**, e1005735 (E. Cascales, Ed.). Public Library of Science.

Connolly, J. P. R., Gabrielsen, M., Goldstone, R. J., Grinter, R., Wang, D., Cogdell, R. J., Walker, D., Smith, D. G. E. & Roe, A. J. (2016). A Highly Conserved Bacterial D-Serine Uptake System Links Host Metabolism and Virulence. *PLoS Pathog* **12**, e1005359. Public Library of Science.

Connolly, J. P., Goldstone, R. J., Burgess, K., Cogdell, R. J., Beatson, S. A., Vollmer, W., Smith, D. G. & Roe, A. J. (2015). The host metabolite D-serine contributes to bacterial niche specificity through gene selection. *ISME J* **9**, 1039–1051. Nature Publishing Group.

Corrigan, R. M., Abbott, J. C., Burhenne, H., Kaever, V. & Gründling, A. (2011). c-di-AMP Is a New Second Messenger in Staphylococcus aureus with a Role in Controlling Cell Size and Envelope Stress. *PLoS Pathog* **7**, e1002217 (A. Cheung, Ed.). Public Library of Science.

Gawthorne, J. A., Audry, L., McQuitty, C., Dean, P., Christie, J. M., Enninga, J. & Roe, A. J. (2016). Visualizing the Translocation and Localization of Bacterial Type III Effector Proteins by Using a Genetically Encoded Reporter System. *Appl Environ Microbiol* **82**, 2700–2708 (T. E. Besser, Ed.). American Society for Microbiology.

Le Guyon, S., Simm, R., Rehn, M. & Römling, U. (2015). Dissecting the cyclic di-guanylate monophosphate signalling network regulating motility in *Salmonella enterica* serovar Typhimurium. *Environ Microbiol* **17**, 1310–1320.

Kirkpatrick, C. L., Martins, D., Redder, P., Frandi, A., Mignolet, J., Chapalay, J. B., Chambon, M., Turcatti, G. & Viollier, P. H. (2016). Growth control switch by a DNA-damage-inducible toxin–antitoxin system in Caulobacter crescentus. *Nat Microbiol* 16008. Nature Publishing Group.

315 **Krachler, A. M., Ham, H. & Orth, K. (2011).** Outer membrane adhesion factor multivalent
316 adhesion molecule 7 initiates host cell binding during infection by Gram-negative
317 pathogens. *Proc Natl Acad Sci* **108**, 11614–11619. National Academy of Sciences.

318 **Lee, C., Wigren, E., Trček, J., Peters, V., Kim, J., Hasni, M. S., Nimtz, M., Lindqvist,**
319 **Y., Park, C. & other authors**. **(2015).** A novel protein quality control mechanism
320 contributes to heat shock resistance of worldwide-distributed *P seudomonas aeruginosa*
321 clone C strains. *Environ Microbiol* **17**, 4511–4526.

322 **Mars, R. A. T., Nicolas, P., Ciccolini, M., Reilman, E., Reder, A., Schaffer, M., Mäder,**
323 **U., Völker, U., van Dijl, J. M. & other authors**. **(2015).** Small Regulatory RNA-
324 Induced Growth Rate Heterogeneity of Bacillus subtilis. *PLOS Genet* **11**, e1005046 (D.
325 B. Kearns, Ed.). Public Library of Science.

326 **Pacheco, A. R., Curtis, M. M., Ritchie, J. M., Munera, D., Waldor, M. K., Moreira, C.**
327 **G. & Sperandio, V. (2012).** Fucose sensing regulates bacterial intestinal colonization.
328 *Nature* **492**, 113–7. NIH Public Access.

329 **Paino, A., Lohermaa, E., Sormunen, R., Tuominen, H., Korhonen, J., Pöllänen, M. T. &**
330 **Ihalin, R. (2012).** Interleukin-1β is internalised by viable Aggregatibacter
331 actinomycetemcomitans biofilm and locates to the outer edges of nucleoids. *Cytokine*
332 **60**, 565–574.

333 **Paino, A., Ahlstrand, T., Nuutila, J., Navickaite, I., Lahti, M., Tuominen, H., Välimaa,**
334 **H., Lamminmäki, U., Pöllänen, M. T. & other authors**. **(2013).** Identification of a
335 Novel Bacterial Outer Membrane Interleukin-1B-Binding Protein from Aggregatibacter
336 actinomycetemcomitans. *PLoS One* **8**, e70509 (J. A. Bengoechea, Ed.). Public Library
337 of Science.

338 **Planamente, S., Salih, O., Manoli, E., Albesa-Jové, D., Freemont, P. S., Filloux, A.,**

Aksyuk, A., Leiman, P., Kurochkina, L. & other authors. **(2016).** TssA forms a gp6-like ring attached to the type VI secretion sheath. *EMBO J* **28**, e201694024. EMBO Press.

Ramsey, M. M. & Whiteley, M. (2009). Polymicrobial interactions stimulate resistance to host innate immunity through metabolite perception. *Proc Natl Acad Sci* **106**, 1578–1583. National Academy of Sciences.

Ryan, R. P., Mccarthy, Y., Andrade, M., Farah, C. S., Armitage, J. P., Dow, J. M. & Lindow, S. E. (2010). Cell–cell signal-dependent dynamic interactions between HD-GYP and GGDEF domain proteins mediate virulence in Xanthomonas campestris. *Proc Natl Acad Sci* **107**, 5989–5994.

Ryan, R. P., Romeo, T., De Keersmaecker, S. C. J. & Coulthurst, S. J. (2009). Nurturing scientific mutualism: A report from the 'Young Microbiologists Mini-Symposium on microbe signalling, organisation and pathogenesis'. *Mol Microbiol* **73**, 760–774. Blackwell Publishing Ltd.

Stacy, A., McNally, L., Darch, S. E., Brown, S. P. & Whiteley, M. (2015). The biogeography of polymicrobial infection. *Nat Rev Microbiol* **14**, 93–105. Nature Research.

Stano, P., Rampioni, G., Carrara, P., Damiano, L., Leoni, L. & Luisi, P. L. (2012). Semi-synthetic minimal cells as a tool for biochemical ICT. *Biosystems* **109**, 24–34.

Turner, K. H., Wessel, A. K., Palmer, G. C., Murray, J. L. & Whiteley, M. (2015). Essential genome of *Pseudomonas aeruginosa* in cystic fibrosis sputum. *Proc Natl Acad Sci* **112**, 4110–4115. National Academy of Sciences.

362