



University of Dundee

Targeting BRAF-mutant tumours with TGFBR1 inhibitors

Spender, Lindsay C.; Inman, Gareth J.

Published in:
Aging

DOI:
[10.18632/aging.101169](https://doi.org/10.18632/aging.101169)

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Spender, L. C., & Inman, G. J. (2017). Targeting BRAF-mutant tumours with TGFBR1 inhibitors. *Aging*, 9(1), 5-6. <https://doi.org/10.18632/aging.101169>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Targeting BRAF-mutant tumours with TGFBR1 inhibitors

Lindsay C Spender and Gareth J Inman

Transforming growth factor beta (TGF β) is a major contributing factor to many diseases including progressive cancers [1]. Given the normal homeostatic functions of TGF β in controlling cell proliferation, immune surveillance and angiogenesis, however, it is much less clear which cancer patients would derive clinical benefit from the administration of anti-TGF β targeted therapy. The use of these therapies is complicated not only by toxicity issues, but also by concerns that interfering with tumour suppressive functions of TGF β may ultimately exacerbate disease. Thus, identifying when and where TGF β signalling promotes tumour progression [2] is key to the success of numerous approaches to block TGF β signalling using a range of humanised antibodies, small molecule inhibitors (SMIs) and antisense oligonucleotides currently being evaluated in clinical trials [3]. Our recent data now provides insight into when these therapies might succeed [4].

To investigate when anti-TGF β targeted therapies might be used appropriately, we screened cancer cell lines carrying defined cancer-associated genetic abnormalities in the RAS/RAF/MEK/ERK signalling pathway for sensitivity to SMIs of the TGF β type 1 receptor (TGFBR1). In assays specifically designed to measure cancer-stem cell like properties including clonogenicity and anchorage independent growth assays, we found that cells carrying mutant BRAF were dependent on TGF β signalling for growth. Although cells carrying mutant KRAS (n=11), NRAS (n=4) and BRAF (n=7) all exhibit constitutive activation of the MAPK pathway, only mutant BRAF cells were universally inhibited by the TGFBR1 inhibitor. Our data were confirmed by siRNA knockdown of the receptor to ensure that the effects of the inhibitor were on-target. Additionally, the dependence of mutant BRAF cells on secreted autocrine TGF β signalling for formation of xenograft tumours was demonstrated by stably expressing short-hairpin shRNA targeting TGF β RNA in the cancer cells. The ability of these cells to establish tumours in mice was significantly impaired by TGF β knockdown. The data suggests, therefore, that mutant BRAF cells are 'hard-wired' to depend on autocrine TGF β signalling for growth in stressful conditions. The data also implies that the presence of mutant BRAF in

sequenced tumour biopsies could act as a biomarker for stratifying patients for anti-TGF β therapy. The mechanism of TGF β -mediated tumour cell growth promotion remains elusive, but appears to be independent of canonical TGF β signalling pathways involving the SMAD transcription factors and may involve activation of the small GTPase RHO-A. It might be misleading, therefore, to use the phosphorylation of SMADs downstream of TGF β receptor activation as an additional biomarker in tumour tissue biopsies to indicate the potential for TGFBR1 inhibitor use in cancer.

The patients most likely to benefit include melanoma cancer patients since mutation of BRAF is detected in approximately 50% of melanomas [5]. Melanoma is particularly prevalent, and mortality highest, in the aging population with more than 80% of deaths from melanoma occurring in people aged over 50 (www.cancerresearch.org). The discovery of the genetic abnormalities associated with melanoma has resulted in an astonishing effort from researchers and the pharmaceutical industry alike to develop and test small molecule inhibitors of mutant BRAF (BRAFi). While these drugs (e.g. vemurafenib) have revolutionised treatment and increased patient survival, chemo-resistance remains a significant clinical issue and the majority of patients relapse and die from drug-resistant metastatic disease. Our analysis of the effect of TGF β inhibitors in both drug-naïve and vemurafenib-resistant patient derived cells therefore provides valuable information relating to when and where TGF β inhibitors might be effective.

We found that TGFBR1 inhibitors remained effective against vemurafenib-resistant patient derived cells. Furthermore, TGF β inhibitors prevented the enhanced cell growth caused by paradoxical activation of the MAPK pathway seen in cells treated with BRAFi. Taken together targeting TGF β signalling in mutant BRAF melanoma is predicted to inhibit tumour growth even in drug-resistant disease, however, the tumour cell microenvironment appears to impact significantly on inhibitor efficacy. We found that cell density reduced the ability of TGFBR1 inhibitors to prevent melanoma growth. In these less stressful conditions, melanoma cells were not dependent on TGF β signalling to the

same extent. The addition melanoma cells have to TGF β signalling correlates with isolation and anchorage independence - situations encountered during tumour spread and outgrowth of metastasis. Using zebrafish xenograft models of melanoma cell metastasis we were able to track single fluorescently tagged cells during the process of migration into tail-fin tissue. As predicted, TGFBR1 inhibitors and TGFBR1 shRNA both reduced melanoma cell migration. In conclusion we envisage that TGFBR1 inhibitors are more likely to be effective in targeting cancer stem cell like properties and metastatic outgrowth, rather than in reducing the size of established tumours and the experience, so far, of investigators working on the clinical development of the TGFBR1 inhibitors supports this idea [6]. To maximise the clinical potential of TGFBR1 inhibitors, therefore, we propose that investigations should focus on their use in the prevention of metastatic recurrence after reducing the established tumour burden by conventional surgical and chemotherapeutic approaches.

REFERENCES

1. Blobe GC, et al. N Engl J Med. 2000; 342:1350–58. doi: 10.1056/NEJM200005043421807.
2. Inman GJ. Curr Opin Genet Dev. 2011; 21:93–99. doi: 10.1016/j.gde.2010.12.004
3. Akhurst RJ and Hata A. Nat Rev Drug Discov. 2012; 11:790–811. doi: 10.1038/nrd3810
4. Spender LC, et al. Oncotarget. 2016. doi: 10.18632/oncotarget.13226
5. Davies H, et al. Nature. 2002; 417:949–54. doi: 10.1038/nature00766
6. Herbertz S, et al. Drug Des Devel Ther. 2015; 9:4479–99. doi: 10.2147/DDDT.S86621

Gareth J Inman: Division of Cancer Research, School of Medicine, University of Dundee, Dundee, UK

Correspondence: Gareth J Inman

Email: g.j.inman@dundee.ac.uk

Keywords: TGF β , BRAF, melanoma, therapy

Received: January 24, 2017

Published: January 31, 2017