



University of Dundee

Knowledge of vitamin D and practices of vitamin D supplementation in a Scottish adult population

Zaremba, Suzanne M.M.; Conduit-Turner, Karen

DOI:

[10.1177/02601060241238824](https://doi.org/10.1177/02601060241238824)

Publication date:

2024

Licence:

CC BY

Document Version

Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Zaremba, S. M. M., & Conduit-Turner, K. (2024). Knowledge of vitamin D and practices of vitamin D supplementation in a Scottish adult population: A cross-sectional study. *Nutrition and Health*. Advance online publication. <https://doi.org/10.1177/02601060241238824>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Knowledge of vitamin D and practices of vitamin D supplementation in a Scottish adult population: A cross-sectional study

Nutrition and Health

1–14

© The Author(s) 2024



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/02601060241238824

journals.sagepub.com/home/nah



Suzanne M.M. Zaremba¹  and Karen Conduit-Turner²

Abstract

Background: Vitamin D supplementation practices (dose and frequency) are relatively unknown in the Scottish population, with no recent up-to-date data available. Reassessing current knowledge, practices, and awareness of vitamin D supplementation following a national health campaign in 2020 by Food Standards Scotland on vitamin D is warranted. **Aim:** This article aims to present the knowledge and awareness of vitamin D, and current vitamin D supplementation practices in adults living in Scotland. **Methods:** A cross-sectional study was performed between June and July 2022 using an online survey adapted from previous work on assessing knowledge of vitamin D in adults. Participants aged 18+, living in Scotland for ≥ 6 months were eligible to participate. Scores for knowledge were calculated as a percentage. Univariate associations between demographic and supplement use were established by χ^2 -test and logistic regression performed to predict factors associated with daily vitamin D intake. **Results:** Four hundred and three participants (72.7% female), mean age 36.4 (± 14.2 years), completed the study. Awareness of vitamin D was very high (99.5%) but the mean overall knowledge score was poor ($31.4 \pm 15.3\%$), with those with a university degree more likely to have knowledge scores at/above the mean compared with those with lower levels of education, $\chi^2(1, N = 393) 10.7, p = 0.001$, odds ratio (OR) = 2.1 (95% confidence interval (CI) 1.7–2.7). Finally, 64.3% took vitamin D supplements, of which 37.5% took them daily during winter months, with only 7.4% taking the recommended daily dose. **Conclusion:** The current study highlights the need to improve both knowledge of vitamin D and practices of vitamin D supplementation during the autumn and winter months in Scotland.

Keywords

Vitamin D, supplementation, knowledge, awareness, Scotland

Introduction

Vitamin D (cholecalciferol) is a fat-soluble vitamin that is synthesised primarily in the skin via activation of 7-dehydrocholesterol by ultraviolet (UVB type) sunlight. Vitamin D can be acquired through dietary sources, including oily fish, eggs and dairy, however these are limited, and meeting requirements can be challenging through dietary means alone (British Dietetic Association, 2019). Therefore, supplementation is the focus for most policies aiming to improve vitamin D levels in the UK population. Vitamin D deficiency, defined as plasma 25-hydroxyvitamin-D < 25 nmol/L (Scientific Advisory Committee on Nutrition, 2016), is problematic in Scotland with nearly half of those in the most deprived areas (47%) with insufficient vitamin D to protect their musculoskeletal health, compared with a quarter of those in the least deprived areas (Purdon et al., 2013). Prevalence of vitamin D deficiency in Scotland has been reported to be higher than other UK nations (Hyppönen and Power, 2007). Due to the high northern latitude of Scotland (55° to

59° North) and cloudy weather, conditions do not often favour vitamin D synthesis via UVB mechanisms (Rhodes et al., 2010). In addition, spending time indoors, obesity, high alcohol intake, and poor diet adversely affect vitamin D status.

Not only is Scottish data lacking with regard to up-to-date data on the prevalence of vitamin D deficiency, there is limited understanding of vitamin D supplementation practices in adults living in Scotland. The Scottish Government recommendation, in line with SACN guidance,

¹ Division of Population Health & Genomics, Ninewells Hospital & Medical School, University of Dundee, Dundee, UK

² Public Health, NHS Tayside, Dundee, UK

Corresponding author:

Suzanne M.M. Zaremba, Division of Population Health & Genomics, Ninewells Hospital & Medical School, University of Dundee, Dundee DD1 9SY, UK.

Email: s.zaremba@dundee.ac.uk

is for adults to consider taking 10 micrograms (μg) of vitamin D daily between October to March (Scientific Advisory Committee on Nutrition, 2016; Scottish Government, 2022a). This advice relates to public health advice for vitamin D supplementation in the general adult population. This advice does not apply to specific medical cases or elderly patients where high loading doses to correct for deficiency status may be relevant. This research is focussed on the general adult population only, in whom $10\mu\text{g}$ is an appropriate daily dose with very low risk of toxicity.

UK survey data published in 2021 by the British Nutrition Foundation showed that 26% took vitamin D supplements all year round, with 8% of British adults taking vitamin D supplements most of the year and 8% taking them during autumn and winter (British Nutrition Foundation, 2021). Over a third of people (39%) reported never taking vitamin D supplements. However, the reported figures did not detail whether these were the recommended dose ($10\mu\text{g}$) or taken daily. A cross-sectional survey by O'Connor et al. (2018) showed similar levels of supplementation within the UK population, 43.5% of individuals took vitamin D regularly but again data did not report if supplementation practices were daily during autumn-winter or the percentage of those taking the recommended dose.

According to Scottish Health Survey (2018) data 14% of Scottish men and 23% of Scottish women take vitamin D supplements, however dose and frequency were not reported. In 2020 Food Standards Scotland (FSS) conducted a survey to assess public adherence to the vitamin D recommendation. They reported that 58% of participants ($n=1465$) were not taking vitamin D supplements, to which the main reason was not being aware of the guidance or what the recommended dose is (Food Standards Scotland, 2020). Subsequently, in 2021–2022 Food Standards Scotland ran a campaign which publicised the Scottish Government recommendation for vitamin D supplementation alongside health information regarding vitamin D. The campaign was run on TV, social media/online, and pop-up field marketing across Scotland (Food Standards Scotland, 2022a). With the UK Government's recent call for evidence and research on vitamin D (UK Government, 2022) it was timely to investigate whether knowledge of vitamin D has improved following the national FSS campaign, assess what barriers to supplementation still exist and identify what the current supplementation practices are in the Scottish adult population. To the researchers' knowledge, this is the first study to quantitatively investigate vitamin D knowledge and supplementation practices, including amount and frequency, in adults living in Scotland.

Materials and methods

Study design, population and timeline

An observational, cross-sectional study was carried out from 15 June to 6 July 2022. Eligibility criteria included male and female adults (≥ 18 years). In order to assess

knowledge of vitamin D and practices of vitamin D supplementation following the winter Food Standards Scotland campaign (run nationally across Scotland from October 2021 to February 2022; Food Standards Scotland, 2021) eligible adults were required to have been resident in Scotland for a minimum of 6 months or longer for potential exposure to the campaign.

Questionnaire

Knowledge and awareness of vitamin D and vitamin D supplementation practice in Scotland were assessed using an online questionnaire, administered via Jisc (Jisc, 2022). Potential participants in this study were invited to participate via social media platforms (Twitter, Facebook, LinkedIn and Instagram), which included a direct link to the questionnaire, study information and consent. Participants were also recruited via University of Dundee newsletters, University of Dundee student societies and Blackboard announcements for staff and students.

The questionnaire was developed using previous studies investigating vitamin D knowledge (Boland et al., 2015; O'Connor et al., 2018), and awareness of vitamin D (Food Standards Scotland, 2021). Questions on vitamin D practices were adapted from a previous national survey on vitamin D (Food Standards Scotland, 2021). Questions from the Scottish Census were used to gather demographic data (Scottish Government, 2022b). Postcodes were used to determine Scottish Index of Multiple Deprivation (SIMD), using the SIMD 2020 postcode lookup tool (Scottish Government, 2020). The questionnaire consisted of 33 questions, containing five sections; demographics, vitamin D awareness, knowledge of vitamin D, vitamin D practices and vitamin D supplements (Appendix 1). To ensure face validity, the researchers reviewed each question in the survey, and knowledgeable individuals ($n=6$) in the field of vitamin D conducted an expert review to assess the appropriateness of the survey questions and response options. Additionally, piloting of the questionnaire was undertaken in a sample of University of Dundee staff and students ($n=12$) for content validity.

Knowledge scoring was performed as per Boland et al. (2015) with each correct answer worth one point, producing a maximum score of seven points. Questions that required only one response were marked as correct, and gained one point, or incorrect for zero points. For questions that included multiple correct responses, each correct response was worth an equal fraction of the overall question score. Questions related to effects and sources of vitamin D both had eight correct answers meaning each one selected was worth $1/8$, for factors that decrease vitamin D each was worth $1/13$, each correct month of the year was worth $1/6$. Equally weighted fractions were deducted for incorrect answers. This penalty for guessing was implemented to prevent participants from scoring 100% on multiple response questions by selecting all possible responses to that question. For this reason, the response "I don't know" was not penalised.

The final knowledge score was recorded as a percentage. Scores were then stratified into less than average and better than average to perform univariate association analysis between categorical variables and knowledge scores. The mean knowledge score was $31.4 \pm 15.3\%$.

For both the recommended daily dose of vitamin D knowledge question and the entry of actual daily dose taken by participants the data entered was inconsistent. Despite the survey question requesting that the participant enters the dose in μg , some participants entered in international units (IU) and identified it as such, however others did not specify a unit. In order to interpret whether the dose was correct or incorrect an assumption was made that doses >100 referred to IU and anything below that referred to μg , unless otherwise stated. The rationale being that most daily vitamin D supplements are $\leq 100 \mu\text{g}$ and this is the upper tolerable limit (European Food Safety Authority, 2016; Rizzoli, 2021).

Statistical analysis

The Statistical Package for Social Science (IBM SPSS software; version 26, IBM, NY, USA) was used to analyse all data.

Descriptive statistics were used to present demographic data and to evaluate knowledge, awareness and practices of vitamin D supplementation. For data that was normally distributed, mean and standard deviations are reported, the median and interquartile range are reported for data that did not have a normal distribution. Univariate associations between categorical variables (age, gender, employment, SIMD, income, ethnicity, and nationality) and supplement use were established by χ^2 -test. To establish factors contributing to daily vitamin D supplement use logistic regression analysis was performed. Factors with a p -value < 0.1 were included in the regression model. Results are expressed from the logistic regression using odds ratios, 95% confidence intervals and a p value < 0.05 was considered statistically significant.

The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines for the presentation and writing of this manuscript were followed (Von Elm et al., 2007).

Results

Demographics

In total 403 participants completed the survey. The mean age was 36.4 (± 14.2 years), with one datapoint excluded due to an invalid age entry. The participants were predominantly female (72.7%), of which 2.0% were pregnant or breastfeeding. Most participants were in full-time employment (43.4%), with 45.6% earning ≥ 700 per week/ $\geq 36,400$ per annum or above. Over a quarter of respondents were from the highest SIMD, quintile 5. Education level was high with 60.5% participants educated to degree level or

above and 15.9% had obtained school completion qualification equivalent or above such as Scottish Highers/GCSEs. The median time living in Scotland was 22.3 years with a range from 0.5 to 77.5 years. In terms of ethnicity 83.4% identified as white and for nationality although Scottish was in the majority when comparing individual nationalities, it made up less than 50% overall. Participants' demographics are summarised in Table 1.

Awareness of vitamin D and food standards Scotland campaign

All participants (99.5%), apart from three (0.5%), had heard of vitamin D. The commonest source of vitamin D information was school/college/university (56.4%), followed by family (45.1%), or from a doctor or nurse (37.9%). Other sources included friends (29.4%), social media (28.9%), television (28.2%) and less frequent sources were books and newspapers. When asked if they were aware of the current recommendation to take a daily supplement in autumn and winter, 65.5% answered 'yes', with 31.5% ($n = 127$) then answering what they believed the recommended dose was. Only 8.2% of participants recalled seeing the FSS vitamin D campaign adverts, and 10% could not recall it. The commonest sources for those who had seen the campaign were social media 37.8% ($n = 17$), television 20.0% ($n = 9$), 15.6% poster/billboard ($n = 7$), 8.9% newspaper ($n = 4$) and 4.4% radio ($n = 2$). 'Other' responses (13.3%) included at work, in a store, health centre and the internet/website advertisement as the source.

Vitamin D knowledge

The mean vitamin D knowledge score (\pm SD) was 31.4 ($\pm 15.3\%$) with 43.4% ($n = 174$) scoring above average. There was a statistically significant association between knowledge scores (stratified into $<$ and $>$ mean of 31.4%) and education level. Those who were educated to a degree level or above were more likely to have scored higher than average (OR = 2.1, 1.7–2.7), and those who were educated below university degree level were more likely to score less than average, χ^2 (1, $N = 393$) = 10.7, $p = 0.002$. There were no statistically significant associations between knowledge and any other demographic factors, including age, gender, income, SIMD, employment status, ethnicity, nationality, pregnancy status and length of residency in Scotland ($p > 0.05$). Participants' knowledge of vitamin D is summarised in Table 2.

Vitamin D supplementation practices

Vitamin D supplements were taken by 64.3% of participants, of which 37.5% took a vitamin D supplement every day and 25.8% less frequently (Table 3).

Of those who took vitamin D supplements, the recommended dose of 10 $\mu\text{g}/400\text{IU}$ was taken daily by 22.8%

Table 1. Demographic patterns of participants (n = 403).

Demographic	Group	n	Percent
Age	18–24	118	29.3
	25–34	86	21.3
	35–49	112	27.8
	50–64	71	17.6
	65+	15	3.7
Gender	Missing	1	0.2
	Male	100	24.8
	Female	293	72.7
	Prefer not to answer	2	0.5
	Other	6	1.5
Pregnant/ breastfeeding	Missing	2	0.5
	Yes	8	2
	No	392	97.3
	Prefer not to answer	1	0.2
Employment	Missing	2	0.5
	Full-time employment	175	43.4
	Part-time employment	63	15.6
	Not currently employed	18	4.5
	Student	123	30.5
	Retired	18	4.5
	Prefer not to answer	2	0.5
Income	Missing	4	1
	Less than £100 pw OR less than £5200 pa	18	4.5
	£100–199 pw or £5200 to £10,399 pa	27	6.7
	£200–299 pw or £10,400 to £15,499 pa	28	6.9
	£300–399 pw or £15,500 to £20,799 pa	17	4.2
	£400–499 pw or £20,800 to £25,999 pa	19	4.7
	£500–699 pw or £26,000 to £36,399 pa	43	10.7
	£700–999 pw or £36,400 to £51,999 pa	61	15.1
	£1000–1499 pw or £52,000 to £77,999 pa	66	16.4
	£1500 pw or more to £78,000 or more pa	57	14.1
	Prefer not to answer	63	15.6
	Missing	4	1
	SIMD quintile	1 (most deprived)	47
2		73	18.1
3		79	19.6
4		96	23.8
5 (least deprived)		104	25.8
Nationality	Scottish	162	40.2
	Non-Scottish	233	57.8
	Missing	8	2
Ethnicity	White	336	83.4
	Mixed or multiple ethnic groups	5	1.2
	Asian, Scottish Asian or British Asian	39	9.7
	Missing	13	3.2

(continued)

Table 1. (continued)

Demographic	Group	n	Percent
	African, Scottish African or British African		
	Caribbean or Black	0	0
	Other ethnic group Arab, Scottish Arab or British Arab, Other	7	1.7
	Missing	3	0.7

pa: per annum; pw: per week; SIMD: Scottish Index of Multiple Deprivation.

(n = 30) participants (Table 4). The majority of vitamin D supplement takers took more than 10 µg/400IU per day (62.9%). A few participants took less than 10 µg/400IU per day (5.3%), with 9.1% unsure of the dose taken. Within the whole sample, 7.4% of participants took a 10 µg/400IU vitamin D supplement daily during the autumn/winter months.

Participants who scored above average for vitamin D knowledge were statistically more likely to take vitamin D supplements daily compared to participants with below average knowledge scores who were less likely to supplement daily (OR = 1.9, 1.5–2.3), $X^2(1, N=401)=6.7$, $p=0.009$ and at the correct dose $X^2(1, N=120)=4.1$, $p=0.043$.

There was a statistically significant association between age and frequency of vitamin D supplementation (stratified into daily supplementation and less frequently). Those who were aged 18–24 and aged ≥65 years were less likely to take a vitamin D supplement daily, and those who were aged 25–64 years were more likely to take a vitamin D supplement every day, $X^2(4, N=402)=44.6$, $p<0.001$. There was no association between age and taking the recommended dose of vitamin D supplements ($p>0.05$).

When stratified into ‘white Scottish’, ‘non-white not Scottish’ and ‘white not Scottish’, participants who were white Scottish were more likely to take the recommended dose of vitamin D, $X^2(2, N=118)=28.5$, $p<0.001$. However, the frequency of dosages was not statistically significant ($p>0.05$).

There were no associations between other demographic factors, gender, education, SIMD, employment or income, with frequency of vitamin D supplementation or supplementing the recommended dose of vitamin D ($p>0.05$).

For those who did not take a vitamin D supplement daily during autumn/winter (n = 252), participants were advised of the current guidance in the survey, of which 76.6% considered themselves as either ‘likely’ or ‘very likely’ to start adhering to the recommendation.

Predicting vitamin D supplement use

Supplement use logistical regression results are summarised in Table 5. There were no factors identified to predict daily vitamin D supplement use ($p>0.05$).

Table 2. Participants' knowledge of vitamin D; questions used in vitamin D knowledge score calculation (n = 401).

Question topic		✓/×	N (% participants who selected answer)	
Benefits of vitamin D	Bone health	✓	325 (80.6)	
	Immune health	✓	283 (70.2)	
	Prevention of Rickets	✓	234 (58.1)	
	Vision health	×	81 (20.1)	
	Hair growth	×	99 (24.6)	
	Prevention of diabetes	✓	53 (13.2)	
	Cardiovascular health	✓	97 (24.1)	
	Cognitive health	✓	118 (29.3)	
	Cancer prevention	✓	75 (18.6)	
	Skin softness	×	41 (10.2)	
	Calcium absorption	✓	242 (60.0)	
	None of the above	×	1 (0.2)	
	I don't know	-	30 (7.4)	
	Other	-	12 (3.0)	
	Sources	Fruits	×	97 (24.1)
		Vegetables	×	124 (30.8)
		Fatty fish	✓	176 (43.7)
Vitamin D supplements		✓	330 (81.9)	
Select cereals		✓	133 (33.0)	
Milk/dairy		✓	138 (34.2)	
Nuts		×	58 (14.4)	
Cod liver		✓	133 (33.0)	
Sun		✓	338 (83.9)	
Eggs		✓	121 (30.0)	
Mushrooms		✓	66 (16.4)	
I don't know		-	9 (2.2)	
Factors that decrease absorption		Skin pigment	✓	212 (52.6)
		Pregnancy/lactation	✓	113 (28.0)
		Dairy allergy	✓	92 (22.8)
		Shade/ clouds	✓	235 (58.3)
		Fatty diets	×	47 (11.7)
	Pollution	✓	126 (31.3)	
	Time of day	✓	206 (51.1)	
	Sunscreen use	✓	155 (38.5)	
	Wind	×	9 (2.2)	
	Latitude	✓	141 (35.0)	
	Vegan diet	✓	124 (30.8)	
	Smoking	×	74 (18.4)	
	Season	✓	210 (52.1)	
	Vegetarian diets	✓	82 (20.3)	
	Age	✓	122 (30.3)	
	Lactose intolerance	✓	53 (13.2)	
	Months of the year when sunshine is sufficient for vitamin D production	I don't know	-	30 (7.4)
January		×	16 (4.0)	
February		×	15 (3.7)	
March		×	43 (10.7)	
April		✓	150 (37.2)	
May		✓	279 (69.2)	
June		✓	365 (90.6)	
July	✓	377 (93.5)		

(continued)

Table 2. (continued)

Question topic		✓/×	N (% participants who selected answer)		
Average daily sun exposure required for fair-skinned to get enough vitamin D	August	✓	352	(87.3)	
	September	✓	184	(45.7)	
	October	×	35	(8.7)	
	November	×	13	(3.2)	
	December	×	12	(3.0)	
	Less than 9minutes per day	×	21	(5.2)	
	9–14minutes per day	✓	66	(16.4)	
	15–20minutes per day	×	123	(30.5)	
	More than 20minutes per day	×	86	(21.3)	
	I don't know	-	107	(26.6)	
Average daily sun exposure required for non-fair-skinned person would need to spend in the sun to get enough vitamin D	Less than 15minutes per day	×	52	(12.9)	
	15–24minutes per day	✓	67	(16.6)	
	About 25–38minutes per day	×	76	(18.9)	
	More than 38minutes per day	×	74	(18.4)	
	I don't know	-	133	(33.0)	
	Missing	×	1	(0.2)	
	Recommended daily dose	Value entered was below recommended	×	10	2.3
		Correct dose entered	✓	61	15.1
		Value entered was above recommended	×	33	13.3

Ticks ✓ denote the correct answers and crosses × indicate an incorrect answer.

Vitamin D supplement purchasing

In terms of where vitamin D supplements were purchased, the majority were from supermarkets (37.1%) and chemists/pharmacies (24.9%). Most participants took vitamin D supplements in tablet form (Table 6). There was a wide variation in vitamin D supplement brands purchased, with high street stores being most frequent. The median (25th and 75th) spend on vitamin D supplements was £3.00 (1.00, 5.00) per month (n = 128).

Barriers to vitamin D supplementation

The most frequent reasons as to why daily vitamin D supplementation was not followed during autumn/winter months were forgetfulness (30.1%) and lack of awareness

of the recommendations (22.1%). Twelve percent of individuals preferred to get vitamins from food and 8% did not see the need for taking vitamin D. Of those who answered 'other', responses ranged from medical conditions that contraindicated vitamin D supplementation advice, to beliefs that sufficient vitamin D could be obtained from food or sunlight (Table 7).

Table 3. Average frequency participants took a vitamin D supplement during the winter (n = 403).

Frequency	N	Percent
Every day	151	37.5
Several times a week	57	14.1
Once a week	20	5.0
Every couple of weeks	22	5.5
Monthly	9	2.2
Never	144	35.7

Table 4. Doses of vitamin D taken daily (n = 132). Ticks ✓ denote the correct answers, crosses × indicate an incorrect answer.

Units	Dose	✓/×	N	Percent of respondents
Micrograms	Below recommended 10µg	×	6	4.5
	10 µg	✓	29	22.0
	Above recommended 10 µg	×	69	52.3
IUs	Below recommended 400 IUs	×	1	0.8
	400 IUs	✓	1	0.8
	Above recommended 400 IUs	×	14	10.6
	Don't know	-	12	9.1

Table 5. Factors contributing to vitamin D supplementation use.

	B	SE	Wald	df	p	OR	95% CI for OR	
							Lower	Upper
Knowledge score	-0.36	0.11	10.62	1	0.09	0.70	0.57	0.87
Gender	-0.75	0.99	0.56	1	0.45	0.47	0.07	3.33
Males								
Gender	-0.38	-0.29	1.77	1	0.19	0.69	0.39	1.20
Females								
Educated to degree level or above	0.09	0.25	0.13	1	0.72	1.10	0.67	1.78
Age	-0.05	0.01	34.90	1	0.06	0.95	0.93	0.97
SIMD 1	-0.76	0.46	2.80	1	0.09	0.47	0.19	1.14
SIMD 2	-0.55	0.46	1.41	1	0.24	0.58	0.23	1.43
SIMD 3	-0.42	0.45	0.87	1	0.35	0.66	0.28	1.56
SIMD 4	-0.22	0.44	0.26	1	0.61	0.80	0.39	1.90
SIMD 5								
Constant	4.15	0.65	40.71	1	0.073	63.43		

Logistic regression predicting likelihood of supplement use on relevant factors (n = 393, n = 11 missing data).

B: B coefficient; CI: confidence interval; df: degrees of freedom; OR: odds ratio; p: p value; SE: standard error of mean; SIMD: Scottish Index of Multiple Deprivation; Wald: Wald chi-square test.

Discussion

This is the first study to quantitatively investigate the knowledge of vitamin D, practices of vitamin D supplementation and identify barriers to vitamin D supplement use in adults living in Scotland. The results suggest that overall knowledge scores of vitamin D are low (31.4%) in the Scottish population, however identification of vitamin D's role in bone health (80.6%) and immunity (70.2%) along with sources of vitamin D, including sun (83.9%) and vitamin D supplements (81.9%), were good. There was poorer knowledge of dietary sources of vitamin D, which is concerning considering that only 37.5% of the individuals in the current study reported to take vitamin D supplements daily during autumn and winter months.

Vitamin D knowledge

In the current study awareness of vitamin D was good at 99.5%, which corroborates the 99% reported in the UK-based study by O'Connor et al. (2018). However, knowledge of vitamin D was poor with an average score of only 31.4%, another similar trend reported in the literature (Kung and Lee, 2006; Boland et al., 2015). This result was not dissimilar to the findings by Boland et al. (2015) where the mean score was 29% in an adult Canadian population.

For identifying sources of vitamin D the participants in the current study outperformed the respondents in research conducted by Kung and Lee (2006) and Toher et al. (2014), particularly in recognition of sunlight (83.9% vs 52.6–74.0%) and supplements (81.9% vs 10.0–38.4%). Source knowledge results were more aligned with the reported findings of O'Connor et al. (2018) and Vu et al. (2010), although overall knowledge remains lower than that seen in the published findings by O'Connor et al.

Table 6. Vitamin D supplement purchase location (n = 205) and form (n = 163).

Question	Response	N	Percent
Place of purchase	Supermarket	76	37.1
	Health food store	37	18.0
	Chemist/Pharmacy	51	24.9
	Online	32	15.6
	Other	9	4.4
Form of supplement	Tablets	119	73.0
	Caplets	18	11.0
	Spray	7	4.3
	Drops	3	1.8
	Jellies	11	6.7
	Chewable tablets	5	3.1

Table 7. Barriers to following the recommended daily vitamin D supplement advice (n = 362).

Reasons	N	Percentage of responses
I'm unaware of the recommended advice	80	22.1
I'm confused by the recommended advice	20	5.5
I don't see the need	32	8.8
I forget	109	30.1
I prefer to get vitamins from food	44	12.2
I can't afford supplements	13	3.6
I don't know where to buy them	6	1.7
I don't like taking supplements	26	7.2
I don't agree with taking tablets	5	1.4
I find it hard to swallow tablets	8	2.2
Other	19	5.2

(2018) where the mean score was 56.6%. However, it should be mentioned that the higher knowledge score reported by O'Connor et al. (2018) is likely to have been influenced by the high level of nutrition students who participated in their research. Lack of knowledge of vitamin D, including dietary sources and government recommendations, was apparent in published findings by Kotta et al. (2015), who conducted qualitative interviews with adults living in England. However, knowledge was not quantitatively scored in the aforementioned study. In the current study, there was a high proportion of females (72.7%) who have been reported elsewhere as having better knowledge of vitamin D (Kung and Lee, 2006; Boland et al., 2015). Miller and Russell (2004) report that American females of childbearing age have good knowledge of dietary supplements, however there was no difference between gender and knowledge of vitamin D in the current study.

Vitamin D supplements

In terms of understanding the vitamin D guidance, only 130 participants attempted to identify the recommended daily dose in the current study, of which 61 identified the

recommended dose, which represents only 15.1% of the total participants. Although this is low, it is an improvement on the 5% of participants who correctly identified the reference nutrient intake (RNI) of vitamin D in the FSS survey (Food Standards Scotland, 2020).

Only 7.4% of participants in the current study took the recommended dose of vitamin D daily during autumn-winter months (10 µg/400IUs). This is a slight improvement on the 5% seen by FSS (2020), which may indicate the FSS campaign had a minor impact. Almost 20% reported consuming doses of vitamin D above the RNI which could raise some concerns about potential toxicity for those with medical conditions, however health status was not monitored in the current work and therefore doses of up to 100 µg are deemed tolerable as per general population advice. The evidence on vitamin D toxicity tends to be based on low-quality case studies, which can be useful to identify risk factors for over-supplementation (Agbabiaka et al., 2018) and potential side effects (De Vincentis et al., 2021). However, they do not offer robust evidence of the impact of over-supplementation, which might lead to nothing more than abnormal blood results (Kilbane et al., 2014). Reported symptoms may potentially be due to co-administration of calcium leading to hypercalcaemia, rather than from vitamin D alone (European Food Safety Authority, 2016).

A potential source of confusion leading to under-supplementation, seen in around 5% of those who took <10 µg daily vitamin D in the current study, may be inconsistencies in RNIs on supplement labels. Some supplement labels state the RNI for vitamin D as 5 µg per day as food labelling is standardised at EU level. Products sold in the UK are required to be labelled in accordance with the EU based RNI for labelling, which is 5 µg (Food Standards Scotland, 2022b). Recommending a daily supplement year-round in Scotland may improve adherence by simplifying the regime (Vlasnik et al., 2005) as 30.1% of the adults included in the current work report that forgetfulness was a common barrier to following the vitamin D supplement recommendation. It seems reasonable to suggest remembering to restart taking vitamin D supplements in autumn-winter may also be an issue. Daily use would also avoid issues with factors that decrease absorption from sunlight and may give better population confidence by removing ambiguity. In addition, supplementation may also be a more advantageous method to optimise vitamin D levels given that only 29% and 33% of Scottish men and women, respectively, are reported to consume oily fish once a week or more, a key source of dietary vitamin D (Scottish Health Survey, 2020).

The median spend on vitamin D supplements was £3.00 per month, which can be considered a reasonable cost given that 45.6% of the participants in the current study had a household income of £700 per week/£36,400 per annum or above. It is important that vitamin D supplements remain affordable and accessible to the public, especially at present with food and energy price inflation.

Signposting individuals to locations where safe and low-priced vitamin D supplements can be purchased would be advantageous, particularly those on low incomes who are not eligible for the free Scottish Vitamins Scheme or Healthy Start initiative (pregnant and breastfeeding females and children under 3 only). This could be a consideration for future vitamin D campaigns, educating the public not only on the health benefits of vitamin D but also where low-cost vitamin D supplements can be found. Vitamin D supplements from supermarkets, particularly own brand products, could be recommended. Considering that brand familiarity can impact purchase intention (Castro et al., 2018), consumers may be more likely to buy branded vitamin supplements than supermarket own-brand products due to consumer belief that branded supplements are of higher quality and more effective than supermarket own-brand products.

Public health campaigns

Only 8.2% of participants in this study recalled the FSS vitamin D campaign (2021–2022). It can therefore be suggested that awareness of the recommendations has more likely come from other sources, namely educational environments (60.5% educated to degree level or above), family and health professionals, as reported in our findings. The current study had a demographic with >10.9% participants resident in Scotland for less than a year, which means they may not have been resident in Scotland for the entire duration of the FSS campaign.

In terms of where the campaign was viewed, social media was the most common avenue with just over half of those who recalled the campaign naming it as a source, with TV and radio significantly lower at 27.0% and 21.0%, respectively. This may indicate declining impact from traditional forms of media to gain health information (Stellefson et al., 2020). Although, this could have been confounded by the relatively young cohort mean age 36.4 years, with younger adults more likely to use social media (Khoros, 2022). However, knowledge alone does not guarantee behaviour change (Baker and Murdoch 2008) and some interventions can unintentionally have negative impacts on behaviours (Nyhan et al., 2014; Stead et al., 2019). There is a need for research on the impact of 'new media' (e.g. liking, sharing or commenting on content and downloading campaign apps) and the cost effectiveness of messages related to health, in order to improve public health interventions (Stead et al., 2019).

From a theoretical point of view, the Health Belief Model (Becker, 1974) and Expanded Health Belief Model (Strecher and Rosenstock, 1997) assume that people will perform a health-promoting behaviour (such as vitamin intake) if they perceive a serious health threat, believe that the health behaviour will reduce the health threat, perceive minimal barriers and believe they can effectively execute the health behaviour. Accordingly, if Scottish adults are aware of the importance of taking vitamin D to

support health but do not know where to get vitamin D other than from the sun or dietary supplements (a barrier that reduces feelings of self-efficacy) they are less likely to seek out these sources and engage with the desired health behaviour. Perhaps the noted lack of vitamin D knowledge from dietary sources, number of individuals who do not take daily vitamin D supplements (over a third of the current cohort never take vitamin D supplements) and high northern latitude of Scotland combined may contribute to the prevalence of vitamin D deficiency in Scotland. However, we acknowledge the need for updated deficiency data, with the most recent vitamin D deficiency data based on Scottish Health Survey data from 2010 to 2011. Ten micrograms of vitamin D is considered a safe daily dose for the general population (SACN, 2016) and given that there is evidence to suggest that Scottish individuals may benefit from daily vitamin D supplementation all year round (Zgaga et al., 2011) we suggest that public health guidance should consider updating advice for Scottish adults to supplement with 10 µg daily, not only during autumn-winter months and not only for high-risk groups.

Limitations and future research

As with all cross-sectional studies it is not possible to comment on the direction of associations. As the majority of the data was quantitative it is difficult to interpret why people are not currently taking vitamin D, aside from the limited number of options offered in the survey. Hard copies of the questionnaire should have been utilised to reach older adults who may be less likely to use the internet, therefore future research should allow the option for electronic and hard copy questionnaires to be used.

The use of convenience sampling is likely to have introduced a degree of bias; this approach meant that individuals associated with a Higher Education institution were predominantly recruited (staff and students) and this is likely to have led to a skewed dataset towards a higher level of education and younger age group, impacting on the generalisability of the results. In terms of age it should be noted that 29% of our study population were <25 and just 4% were >65 compared to 12% and 21%, respectively, reported by FSS. At risk population groups were underrepresented in the current work, and hence studies focussing on pregnant/breastfeeding females, children, older adults >65 years and individuals with darker skin are warranted. This is important as recent evidence from Tanna et al. (2023) report that increasing age is negatively associated with vitamin D supplement intake in British adults. Although the current study did have a larger sample included than other similar work (O'Connor et al. 2018; Toher et al., 2014), a larger sample would enable better understanding of the diverse Scottish population.

Data were stratified using mean and medians, therefore it is important to acknowledge the drawbacks of taking such approach, including sensitivity to outliers, loss of precision

and potential impact on statistical testing, that is, when assumptions of normal distribution are required (Field, 2017). Scoring of the questionnaire followed the protocol adopted by Boland et al. (2015) which allowed for direct comparison of scores with the current data, however this scoring technique was perhaps cumbersome with some correct answers weighted for as little as 1/7 or 1/13 of a mark.

Logistic regression was performed to determine whether age, gender, SIMD or knowledge were predictors of daily vitamin D supplementation, which were not shown to be significant in the current study. This may have been contributed to by the demographics of the group being skewed towards a younger, more educated group and the limitations of SIMD as an indicator of relative affluence, particularly in a student population. This may be an avenue for further research.

A further limitation of this study was that some of the deviation in reported dosing regimens for vitamin D may have been due to individualised medical advice. Around 38% of respondents reported hearing about vitamin D from a doctor or nurse, and although they may have received generic advice around the public health guidance (10 µg daily) it is also possible that they were given specific regimens due to certain health conditions that would have required either high loading doses followed by maintenance, or reduced doses due to concerns of toxicity. It would be important in any future research to clearly identify cases with medical conditions that may require altered dosing regimes as per national guidance (NICE, 2022). Furthermore, during the COVID pandemic residents in Scotland, as well as other parts of the UK, were provided free vitamin D 10 µg supplements over the wintertime which may have affected respondent awareness. It is reported that 40% of Scottish shielders took up the offer of free vitamin D (Scottish Government, 2021). The population studied had very good knowledge of vitamin D's role in immunity, the prevention of rickets and bone health yet over 63% of the sample who took daily vitamin D supplements during autumn-winter took more than the recommended dose to support skeletal on bone health (>10 µg). This highlights a future avenue for exploration with regard to understanding knowledge and practices of at-risk individuals with impaired immune function and rickets.

Conclusion

In conclusion, the current study highlights the need to improve both knowledge of vitamin D and practice of vitamin D supplementation during the autumn and winter months in Scotland. It is clear that despite the effort of FSS to raise awareness of vitamin D, there is still a significant gap in the Scottish population's knowledge of vitamin D and improvements need to be made to supplementation practices. The true extent to which poor supplementation practices translate into vitamin D deficiency is currently

unknown, especially since deficiency data in Scottish adults has not been updated in over a decade.

Acknowledgements

We would like to thank all participants who took part in this survey. We are grateful of their time to make this research possible. We would also like to thank Alana McDonald and Dr Gillian Purdon (Food Standards Scotland) for their expert guidance during the preparation of the survey.

Availability of data and materials

Data are available from the corresponding author upon request.

Author contributions

SZ and KC-T contributed to conceptualisation, methodology, investigation, data curation formal analysis, visualisation, writing-original draft preparation, and writing-review and editing. KC-T was involved in project administration and SZ in supervision.

Consent for publication

All authors approved the submission of the manuscript and consented to the publication of this manuscript.

Declaration of conflicting interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Ethical approval

The study was conducted in accordance with the Declaration of Helsinki and approved by University of Dundee School of Medicine Research Ethics Committee with the reference 22/58 on the 9 June 2022. Written informed consent was obtained from all subjects involved in the study.

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Suzanne M.M. Zaremba  <https://orcid.org/0000-0003-1479-9653>

Supplemental material

Supplemental material for this article is available online.

References

- Agbabiaka TB, Spencer NH, Khanom S, et al. (2018) Prevalence of drug-herb and drug-supplement interactions in older adults: a cross-sectional survey. *The British Journal of General Practice* 68(675): e711–e717.
- Baker H and Murdoch IE (2008) Can a public health intervention improve awareness and health-seeking behaviour for glaucoma? *British Journal of Ophthalmology* 92(12): 1671–1675.
- Becker MH (1974) The health belief model and personal health behavior. *Health Education Monographs* 2: 324–508.

- Boland S, Irwin JD and Johnson AM (2015) A survey of university students' vitamin D-related knowledge. *Journal of Nutrition Education and Behavior* 47: 99–103.
- British Dietetic Association (2019) Vitamin D: Food Fact Sheet. Available at: <https://www.bda.uk.com/resource/vitamin-d.html> (accessed 31 May 2023).
- British Nutrition Foundation (2021) British Nutrition Foundation survey reveals 49% adults unaware of UK Government guidelines for vitamin D. Available at: <https://www.nutrition.org.uk/news/2021/british-nutrition-foundation-survey-reveals-49-adults-unaware-of-uk-government-guidelines-for-vitamin-d/> (accessed 31 May 2023).
- Castro IA, Majmundar A, Williams C, et al. (2018) Customer purchase intentions and choice in food retail environments: a scoping review. *International Journal of Environmental Research and Public Health* 15(11): 2493.
- De Vincentis S, Russo A, Milazzo M, et al. (2021) How much vitamin D is too much? A case report and review of the literature. *Endocrine, Metabolic & Immune Disorders Drug Targets* 21(9): 1653–1659.
- European Food Safety Authority (2016) Dietary reference values for vitamin D. *EFSA Journal* 14(10): e04547.
- Field A (2017) *Discovering Statistics using IBM SPSS Statistics*. London, England: Sage Publications.
- Food Standards Scotland (2020) Vitamin D Omnibus Summary of research results November 2020. Available at: <https://www.foodstandards.gov.scot/publications-and-research/publications/vitamin-d-omnibus-survey> (accessed 31 May 2023).
- Food Standards Scotland (2021) Vitamin D Partner Toolkit 2021. Available at: https://www.foodstandards.gov.scot/downloads/fss_vitamin_d_toolkit.pdf (accessed 31 May 2023).
- Food Standards Scotland (2022a) Vitamin D Campaign evaluation. Available at: https://www.foodstandards.gov.scot/downloads/FSS_Vitamin_D_Campaign_Research_202122.pdf (accessed 31 May 2023).
- Food Standards Scotland (2022b) Your questions on vitamin D answered by Food Standards Scotland's expert. Available at: <https://www.foodstandards.gov.scot/news-and-alerts/your-questions-on-vitamin-d-answered-by-food-standards-scotlands-expert> (accessed 31 May 2023).
- Hypönen E and Power C (2007) Hypovitaminosis D in British adults at age 45 y: nationwide cohort study of dietary and lifestyle predictors. *The American Journal of Clinical Nutrition* 85(3): 860–868.
- Jisc (2022) Online surveys. Available at: <https://www.onlinesurveys.ac.uk/> (accessed 31 May 2023).
- Khoros (2022) The 2022 Social Media Demographics Guide. Available at: <https://khoros.com/resources/social-media-demographics-guide> (accessed 31 May 2023).
- Kilbane MT, O'Keane M, Morrin M, et al. (2014) The double-edged sword of vitamin D in Ireland: the need for public health awareness about too much as well as too little. *Irish Journal of Medical Science* 183(3): 485–487.
- Kotta S, Gadhvi D, Jakeways N, et al. (2015) Test me and treat me"—attitudes to vitamin D deficiency and supplementation: a qualitative study. *BMJ Open* 5(7): e007401.
- Kung AW and Lee KK (2006) Knowledge of vitamin D and perceptions and attitudes toward sunlight among Chinese middle-aged and elderly women: a population survey in Hong Kong. *BMC Public Health* 6: 226.
- Miller CK and Russell T (2004) Knowledge of dietary supplement label information among female supplement users. *Patient Education and Counseling* 52(3): 291–296.
- National Institute for Health and Care Excellence (2022) Vitamin D deficiency in adults: management of vitamin D deficiency or insufficiency. Available at: <https://cks.nice.org.uk/topics/vitamin-d-deficiency-in-adults/management/management/#how-to-treat> (accessed 19 December 2023)
- Nyhan B, Reifler J, Richey S, et al. (2014) Effective messages in vaccine promotion: a randomized trial. *Pediatrics* 133(4): e835–e842.
- O'Connor C, Glatt D, White L, et al. (2018) Knowledge, attitudes and perceptions towards vitamin D in a UK adult population: a cross-sectional study. *International Journal of Environmental Research and Public Health* 15(11): 2387.
- Purdon G, Comrie F, Rutherford L, et al. (2013) Vitamin D status of Scottish adults: results from the 2010 & 2011 Scottish Health Surveys. Available at: https://www.foodstandards.gov.scot/downloads/Report_Final.pdf (accessed 31 May 2023).
- Rhodes LE, Webb AR, Fraser HI, et al. (2010) Recommended summer sunlight exposure levels can produce sufficient (>=20 ng ml(-1)) but not the proposed optimal (>=32 ng ml(-1)) 25(OH)D levels at UK latitudes. *Journal of Investigative Dermatology* 130: 1411–1418.
- Rizzoli R (2021) Vitamin D supplementation: upper limit for safety revisited? *Aging Clinical and Experimental Research* 33: 19–24.
- Scientific Advisory Committee on Nutrition (2016) SACN vitamin D and health report. Available at: <https://www.gov.uk/government/publications/sacn-vitamin-d-and-health-report> (accessed 31 May 2023).
- Scottish Government (2020) Scottish Index of Multiple Deprivation 2020v2 postcode lookup file. Available at: <https://www.gov.scot/publications/scottish-index-of-multiple-deprivation-2020v2-postcode-look-up/> (accessed 31 May 2023).
- Scottish Government (2021) Coronavirus (COVID-19): Vitamin D take up by the shielding group – data summary. Available at: <https://www.gov.scot/publications/coronavirus-covid-19-vitamin-d-take-up-by-the-shielding-group/> (accessed 19 December 2023).
- Scottish Government (2022a) Vitamin D: advice for all age groups. Available at: <https://www.gov.scot/publications/vitamin-d-advice-for-all-age-groups/> (accessed 31 May 2023).
- Scottish Government (2022b) Scotland's Census 2022 paper questionnaires. Available at: <https://www.scotlandscensus.gov.uk/documents/scotland-s-census-2022-paper-questionnaires/> (accessed 31 May 2023).
- Scottish Health Survey (2018) Scottish Health Survey 2017 - volume one: main report. Available at: <https://www.gov.scot/publications/scottish-health-survey-2017-volume-1-main-report/pages/82/> (accessed 31 May 2023).
- Scottish Health Survey (2020) Scottish Health Survey 2018: main report - revised 2020. Available at: <https://www.gov.scot/publications/scottish-health-survey-2018-volume-1-main-report/> (accessed 31 May 2023).
- Stead M, Angus K, Langley T, et al. (2019) Mass media to communicate public health messages in six health topic areas: a systematic review and other reviews of the evidence. *Public Health Research* 7(8). DOI: 10.3310/phr07080.
- Stellefson M, Paige SR, Chaney BH, et al. (2020) Evolving role of social media in health promotion: updated responsibilities for health education specialists. *International Journal of Environmental Research and Public Health* 17(4): 1153.
- Strecher V and Rosenstock I (1997) The health belief model. In: Glanz K, Lewis FM and Rimer BK (eds) *Health Behavior*

and Health Education: Theory, Research, and Practice. San Francisco: Jossey-Bass, 41–58.

- Tanna NK, Karki M, Webber I, et al. (2023) Knowledge, attitudes, and practices associated with vitamin D supplementation: a cross-sectional online community survey of adults in the UK. *PLoS ONE* 18(8): e0281172.
- Toher C, Lindsay K, McKenna M, et al. (2014) Relationship between vitamin D knowledge and 25-hydroxyvitamin D levels amongst pregnant women. *Journal of Human Nutrition and Dietetics* 27(3): 261–269.
- UK Government (2022) Vitamin D Call for evidence. Available at: <https://www.gov.uk/government/consultations/vitamin-d-call-for-evidence/vitamin-d-call-for-evidence> (accessed 31 May 2023).
- Vlasnik JJ, Aliotta SL and DeLor B (2005) Medication adherence: factors influencing compliance with prescribed medication plans. *The Case Manager* 16(2): 47–51.
- von Elm E, Altman DG, Egger M, et al. (2007) The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet* 370(9596): 1453–1457.
- Vu LH, van der Pols J, Whiteman DC, et al. (2010) Knowledge and attitudes about vitamin D and impact on sun protection practices among urban office workers in Brisbane, Australia. *Cancer Epidemiology, Biomarkers & Prevention* 19(7): 1784–1789.
- Zgaga L, Theodoratou E, Farrington SM, et al. (2011) Diet, environmental factors, and lifestyle underlie the high prevalence of vitamin D deficiency in healthy adults in Scotland, and supplementation reduces the proportion that are severely deficient. *Journal of Nutrition* 141(8): 1535–1542.

Appendix

Demographics

1. Age in years _____

2. Gender:

- Male
- Female
- Other (with a blank entry field for the participant to self-identify)
- Prefer not to answer

3. Are you currently pregnant or breastfeeding?

- Yes
- No
- Prefer not to answer

4. Employment:

- F/T employment
- P/T employment
- Not currently employed
- Student
- Retired

Household Income:

5. What was your household's total income from all sources over the last 12 months* **:

*Count income from every person in the household including benefits.

**Do not deduct tax/National Insurance/Health insurance payments/Superannuation

- Less than £100 per week OR less than £5200 per year
- £100–199 per week or £5200 to £10,399 per year
- £200–299 per week or £10,400 to £15,499 per year
- £300–399 per week or £15,500 to £20,799 per year
- £400–499 per week or £20,800 to £25,999 per year
- £500–699 per week or £26,000 to £36,399 per year
- £700–999 per week or £36,400 to £51,999 per year
- £1000–1499 per week or £52,000 to £77,999 per year
- £1500 per week or more to £78,000 or more per year
- Prefer not to answer

6. Highest level of education

- O Grade, Standard Grade, National 3, 4 or 5, Intermediate 1 or 2, GCSE, CSE or equivalent
- Higher, Advanced Higher, SCE Higher Grade, CSYS, A Level, AS Level or equivalent
- Apprenticeship (trade or equivalent)
- Apprenticeship (Foundation or equivalent)
- Apprenticeship (Modern or equivalent)
- Apprenticeship (Graduate or equivalent)
- GSVQ Foundation or Intermediate, SVQ level 1 or 2, SCOTVEC Module, City and Guilds Craft or equivalent
- GSVQ Advanced, SVQ level 3, ONC, OND, SCOTVEC National Diploma, City and Guilds Advanced Craft or equivalent
- HNC, HND, SVQ level 4 or equivalent
- Other school qualifications not already mentioned (including foreign qualifications)
- Other post-school but pre-Higher Education qualifications not already mentioned (including foreign qualifications)
- Degree, Postgraduate qualifications, Masters, PhD, SVQ level 5 or equivalent
- Professional qualifications (for example, teaching, nursing, accountancy) Other Higher Education qualifications not already mentioned (including foreign qualifications)
- No qualifications
- Prefer not to answer

Nationality

7. What is your nationality? _____

Residence

8. How long have you lived in Scotland: _____ years
_____ months

9. Postcode _____ (Please include full postcode e.g. DD1 1HG)

Ethnicity:**10. How would you describe your ethnic origin? (Tick one box only)**

What is your ethnic group? Choose ONE section from A to F, then tick ONE box which best describes your ethnic group or background:

A White:

- Scottish
 Other British
 Irish
 Polish
 Gypsy / Traveller
 Roma
 Showman / Showwoman

Other white ethnic group, please write in: _____

B Mixed or multiple ethnic groups:

- Any mixed or multiple ethnic groups, please write in: _____

C Asian:

- Scottish Asian or British Asian Pakistani, Scottish Pakistani or British

Pakistani:

- Indian, Scottish Indian or British Indian
 Bangladeshi, Scottish Bangladeshi or British Bangladeshi
 Chinese, Scottish Chinese or British Chinese
 Other, please write in: _____

D African, Scottish African or British African:

Please write in (for example, NIGERIAN, SOMALI): _____

E Caribbean or Black:

Please write in (for example, SCOTTISH CARIBBEAN, BLACK SCOTTISH): _____

F Other ethnic group

Arab, Scottish Arab or British Arab Other, please write in (for example, SIKH, JEWISH) _____

Vitamin D awareness**11. Have you heard of vitamin D?**

- Yes
 No

If NO reroute to Information-point 16

12. Where did you hear about vitamin D (tick all that apply)?

- Doctor/ Nurse
 Other Health Professional
 School / college / university
 Newspaper
 Family
 Friend
 Television
 Radio
 Poster/ Billboard
 Magazine
 Book
 Social media
 I don't know
 Other _____

Knowledge of vitamin D**13. Vitamin D helps with which of the following health effects? (Check all that apply)**

- Bone health
 Immune health
 Prevention of Rickets
 Vision Health
 Hair Growth
 Prevention of diabetes
 Cardio-vascular health
 Cognitive health
 Cancer prevention
 Skin softness
 Calcium absorption
 None of the above
 Other _____
 I don't know

14. Where do you think vitamin D comes from? (Check all that are correct)

- Fruits
 Vegetables
 Fatty fish
 Vitamin D supplements
 Select cereals
 Milk/ Dairy
 Nuts
 Cod liver
 Sun
 Eggs
 Mushrooms
 I don't know

15. Factors that can decrease the amount of vitamin D a person can get are (check all that apply):

- | | | |
|--|---|--|
| <input type="checkbox"/> Skin pigment | <input type="checkbox"/> Pregnancy/ lactation | <input type="checkbox"/> Dairy allergy |
| <input type="checkbox"/> Shade/ clouds | <input type="checkbox"/> Fatty diets | <input type="checkbox"/> Pollution |
| <input type="checkbox"/> Time of day | <input type="checkbox"/> Sunscreen use | <input type="checkbox"/> Wind |
| <input type="checkbox"/> Latitude | <input type="checkbox"/> Vegan diet | <input type="checkbox"/> Smoking |
| <input type="checkbox"/> Season | <input type="checkbox"/> Vegetarian diets | <input type="checkbox"/> I don't know |
| <input type="checkbox"/> Age | <input type="checkbox"/> Lactose intolerance | |

16. Information page with no questions will be displayed on JISC with the following text:

Although some foods can be a source of vitamin D it is incredibly difficult to get enough from food alone. The most effective sources, other than from vitamin D supplements is the sun.

17. How much time would the average fair-skinned person need to spend in the sun to get enough vitamin D, if their bare legs and arms were exposed and without sunscreen?

- Less than 9minutes per day
 9–14minutes per day
 15–20minutes per day
 More than 20minutes per day
 I don't know

18. How much time would the average non-fair-skinned (i.e. non-Caucasian) person need to spend in the sun to get enough vitamin D, if their bare legs and arms were exposed?

- Less than 15minutes per day
 15–24 minutes per day
 About 25–38minutes per day
 More than 38minutes per day
 I don't know

19. In Scotland, during which months of the year is the sunshine adequate to enable adults to make vitamin D, if their bare legs and arms were exposed? (Tick all months that apply)

- January
 February
 March
 April
 May
 June
 July
 August
 September
 October
 November
 December

20. Are you aware of the recommendation for adults to consider taking a daily vitamin D supplement between October and March each year in Scotland?

- Yes
 No

21. How many micrograms of vitamin D should you take as a daily supplement living in Scotland?

_____ micrograms

- Don't know

Vitamin D practices

22. On average how often, if at all, do you take a vitamin D supplement during the winter months?

- Every day (route to Q23)
 Several times a week (route to Q24)
 Once a week (route to Q24)
 Every couple of weeks (route to Q24)
 Monthly (route to Q24)
 Never (route to Q24)

23. What dose of vitamin D do you take in the winter months? _____ micrograms per day

(to convert international units to micrograms take the IU and divide it by 40. For example: 800 IU vitamin D / 40 = 20 micrograms vitamin D) (route to Q27)

24. If you don't take a supplement every day, what is the reason? Please select all that apply

- I'm unaware of the recommended advice
 I'm confused by the recommended advice
 I don't see the need
 I forget
 I prefer to get vitamins from food
 I can't afford supplements
 I don't know where to buy them
 I don't like taking supplements
 I don't agree with taking tablets
 I find it hard to swallow tablets
 Other, please state (_____)

25. Information page with no questions will be displayed on JISC with the following text:

In Scotland we only get enough of the right kind of sunlight for our bodies to make vitamin D between April and September, mostly between 11am and 3pm. Some people, including people from minority ethnic groups with darker skin, are at higher risk of vitamin D deficiency and should take a daily supplement all year round. Taking a daily 10 microgram vitamin D supplement, particularly between October and March, supports bone and muscle health and reduces our risk of vitamin D deficiency.

26. Knowing these benefits now, how likely are you to consider taking a daily 10 microgram vitamin D supplement between October and March?

- Very likely
- Likely
- Neither likely nor unlikely
- Unlikely
- Very unlikely

Supplements

27. Where do you buy your vitamin D supplements?

- Supermarket
- Health food store
- Chemist/Pharmacy
- Online
- Other, please state _____

28. What form of vitamin D supplement do you use?

- Tablets
- Caplets
- Spray
- Drops
- Jellies
- Chewable tablets

29. Which brand of vitamin D supplement do you currently use? _____

- Don't know the brand

30. On average, how much do you spend on vitamin D supplements per month? £ _____

- Don't know

31. Please indicate how many times, on average, you have taken the following supplements during the last 3 months:

	Never or <1 a month	1-3 a month	Once a week	2-4 a week	5-6 a week	Once a day	2-3 a day	4-5 a day	6 +
Multivitamin									
Calcium									
Calcium + Vitamin D									

FSS Campaign

32. Did you see/hear about the Food Standards Scotland campaign about vitamin D over the winter months?

- Yes (if yes route to Q33)
- No (route to point 34)
- Can't remember (route to point 34)



33. Where did you see/hear about the FSS campaign (please tick all that apply)

- Newspaper
- Television
- Radio
- Poster/ Billboard
- Magazine
- Social media
- Other _____

34. End of survey. Thank you for taking the time to complete this survey