

University of Dundee

DOCTOR OF PHILOSOPHY

Skin barrier dysfunction in common genetic disorders

Chen, Huijia

Award date:
2011

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

DOCTOR OF PHILOSOPHY

Skin barrier dysfunction in common genetic disorders

Huijia Chen

2011

University of Dundee

Conditions for Use and Duplication

Copyright of this work belongs to the author unless otherwise identified in the body of the thesis. It is permitted to use and duplicate this work only for personal and non-commercial research, study or criticism/review. You must obtain prior written consent from the author for any other use. Any quotation from this thesis must be acknowledged using the normal academic conventions. It is not permitted to supply the whole or part of this thesis to any other person or to post the same on any website or other online location without the prior written consent of the author. Contact the Discovery team (discovery@dundee.ac.uk) with any queries about the use or acknowledgement of this work.

INTRODUCTION	1
1.1 THE STRUCTURE OF SKIN	2
1.1.1 The epidermis	2
1.1.2 The dermis	6
1.2 FORMATION OF A SELECTIVELY PERMEABLE SKIN BARRIER	8
1.2.1 Structural protein components of the skin barrier	9
1.2.2.1 Keratins	9
1.2.1.2 Filaggrin	10
1.2.1.3 Involucrin	11
1.2.1.4 Loricrin	12
1.2.1.5 Small proline rich proteins (SPRRs)	12
1.2.1.6 Late cornified envelope proteins (LCEs)	13
1.2.1.7 Corneodesmosomes	14
1.2.2 Lipid components of the skin barrier	16
1.2.2.1 Ceramides	16
1.2.2.2 Free fatty acids	18
1.2.2.3 Cholesterol	18
1.2.3 Biochemical components involved in skin barrier formation	19
1.2.3.1 Calcium ions	19
1.2.3.2 Enzymes relevant in barrier formation	21
1.2.3.3 Anti-microbial peptides	22
1.3 THE GENETICS OF SKIN DISEASES	24
1.3.1 Methods for finding causative genes in skin disorders	24

1.3.1.1 Genetic linkage analysis	24
1.3.1.2 Candidate gene approach	26
1.3.1.3 Sanger sequencing	26
1.3.2 Mendelian skin disorders	27
1.3.2.1 Modes of inheritance	27
1.3.3 Complex genetic traits of the skin	32
1.3.4 Gene interactions in skin disorders	33
1.4 AIMS OF THESIS	35

**CHARACTERISATION OF FILAGGRIN GENE (*FLG*)-
NULL MUTATIONS IN COMMON GENODERMATOSES**

	36
2.1 INTRODUCTION	37
2.1.1 The biology of filaggrin	37
2.1.1.1 The role of filaggrin in skin barrier formation	37
2.1.1.1 The genetic structure of the filaggrin gene (<i>FLG</i>)	39
2.1.1.3 Expression control of profilaggrin	41
2.1.1.4 Post-translational processing of profilaggrin	42
2.1.1.5 Final degradation of filaggrin	48
2.1.2 Common genodermatoses associated with filaggrin	50
2.1.2.1 Ichthyosis vulgaris	50
2.1.2.2 Atopic dermatitis	51
2.1.3 Challenges in studying the global genetics of <i>FLG</i>	55
2.1.3 Aims of this chapter	57
2.2 MATERIALS AND METHODS	58

2.2.1 Clinical networking for dermatology research in Singapore	58
2.2.2 Clinical material and data collection	59
2.2.2.1 Pilot study of eight Singaporean Chinese ichthyosis vulgaris patients	59
2.2.2.2 FLG-null mutation discovery cohort	59
2.2.2.1 Childhood-adolescent Singaporean Chinese atopic dermatitis cohort	61
2.2.2.4 Population controls	61
2.2.2.5 Exclusion criteria and ethnicity of all cases and controls	62
2.2.3 Immunohistochemistry	63
2.2.4 Comprehensive sequence analysis of <i>FLG</i>	63
2.2.5 Medium-high throughput screening methodologies for <i>FLG</i>-null mutations	66
2.2.5.1 Fluorescent PCR and allelic size discrimination	66
2.2.5.2 Restriction digest and gel pattern analysis	69
2.2.5.3 Direct sequencing of small PCR fragments	71
2.2.6 Statistical analysis	71
2.3 RESULTS	74
2.3.1 Pilot study: <i>FLG</i>-null mutations in eight Singaporean-Chinese patients with ichthyosis vulgaris	74
2.3.2 Mining for novel <i>FLG</i>-null mutations in the discovery cohort	78
2.3.3 Observation of overlapping clinical features in ichthyosis vulgaris and X-linked ichthyosis	83
2.3.4 The association of <i>FLG</i>-null mutations with atopic dermatitis in Singapore	83

2.3.5 The Singaporean Chinese atopic dermatitis <i>FLG</i> -null mutation landscape is distinct from that in Europe	88
2.3.6 Palmar hyperlinearity and keratosis pilaris are clinical markers of <i>FLG</i> -null mutations	88
2.4 DISCUSSION	93
2.4.1 Population specific <i>FLG</i> -null mutations cause ichthyosis vulgaris in Singaporean Chinese	93
2.4.2 A wide spectrum of <i>FLG</i> -null mutations is present in Asia	95
2.4.3 A smaller proportion of Asian atopic dermatitis patients carry <i>FLG</i> -null mutations	98
2.4.4 The <i>FLG</i> -null mutation heterozygote advantage hypothesis in Europe and Asia	100
2.4.5 Towards the non-invasive diagnosis of <i>FLG</i> -null mutations	101
2.5 CONCLUSIONS	103
<u>THE FILAGGRIN GENE IN OTHER ALLERGIC CONDITIONS AND GENODERMATOSES</u>	105
3.1 INTRODUCTION	106
3.1.1 Secondary atopic conditions related to atopic dermatitis	106
3.1.1.1 Peanut allergy	107
3.1.2 Dry skin and acne vulgaris	108
3.1.3 Aims of this chapter	111
3.2 MATERIALS AND METHODS	112
3.2.1 Study populations	112

3.2.1.1 Childhood-adolescent Singaporean Chinese atopic dermatitis cohort (sub-group)	112
3.2.1.2 Peanut sensitisation pilot study	113
3.2.1.3 Adult Singaporean Chinese acne vulgaris cohort	114
3.2.2. Screening for Asian specific <i>FLG</i>-null mutations	114
3.2.3 Statistical analysis	115
3.3 RESULTS AND DISCUSSION	117
3.3.1 <i>FLG</i>-null mutations increase the susceptibility of recurrent infections in the presence of atopic dermatitis	117
3.3.2 <i>FLG</i>-null mutations are not significantly associated with peanut sensitisation in 27 Singaporean Chinese children	122
3.3.3 <i>FLG</i>-null mutations do not protect against acne vulgaris	124
3.4 CONCLUSIONS	127
 <u>INTERACTION OF SKIN BARRIER GENES IN GENODERMATOSES</u>	 <u>128</u>
4.1 INTRODUCTION	129
4.1.1 Interaction of different genes in genodermatoses	129
4.1.2 The role of <i>KLK7</i> and <i>SPINK5</i> in the skin barrier	130
4.1.3 The importance of the cholesterol synthesis pathway	131
4.1.3.1 The role of steroid sulphatase in the cholesterol synthesis pathway	131
4.1.3.2 Null mutations in the steroid sulphatase gene (STS) cause X-linked ichthyosis	133
4.1.4 Aims of this chapter	137
4.2 MATERIALS AND METHODS	138

4.2.1 Study populations and clinical material	138
4.2.1.1 Cases and cohorts for the SPINK5, KLK7 and FLG variants study	138
4.2.1.2 Indian and Singaporean Chinese cases of X-linked ichthyosis	139
4.2.2 Genotyping of <i>KLK7</i>, <i>SPINK5</i>, <i>STS</i> and <i>FLG</i>	139
4.2.2.1 Analysis of KLK7 by fluorescent PCR and allelic size discrimination	140
4.2.2.2 SNP analysis of SPINK5 by Taqman Genotyping Assay	140
4.2.2.3 Direct sequencing of STS	141
4.2.2.4 Sequencing of FLG and screening of novel FLG-null mutation	141
4.2.3 Detection of <i>STS</i> with fluorescent <i>in situ</i> hybridisation (FISH) and steroid sulphatase activity assay	143
4.2.4 Statistical analysis	144
4.3 RESULTS AND DISCUSSION	146
4.3.1 <i>KLK7</i> and <i>SPINK5</i> show no association with atopic dermatitis	146
4.3.2 Maternally-inherited <i>SPINK5</i> demonstrate weak interaction with <i>FLG</i>-null mutations	149
4.3.3 Exacerbation of X-linked ichthyosis phenotype due to <i>STS</i> and <i>FLG</i> interaction	151
4.3.3.1 Unusually severe X-linked ichthyosis in an Indian female	151
4.3.3.2 Homozygous STS and heterozygous FLG mutations in severe X-linked ichthyosis	156
4.3.4 Overlapping clinical phenotype between ichthyosis vulgaris and X-linked ichthyosis in Singaporean Chinese patients with atopic dermatis	160

4.3.4.1 Clinical features of atopic dermatitis patients with no detected FLG-null mutations	161
4.3.4.2 STS mutations detected in two Singaporean Chinese diagnosed with ichthyosis vulgaris and atopic dermatitis	164
4.4 CONCLUSIONS	167

THE EPIDERMAL DIFFERENTIATION COMPLEX (EDC)
CONTAINS PSORIASIS SUSCEPTIBILITY GENES **168**

5.1 INTRODUCTION	169
5.1.1 Psoriasis is caused by genetic changes in the immune system and skin	169
5.1.2 Genome-wide association studies identify psoriasis susceptibility genes in the EDC	171
5.1.3 Aims of this chapter	175
5.2 METHODS	176
5.2.1 Study population	176
5.2.2 Selection of EDC candidate genes and SNPs for genotyping	177
5.2.3 Genotyping with Sequenom[®] and Infinium[™] platforms	179
5.2.4 Statistical and stratification analyses	180
5.3 RESULTS	182
5.3.1 Fine mapping of the EDC reveal significant association of psoriasis with the involucrin gene (<i>IVL</i>) region	182
5.3.2 Expanded analysis further reveal significant association of psoriasis with an additional EDC gene region	185
5.4 DISCUSSION AND CONCLUSION	190

5.4.1 The EDC is an important psoriasis susceptibility locus in Asia 190

**5.4.2 Candidate genes for psoriasis susceptibility lie in close proximity
with *IVL* or small proline rich repeat genes (*SPRRs*) 191**

OVERALL CONCLUSIONS AND DISCUSSION 195

REFERENCES 202

APPENDIX I: PUBLICATIONS

**APPENDIX II: COMPREHENSIVE TABLES OF SNPS
GENOTYPED USING SEQUENOM AND INFINIUM
PLATFORMS**