



University of Dundee

Protein Kinases in Pluripotency - Beyond the Usual Suspects

Fernandez-Alonso, Rosalia; Bustos, Francisco; Williams, Charles A. C.; Findlay, Greg M.

Published in:
Journal of Molecular Biology

DOI:
[10.1016/j.jmb.2017.04.013](https://doi.org/10.1016/j.jmb.2017.04.013)

Publication date:
2017

Licence:
CC BY

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Fernandez-Alonso, R., Bustos, F., Williams, C. A. C., & Findlay, G. M. (2017). Protein Kinases in Pluripotency - Beyond the Usual Suspects. *Journal of Molecular Biology*, 429(10), 1504-1520.
<https://doi.org/10.1016/j.jmb.2017.04.013>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Protein Kinases in Pluripotency—Beyond the Usual Suspects

Rosalia Fernandez-Alonso, Francisco Bustos,
Charles A.C. Williams and Greg M. Findlay

The MRC Protein Phosphorylation and Ubiquitylation Unit, School of Life Sciences, The University of Dundee, Dundee DD1 5EH, UK

Correspondence to Greg M. Findlay: g.m.findlay@dundee.ac.uk

<http://dx.doi.org/10.1016/j.jmb.2017.04.013>

Edited by Patrick Griffin

Abstract

Post-translational modification of proteins by phosphorylation plays a key role in regulating all aspects of eukaryotic biology. Embryonic stem cell (ESC) pluripotency, defined as the ability to differentiate into all cell types in the adult body, is no exception. Maintenance and dissolution of pluripotency are tightly controlled by phosphorylation. As a result, key signalling pathways that regulate pluripotency have been identified and their functions well characterised. Amongst the best studied are the fibroblast growth factor (FGF)-ERK1/2 pathway, PI3K-AKT, the leukemia inhibitory factor (LIF)-JAK-STAT3 axis, Wnt-GSK3 signalling, and the transforming growth factor (TGF) β family. However, these kinase pathways constitute only a small proportion of the protein kinase complement of pluripotent cells, and there is accumulating evidence that diverse phosphorylation systems modulate ESC pluripotency. Here, we review recent progress in understanding the overarching role of phosphorylation in mediating communication from the cellular environment, metabolism, and cell cycle to the core pluripotency machinery.

© 2017 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Pluripotency is a fundamental of metazoan biology and is defined as the theoretical capability of a single cell to differentiate into any lineage in the developing organism [1]. The potential of pluripotent cells to form any tissue or organ in the body has pushed pluripotency research to the fore in the field of regenerative medicine. Pluripotent cells were initially isolated from developing embryos as embryonic stem cells (ESCs) [2–4]. More recently, induced pluripotent stem cells (iPSCs) have been derived by developmentally reprogramming somatic cells, and these closely resemble ESCs at the molecular level [5–7].

Pluripotency comprises at least two molecularly distinct states, which differ according to species and developmental context [8]. Naïve pluripotency is a developmental “ground state” characteristic of cells from the preimplantation mouse embryonic epiblast [mouse ESCs (mESCs)]. Primed pluripotency is

characteristic of post-implantation mouse epiblast stem cells (EpiSCs) and ESCs isolated from human embryos (hESCs), although human naïve pluripotent cells were recently derived from primed hESCs or human embryos [9–16]. An intermediate state termed formative pluripotency was recently described [17], which represents the initial acquisition of developmental characteristics by naïve cells. Molecular distinctions and definitions of pluripotent states have been extensively reviewed elsewhere [18].

Pluripotency acquisition and maintenance are intrinsically linked to the expression of a pluripotency gene regulatory network, particularly the core transcription factor triumvirate of OCT4/PO5F1, SOX2, and NANOG [19–21]. OCT4 is a “governor” of pluripotency and is expressed in both naïve and primed pluripotent states via distinct enhancer elements [22]. OCT4 also controls lineage allocation following pluripotent exit [23]. SOX2 and NANOG are key regulators of naïve and primed pluripotency,

although Nanog is expressed at a reduced level in primed cells. Other transcriptional regulators of pluripotency are differentially expressed in naïve and primed cells. The naïve state is marked by the expression of the key NANOG effector *Esrrb*, Krueppel-like factors (*Klf2/4*), *Rex1*, *Fgf4*, and *Nr0b1* [24]. As the term suggests, primed pluripotent cells express markers of lineage priming, including the de novo DNA methyltransferases *Dnmt3a/b* [25], the embryonic epiblast marker *Fgf5*, and lineage-specific transcription factors such as *Brachyury* [24].

Pluripotency gene regulatory networks are under strict control of extrinsic and intrinsic signalling networks. Dynamic flow of signalling information between and within cells is dependent upon networks of reversible post-translational modifications, particularly protein phosphorylation [26]. Thus, pluripotent states can be accessed and stabilised by modulating activities of protein kinases. Specifically, cytokines, growth factors, and selective protein kinases inhibitors can be exploited to manipulate pluripotency pathways. Pluripotent mESCs were initially captured [24] using a combination of bone morphogenetic protein (BMP) to activate the SMAD1 inhibitor of differentiation pathway and leukemia inhibitory factor (LIF) to activate JAK-STAT3 signalling. Accessing naïve “ground state” mESC pluripotency, however, requires the inhibition of two protein kinases: MEK1/2, which phosphorylates and activates ERK1/2 MAP kinase; and GSK3, an antagonist of Wnt signalling [27,28]. Primed pluripotency in human and mouse is supported by fibroblast growth factor (FGF) and Activin [29–31], whilst human naïve pluripotency can be accessed using distinct combinations of growth factors and inhibitors [9–16].

In this review, we train our focus away from these well-understood pluripotency signalling pathways to explore the role of emerging signalling networks and their impact upon the maintenance and dissolution of pluripotent states. Furthermore, we shed light on exciting but poorly appreciated roles for cell cycle, environmental, metabolic, structural, and stress-regulated phosphorylation networks in pluripotency regulation.

Novel Pluripotency Signalling Pathways

The human protein kinome consists of 538 kinases[†] [32] and includes some of the most studied enzymes in biology. However, vast swathes of the kinome have not yet been investigated, and understudied kinases likely perform key functions in biological processes such as pluripotency maintenance and dissolution [33]. Indeed, total and phosphoproteome analysis indicates that at least 300–400 kinases are expressed in pluripotent mESCs (Jens Hukelman and G.M.F., unpublished data) and hESCs [34]. In this section, we discuss technologies that have been employed to uncover new pluripotency kinase signalling pathways.

Phosphoproteomic profiling is a powerful method to identify novel mechanisms by which phosphorylation modulates pluripotency. Initial studies focussed on identifying novel targets of well-understood signalling pathways (e.g., FGF2 in hESCs) [35,36]. However, unbiased comparison of the phosphoproteomes from pluripotent hESCs and those from differentiating or somatic cells has uncovered new phosphorylated targets relevant for pluripotency regulation. In this manner, multiple receptor tyrosine kinases with previously unappreciated roles in hESC pluripotency were identified [37]. Furthermore, kinase-substrate motif prediction analysis of differentially phosphorylated proteins indicates that distinct families of kinases are active in hESCs compared to somatic cells [38]. Importantly, this approach suggests novel roles for cyclin-dependent kinases (CDKs), Aurora, p38, and c-Jun N-terminal kinases (JNKs) in pluripotency regulation [34] and reveals key hESC phosphorylation modules centred on CDK1/2 [39] and DNA methyltransferases [40]. Perhaps most excitingly, phosphoproteomic studies have the potential to reveal new pluripotency kinases, affirming phosphoproteomics as a core tool for systematic identification of novel mechanisms of pluripotency regulation.

Functional screening has also proven invaluable to uncover new pluripotency kinases. Global RNA interference screens have identified kinases that block mESC pluripotency maintenance [41] and reprogramming to pluripotency [42]. Interestingly, kinases such as TESK1 and LIMK2 identified in these studies have no previously described role in pluripotency or reprogramming. Future research will explore the molecular functions and regulation of these kinases in pluripotent cells. Similarly, cellular screening of kinase inhibitor libraries has elucidated new pluripotency pathways. A survey of selective kinase inhibitors uncovered a novel role for the ERK5 MAP kinase pathway in modulating mESC naïve-primed pluripotent transition [43]. In mESCs, ERK5 promotes the expression of a key network of naïve pluripotency factors, including *Klf2*, *Esrrb*, and *Rex1* [43,44], which suppresses the transition of naïve cells to the primed state. Interestingly, both the ERK5 kinase and a C-terminal transcriptional activation domain are required for naïve maintenance [43]. ERK5 may therefore function in concert with transcription factors SP1 and MEF2 [45–47], which are required for *Klf2/4* expression in other developmental systems [44,46,48]. As with many emerging pluripotency pathways, identification of novel ERK5 substrates would shed light on the mechanisms by which ERK5 controls pluripotency. In addition, identifying factors that specifically activate ERK5 may have utility in capturing naïve pluripotency and/or reprogramming somatic cells. In this regard, BMP, LIF, and FGF activate ERK5 in mESCs and other cell types [44,49,50].

Identification of novel pluripotency kinase pathways thus represents an exciting niche within the ESC

arena. Increasingly potent and selective tool compounds, including high-value collections from pharmaceutical companies and academic consortia [51,52] in combination with genome editing technologies, will provide unique opportunities to specifically disrupt kinase function in pluripotent cells [33]. Ultimately, we envision that pathways identified using these cutting-edge approaches will influence pluripotent stem cell-based clinical applications. In the meantime, a number of more prominent kinase signalling networks have relatively poorly understood roles in pluripotency regulation, which we now discuss.

Cell Cycle and DNA Damage Response (DDR) Kinases in Pluripotency Regulation

Cell cycle progression is governed by a complex kinase network centred on the CDKs. Cell cycle phase is intrinsically linked to pluripotency maintenance and lineage commitment, with G1 phase providing a key window for developmental decision-making. A short G1 is deterministic for pluripotency maintenance [53], whilst lengthening G1 promotes differentiation [54], implying that kinases that govern cell cycle progression couple directly to the pluripotency machinery.

Cell cycle kinases and pluripotency

Phosphoproteomic analysis pinpoints CDK1/2 as a pluripotency hub in hESCs [39], and several themes are now emerging with regard to the mechanisms by which CDKs modulate pluripotency. CDKs directly regulate phosphorylation and/or expression of pluripotency factors or couple developmental decision-making to cell cycle phase. CDK activity can also pattern the activation of signalling pathways that regulate pluripotency and differentiation.

In mESCs, CDK1 kinase activity is essential for pluripotency maintenance [55], which is underpinned at least in part by the modulation of core pluripotency factors. CDK1 and OCT4 appear to cooperate in an unknown way to repress differentiation [56]. However, direct phosphorylation of SOX2 by CDKs promotes pluripotency acquisition [57], whilst NANOG is phosphorylated by CDK1 *in vitro* [58]. Although the mechanisms and functional significance remain unclear, these studies set a precedent for the direct modulation of pluripotency factors by CDK-Cyclin activities. Comprehensive analysis of CDK-dependent phosphorylation of core pluripotency transcription factors will determine the generality of this mechanism. In addition to direct phosphorylation of pluripotency factors, transcriptional regulation of pluripotency by CDK-Cyclins has been described. In hESCs, Cyclin D forms specific transcriptional complexes to control lineage specific genes [59], although this is independent of CDK kinase activity. CDK2 activity also

promotes lineage-specific gene expression in hESCs by phosphorylating and activating the histone methyltransferase MLL2 [60].

Perhaps unsurprisingly, CDK-Cyclin activities also control pluripotency via cell cycle phase, and as discussed previously, this may relate to G1 length [61]. Gonzales and colleagues describe a molecular mechanism underpinning G1 as a critical cell cycle phase during which mESC pluripotency is maintained or dissolved [62]. Elevated CDK-Cyclin B and ataxia telangiectasia mutated (ATM)/ATR kinase activities during S and G2 cell cycle phases of the mESC cycle protect against pluripotent exit. However, in G1, these activities are less prominent, providing a window for differentiation [62].

CDK activity also modulates pluripotency more indirectly by patterning the signalling landscape. CDK4/6 phosphorylates SMAD2/3 at a distinct regulatory region to suppress transforming growth factor beta (TGF β)-SMAD2/3-dependent expression of lineage-specific genes [63], indicating that CDK-Cyclin activities not only directly modulate expression and function of pluripotency genes but also impact the interpretation of developmental signals. This may be an important mechanism by which CDKs coordinate pluripotency, ensuring that cells respond to developmental cues only within the appropriate cell cycle phase.

Finally, other cell cycle kinases have been uncovered as key pluripotency regulators. In mESCs, a kinome-wide RNAi screen showed that mitotic Aurora kinases regulate phosphorylation and degradation of p53 to promote pluripotency [41]. Furthermore, Aurora modulates OCT4 function in mESCs [64], exemplifying the increasingly important molecular connections between cell cycle protein kinases and pluripotency. ESCs have therefore evolved a variety of mechanisms to tightly couple pluripotency and cell cycle phase, and these await further discovery and investigation.

DNA damage signalling in pluripotent cells

The unique ESC cell cycle, with its short G1 phase, subjects the genome to significant replication stress [65]. This can drive pluripotency exit [66], presumably to prevent the potential transmission of mutations to future somatic lineages and the germline. As a result, the DDR is “hard-wired” to the pluripotency machinery to ensure that DNA is efficiently repaired in pluripotent cells [66]. As in somatic cells, the DDR centres on CHEK1/2 and ATM/ATR kinases. Activation of CHEK1 [67] and ATM [68] is required for reprogramming and iPSC genome stability, establishing the importance of DDR kinases in acquisition and maintenance of pluripotency. Interestingly, mESCs can activate ATM without inducing p53 [65] in order to mount an efficient DDR without promoting differentiation.

Although ESCs use the same conserved core DDR machinery as somatic cells, specific factors

orchestrate DDR signalling to maintain ESC genome integrity. In mESCs, the pluripotency factor SALL4 recruits the Rad50/MRN DDR complex to specific genomic regions, which activates ATM to promote DNA repair [69]. Furthermore, mESCs specifically utilise the scaffold FILIA/KHDC3 to activate DDR kinases [70]. FILIA is phosphorylated following DNA damage, which activates ATM and CHEK2, promoting DNA repair and genome stability [70]. A major unresolved question concerns why ESCs utilise unique factors to coordinate the DDR.

Nutrition and energy sensing kinases in pluripotency

As exemplified by the cell cycle and DDR, fundamental cellular processes profoundly influence pluripotency acquisition and/or maintenance. Cellular nutrition and energy status continue along this theme (See Fig. 1). As in other cell types, ESC nutrient responses are controlled by the PI3K-related protein kinase mechanistic target of rapamycin (mTOR) [71].

mTOR is the kinase in two distinct complexes: mTORC1, defined by mTOR interaction with regulatory-associated protein of mTOR (RAPTOR) [72,73], and mTORC2, defined by mTOR interaction with rapamycin-insensitive companion of mTOR (RICTOR) [74]. In response to nutrients, mTORC1 phosphorylates and activates p70 S6 kinase 1 (which phosphorylates ribosomal S6 protein [75]) and eIF4E binding protein 1 [76]. mTORC1 integrates nutritional signals to enhance global and cap-dependent translation, ribosome biosynthesis, gene expression, and autophagy [77,78].

mTOR is a central regulator of ESC pluripotency

Consistent with its role as a major nutrient sensor, mTOR is critical for ESC growth and proliferation [79], early embryonic development [80], and pluripotency [79,81–85] in both mouse and human. In hESCs, mTORC1 activity is required for long-term pluripotency maintenance and promotes the expression of pluripotency factors OCT4, SOX2, and NANOG [85]. Furthermore, mTORC1 inhibition results in hESC




Figure 1. Metabolism and energetic signaling in pluripotency maintenance, acquisition, and exit. (Left) Primed mESC, hESC, and iPSC primarily utilise glycolytic metabolism. Under these metabolic conditions, AMPK signaling phosphorylates and activates ULK1 to stimulate mitophagy. Pluripotency factors such as OCT4, SOX2, and NANOG are highly expressed, and SOX2 recruits NuRD repressor complex to restrict mTOR expression. However, mTOR signaling supports pluripotency and protein translation. (Right) Differentiation induces a metabolic switch to mitochondrial oxidative respiration-based metabolism. mTOR expression and activation increase to elevate global translation. High mTOR activity promotes ULK1 inhibitory phosphorylation, which suppresses mitophagy, allowing mitochondrial accumulation.

differentiation to mesendoderm [83], suggesting an involvement of mTORC1 activity in pluripotency and restraining lineage specification.

mTOR complexes display highly specific signalling dynamics in pluripotent and differentiating cells. Although active in mouse and human ESCs, mTORC1&2 activity further increases during differentiation [81,84,86], which may require ERK1/2 activation of p90 ribosomal S6 kinase [81]. Interestingly, an mTOR-specific protein inhibitor, DEPTOR, acts to maintain mTOR kinase activity below a threshold level in both hESCs and mESCs [87]. DEPTOR suppression may thereby promote the increased mTORC1&2 activity observed upon ESC differentiation.

As expected from these studies, mTOR also displays complex signalling and expression dynamics during somatic cell reprogramming. In the mouse, mTORC1 and mTORC2 activities decrease during reprogramming via SOX2-dependent recruitment of the NuRD epigenetic remodelling complex, which shuts down the *mTOR* promoter [88]. Congruently, mTOR kinase inhibition enhances somatic cell reprogramming [76], although restoration of mTOR activity is critical for the final steps of iPSC generation, presumably because mTOR is required for pluripotency maintenance [81,84].

Most reports therefore suggest that different thresholds of mTOR signalling specify distinct developmental identities. Consistent with this notion, exciting recent work shows that partial inhibition of mTORC1 activity maintains a state of “paused pluripotency” similar to diapause, a developmental arrest induced by nutrient starvation [89]. Why restrained mTOR activity is more compatible with pluripotency is not currently understood. However, mTOR drives global translation, and mESCs have lower translation rates than differentiated cells [84,86], which may buffer against spurious expression of lineage-specific factors and differentiation.

Amino acid signalling in ESC pluripotency and self-renewal

A key function of the mTOR pathway is to mount cellular responses to nutrient availability, primarily amino acid levels. In contrast to somatic cells, where mTORC1 is activated in response to branched chain amino acids leucine and arginine [90], mESCs are exquisitely dependent upon threonine and show little requirement for leucine/arginine for growth/proliferation [91]. Intriguingly, threonine reportedly regulates activity of both mTORC1&2 complexes in mESCs to maintain proliferation and *Oct4/Pou5f1* expression [92]. How threonine might control mTORC1&2 is not known, although evidence suggests that this mechanism is distinct from the canonical branched chain amino acid sensing by RAG GTPases [93]. Thus, elucidating the molecular basis of this unique ESC nutrient-sensing system remains an important unresolved question.

Kinase Signalling and Metabolic Programming in Pluripotency

As is the case with the nutrient-sensing machinery, metabolic status and pluripotency are fundamentally intertwined. Distinct pluripotent states differentially utilise glycolytic and oxidative energy production systems [94], which play a key role in pluripotency maintenance. Whilst naïve ESCs and somatic cells produce energy via mitochondrial oxidative phosphorylation, primed hESCs and mESCs obtain their ATP from glycolysis [95]. Transition between pluripotent states, pluripotency acquisition, and differentiation therefore involves a metabolic conversion [94,96–99], and mTOR plays several distinct roles in metabolic regulation in ESCs.

Autophagy/mitophagy in metabolic switching and pluripotency

mTOR supports glycolytic metabolism and pluripotency in primed hESCs and mouse EpiSCs [100], at least in part by promoting the expression of pentose phosphate pathway genes [101,102]. In mESCs, mTOR also controls a unique GALECTIN-1 system for glucose uptake, which is essential for proliferation [103]. However, the best understood mTOR function in metabolic conversion from oxidative to glycolytic metabolism is via mitochondrial removal by the process of mitochondrial autophagy (mitophagy) [104]. Autophagy recycles proteins and organelles via the formation of intracellular autophagosomes targeted for lysosomal degradation [105]. This is critical for mESC pluripotency maintenance, as oxidative metabolism drives pluripotent exit [104]. Conversely, mESC differentiation increases mitochondrial synthesis to support oxidative metabolism [106]. In a similar vein, active mitophagy ensures that hESCs have relatively few mitochondria compared to somatic cells [7,107], which restrict differentiation.

In addition to mTOR, autophagy/mitophagy is tightly regulated by the activities of several key protein kinases. A complex containing UNC51-like kinase-1 (ULK1) controls isolation membrane (phagophore) assembly, autophagosome nucleation, and expansion [108]. ULK1 integrates opposing signals from AMP-activated protein kinase (AMPK) and mTORC1 to control phagophore assembly. AMPK phosphorylates ULK1 to drive autophagy, whilst mTOR phosphorylates a distinct site on ULK1 to suppress autophagy [109,110]. Elevated mTORC1 activity observed during differentiation thereby reduces autophagy in somatic cells [81].

Autophagy has been extensively studied during reprogramming to pluripotency. Canonical (ATG3-dependent [88]) and non-canonical autophagy pathways [111,112] are activated during

reprogramming to drive oxidative to glycolytic conversion [76]. AMPK-induced autophagy is required for pluripotency acquisition in both mouse and human [76,88,113,114], and pharmacological AMPK activation or mTOR inhibition enhances cell reprogramming by modulating autophagy [76]. Furthermore, increased *mTOR* transcription and mTOR kinase activity during the final stages of reprogramming drive autophagy, which aids pluripotency acquisition [88]. Specifically manipulating mitophagy and metabolic reprogramming using kinase inhibitors would add further valuable insight into the role of these processes in pluripotency.

Cellular and Environmental Stress Signalling in Pluripotency

Like nutrient and metabolic status, environmental stress profoundly impacts pluripotency, although ESCs have high stress tolerance compared to differentiated cells [115,116]. Responses to environmental stress are mediated by classical MAP kinase cascades [117] and other kinase signalling pathways. These pathways therefore perform key functions in pluripotent cells (See Fig. 2).




Figure 2. Multiple stress signalling pathways modulate pluripotency and differentiation. Oxidative stress signalling promotes the expression of lineage-specific factors by activating multiple signalling pathways. Reactive oxygen species (ROS) produced by oxidative metabolism promotes the activation of ERK1/2 via inhibition of protein tyrosine phosphatases (PTPs). ROS also results in the dissociation of TRX and ASK1 to enable JNK1 and p38 activation. These pathways are thought to promote pluripotent exit and differentiation in response to ROS. Hypoxic conditions also increase ROS production and endoplasmic reticulum (ER) stress, which induces unfolded protein response (UPR) genes, VEGF induction, and differentiation. Hyperosmotic activation of p38 promotes Oct4 expression, whilst JNK inhibits KLF4 and OCT4. In contrast, shear stress (SS) promotes the expression of pluripotency factors including OCT4 and REX1. The ERK5 signalling pathway may be a key mechanism to integrate SS signals with the core pluripotency gene regulatory network.

Oxidative stress

As discussed above, primed pluripotent ESCs rely primarily on glycolysis for ATP generation [99], which minimises oxidative stress generated by reactive oxygen species (ROS) and oxidative metabolism [118]. In hESCs, inhibition of oxidative phosphorylation enhances pluripotency [119], and culture in low oxygen/hypoxia improves pluripotency maintenance and reduces chromosomal abnormalities [120,121]. Although acute oxidative stress does not directly impact pluripotency gene expression [122], it can influence lineage choice during differentiation [41,123,124]. ESCs therefore appear to be metabolically programmed to avoid differentiation induced by oxidative stress.

Oxidative stress activates the ERK1/2, JNK, and p38 MAP kinase signalling cascades, the phosphoinositide 3-kinase (PI(3)K)/AKT pathway, and the nuclear factor (NF)- κ B signalling pathways [125]. ERK1/2 and PI(3)K/AKT activation occurs largely by protein tyrosine phosphatase inhibition and receptor tyrosine kinase phosphorylation [125], whilst the p38 and JNK MAP kinase cascades are activated by apoptosis signal-regulating kinase 1 (ASK1). The redox regulatory protein thioredoxin (TRX) normally inhibits ASK1, but oxidative stress dissociates the TRX-ASK1 complex, facilitating JNK and p38 phosphorylation and activation [126,127]. NF- κ B is also activated by several stress pathways, including I κ B kinase (IKK) α/β and ERK1/2 [128].

In mESCs, NF- κ B transcription promotes differentiation [129], and NANOG maintains pluripotency at least in part by suppressing NF- κ B signalling [130]. NF- κ B transcription is also activated during hESC differentiation [131], although intriguingly, the key NF- κ B kinase activator IKK α/β promotes OCT4 and NANOG expression in mESCs [132], suggesting that the role of the NF- κ B signalling pathway in pluripotency is not yet fully understood. ERK1/2, JNK, and p38 MAP kinase signalling pathways also promote mESC pluripotent exit [133–135], although whether these are oxidative stress-specific responses remains to be determined. Furthermore, as we discuss below, the role of MAP kinases in pluripotency may depend on both cellular context and type of stress response. Nevertheless, these data indicate that signalling responses to oxidative stress have a central function in pluripotency maintenance and gene expression.

Hyperosmotic stress

Hyperosmotic stress occurs when external osmolarity exceeds the intracellular physiological range and has been linked to pluripotency regulation. The WNK1–4 family kinases have emerged as key modulators of the cellular response to hyperosmotic conditions by regulating ion transport [136,137]. However, investigation of WNK kinase functions in

pluripotent cells awaits the development of selective tool compounds. To date, most studies into kinase signalling in hyperosmotic stress responses focus on MAP kinase signalling pathways p38 and JNK.

In contrast to oxidative stress, hyperosmotic activation of p38 actively promotes pluripotency [138] and facilitates somatic reprogramming to iPSCs [138,139]. During reprogramming, p38 specifically promotes the demethylation of pluripotency gene promoters [138] via an undefined mechanism. In addition, p38 kinase inhibitors specifically suppress OCT4 transcriptional activity under hyperosmotic conditions [140]. In contrast, JNK activity promotes pluripotent exit following acute hyperosmotic stress by suppressing OCT4 expression [141] and phosphorylating KLF4 to suppress transcriptional activity [142]. Therefore, a somewhat confusing picture of pathway-specific stress responses in pluripotent cells is emerging. Dissecting the roles of individual kinases under specific stress conditions using selective kinase inhibitors should provide clarity to this field.

Signalling responses to other environmental and cellular stresses

A further stress exerted on pluripotent cells is shear stress (SS), caused by fluid movement over the cell surface. First reported in endothelial cells [143], SS occurs in response to load-bearing or surface laminar flow [144] and can be lethal in preimplantation embryos [145]. ESCs reportedly mount differential responses depending on the strength and duration of SS. In pluripotent mESCs, SS stabilises the expression of *Oct4* and the naïve pluripotency marker *Rex1* [146]. In contrast, acute SS promotes ectodermal fate in differentiating mESCs, whilst prolonged SS results in mesendoderm specification [147]. Thus, it may be desirable to exploit culture conditions that generate a certain threshold of SS to efficiently propagate naïve pluripotent cells.

SS signals are primarily transduced by mechanosensors such as integrins, receptor tyrosine kinases (particularly VEGFR2), G-protein-coupled receptors, ion channels, and intercellular junction proteins [148–152]. These trigger kinase signalling cascades that modulate the transcription of SS response genes. In particular, the ERK5 pathway, which promotes naïve pluripotency in mESCs [43], plays a key role in SS responses. ERK5 is thought to mediate SS responses via KLF2 induction [153] or NRF2 activation [154] and therefore may be the missing link that connects SS signals to the core pluripotency machinery.

Finally, endoplasmic reticulum (ER) homeostasis and the unfolded protein response (UPR) are crucial for cell viability and are regulated by protein kinases IRE1 and PERK [155]. In ESCs, ER stress promotes Vascular Endothelial Growth Factor (VEGF) expression and vascular differentiation [156], and inhibiting ER stress reduces VEGF expression and supports

mESC pluripotency [157,158]. However, hESCs have a relatively active UPR [159], suggesting that pluripotent cells nevertheless rely on a threshold level of UPR for survival and/or to maintain pluripotency. Thus, a key question is establishing how ESCs effectively balance the UPR to prevent spurious differentiation.

Mechanical and Structural Signalling in Pluripotency

As we have seen, eliciting appropriate stress responses is critical to maintain pluripotency and survival at the level of individual cells. However,

metazoan development also requires that cells sense the topography of their immediate environment. As a result, intercellular forces and mechanical signals from the extracellular matrix (ECM) profoundly influence pluripotency (See Fig. 3). Understanding the molecular pathways involved in ESC mechanotransduction will be invaluable to develop effective cell culture strategies to exploit pluripotent cells for tissue regeneration [160].

Cell-cell interactions: signaling downstream of cadherins in pluripotency

The mammalian genome encodes more than 100 cadherins [161], which are key components of




Figure 3. Mechanical and structural signaling in pluripotency. (1) E-cadherin (CDH1) is a key component of adherens junctions and forms a ternary complex with the LIF receptor and gp130 in mESCs. This complex is required for efficient JAK-STAT3 signalling and pluripotency factor expression. The STAT3 target KLF4 activates *Cdh1* expression, generating a positive feedback loop to stabilise naïve pluripotency. (2) Integrins couple the extracellular matrix (ECM) to intracellular signaling. Integrins activate focal adhesion kinase (FAK) to support pluripotency via activation of PI3K-AKT and MDM2. Downstream, the Src family kinase (SFK) YES phosphorylates the transcription factor YAP to promote pluripotency. (3) The Hippo pathway provides a further link between cellular mechanics and pluripotency. The Ser/Thr kinases MST1/2 and LATS1/2 form a kinase cascade that phosphorylates YAP (at a site distinct from that phosphorylated by YES), which promotes 14-3-3 binding and cytoplasmic retention. Nuclear YAP/TAZ/TEAD promotes the transcription of pluripotency genes including *Oct4*.

adherens junctions. E-cadherin (CDH1) is a master regulator of mESC and hESC biology [162] and underpins the compact mESC morphology [163]. A key signalling function of CDH1 in mESCs is to support the activation of the LIF signalling pathway. *Cdh1*^{-/-} mESCs respond poorly to LIF stimulation [164–166], displaying reduced STAT3 activation and pluripotency gene expression [167]. CDH1 forms a ternary complex with LIF receptor and gp130, and this interaction is required for efficient LIF signalling [164]. Interestingly, this system is stabilised by a positive feedback loop, whereby the STAT3 target *Klf4* activates the *Cdh1* promoter [168], promoting cadherin-based cell–cell adhesion and LIF signalling to maintain pluripotency.

In primed ESCs, the cadherin picture becomes more complex. CDH1 depletion converts naïve mESCs into primed EpiSCs, whereupon N-cadherin (CDH2) is upregulated [167] to support pluripotency in this context [169]. However, CDH1 retains a critical signalling role in primed cells, by potentiating ACTIVIN-SMAD2/3 signalling and *Nanog* expression in EpiSCs [165], and PI3K-AKT signalling and *NANOG* and *OCT4* expression in hESCs [170]. Therefore, cadherins have clear signalling functions in both naïve and primed ESCs, and these are required for pluripotency maintenance.

Cell–ECM interactions: signalling downstream of integrins in pluripotency

Integrins are heterodimeric transmembrane receptors that couple the ECM to intracellular signalling networks via cytoskeletal adaptor proteins [171]. Although integrin engagement with ECM substrates supports self-renewal and pluripotency of hESCs [172–176], the mechanisms are poorly understood in comparison to the cadherins. Nevertheless, we explore the mechanisms by which integrins signal to the core pluripotency gene regulatory network.

Focal adhesion kinase (FAK) is a non-receptor tyrosine kinase activated by focal adhesion formation and integrin activation. FAK activation generates phosphotyrosine docking sites for adaptor/scaffolding proteins and signalling molecules [177], which include Src family kinases (SFKs), the scaffold p130CAS, the GRB2 adaptor, and PI3K. These activate downstream signalling pathways including ERK1/2 and AKT [178]. In hESCs, integrin signalling to FAK supports pluripotency, and FAK inhibition results in differentiation and anoikis [179]. Integrin activation of FAK activates PI3K-AKT and MDM2 to suppress p53 activity [179], which supports cell survival and pluripotency [180,181]. However, FAK is also activated following hESC differentiation and inactivated during the reprogramming of fibroblasts to iPSCs [182]. Furthermore, mESC pluripotency maintenance inversely correlates with integrin activation [183]. Therefore, there is contradictory evidence regarding the role of integrin signalling in pluripotency, which will only be

resolved by developing strategies to disentangle the pleiotropic functions of integrins and FAK in ESCs.

SFKs are membrane-associated non-receptor tyrosine kinases, which transduce signals from integrins and other cell surface receptors to the actin cytoskeleton [184]. There are eight mammalian SFKs [185], many of which are expressed in mESCs. Paradoxically, inhibition of all SFKs in mESCs promotes pluripotency [186], whilst inhibition of specific isoforms can induce differentiation [187]. This conundrum was elegantly tackled using inhibitor-resistant SFK mutants, which showed that selective c-SRC activation induces primitive endoderm differentiation [188]. In contrast, specific YES knockdown suppresses *Nanog* and *Oct4*, inducing mESC differentiation [189]. Interestingly, the differentiation-promoting activity of c-SRC is antagonised by YES in mESCs [189] and hESCs [190], indicating that these kinases may directly inhibit or compete with each other.

To further complicate matters, SFKs isoforms are activated by multiple stimuli. In addition to activation by integrins and receptor tyrosine kinases such as FGFRs [35], YES and HCK are activated by LIF in mESCs via recruitment to gp130 [186,191–193], consistent with a role in pluripotency signalling. YES kinase activity is suppressed following mESC differentiation [189], whilst HCK activation reduces the LIF requirement for mESC self-renewal [191]. Additionally, specific SFK functions may be underpinned by distinct developmental expression profiles. OCT4 positively regulates Yes expression in mESCs [194], ensuring that YES is abundant and activated in pluripotent cells. YES specifically phosphorylates YAP, a transcription factor inhibited by the Hippo pathway (see section below), which drives YAP-TEAD-dependent *Oct4* transcription and mESC pluripotency [192]. Accumulating evidence therefore suggests that c-SRC drives differentiation, whilst YES and HCK function to maintain pluripotency. However, lack of specific tools to study SFKs has hindered efforts to identify the molecular mechanisms underpinning their functional specificities in ESCs.

Hippo signalling connects cellular mechanics to the pluripotency machinery

A final key cellular mechanosensing system centres on the conserved Hippo kinase signalling pathway, which coordinates cell growth, proliferation, and fate in response to cell–cell contact and polarity. The core Hippo pathway consists of the MST1/2 (Hpo in *Drosophila*) and LATS1/2 (Wts) kinases, which form a cascade that phosphorylates YAP (Yki) and TAZ transcriptional cofactors. YAP phosphorylation promotes 14-3-3 binding and prevents YAP/TAZ accumulation in the nucleus [195,196]. Hippo activation therefore restricts YAP/TAZ-dependent transcription of genes required for cell growth and proliferation and is emerging as a major regulator of pluripotency [197].

Hippo signalling is activated by mechanical inputs from adherens junctions, tight junctions, apical–basal polarity complexes, and the actin cytoskeleton [198]. F-actin stabilization results in YAP/TAZ activation, whilst F-actin disruption drives Hippo activation and inactivation of YAP/TAZ [199]. G-protein-coupled receptors also respond to mechanical signals to modulate RHO family GTPase activation and actin dynamics [200]. In hESCs, the guanine nucleotide exchange factor AKAP-LBC activates RHOA signalling by modulating actin microfilament organization, and this is required to sustain the nuclear localisation of YAP/TAZ [201].

YAP and TEAD cofactors are highly expressed in ESCs [202] and, along with TAZ, are required for maintenance of mouse and human pluripotency [203–205]. Elevated YAP/TAZ activity also maintains and expands tissue-specific stem cell compartments [206–213]. Overexpression of YAP along with OCT4, SOX2, and KLF4 drives the reprogramming of mouse fibroblasts to pluripotency [205]. Conversely, activation of Hippo signalling is a barrier to reprogramming [214,215], although LATS2 appears to suppress reprogramming by inhibiting TAZ but not YAP [214]. Nevertheless, The YAP/TAZ/TEAD transcriptional module is a crucial determinant of pluripotency, and its function is directly opposed by the Hippo signalling pathway.

Mechanistically, YAP/TAZ/TEAD drives mESC pluripotency by directly inducing *Oct4* and *Nanog* expression [192]. However, YAP also patterns and integrates signals from other pluripotency signalling pathways. Indeed, YAP may promote naïve pluripotency in part by suppressing differentiation-inducing effects of GSK3 inhibition in hESCs [216]. As discussed previously, YAP phosphorylation by the SFK YES increases TEAD2 transcriptional activity at the *Oct4* and *Nanog* promoters in mESCs [192]. In addition, cross-talk between Hippo and TGF β /BMP signalling also promotes pluripotency. In hESCs, TAZ associates with SMAD2/3 to maintain nuclear localisation and potentiate *OCT4* and *NANOG* expression in response to TGF β [203]. Furthermore, Beyer *et al.* identified a regulatory complex composed of TAZ/YAP with TEADs with SMAD2/3 and OCT4 (termed TSO). TSO acts to suppress the expression of differentiation markers whilst supporting the expression of core pluripotency genes, thereby maintaining pluripotency [217]. Additionally, in mESCs, YAP has been shown to function in the BMP pathway via SMAD1-dependent recruitment to BMP-responsive enhancers to block neural differentiation [204]. Therefore, the Hippo signalling pathway and its YAP/TAZ/TEAD transcriptional module integrate diverse signals to couple the physical environment to the transcriptional regulation of pluripotency.

Perspectives

Biological function has been ascribed to a relatively small fraction of the kinome, and evidence suggests that understudied kinases and signalling pathways play critical roles in key biological processes. In this review, we explore newly uncovered pluripotency kinases and more established kinase signalling pathways with emerging roles in pluripotency regulation. A general theme is that environmental and cellular conditions modulate diverse kinase networks, which profoundly impact the expression and function of pluripotency factors. These molecular connections ensure that pluripotency is either maintained or dissolved depending on the cellular environment. Although much progress has been made, a more complete understanding of how cell cycle, DNA damage signalling, metabolism, stress, and mechanical factors modulate pluripotency is required to exploit these processes in pluripotent cell technologies. Furthermore, recent advances in our understanding of distinct pluripotent states require that we investigate kinase functions in pluripotent cells within the framework of the naïve-formative-primed pluripotency paradigm.

A significant future challenge will be to elucidate key mechanisms and substrates by which protein kinases control pluripotency. In this regard, phosphoproteomic studies and “functional kinomics” enable novel phosphorylation networks to be comprehensively mapped and interrogated. We propose that kinase signalling pathways identified using these unbiased approaches will elaborate the most exciting new molecular targets to be exploited in pluripotent stem cell applications.

Acknowledgements

The authors would like to apologise to those whose work could not be cited due to space limitations. C.A.C.W. is a recipient of an MRC PhD studentship, and G.M.F. is supported in part by a Medical Research Council New Investigator Award (MR/N000609/1) and a Tenovus Scotland research grant (T15/11).

Received 7 November 2016;
Received in revised form 21 April 2017;
Accepted 21 April 2017
Available online 26 April 2017

Keywords:

pluripotency;
phosphorylation;
kinase;
embryonic stem cells;
signalling networks

Abbreviations used:

ESC, embryonic stem cell; iPSC, induced pluripotent stem cell; mESC, mouse ESC; EpiSC, epiblast stem cell; hESC, human ESC; BMP, bone morphogenetic protein; LIF, leukemia inhibitory factor; FGF, fibroblast growth factor; CDK, cyclin-dependent kinase; JNK, c-Jun N-terminal kinase; DDR, DNA damage response; ATM, ataxia telangiectasia mutated; mTOR, mechanistic target of rapamycin; ULK1, UNC51-like kinase-1; AMPK, AMP-activated protein kinase; ROS, reactive oxygen species; NF, nuclear factor; ASK1, apoptosis signal-regulating kinase 1; TRX, thioredoxin; SS, shear stress; ER, endoplasmic reticulum; ECM, extracellular matrix; FAK, focal adhesion kinase; SFK, Src family kinase; TGF β , transforming growth factor beta; RAPTOR, regulatory associated protein of mTOR; RICTOR, rapamycin insensitive companion of mTOR.

References

- [1] C.E. Murry, G. Keller, Differentiation of embryonic stem cells to clinically relevant populations: lessons from embryonic development, *Cell* 132 (2008) 661–680.
- [2] M.J. Evans, M.H. Kaufman, Establishment in culture of pluripotential cells from mouse embryos, *Nature* 292 (1981) 154–156.
- [3] G.R. Martin, Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells, *Proc Natl Acad Sci U S A.* 78 (1981) 7634–7638.
- [4] J.A. Thomson, J. Itskovitz-Eldor, S.S. Shapiro, M.A. Waknitz, J.J. Swiergiel, V.S. Marshall, et al., Embryonic stem cell lines derived from human blastocysts, *Science* 282 (1998) 1145–1147.
- [5] K. Takahashi, K. Tanabe, M. Ohnuki, M. Narita, T. Ichisaka, K. Tomoda, et al., Induction of pluripotent stem cells from adult human fibroblasts by defined factors, *Cell* 131 (2007) 861–872.
- [6] K. Takahashi, S. Yamanaka, Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors, *Cell* 126 (2006) 663–676.
- [7] J. Yu, M.A. Vodyanik, K. Smuga-Otto, J. Antosiewicz-Bourget, J.L. Frane, S. Tian, et al., Induced pluripotent stem cell lines derived from human somatic cells, *Science* 318 (2007) 1917–1920.
- [8] T. Kalkan, N. Olova, M. Roode, C. Mulas, H.J. Lee, I. Nett, et al., Tracking the embryonic stem cell transition from ground state pluripotency, *Development* 144 (7) (2017) 1221–1234.
- [9] H. Chen, I. Aksoy, F. Gonnot, P. Osteil, M. Aubry, C. Hamela, et al., Reinforcement of STAT3 activity reprogrammes human embryonic stem cells to naïve-like pluripotency, *Nature Commun.* 6 (2015) 7095.
- [10] G. Duggal, S. Warriar, S. Ghimire, D. Broekaert, M. Van der Jeught, S. Lierman, et al., Alternative routes to induce naïve pluripotency in human embryonic stem cells, *Stem Cells.* 33 (2015) 2686–2698.
- [11] O. Gafni, L. Weinberger, A.A. Mansour, Y.S. Manor, E. Chomsky, D. Ben-Yosef, et al., Derivation of novel human ground state naïve pluripotent stem cells, *Nature* 504 (2013) 282–286.
- [12] G. Guo, F. von Meyenn, F. Santos, Y. Chen, W. Reik, P. Bertone, et al., Naïve pluripotent stem cells derived directly from isolated cells of the human inner cell mass, *Stem Cell Reports* 6 (2016) 437–446.
- [13] Y. Takashima, G. Guo, R. Loos, J. Nichols, G. Ficz, F. Krueger, et al., Resetting transcription factor control circuitry toward ground-state pluripotency in human, *Cell* 158 (2014) 1254–1269.
- [14] T.W. Theunissen, B.E. Powell, H. Wang, M. Mitalipova, D.A. Faddah, J. Reddy, et al., Systematic identification of culture conditions for induction and maintenance of naïve human pluripotency, *Cell Stem Cell* 15 (2014) 471–487.
- [15] M. Van der Jeught, J. Taelman, G. Duggal, S. Ghimire, S. Lierman, S.M. Chuva de Sousa Lopes, et al., Application of small molecules favoring naïve pluripotency during human embryonic stem cell derivation, *Cell Reprogram.* 17 (2015) 170–180.
- [16] C.B. Ware, A.M. Nelson, B. Mecham, J. Hesson, W. Zhou, E.C. Jonlin, et al., Derivation of naïve human embryonic stem cells, *Proc Natl Acad Sci U S A.* 111 (2014) 4484–4489.
- [17] T. Kalkan, A. Smith, Mapping the route from naïve pluripotency to lineage specification, *Philos Trans R Soc Lond B Biol Sci.* 369 (1657) (2014).
- [18] L. Weinberger, M. Ayyash, N. Novershtern, J.H. Hanna, Dynamic stem cell states: naïve to primed pluripotency in rodents and humans, *Nat Rev Mol Cell Biol.* 17 (2016) 155–169.
- [19] L.A. Boyer, T.I. Lee, M.F. Cole, S.E. Johnstone, S.S. Levine, J.P. Zucker, et al., Core transcriptional regulatory circuitry in human embryonic stem cells, *Cell* 122 (2005) 947–956.
- [20] K. Mitsui, Y. Tokuzawa, H. Itoh, K. Segawa, M. Murakami, K. Takahashi, et al., The homeoprotein Nanog is required for maintenance of pluripotency in mouse epiblast and ES cells, *Cell* 113 (2003) 631–642.
- [21] J. Nichols, B. Zevnik, K. Anastasiadis, H. Niwa, D. Klewe-Nebenius, I. Chambers, et al., Formation of pluripotent stem cells in the mammalian embryo depends on the POU transcription factor Oct4, *Cell* 95 (1998) 379–391.
- [22] C. Buecker, R. Srinivasan, Z. Wu, E. Calo, D. Acampora, T. Faial, et al., Reorganization of enhancer patterns in transition from naïve to primed pluripotency, *Cell Stem Cell* 14 (2014) 838–853.
- [23] H. Niwa, J. Miyazaki, A.G. Smith, Quantitative expression of Oct-3/4 defines differentiation, dedifferentiation or self-renewal of ES cells, *Nat Genet.* 24 (2000) 372–376.
- [24] J. Nichols, A. Smith, Naïve and primed pluripotent states, *Cell Stem Cell.* 4 (2009) 487–492.
- [25] G. Ficz, M.R. Branco, S. Seisenberger, F. Santos, F. Krueger, T.A. Hore, et al., Dynamic regulation of 5-hydroxymethylcytosine in mouse ES cells and during differentiation, *Nature* 473 (2011) 398–402.
- [26] P. Cohen, The origins of protein phosphorylation, *Nat Cell Biol.* 4 (2002) E127–E130.
- [27] Q.L. Ying, J. Wray, J. Nichols, L. Battle-Morera, B. Doble, J. Woodgett, et al., The ground state of embryonic stem cell self-renewal, *Nature* 453 (2008) 519–523.
- [28] J. Nichols, J. Silva, M. Roode, A. Smith, Suppression of Erk signalling promotes ground state pluripotency in the mouse embryo, *Development* 136 (2009) 3215–3222.
- [29] I.G. Brons, L.E. Smithers, M.W. Trotter, P. Rugg-Gunn, B. Sun, S.M. Chuva de Sousa Lopes, et al., Derivation of pluripotent epiblast stem cells from mammalian embryos, *Nature* 448 (2007) 191–195.

- [30] P.J. Tesar, J.G. Chenoweth, F.A. Brook, T.J. Davies, E.P. Evans, D.L. Mack, et al., New cell lines from mouse epiblast share defining features with human embryonic stem cells, *Nature* 448 (2007) 196–199.
- [31] L. Vallier, M. Alexander, R.A. Pedersen, Activin/Nodal and FGF pathways cooperate to maintain pluripotency of human embryonic stem cells, *J Cell Sci.* 118 (2005) 4495–4509.
- [32] G. Manning, D.B. Whyte, R. Martinez, T. Hunter, S. Sudarsanam, The protein kinase complement of the human genome, *Science* 298 (2002) 1912–1934.
- [33] O. Fedorov, S. Muller, S. Knapp, The (un)targeted cancer kinome, *Nat Chem Biol.* 6 (2010) 166–169.
- [34] I. Singec, A.M. Crain, J. Hou, B.T. Tobe, M. Talantova, A.A. Winquist, et al., Quantitative analysis of human pluripotency and neural specification by in-depth (phospho)proteomic profiling, *Stem Cell Reports* 7 (2016) 527–542.
- [35] V.M. Ding, P.J. Boersema, L.Y. Foong, C. Preisinger, G. Koh, S. Natarajan, et al., Tyrosine phosphorylation profiling in FGF-2 stimulated human embryonic stem cells, *PLoS One* 6 (2011) e17538.
- [36] A.D. Zoumaro-Djajoon, V. Ding, L.Y. Foong, A. Choo, A.J. Heck, J. Munoz, Investigating the role of FGF-2 in stem cell maintenance by global phosphoproteomics profiling, *Proteomics* 11 (2011) 3962–3971.
- [37] L.M. Brill, W. Xiong, K.B. Lee, S.B. Ficarro, A. Crain, Y. Xu, et al., Phosphoproteomic analysis of human embryonic stem cells, *Cell Stem Cell* 5 (2009) 204–213.
- [38] D.H. Phanstiel, J. Brumbaugh, C.D. Wenger, S. Tian, M.D. Probasco, D.J. Bailey, et al., Proteomic and phosphoproteomic comparison of human ES and iPS cells, *Nat Methods* 8 (2011) 821–827.
- [39] D. Van Hoof, J. Munoz, S.R. Braam, M.W. Pinkse, R. Linding, A.J. Heck, et al., Phosphorylation dynamics during early differentiation of human embryonic stem cells, *Cell Stem Cell* 5 (2009) 214–226.
- [40] K.T. Rigbolt, T.A. Prokhorova, V. Akimov, J. Henningsen, P.T. Johansen, I. Kratchmarova, et al., System-wide temporal characterization of the proteome and phosphoproteome of human embryonic stem cell differentiation, *Science Signaling* 4 (2011) rs3.
- [41] D.F. Lee, J. Su, Y.S. Ang, X. Carvajal-Vergara, S. Mulero-Navarro, C.F. Pereira, et al., Regulation of embryonic and induced pluripotency by aurora kinase-p53 signaling, *Cell Stem Cell* 11 (2012) 179–194.
- [42] K. Sakurai, I. Talukdar, V.S. Patil, J. Dang, Z. Li, K.Y. Chang, et al., Kinome-wide functional analysis highlights the role of cytoskeletal remodeling in somatic cell reprogramming, *Cell Stem Cell* 14 (2014) 523–534.
- [43] C.A.C. Williams, R. Fernandez-Alonso, J. Wang, R. Toth, N.S. Gray, G.M. Findlay, Erk5 is a key regulator of naïve-primed transition and embryonic stem cell identity, *Cell Reports* 16 (2016) 1820–1828.
- [44] M. Morikawa, D. Koinuma, A. Mizutani, N. Kawasaki, K. Holmborn, A. Sundqvist, et al., BMP sustains embryonic stem cell self-renewal through distinct functions of different kruppel-like factors, *Stem Cell Reports* 6 (2016) 64–73.
- [45] Y. Kato, BMK1/ERK5 regulates serum-induced early gene expression through transcription factor MEF2C, *The EMBO Journal.* 16 (1997) 7054–7066.
- [46] K. Sunadome, T. Yamamoto, M. Ebisuya, K. Kondoh, A. Sehara-Fujisawa, E. Nishida, ERK5 regulates muscle cell fusion through Klf transcription factors, *Developmental Cell* 20 (2011) 192–205.
- [47] C. Yan, H. Luo, J.D. Lee, Abe Ji, B.C. Berk, Molecular cloning of mouse ERK5/BMK1 splice variants and characterization of ERK5 functional domains, *J. Biol. Chem.* 276 (2001) 10,870–10,878.
- [48] K.M. Parmar, H.B. Larman, G. Dai, Y. Zhang, E.T. Wang, S.N. Moorthy, et al., Integration of flow-dependent endothelial phenotypes by Kruppel-like factor 2, *Journal of Clinical Investigation* 116 (2006) 49–58.
- [49] K. Kesavan, K. Lobel-Rice, W. Sun, R. Lapadat, S. Webb, G.L. Johnson, et al., MEKK2 regulates the coordinate activation of ERK5 and JNK in response to FGF-2 in fibroblasts, *J Cell Physiol.* 199 (2004) 140–148.
- [50] Y. Nakaoka, K. Nishida, Y. Fujio, M. Izumi, K. Terai, Y. Oshima, et al., Activation of gp130 transduces hypertrophic signal through interaction of scaffolding/docking protein Gab1 with tyrosine phosphatase SHP2 in cardiomyocytes, *Circ Res.* 93 (2003) 221–229.
- [51] J.M. Elkins, V. Fedele, M. Szklarz, K.R. Abdul Azeez, E. Salah, J. Mikolajczyk, et al., Comprehensive characterization of the published kinase inhibitor set, *Nat Biotechnol.* 34 (2016) 95–103.
- [52] E. Jacoby, G. Tresadern, S. Bembek, B. Wroblowski, C. Buyck, J.M. Neefs, et al., Extending kinome coverage by analysis of kinase inhibitor broad profiling data, *Drug Discov Today* 20 (2015) 652–658.
- [53] D. Coronado, M. Godet, P.Y. Bourillot, Y. Tapponnier, A. Bernat, M. Petit, et al., A short G1 phase is an intrinsic determinant of naïve embryonic stem cell pluripotency, *Stem Cell Res.* 10 (2013) 118–131.
- [54] A. Calder, I. Roth-Albin, S. Bhatia, C. Pilquill, J.H. Lee, M. Bhatia, et al., Lengthened G1 phase indicates differentiation status in human embryonic stem cells, *Stem Cells and Dev.* 22 (2013) 279–295.
- [55] W.W. Zhang, X.J. Zhang, H.X. Liu, J. Chen, Y.H. Ren, D.G. Huang, et al., Cdk1 is required for the self-renewal of mouse embryonic stem cells, *J. Cell. Biochem.* 112 (2011) 942–948.
- [56] L. Li, J. Wang, J. Hou, Z. Wu, Y. Zhuang, M. Lu, et al., Cdk1 interplays with Oct4 to repress differentiation of embryonic stem cells into trophoblast, *FEBS Letters* 586 (2012) 4100–4107.
- [57] J. Ouyang, W. Yu, J. Liu, N. Zhang, L. Florens, J. Chen, et al., Cyclin-dependent kinase-mediated Sox2 phosphorylation enhances the ability of Sox2 to establish the pluripotent state, *J. Biol. Chem.* 290 (2015) 22,782–22,794.
- [58] J. Brumbaugh, J.D. Russell, P. Yu, M.S. Westphall, J.J. Coon, J.A. Thomson, NANOG is multiply phosphorylated and directly modified by ERK2 and CDK1 *in vitro*, *Stem Cell Reports* 2 (2014) 18–25.
- [59] S. Pauklin, P. Madrigal, A. Bertero, L. Vallier, Initiation of stem cell differentiation involves cell cycle-dependent regulation of developmental genes by cyclin D, *Genes & Dev.* 30 (2016) 421–433.
- [60] A.M. Singh, Y. Sun, L. Li, W. Zhang, T. Wu, S. Zhao, et al., Cell-cycle control of bivalent epigenetic domains regulates the exit from pluripotency, *Stem Cell Reports* 5 (2015) 323–336.
- [61] C. Lange, F. Calegari, Cdks and cyclins link G1 length and differentiation of embryonic, neural and hematopoietic stem cells, *Cell Cycle* 9 (2010) 1893–1900.
- [62] K.A. Gonzales, H. Liang, Y.S. Lim, Y.S. Chan, J.C. Yeo, C.P. Tan, et al., Deterministic restriction on pluripotent state dissolution by cell-cycle pathways, *Cell* 162 (2015) 564–579.

- [63] S. Pauklin, L. Vallier, The cell-cycle state of stem cells determines cell fate propensity, *Cell* 155 (2013) 135–147.
- [64] J. Shin, T.W. Kim, H. Kim, H.J. Kim, M.Y. Suh, S. Lee, et al., Aurkb/PP1-mediated resetting of Oct4 during the cell cycle determines the identity of embryonic stem cells, *eLife* 5 (2016) e10877.
- [65] I.A. Chuykin, M.S. Lianguzova, T.V. Pospelova, V.A. Pospelov, Activation of DNA damage response signaling in mouse embryonic stem cells, *Cell Cycle* 7 (2008) 2922–2928.
- [66] A.K. Ahuja, K. Jodkowska, F. Teloni, A.H. Bizard, R. Zellweger, R. Herrador, et al., A short G1 phase imposes constitutive replication stress and fork remodelling in mouse embryonic stem cells, *Nature Commun.* 7 (2016) 10,660.
- [67] S. Ruiz, A.J. Lopez-Contreras, M. Gabut, R.M. Marion, P. Gutierrez-Martinez, S. Bua, et al., Limiting replication stress during somatic cell reprogramming reduces genomic instability in induced pluripotent stem cells, *Nature Commun.* 6 (2015) 8036.
- [68] T. Kinoshita, G. Nagamatsu, T. Kosaka, K. Takubo, A. Hotta, J. Ellis, et al., Ataxia-telangiectasia mutated (ATM) deficiency decreases reprogramming efficiency and leads to genomic instability in iPS cells, *Biochemical and Biophysical Res. Commun.* 407 (2011) 321–326.
- [69] J. Xiong, D. Todorova, N.Y. Su, J. Kim, P.J. Lee, Z. Shen, et al., Stemness factor Sall4 is required for DNA damage response in embryonic stem cells, *J. Cell Biol.* 208 (2015) 513–520.
- [70] B. Zhao, W.D. Zhang, Y.L. Duan, Y.Q. Lu, Y.X. Cun, C.H. Li, et al., Filia is an ESC-specific regulator of DNA damage response and safeguards genomic stability, *Cell Stem Cell* 16 (2015) 684–698.
- [71] E.J. Brown, M.W. Albers, T.B. Shin, K. Ichikawa, C.T. Keith, W.S. Lane, et al., A mammalian protein targeted by G1-arresting rapamycin-receptor complex, *Nature* 369 (1994) 756–758.
- [72] K. Hara, Y. Maruki, X. Long, Yoshino Ki, N. Oshiro, S. Hidayat, et al., Raptor, a binding partner of target of rapamycin (TOR), mediates TOR action, *Cell* 110 (2002) 177–189.
- [73] D.H. Kim, D.D. Sarbassov, S.M. Ali, J.E. King, R.R. Latek, H. Erdjument-Bromage, et al., mTOR interacts with raptor to form a nutrient-sensitive complex that signals to the cell growth machinery, *Cell* 110 (2002) 163–175.
- [74] D.S. Dos, S.M. Ali, D.H. Kim, D.A. Guertin, R.R. Latek, H. Erdjument-Bromage, et al., Rictor, a novel binding partner of mTOR, defines a rapamycin-insensitive and raptor-independent pathway that regulates the cytoskeleton, *Current Biol.* 14 (2004) 1296–1302.
- [75] G. Thomas, The S6 kinase signaling pathway in the control of development and growth, *Biol Res.* 35 (2002) 305–313.
- [76] X.M. Ma, J. Blenis, Molecular mechanisms of mTOR-mediated translational control, *Nature Reviews Mol. Cell Biol.* 10 (2009) 307–318.
- [77] Q. Yang, K.-L. Guan, Expanding mTOR signaling, *Cell Res.* 17 (2007) 666–681.
- [78] R. Zoncu, A. Efeyan, D.M. Sabatini, mTOR: from growth signal integration to cancer, diabetes and ageing, *Nature Reviews Mol. Cell Biol.* 12 (2011) 21–35.
- [79] Gangloff Y-G, M. Mueller, S.G. Dann, P. Svoboda, M. Sticker, J.-F. Spetz, et al., Disruption of the mouse mTOR gene leads to early postimplantation lethality and prohibits embryonic stem cell development disruption of the mouse mTOR gene leads to early postimplantation lethality and prohibits embryonic stem cell development, *Development* 131 (2004) 9508–9516.
- [80] M. Murakami, T. Ichisaka, M. Maeda, N. Oshiro, K. Hara, F. Edenhofer, et al., mTOR is essential for growth and proliferation in early mouse embryos and embryonic stem cells, *Mol Cell Biol.* 24 (2004) 6710–6718.
- [81] M.Y. Cherepkova, G.S. Sineva, V.A. Pospelov, Leukemia inhibitory factor (LIF) withdrawal activates mTOR signaling pathway in mouse embryonic stem cells through the MEK/ERK/TSC2 pathway, *Nature Publishing Group* (2016) 1–10.
- [82] K.-W. Lee, J.-Y. Yook, M.-Y. Son, M.-J. Kim, D.-B. Koo, Y.-M. Han, et al., Rapamycin promotes the osteoblastic differentiation of human embryonic stem cells by blocking the mTOR pathway and stimulating the BMP/Smad pathway, *Stem Cells and Dev.* 19 (2010) 557–568.
- [83] E.J.P. Nazareth, N. Rahman, T. Yin, P.W. Zandstra, A multi-lineage screen reveals mTORC1 inhibition enhances human pluripotent stem cell mesoderm and blood progenitor production, *Stem Cell Reports* 6 (2016) 679–691.
- [84] P. Sampath, D.K. Pritchard, L. Pabon, H. Reinecke, S.M. Schwartz, D.R. Morris, et al., A hierarchical network controls protein translation during murine embryonic stem cell self-renewal and differentiation, *Cell Stem Cell* 2 (2008) 448–460.
- [85] J. Zhou, P. Su, L. Wang, J. Chen, M. Zimmermann, O. Genbacev, et al., mTOR supports long-term self-renewal and suppresses mesoderm and endoderm activities of human embryonic stem cells, *Proceedings of the National Academy of Sciences of the United States of America.* 106 (2009) 7840–7845.
- [86] Easley Ca, A. Ben-Yehudah, C.J. Redinger, S.L. Oliver, S.T. Varum, V.M. Eisinger, et al., mTOR-mediated activation of p70 S6K induces differentiation of pluripotent human embryonic stem cells, *Cellular Reprogramming (Formerly "Cloning and Stem Cells")* 12 (2010) 263–273.
- [87] P. Agrawal, J. Reynolds, S. Chew, D.A. Lamba, R.E. Hughes, DEPTOR is a stemness factor that regulates pluripotency of embryonic stem cells, *J. Biol. Chem.* 289 (2014) 31,818–31,826.
- [88] S. Wang, P. Xia, B. Ye, G. Huang, J. Liu, Z. Fan, Transient activation of autophagy via Sox2-mediated suppression of mTOR is an important early step in reprogramming to pluripotency, *Cell Stem Cell* 13 (2013) 617–625.
- [89] A. Bulut-Karslioglu, S. Biechele, H. Jin, T.A. Macrae, M. Hejna, M. Gertsenstein, et al., Inhibition of mTOR induces a paused pluripotent state, *Nature* 540 (2016) 119–123.
- [90] K. Hara, K. Yonezawa, Q.P. Weng, M.T. Kozlowski, C. Belham, J. Avruch, Amino acid sufficiency and mTOR regulate p70 S6 kinase and eIF-4E BP1 through a common effector mechanism, *J. Biol. Chem.* 273 (1998) 14,484–14,494.
- [91] J. Wang, P. Alexander, L. Wu, R. Hammer, O. Cleaver, S.L. McKnight, Dependence of mouse embryonic stem cells on threonine catabolism, *Science* 325 (2009) 435–439.
- [92] J.M. Ryu, H.J. Han, L-threonine regulates G1/S phase transition of mouse embryonic stem cells via PI3K/Akt, MAPKs, and mTORC pathways, *J. Biol. Chem.* 286 (2011) 23,667–23,678.
- [93] Y. Sancak, T.R. Peterson, Y.D. Shaul, R.A. Lindquist, C.C. Thoreen, L. Bar-Peled, et al., The Rag GTPases bind raptor and mediate amino acid signaling to mTORC1, *Science* 320 (2008) 1496–1501.
- [94] W. Zhou, M. Choi, D. Margineantu, L. Margaretha, J. Hesson, C. Cavanaugh, et al., HIF1 α induced switch from bivalent to exclusively glycolytic metabolism during ESC-to-EpiSC/hESC transition, *EMBO J.* 31 (2012) 2103–2116.

- [95] S. Varum, A.S. Rodrigues, M.B. Moura, O. Momcilovic, C.A. Easley Iv, J. Ramalho-Santos, et al., Energy metabolism in human pluripotent stem cells and their differentiated counterparts, *PLoS ONE* 6 (2011) e20914.
- [96] C.D.L. Folmes, T.J. Nelson, A. Martinez-Fernandez, D.K. Arrell, J.Z. Lindor, P.P. Dzeja, et al., Somatic oxidative bioenergetics transitions into pluripotency-dependent glycolysis to facilitate nuclear reprogramming, *Cell Metabolism* 14 (2011) 264–271.
- [97] L.A. Sena, N.S. Chandel, Physiological roles of mitochondrial reactive oxygen species, *48* (2) (2012) 158–167.
- [98] S.T. Suhr, E.A. Chang, J. Tjong, N. Alcasid, G.A. Perkins, M.D. Goissis, et al., Mitochondrial rejuvenation after induced pluripotency, *PLoS ONE* 5 (2010) e14095.
- [99] J. Zhang, E. Nuebel, G.Q. Daley, C.M. Koehler, M.A. Teitell, Metabolic regulation in pluripotent stem cells during reprogramming and self-renewal, *Cell Stem Cell* 11 (2012) 589–595.
- [100] W. Gu, X. Gaeta, A. Sahakyan, B. Chan Alanna, S. Hong Candice, R. Kim, et al., Glycolytic metabolism plays a functional role in regulating human pluripotent stem cell state, *Cell Stem Cell* (2016) 1–15.
- [101] K. Düvel, J.L. Yecies, S. Menon, P. Raman, A.I. Lipovsky, A.L. Souza, et al., Activation of a metabolic gene regulatory network downstream of mTOR complex 1, *Mol. Cell* 39 (2010) 171–183.
- [102] J.G. Ryall, T. Cliff, S. Dalton, V. Sartorelli, Metabolic reprogramming of stem cell epigenetics, *Cell Stem Cell* 17 (6) (2015) 651–662.
- [103] M.Y. Lee, H.J. Han, Galectin-1 upregulates glucose transporter-1 expression level via protein kinase C, phosphoinositol-3 kinase, and mammalian target of rapamycin pathways in mouse embryonic stem cells, *The International J. Biochem. & Cell Biol.* 40 (2008) 2421–2430.
- [104] K. Liu, Q. Zhao, P. Liu, J. Cao, J. Gong, C. Wang, et al., ATG3-dependent autophagy mediates mitochondrial homeostasis in pluripotency acquirement and maintenance, *Autophagy* (2016) 1–9.
- [105] M. Kundu, C.B. Thompson, Autophagy: basic principles and relevance to disease, *Annual Review of Pathology: Mechanisms of Disease.* 3 (2008) 427–455.
- [106] J.M. Facucho-Oliveira, J. Alderson, E.C. Spikings, S. Egginton, J.C. St John, Mitochondrial DNA replication during differentiation of murine embryonic stem cells, *J. Cell Sci.* 120 (2007) 4025–4034.
- [107] J.C. St. John, J. Ramalho-Santos, H.L. Gray, P. Petrosko, V.Y. Rawe, C.S. Navara, et al., The expression of mitochondrial DNA transcription factors during early cardiomyocyte, *Cloning and Stem Cells* 7 (2005) 141–153.
- [108] J. Kaur, J. Debnath, Autophagy at the crossroads of catabolism and anabolism, *Nature Reviews Mol. Cell Biol.* 16 (2015) 461–472.
- [109] D.F. Egan, D.B. Shackelford, M.M. Mihaylova, S. Gelino, R.A. Kohnz, W. Mair, et al., Phosphorylation of ULK1 (hATG1) by AMP-activated protein kinase connects energy sensing to mitophagy, *Science* 331 (2011) 456–461.
- [110] J. Kim, M. Kundu, B. Viollet, K.-L. Guan, AMPK and mTOR regulate autophagy through direct phosphorylation of Ulk1, *Nature Cell Biol.* 13 (2011) 132–141.
- [111] S. Honda, S. Arakawa, Y. Nishida, H. Yamaguchi, E. Ishii, S. Shimizu, Ulk1-mediated Atg5-independent macroautophagy mediates elimination of mitochondria from embryonic reticulocytes, *Nature Commun* 5 (2014) 4004.
- [112] Y. Nishida, S. Arakawa, K. Fujitani, H. Yamaguchi, T. Mizuta, T. Kanaseki, et al., Discovery of Atg5/Atg7-independent alternative macroautophagy, *Nature* 461 (2009) 654–658.
- [113] T.-C. Kuo, C.-T. Chen, D. Baron, T.T. Onder, S. Loewer, S. Almeida, et al., Midbody accumulation through evasion of autophagy contributes to cellular reprogramming and tumorigenicity, *Nature Cell Biol.* 13 (2011) 1214–1223.
- [114] S. Varum, O. Momcilović, C. Castro, A. Ben-Yehudah, J. Ramalho-Santos, C.S. Navara, Enhancement of human embryonic stem cell pluripotency through inhibition of the mitochondrial respiratory chain, *Stem Cell Res.* 3 (2009) 142–156.
- [115] G. Saretzki, Stress defense in murine embryonic stem cells is superior to that of various differentiated murine cells, *Stem Cells* 22 (2004) 962–971.
- [116] G. Saretzki, T. Walter, S. Atkinson, J.F. Passos, B. Bareth, W.N. Keith, et al., Downregulation of multiple stress defense mechanisms during differentiation of human embryonic stem cells, *Stem Cells* 26 (2008) 455–464.
- [117] L.A. Tibbles, J.R. Woodgett, The stress-activated protein kinase pathways, *Cell Mol Life Sci.* 55 (1999) 1230–1254.
- [118] K.J. Davies, Oxidative stress, antioxidant defenses, and damage removal, repair, and replacement systems, *IUBMB Life* 50 (2000) 279–289.
- [119] S. Varum, O. Momčilović, C. Castro, A. Ben-Yehudah, J. Ramalho-Santos, C.S. Navara, Enhancement of human embryonic stem cell pluripotency through inhibition of the mitochondrial respiratory chain, *Stem Cell Res.* 3 (2009) 142–156.
- [120] T. Ezashi, P. Das, R.M. Roberts, Low O₂ tensions and the prevention of differentiation of hES cells, *Proceedings of the National Academy of Sciences of the United States of America.* 102 (2005) 4783–4788.
- [121] N.R. Forsyth, A. Musio, P. Vezzoni, Simpson AHRW, B.S. Noble, J. McWhir, Physiologic oxygen enhances human embryonic stem cell clonal recovery and reduces chromosomal abnormalities, *Cloning and Stem Cells.* 8 (2006) 16–23.
- [122] Y.-L.L. Guo, S. Chakraborty, S.S. Rajan, R. Wang, F. Huang, Effects of oxidative stress on mouse embryonic stem cell proliferation, apoptosis, senescence, and self-renewal, *Stem Cells and Dev.* 19 (2010) 1321–1331.
- [123] M. Schmelter, B. Ateghang, S. Helmig, M. Wartenberg, H. Sauer, Embryonic stem cells utilize reactive oxygen species as transducers of mechanical strain-induced cardiovascular differentiation, *FASEB J.: Official Publication of the Federation of American Societies for Experimental Biology.* 20 (2006) 1182–1184.
- [124] Q. Xiao, Z. Luo, A.E. Pepe, A. Margariti, L. Zeng, Q. Xu, Embryonic stem cell differentiation into smooth muscle cells is mediated by Nox4-produced H₂O₂, *American J. Physiol. Cell Physiol.* 296 (2009) 711–723.
- [125] T. Finkel, N.J. Holbrook, Oxidants, oxidative stress and the biology of ageing, *Nature* 408 (2000) 239–247.
- [126] C.-C. Hsieh, J. Papaconstantinou, Thioredoxin–ASK1 complex levels regulate ROS-mediated p38 MAPK pathway activity in livers of aged and long-lived Snell dwarf mice, *FASEB J.* 20 (2006) 259–268.
- [127] M. Saitoh, H. Nishitoh, M. Fujii, K. Takeda, K. Tobiume, Y. Sawada, et al., Mammalian thioredoxin is a direct inhibitor of apoptosis signal-regulating kinase (ASK) 1, *EMBO J.* 17 (1998) 2596–2606.
- [128] F. Mercurio, Manning aM. NF-kappaB as a primary regulator of the stress response, *Oncogene* 18 (1999) 6163–6171.

- [129] Y.-E. Kim, H.-B. Kang, J.-A. Park, K.-H. Nam, H.-J. Kwon, Y. Lee, Upregulation of NF- κ B upon differentiation of mouse embryonic stem cells, *BMB Reports* 41 (2008) 705–709.
- [130] J. Torres, F.M. Watt, Nanog maintains pluripotency of mouse embryonic stem cells by inhibiting NF κ B and cooperating with Stat3, *Nature Cell Biol.* 10 (2008) 194–201.
- [131] C. Yang, S.P. Atkinson, F. Vilella, M. Lloret, L. Armstrong, D.A. Mann, et al., Opposing putative roles for canonical and noncanonical NF κ B signaling on the survival, proliferation, and differentiation potential of human embryonic stem cells, *Stem Cells* 28 (2010) 1970–1980.
- [132] P. Lüningschrör, B. Kaltschmidt, C. Kaltschmidt, Knockdown of IKK1/2 promotes differentiation of mouse embryonic stem cells into neuroectoderm at the expense of mesoderm, *Stem Cell Reviews and Reports.* 8 (2012) 1098–1108.
- [133] T. Burdon, C. Stracey, I. Chambers, J. Nichols, A. Smith, Suppression of SHP-2 and ERK signalling promotes self-renewal of mouse embryonic stem cells, *Dev Biol.* 210 (1999) 30–43.
- [134] X. Qi, T.G. Li, J. Hao, J. Hu, J. Wang, H. Simmons, et al., BMP4 supports self-renewal of embryonic stem cells by inhibiting mitogen-activated protein kinase pathways, *Proc Natl Acad Sci U S A.* 101 (2004) 6027–6032.
- [135] P. Xu, K. Yoshioka, D. Yoshimura, Y. Tominaga, T. Nishioka, M. Ito, et al., *In vitro* development of mouse embryonic stem cells lacking JNK/stress-activated protein kinase-associated protein 1 (JSAP1) scaffold protein revealed its requirement during early embryonic neurogenesis, *J. Biol. Chem.* 278 (2003) 48,422–48,433.
- [136] D.R. Alessi, J. Zhang, A. Khanna, T. Hochdorfer, Y. Shang, K.T. Kahle, The WNK-SPAK/OSR1 pathway: master regulator of cation-chloride cotransporters, *Science Signaling* 7 (2014) re3.
- [137] C. Richardson, D.R. Alessi, The regulation of salt transport and blood pressure by the WNK-SPAK/OSR1 signalling pathway, *J Cell Sci.* 121 (2008) 3293–3304.
- [138] X. Xu, Q. Wang, Y. Long, R. Zhang, X. Wei, M. Xing, et al., Stress-mediated p38 activation promotes somatic cell reprogramming, *Cell Res.* 23 (2012) 131–141.
- [139] K. Takahashi, K. Tanabe, M. Ohnuki, M. Narita, T. Ichisaka, K. Tomoda, et al., Induction of pluripotent stem cells from adult human fibroblasts by defined factors, *Cell* 131 (2007) 861–872.
- [140] J.P. Saxe, A. Tomilin, H.R. Schöler, K. Plath, J. Huang, Post-translational regulation of Oct4 transcriptional activity, *PLoS ONE* 4 (2009) e4467–e4477.
- [141] C.R. Amura, L. Marek, R.A. Winn, L.E. Heasley, Inhibited neurogenesis in JNK1-deficient embryonic stem cells, *Mol. and Cellular Biol.* 25 (2005) 10,791–10,802.
- [142] K. Yao, M.O. Ki, H. Chen, Y.Y. Cho, S.H. Kim, D.H. Yu, et al., JNK1 and 2 play a negative role in reprogramming to pluripotent stem cells by suppressing Klf4 activity, *Stem Cell Res.* 12 (2014) 139–152.
- [143] R.P. Franke, M. Gräfe, H. Schnittler, D. Seiffge, C. Mittermayer, D. Drenckhahn, Induction of human vascular endothelial stress fibres by fluid shear stress, *Nature* 307 (1984) 648–649.
- [144] Y. Xie, shear stress induces preimplantation embryo death that is delayed by the zona pellucida and associated with stress-activated protein kinase-mediated apoptosis, *Biol. of Reproduction.* 75 (2006) 45–55.
- [145] H.B. Croxatto, Physiology of gamete and embryo transport through the fallopian tube, *Reprod Biomed Online.* 4 (2002) 160–169.
- [146] T. Gareau, G.G. Lara, R.D. Shepherd, R. Krawetz, D.E. Rancourt, K.D. Rinker, et al., Shear stress influences the pluripotency of murine embryonic stem cells in stirred suspension bioreactors, *Journal of Tissue Engineering and Regenerative Medicine.* 8 (2014) 268–278.
- [147] R.P. Wolfe, J. Leleux, R.M. Nerem, T. Ahsan, Effects of shear stress on germ lineage specification of embryonic stem cells, *Integrative Biol* 4 (2012) 1263.
- [148] S. Jalali, M.A. del Pozo, K.-D. Chen, H. Miao, Y.-S. Li, M.A. Schwartz, et al., Integrin-mediated mechanotransduction requires its dynamic interaction with specific extracellular matrix (ECM) ligands, *Proceedings of the National Academy of Sciences.* 98 (2001) 1042–1046.
- [149] M.J. Kuchan, H. Jo, J.A. Frangos, Role of G proteins in shear stress-mediated nitric oxide production by endothelial cells, *The American J. Physiol.* 267 (1994) C753–C758.
- [150] E. Tzima, M. Irani-Tehrani, W.B. Kiosses, E. DeJana, Schultz Da, B. Engelhardt, et al., A mechanosensory complex that mediates the endothelial cell response to fluid shear stress, *Nature* 437 (2005) 426–431.
- [151] Y. Wang, H. Miao, S. Li, K.-D. Chen, Y.-S. Li, S. Yuan, et al., Interplay between integrins and FLK-1 in shear stress-induced signaling, *American J. Physiol. Cell Physiol.* 283 (2002) C1540–C7.
- [152] K. Yamamoto, T. Sokabe, T. Matsumoto, K. Yoshimura, M. Shibata, N. Ohura, et al., Impaired flow-dependent control of vascular tone and remodeling in P2X4-deficient mice, *Nat Med.* 12 (2006) 133–137.
- [153] A. Young, W. Wu, W. Sun, H.B. Larman, N. Wang, Y.S. Li, et al., Flow activation of AMP-activated protein kinase in vascular endothelium leads to krüppel-like factor 2 expression, *Arteriosclerosis, Thrombosis, and Vascular Biol.* 29 (2009) 1902–1908.
- [154] M. Kim, S. Kim, J.H. Lim, C. Lee, H.C. Choi, C.H. Woo, Laminar flow activation of ERK5 protein in vascular endothelium leads to atheroprotective effect via NF-E2-related factor 2 (Nrf2) activation, *J. Biol. Chem.* 287 (2012) 40,722–40,731.
- [155] I. Tabas, D. Ron, Integrating the mechanisms of apoptosis induced by endoplasmic reticulum stress, *Nature Cell Biol.* 13 (2011) 184–190.
- [156] K. Kratochvilova, L. Moran, S. Padourova, S. Stejskal, L. Tesarova, P. Simara, et al., The role of the endoplasmic reticulum stress in stemness, pluripotency and development, *Eur J Cell Biol.* 95 (2016) 115–123.
- [157] G. Chen, X. Xu, L. Zhang, Y. Fu, M. Wang, H. Gu, et al., Blocking autocrine VEGF signaling by sunitinib, an anti-cancer drug, promotes embryonic stem cell self-renewal and somatic cell reprogramming, *Cell Res.* 24 (2014) 1121–1136.
- [158] K. Kratochvilová, L. Moráň, S. Paďourová, S. Stejskal, L. Tesařová, P. Šimara, et al., The role of the endoplasmic reticulum stress in stemness, pluripotency and development, *European J. Cell Biol.* 95 (2016) 115–123.
- [159] L. Liu, C. Liu, Y. Zhong, A. Apostolou, S. Fang, ER stress response during the differentiation of H9 cells induced by retinoic acid, *Biochemical and Biophysical Res. Commun.* 417 (2012) 738–743.
- [160] W.L. Murphy, T.C. McDevitt, A.J. Engler, Materials as stem cell regulators, *Nature Materials.* 13 (2014) 547–557.

- [161] P. Hulpiau, F. van Roy, Molecular evolution of the cadherin superfamily, *The International J. Biochem. & Cell Biol.* 41 (2009) 349–369.
- [162] D. Li, J. Zhou, L. Wang, M.E. Shin, P. Su, X. Lei, et al., Integrated biochemical and mechanical signals regulate multifaceted human embryonic stem cell functions, *J. Cell Biol.* 191 (2010) 631–644.
- [163] H.L. Spencer, A.M. Eastham, C.L. Merry, T.D. Southgate, F. Perez-Campo, F. Soncin, et al., E-cadherin inhibits cell surface localization of the pro-migratory 5T4 oncofetal antigen in mouse embryonic stem cells, *Mol. Biol. Cell* 18 (2007) 2838–2851.
- [164] I. del Valle, S. Rudloff, A. Carles, Y. Li, E. Liszewska, R. Vogt, et al., E-cadherin is required for the proper activation of the Lifr/Gp130 signaling pathway in mouse embryonic stem cells, *Development* 140 (2013) 1684–1692.
- [165] F. Soncin, L. Mohamet, D. Eckardt, S. Ritson, A.M. Eastham, N. Bobola, et al., Abrogation of E-cadherin-mediated cell-cell contact in mouse embryonic stem cells results in reversible LIF-independent self-renewal, *Stem Cells* 27 (2009) 2069–2080.
- [166] F. Soncin, C.M. Ward, The function of e-cadherin in stem cell pluripotency and self-renewal, *Genes* 2 (2011) 229–259.
- [167] K. Hawkins, L. Mohamet, S. Ritson, C.L. Merry, C.M. Ward, E-cadherin and, in its absence, N-cadherin promotes Nanog expression in mouse embryonic stem cells via STAT3 phosphorylation, *Stem Cells* 30 (2012) 1842–1851.
- [168] J.L. Yori, E. Johnson, G. Zhou, M.K. Jain, R.A. Keri, Kruppel-like factor 4 inhibits epithelial-to-mesenchymal transition through regulation of E-cadherin gene expression, *J. Biol. Chem.* 285 (2010) 16,854–16,863.
- [169] T. Takehara, T. Teramura, Y. Onodera, J. Frampton, K. Fukuda, Cdh2 stabilizes FGFR1 and contributes to primed-state pluripotency in mouse epiblast stem cells, *Sci Rep.* 5 (2015) 14,722.
- [170] T.S. Huang, L. Li, L. Moalim-Nour, D. Jia, J. Bai, Z. Yao, et al., A Regulatory network involving beta-catenin, e-cadherin, PI3k/Akt, and slug balances self-renewal and differentiation of human pluripotent stem cells in response to Wnt Signaling, *Stem Cells* 33 (2015) 1419–1433.
- [171] R.O. Hynes, Integrins: bidirectional, allosteric signaling machines, *Cell* 110 (2002) 673–687.
- [172] M.A. Baxter, M.V. Camarasa, N. Bates, F. Small, P. Murray, D. Edgar, et al., Analysis of the distinct functions of growth factors and tissue culture substrates necessary for the long-term self-renewal of human embryonic stem cell lines, *Stem Cell Res.* 3 (2009) 28–38.
- [173] S.R. Braam, L. Zeinstra, S. Litjens, D. Ward-van Oostwaard, S. van den Brink, L. van Laake, et al., Recombinant vitronectin is a functionally defined substrate that supports human embryonic stem cell self-renewal via alphavbeta5 integrin, *Stem Cells.* 26 (2008) 2257–2265.
- [174] T. Miyazaki, S. Futaki, H. Suemori, Y. Taniguchi, M. Yamada, M. Kawasaki, et al., Laminin E8 fragments support efficient adhesion and expansion of dissociated human pluripotent stem cells, *Nature Commun.* 3 (2012) 1236.
- [175] S. Rodin, L. Antonsson, O. Hovatta, K. Tryggvason, Monolayer culturing and cloning of human pluripotent stem cells on laminin-521-based matrices under xeno-free and chemically defined conditions, *Nature Protocols.* 9 (2014) 2354–2368.
- [176] D. Soteriou, B. Iskender, A. Byron, J.D. Humphries, S. Borg-Bartolo, M.C. Haddock, et al., Comparative proteomic analysis of supportive and unsupportive extracellular matrix substrates for human embryonic stem cell maintenance, *J. Biol. Chem.* 288 (2013) 18,716–18,731.
- [177] M.C. Frame, H. Patel, B. Serrels, D. Lietha, M.J. Eck, The FERM domain: organizing the structure and function of FAK, *Nature Reviews Mol. Cell Biol.* 11 (2010) 802–814.
- [178] J.T. Parsons, Focal adhesion kinase: the first ten years, *J. Cell Sci.* 116 (2003) 1409–1416.
- [179] L. Vitillo, M. Baxter, B. Iskender, P. Whiting, S.J. Kimber, Integrin-associated focal adhesion kinase protects human embryonic stem cells from apoptosis, detachment, and differentiation, *Stem Cell Reports.* 7 (2016) 167–176.
- [180] A.K. Jain, K. Allton, M. Iacovino, E. Mahen, R.J. Milczarek, T.P. Zwaka, et al., p53 regulates cell cycle and microRNAs to promote differentiation of human embryonic stem cells, *PLoS Biol.* 10 (2012) e1001268.
- [181] A.M. Singh, D. Reynolds, T. Cliff, S. Ohtsuka, A.L. Mattheyses, Y. Sun, et al., Signaling network crosstalk in human pluripotent cells: a Smad2/3-regulated switch that controls the balance between self-renewal and differentiation, *Cell Stem Cell* 10 (2012) 312–326.
- [182] L.G. Villa-Diaz, J.K. Kim, A. Laperle, S.P. Palecek, P.H. Krebsbach, Inhibition of focal adhesion kinase signaling by integrin alpha6beta1 supports human pluripotent stem cell self-renewal, *Stem Cells* 34 (2016) 1753–1764.
- [183] S.P. Toya, K.K. Wary, M. Mittal, F. Li, P.T. Toth, C. Park, et al., Integrin alpha6beta1 expressed in ESCs instructs the differentiation to endothelial cells, *Stem Cells* 33 (2015) 1719–1729.
- [184] M.P. Playford, M.D. Schaller, The interplay between Src and integrins in normal and tumor biology, *Oncogene* 23 (2004) 7928–7946.
- [185] S.M. Thomas, J.S. Brugge, Cellular functions regulated by Src family kinases, *Annual Review of Cell and Developmental Biol.* 13 (1997) 513–609.
- [186] C. Anneren, C.A. Cowan, D.A. Melton, The Src family of tyrosine kinases is important for embryonic stem cell self-renewal, *J. Biol. Chem.* 279 (2004) 31,590–31,598.
- [187] M.A. Meyn 3rd, S.J. Schreiner, T.P. Dumitrescu, G.J. Nau, T.E. Smithgall, SRC family kinase activity is required for murine embryonic stem cell growth and differentiation, *Mol. Pharmacology* 68 (2005) 1320–1330.
- [188] M.A. Meyn 3rd, T.E. Smithgall, Chemical genetics identifies c-Src as an activator of primitive ectoderm formation in murine embryonic stem cells, *Science Signaling* 2 (2009) ra64.
- [189] X. Zhang, M.A. Meyn 3rd, T.E. Smithgall, c-Yes tyrosine kinase is a potent suppressor of ES cell differentiation and antagonizes the actions of its closest phylogenetic relative, c-Src, *ACS Chem Biol.* 9 (2014) 139–146.
- [190] X. Zhang, C. Simerly, C. Hartnett, G. Schatten, T.E. Smithgall, Src-family tyrosine kinase activities are essential for differentiation of human embryonic stem cells, *Stem Cell Res.* 13 (2014) 379–389.
- [191] M. Ernst, D.P. Gearing, A.R. Dunn, Functional and biochemical association of Hck with the LIF/IL-6 receptor signal transducing subunit gp130 in embryonic stem cells, *EMBO J.* 13 (1994) 1574–1584.
- [192] C. Tamm, N. Bower, C. Anneren, Regulation of mouse embryonic stem cell self-renewal by a Yes-YAP-TEAD2 signaling pathway downstream of LIF, *J. Cell Sci.* 124 (2011) 1136–1144.
- [193] M. Trouillas, C. Saucourt, B. Guillotin, X. Gauthereau, L. Ding, F. Buchholz, et al., Three LIF-dependent signatures and gene clusters with atypical expression profiles, identified by transcriptome studies in mouse ES cells and early derivatives, *BMC Genomics* 10 (2009) 73.

- [194] Z. Zhang, B. Liao, M. Xu, Y. Jin, Post-translational modification of POU domain transcription factor Oct-4 by SUMO-1, *FASEB J.: Official Publication of the Federation of American Societies for Experimental Biology*. 21 (2007) 3042–3051.
- [195] Q.Y. Lei, H. Zhang, B. Zhao, Z.Y. Zha, F. Bai, X.H. Pei, et al., TAZ promotes cell proliferation and epithelial-mesenchymal transition and is inhibited by the hippo pathway, *Mol. and Cellular Biol.* 28 (2008) 2426–2436.
- [196] B. Zhao, X. Wei, W. Li, R.S. Udan, Q. Yang, J. Kim, et al., Inactivation of YAP oncoprotein by the Hippo pathway is involved in cell contact inhibition and tissue growth control, *Genes & Dev.* 21 (2007) 2747–2761.
- [197] J.S. Mo, H.W. Park, K.L. Guan, The Hippo signaling pathway in stem cell biology and cancer, *EMBO Reports*. 15 (2014) 642–656.
- [198] Z. Meng, T. Moroishi, K.L. Guan, Mechanisms of Hippo pathway regulation, *Genes & Dev.* 30 (2016) 1–17.
- [199] M. Aragona, T. Panciera, A. Manfrin, S. Giullitti, F. Michielin, N. Elvassore, et al., A mechanical checkpoint controls multicellular growth through YAP/TAZ regulation by actin-processing factors, *Cell* 154 (2013) 1047–1059.
- [200] S. Dupont, L. Morsut, M. Aragona, E. Enzo, S. Giullitti, M. Cordenonsi, et al., Role of YAP/TAZ in mechanotransduction, *Nature* 474 (2011) 179–183.
- [201] M. Ohgushi, M. Minaguchi, Y. Sasai, Rho-signaling-directed YAP/TAZ activity underlies the long-term survival and expansion of human embryonic stem cells, *Cell Stem Cell* 17 (2015) 448–461.
- [202] M. Ramalho-Santos, S. Yoon, Y. Matsuzaki, R.C. Mulligan, D.A. Melton, “Stemness”: transcriptional profiling of embryonic and adult stem cells, *Science* 298 (2002) 597–600.
- [203] X. Varelas, R. Sakuma, P. Samavarchi-Tehrani, R. Peerani, B.M. Rao, J. Dembowy, et al., TAZ controls Smad nucleocytoplasmic shuttling and regulates human embryonic stem-cell self-renewal, *Nature Cell Biol.* 10 (2008) 837–848.
- [204] C. Alarcon, A.I. Zaromytidou, Q. Xi, S. Gao, J. Yu, S. Fujisawa, et al., Nuclear CDKs drive Smad transcriptional activation and turnover in BMP and TGF-beta pathways, *Cell* 139 (2009) 757–769.
- [205] I. Lian, J. Kim, H. Okazawa, J. Zhao, B. Zhao, J. Yu, et al., The role of YAP transcription coactivator in regulating stem cell self-renewal and differentiation, *Genes & Dev.* 24 (2010) 1106–1118.
- [206] S. Benhamouche, M. Curto, I. Saotome, A.B. Gladden, C.H. Liu, M. Giovannini, et al., Nf2/Merlin controls progenitor homeostasis and tumorigenesis in the liver, *Genes & Dev.* 24 (2010) 1718–1730.
- [207] F.D. Camargo, S. Gokhale, J.B. Johnnidis, D. Fu, G.W. Bell, R. Jaenisch, et al., YAP1 increases organ size and expands undifferentiated progenitor cells, *Curr Biol.* 17 (2007) 2054–2060.
- [208] X. Cao, S.L. Pfaff, F.H. Gage, YAP regulates neural progenitor cell number via the TEA domain transcription factor, *Genes & Dev.* 22 (2008) 3320–3334.
- [209] K.P. Lee, J.H. Lee, T.S. Kim, T.H. Kim, H.D. Park, J.S. Byun, et al., The Hippo-Salvador pathway restrains hepatic oval cell proliferation, liver size, and liver tumorigenesis, *Proc Natl Acad Sci U S A.* 107 (2010) 8248–8253.
- [210] K. Schlegelmilch, M. Mohseni, O. Kirak, J. Pruszk, J.R. Rodriguez, D. Zhou, et al., Yap1 acts downstream of alpha-catenin to control epidermal proliferation, *Cell* 144 (2011) 782–795.
- [211] N. Zhang, H. Bai, K.K. David, J. Dong, Y. Zheng, J. Cai, et al., The Merlin/NF2 tumor suppressor functions through the YAP oncoprotein to regulate tissue homeostasis in mammals, *Dev Cell* 19 (2010) 27–38.
- [212] D. Zhou, C. Conrad, F. Xia, J.S. Park, B. Payer, Y. Yin, et al., Mst1 and Mst2 maintain hepatocyte quiescence and suppress hepatocellular carcinoma development through inactivation of the Yap1 oncogene, *Cancer Cell* 16 (2009) 425–438.
- [213] D. Zhou, Y. Zhang, H. Wu, E. Barry, Y. Yin, E. Lawrence, et al., Mst1 and Mst2 protein kinases restrain intestinal stem cell proliferation and colonic tumorigenesis by inhibition of Yes-associated protein (Yap) overabundance, *Proc Natl Acad Sci U S A.* 108 (2011) E1312–E1320.
- [214] H. Qin, K. Blaschke, G. Wei, Y. Ohi, L. Blouin, Z. Qi, et al., Transcriptional analysis of pluripotency reveals the Hippo pathway as a barrier to reprogramming, *Human Mol. Genet.* 21 (2012) 2054–2067.
- [215] H. Qin, A. Diaz, L. Blouin, R.J. Lebbink, W. Patena, P. Tanbun, et al., Systematic identification of barriers to human iPSC generation, *Cell* 158 (2014) 449–461.
- [216] H. Qin, M. Hejna, Y. Liu, M. Percharde, M. Wossidlo, L. Blouin, et al., YAP induces human naïve pluripotency, *Cell Reports* 14 (2016) 2301–2312.
- [217] T.A. Beyer, A. Weiss, Y. Khomchuk, K. Huang, A.A. Ogunjimi, X. Varelas, et al., Switch enhancers interpret TGF-beta and Hippo signaling to control cell fate in human embryonic stem cells, *Cell Reports*. 5 (2013) 1611–1624.