



University of Dundee

Loss of Cardioprotective Effects at the ADAMTS7 Locus as a Result of Gene-Smoking Interactions

Saleheen, Danish; Zhao, Wei; Young, Robin; Nelson, Christopher P.; Ho, Weang Kee; Ferguson, Jane F.

Published in:
Circulation

DOI:
[10.1161/CIRCULATIONAHA.116.022069](https://doi.org/10.1161/CIRCULATIONAHA.116.022069)

Publication date:
2017

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Saleheen, D., Zhao, W., Young, R., Nelson, C. P., Ho, W. K., Ferguson, J. F., Rasheed, A., Ou, K., Nurnberg, S. T., Bauer, R. C., Goel, A., Do, R., Stewart, A. F. R., Hartiala, J., Zhang, W., Thorleifsson, G., Strawbridge, R. J., Sinisalo, J., Kanoni, S., ... EPIC-CVD, PROMIS, CARDIoGRAMplusC4D (2017). Loss of Cardioprotective Effects at the *ADAMTS7* Locus as a Result of Gene-Smoking Interactions. *Circulation*, 135(24), 2336-2353. <https://doi.org/10.1161/CIRCULATIONAHA.116.022069>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 Loss of cardio-protective effects at the *ADAMTS7* locus due to gene-smoking interactions

2 Danish Saleheen PhD,^{1,2} Wei Zhao MSc,¹ Robin Young PhD,³ Christopher P Nelson PhD,⁴
3 WeangKee Ho PhD,³ Jane F Ferguson PhD,⁵ Asif Rasheed MBBS,² Kristy Ou BS,⁵ Sylvia T
4 Nurnberg PhD,^{5,7} Robert C. Bauer PhD,¹⁴ Anuj Goel MSc,⁶ Ron Do PhD,^{7,8} Alexandre FR Stewart
5 PhD,⁹ Jaana Hartiala PhD,¹⁰ Weihua Zhang PhD,^{11,12} Gudmar Thorleifsson PhD,^{50,51} Rona J
6 Strawbridge PhD,¹³ Juha Sinisalo PhD,⁶¹ Stavroula Kanoni PhD,¹⁵ Sanaz Sedaghat PhD,¹⁶ Eirini
7 Marouli PhD,^{15,17} Kati Kristiansson PhD,¹⁸ Jing Hua Zhao PhD,¹⁹ Robert Scott PhD,¹⁹ Dominique
8 Gauguier PhD,²⁰ Svati H Shah MD,²¹ Albert Vernon Smith PhD,^{22,36} Natalie van Zuydam PhD,²³
9 Amanda J Cox PhD,²⁴ Christina Willenborg PhD,^{25,26}, Thorsten Kessler MD,^{27,28} Lingyao Zeng
10 PhD,^{27,29} Michael A Province PhD,³⁰ Andrea Ganna PhD,^{31,32} Lars Lind PhD,³³ Nancy L. Pedersen
11 PhD,³⁴ Charles C White PhD,³⁵ Anni Joensuu MSc,^{18,37} Marcus Edi Kleber PhD,³⁸ Alistair S Hall
12 PhD,³⁹ Winfried März PhD,⁴⁰ Veikko Salomaa PhD,¹⁸ Christopher O'Donnell MD,⁴¹ Erik Ingelsson
13 PhD,^{42,43} Mary F Feitosa PhD,³⁰ Jeanette Erdmann PhD,^{25,26} Donald W. Bowden PhD,²⁴ Colin N.A.
14 Palmer PhD,²³ Vilmundur Gudnason PhD,^{22,36} Ulf De Faire PhD,⁴⁴ Pierre Zalloua PhD,⁴⁵ Nicholas
15 Wareham PhD,²⁰ John R Thompson PhD,⁴⁶ Kari Kuulasmaa PhD,¹⁸ George Dedoussis PhD,¹⁷
16 Markus Perola PhD,^{18,37} Abbas Dehghan PhD,¹⁶ John C Chambers PhD,^{11,12,47} Jaspal Kooner
17 MD,^{11,47,48} Hooman Allayee PhD,¹⁰ Panos Deloukas PhD,^{15,49} Ruth McPherson PhD,⁹ Kari
18 Stefansson PhD,^{50,51} Heribert Schunkert MD,^{27,29} Sekar Kathiresan MD,⁵²⁻⁵⁵ Martin Farrall PhD,⁶
19 EPIC-CVD,³ Philippe Marcel Frossard DSC,² Daniel J Rader MD,^{56,57} Nilesh J Samani MD,⁴
20 PROMIS,² CARDIoGRAMplusC4D, Muredach P. Reilly MD.¹⁴

21 Running Title: Gene*Smoking interaction, *ADAMTS7* locus & CHD risk

22 1. Department of Biostatistics and Epidemiology, University of Pennsylvania, Philadelphia, Pennsylvania, USA
23 2. Center for Non-Communicable Diseases, Karachi, Pakistan.
24 3. Department of Public Health and Primary Care, University of Cambridge, UK
25 4. Department of Cardiovascular Sciences, University of Leicester, Leicester, LE3 9QP, UK
26 5. Cardiology Division, Department of Medicine, Vanderbilt University, Nashville, Tennessee, USA
27 6. Division of Cardiovascular Medicine, Radcliffe Department of Medicine & Wellcome Trust Centre for Human
28 Genetics, University of Oxford, Oxford, UK
29 7. The Charles Bronfman Institute of Personalized Medicine, Icahn School of Medicine at Mount Sinai, New
30 York, NY, USA
31 8. Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY,
32 USA
33 9. Ruddy Canadian Cardiovascular Genetics Centre, University of Ottawa Heart Institute, Ottawa, Canada
34 10. Institute for Genetic Medicine and Department of Preventive Medicine, Keck School of Medicine, University
35 of Southern California, Los Angeles, California, 90033, USA
36 11. Department of Epidemiology and Biostatistics, Imperial College London, London, UK
37 12. Department of Cardiology, Ealing Hospital NHS Trust, Middlesex, UK
38 13. Cardiovascular Medicine Unit, Department of Medicine Solna, Karolinska Institutet, Stockholm, Sweden
39 14. Cardiology Division, Department of Medicine and the Irving Institute for Clinical and Translational
40 Research, Columbia University Medical Center, New York, USA
41 15. William Harvey Research Institute, Barts and the London School of Medicine and Dentistry, Queen Mary
42 University of London, London, UK
43 16. Department of Epidemiology, Erasmus University Medical center, Rotterdam, The Netherlands
44 17. Department of Dietetics-Nutrition, Harokopio University, 70 El. VenizelouStr, Athens, Greece
45 18. National Institute for Health and Welfare, Helsinki, Finland
46 19. MRC Epidemiology Unit, Institute of Metabolic Science, University of Cambridge School of Clinical
47 Medicine, Box 285, Cambridge Biomedical Campus, Cambridge, CB2 0QQ, United Kingdom

48 20. INSERM, UMRS1138, Centre de Recherche des Cordeliers, Paris, France
49 21. Division of Cardiology, Department of Medicine, Duke University Medical Center, Durham, North Carolina,
50 27710, USA
51 22. Icelandic Heart Association, Kopavogur, Iceland
52 23. Medical Research Institute, Ninewells Hospital and Medical School, University of Dundee, Dundee, UK
53 24. Center for Genomics and Personalized Medicine Research, Wake Forest School of Medicine, Winston-
54 Salem, North Carolina, USA
55 25. Institut für Integrative und Experimentelle Genomik, Universität zu Lübeck, Lübeck, Germany
56 26. DZHK (German Research Center for Cardiovascular Research) partner site Hamburg–Lübeck–Kiel,
57 Lübeck, Germany
58 27. Deutsches Herzzentrum München, Technische Universität München, München, Germany
59 28. Klinikum rechts der Isar, München, Germany
60 29. DZHK (German Centre for Cardiovascular Research), partner site Munich Heart Alliance, München,
61 Germany
62 30. Department of Genetics, Washington University School of Medicine, St. Louis, Missouri, USA
63 31. Analytic and Translational Genetics Unit, Department of Medicine, Massachusetts General Hospital and
64 Harvard Medical School, Boston, Massachusetts 02114, USA
65 32. Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, 7 Cambridge Center,
66 Cambridge, Massachusetts 02142, USA
67 33. Department of Medical Sciences, Cardiovascular Epidemiology, Uppsala University, Uppsala, Sweden
68 34. Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden
69 35. Department of Biostatistics Boston University School of Public Health Framingham Heart Study 801
70 Massachusetts Avenue, CT3 Boston, Massachusetts, 02118, USA
71 36. Faculty of Medicine, University of Iceland, Reykjavik, Iceland
72 37. University of Helsinki, Institute for Molecular Medicine, Finland (FIMM)
73 38. Department of Medicine, Mannheim Medical Faculty, Heidelberg University, Heidelberg, Germany
74 39. Leeds Institute of Genetics, Health and Therapeutics, University of Leeds, Leeds, UK
75 40. Synlab Academy, Synlab Services GmbH, Mannheim, Germany and Clinical Institute of Medical and
76 Chemical Laboratory Diagnostics, Medical University of Graz, Graz, Austria
77 41. National Heart, Lung, and Blood Institute and the Framingham Heart Study, National Institutes of Health,
78 Bethesda, Maryland, USA
79 42. Department of Medical Sciences, Molecular Epidemiology and Science for Life Laboratory, Uppsala
80 University, Uppsala, Sweden
81 43. Department of Medicine, Division of Cardiovascular Medicine, Stanford University School of Medicine,
82 Stanford, California, USA
83 44. Division of Cardiovascular Epidemiology, Institute of Environmental Medicine, Karolinska Institutet,
84 Stockholm, Sweden
85 45. Lebanese American University, School of Medicine, Beirut 1102 2801, Lebanon
86 46. Department of Health Sciences, University of Leicester, Leicester, UK
87 47. Imperial College Healthcare NHS Trust, London, UK
88 48. Cardiovascular Science, National Heart and Lung Institute, Imperial College London, London, UK
89 49. Princess Al-Jawhara Al-Brahim Centre of Excellence in Research of Hereditary Disorders (PACER-HD),
90 King Abdulaziz University, Jeddah, Saudi Arabia
91 50. deCODE Genetics, Sturlugata 8, IS-101 Reykjavik, Iceland
92 51. University of Iceland, School of Medicine, 101 Reykjavik, Iceland
93 52. Broad Institute of the Massachusetts Institute of Technology and Harvard University, Cambridge,
94 Massachusetts, USA
95 53. Cardiovascular Research Center, Massachusetts General Hospital, Boston, Massachusetts, USA
96 54. Center for Human Genetic Research, Massachusetts General Hospital, Boston, Massachusetts, USA
97 55. Department of Medicine, Harvard Medical School, Boston, Massachusetts, USA
98 56. Department of Genetics, University of Pennsylvania, Philadelphia, Pennsylvania, USA
99 57. Division of Translational Medicine and Human Genetics, Department of Medicine, Perelman School of
100 Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA

101
102
103

104
105
106
107
108
109
110
111
112
113
114
115

116
117
118
119
120
121
122
123
124
125

126

127

128

129

130

131

132

133

134

135

136

137

Corresponding authors:

Danish Saleheen, MBBS, PhD
Assistant Professor of Biostatistics and Epidemiology
University of Pennsylvania
11-134 Translational Research Center
3400 Civic Center Boulevard
Philadelphia, PA 19104
Tel: 215-573-6323
Fax: 215-573-2094
Email: saleheen@mail.med.upenn.edu

Muredach P. Reilly, MBBCH, MSCE
Herbert and Florence Irving Professor of Medicine
Director of the Irving Institute for Clinical and Translational Research
Columbia University
622 West 168th Street, PH10-305
New York, NY 10032
Tel: 212-305-9453
Fax: 212-305-3213
Email: mpr2144@cumc.columbia.edu

138 **ABSTRACT**

139 **Background** Common diseases such as coronary heart disease (CHD) are complex in etiology. The
140 interaction of genetic susceptibility with lifestyle factors may play a prominent role. However, gene-
141 environment interactions for CHD have been difficult to identify. Here, we investigate interaction of
142 smoking behavior, a potent lifestyle factor, with genotypes that have been shown to associate with
143 CHD risk.

144 **Methods** We analyzed data on 60,919 CHD cases and 80,243 controls from 29 studies for gene-
145 smoking interactions for genetic variants at 45 loci previously reported to associate with CHD risk.
146 We also studied 5 loci associated with smoking behavior. Study specific gene-smoking interaction
147 effects were calculated and pooled using fixed-effects meta-analyses. Interaction analyses were
148 declared to be significant at a *P-value* < 1.0×10^{-3} (Bonferroni correction for 50 tests).

149 **Results** We identified novel gene-smoking interaction for a variant upstream of the *ADAMTS7* gene.
150 Every T allele of rs7178051 was associated with lower CHD risk by 12% in never-smokers (*P-value*:
151 1.3×10^{-16}) compared to 5% in ever-smokers (*P-value*: 2.5×10^{-4}) translating to a 60% loss of CHD
152 protection conferred by this allelic variation in people who smoked tobacco (*Interaction P-value*:
153 8.7×10^{-5}). The protective T allele at rs7178051 was also associated with reduced *ADAMTS7*
154 expression in human aortic endothelial cells and lymphoblastoid cell lines. Exposure of human
155 coronary artery smooth muscle cells to cigarette smoke extract led to induction of *ADAMTS7*.

156 **Conclusion** Allelic variation at rs7178051 that associates with reduced *ADAMTS7* expression
157 confers stronger CHD protection in “never-smokers” compared to “ever-smokers”. Increased
158 vascular *ADAMTS7* expression may contribute to the loss of CHD protection in smokers.

159 **Key words:** Gene-smoking interaction, gene-environment interaction, coronary heart disease,
160 *ADAMTS7*, smoking.

161 **Word count: 269**

162

163

164 **Clinical Perspective**

165 1) What is new?

- 166
- 167 • Using data on 60,919 CHD cases and 80,243 controls, this study conducted gene-
168 environment interaction analyses to investigate effect modification by smoking behavior at
169 established CHD and smoking related loci.
 - 170 • Cardio-protective effects associated with allelic variation at the *ADAMTS7* locus were
171 attenuated by 60% in people who smoked tobacco compared to those who did not smoke.
 - 172 • Allelic variation at *ADAMTS7* associated with reduced CHD risk was associated with reduced
173 *ADAMTS7* expression in human aortic endothelial cells and lymphoblastoid cell lines.
 - 174 • Exposure of human coronary artery smooth muscle cells to cigarette smoke extract led to
175 induction of *ADAMTS7*.
- 176

177 2) What are the clinical implications?

- 178
- 179 • These human genomic data provide new insights into potential mechanisms of CHD in
180 cigarette smokers.
 - 181 • Findings from this study also point towards the directional impact of the *ADAMTS7* locus on
182 CHD.
 - 183 • *ADAMTS7* and its substrates have a specific role in cigarette smoking related CHD.
 - 184 • Inhibition of *ADAMTS7* is a novel potential therapeutic strategy for CHD that may have
185 particular benefits in individuals who smoke cigarettes.
- 186

187

188

189

190

191

192

193

194 **INTRODUCTION**

195 Coronary heart disease (CHD) is a complex disorder resulting from the interplay of lifestyle
196 and genetic factors.^{1, 2} Yet, gene-environment interactions for CHD have been difficult to identify.
197 Cigarette smoking is one of the strongest lifestyle risk factors for CHD but the underlying molecular
198 mechanisms of CHD in humans who smoke remain uncertain.³⁻⁵ Cigarette smoking accounts for
199 one-fifth of all CHD events globally and is responsible for ~1.6 million deaths attributable to CHD
200 annually.⁶ Genome-wide association studies (GWAS) have improved our understanding on the
201 genetic predisposition to both CHD and smoking behavior.⁷⁻¹⁰ Joint or interactive effects of genetic
202 variation and smoking behavior in the etiology of CHD, however, remain poorly understood. GWAS
203 can provide new opportunities to investigate gene-smoking interactions.

204 We hypothesized that genetic predisposition to CHD is modified by cigarette smoking at
205 loci discovered by GWAS to be associated with either CHD or smoking behavior. We conducted a
206 focused experiment at 50 main-effect loci (45 for CHD and 5 for smoking behavior) using genetic
207 data and information on smoking behavior in 60,919 CHD cases and 80,243 controls from 29
208 different studies. We report novel findings on gene-smoking interactions in CHD.

209

210

211

212

213

214

215

216

217

218

219

220

221 **METHODS**

222 **Summary of study Design**

223 All studies participating in the CARDIoGRAMplusC4D consortium⁷⁻⁹ that had information
224 available on smoking status, CHD risk and genotypes at the 50 CHD and smoking behavior-
225 associated loci were invited to participate. The current study had five inter-related components
226 (**Supplementary Figure-1**). First, as part of the quality control, we investigated the association of
227 smoking status with CHD risk within each study. Second, we performed an updated analysis of all
228 the SNPs (\pm 50 KB) at the 45 established CHD loci to identify the variant with the strongest CHD
229 association in our study population at each established CHD locus. Effect estimates from each study
230 in association with CHD risk were obtained and pooled to identify the strongest CHD associated
231 variant (“lead variant”). Third, we investigated gene-smoking interactions at these 45 CHD loci and at
232 5 loci related to smoking behavior. Fourth, for loci demonstrating differential CHD associations by
233 smoking status, we mapped the interaction region, examined linkage disequilibrium (LD) across the
234 region and performed conditional analyses to identify independent genetic signals. Finally, for loci
235 exhibiting gene-smoking interaction in CHD, we assessed eQTL data for association of variants with
236 expression of local genes in available datasets and examined expression of these genes in multiple
237 cell types that play prominent roles in smoking-CHD pathobiology.

238 **Harmonization of phenotypes and genotypes**

239 Summary level estimates for each study were shared via a secure FTP site. We used
240 “ever-smoking” as a primary exposure and data were harmonized by uniformly characterizing
241 participants in each study into two categories, “ever-smokers” and “never-smokers”. “Ever-smokers”
242 were defined as those who had smoked more than 100 cigarettes in a lifetime. For case-control
243 studies, information on “ever smoking” status collected at the time of enrollment was used for the
244 current analyses; whereas for prospective cohort studies, information on smoking status obtained at
245 the baseline visit was used for the current investigation. CHD was defined based on evidence from
246 angiography or history of verified myocardial infarction (MI), percutaneous coronary interventions
247 (PCI) or coronary artery bypass grafting (CABG) as published in CARDIoGRAMplusC4D projects.⁷⁻⁹
248 Genotype data generated through GWAS (directly genotyped or imputed) or cardio-metabochip
249 (directly genotyped only) arrays were obtained from each study and all genetic data were aligned
250 using the build-37 reference panel. Imputed SNPs were removed if they had any of the following: (i)
251 a minor allele frequency of <1%; (ii) info score of <0.90; or (iii) confidence score <0.90. For each
252 study, GWAS data were imputed using the Phase II CEU HapMap reference population.¹¹ Standard

253 quality control criteria were applied by each participating study, as described previously.⁷ All
254 participating studies in the CARDIoGRAMplusC4D consortium were approved by their locally
255 relevant institutional review boards and all participants gave written informed consent before their
256 enrollment in each study.⁷⁻⁹

257 **STATISTICAL ANALYSIS**

258 **Gene-smoking interaction analyses**

259 Initial quality control and association of established CHD loci with CHD risk: As part of an initial
260 quality control, effect estimates from each study were obtained for “ever-smoking” status and CHD
261 risk using a case-control logistic regression model adjusted for age and sex. Each participating study
262 also assessed and, if needed, controlled for population stratification by including principal
263 components as covariates in the regression model as described earlier.⁷⁻⁹ To identify variant(s) with
264 the most significant association with CHD risk at established CHD loci in our study population,
265 logistic regression analyses were conducted by each participating study for all the SNPs flanking
266 (± 50 kb) the lead variant previously reported at each CHD locus. Effect estimates and standard
267 errors were obtained and meta-analyzed using a fixed-effects inverse variance approach. All lead
268 variants identified through these analyses were further investigated for gene-smoking interactions in
269 CHD. One lead variant per locus was selected for primary gene-smoking interaction analyses.

270 Investigation of the APOE locus: Although APOE has been recently established as a GWAS locus,⁷
271 previous studies prior to GWAS have suggested that CHD risk is higher among carriers of the $\epsilon 4$
272 allele at the APOE locus in smokers than in non-smokers.¹²⁻¹⁴ Because the $\epsilon 2$, $\epsilon 3$ and $\epsilon 4$ alleles at
273 the APOE locus are not captured by the GWAS platform, we specifically conducted genotyping for
274 rs429358 and rs7412 variants to capture the three epsilon (ϵ) alleles in 13,822 participants (including
275 7,286 first-onset myocardial infarction cases) in the PROMIS study.¹⁵

276 Gene-smoking interaction analyses at CHD and smoking loci: To assess gene-smoking interactions,
277 analyses were conducted within each study, adjusted for age, sex and other study specific
278 covariates (e.g., principal components), and variants were analyzed in association with CHD
279 separately in “ever-smokers” and “never-smokers”. Results from the two groups were then used to
280 test for interaction within each study. For the 50 variants, an interaction test statistic was calculated
281 within each study using the following equation as adapted from Teslovich *TM* et.al.¹⁶

282
$$\frac{(\beta_n - \beta_e)}{\sqrt{SE_n^2 + SE_e^2}}$$

283 where β_n and β_e are the beta coefficients for the SNP in never-smokers and ever-smokers
284 respectively, SE_n and SE_e are the standard errors for the log-ORs estimated for never-smokers and
285 ever-smokers, respectively. Study specific interaction beta(s) and se(s) were calculated within each
286 study and were pooled across studies using a fixed-effects meta-analysis. Interaction analyses were
287 declared to be significant at a P-value of $<1.0 \times 10^{-3}$ (Bonferroni correction for 50 tests).

288 Conditional analyses on chr.15q25.1: At chr.15q25.1, we observed two variants exhibiting gene-
289 smoking interactions for CHD. The proximity of these two signals raised the possibility that the
290 observed interactions may represent a single interaction locus across the entire region. To
291 investigate this possibility we undertook conditional analyses using an approximate conditional and
292 joint analyses approach, also known as GCTA (Genome-wide Complex Trait Analysis), as described
293 previously.¹⁷⁻²² Briefly, this method leverages summary-level statistics from a meta-analysis and uses
294 LD corrections between SNPs estimated from a reference sample. Such an approach has been
295 shown to yield similar results to that obtained from conditional analyses conducted on individual
296 participant data and has been successfully implemented in several other studies that have fine-
297 mapped loci for other complex traits.¹⁷⁻²² Using this approach, we first conducted separate
298 conditional analyses at the chr.15q25.1 locus to identify main-effect variant(s) independently
299 associated with CHD and smoking behavior, respectively. We used the meta-analyzed data for CHD
300 main effects in the CARDIoGRAMplus4D consortium to identify SNPs independently associated with
301 CHD risk and we used the genetic meta-analysis data from the Tobacco and Genetics Consortium
302 (TGC) in 140,000 participants to identify variants independently associated with smoking behavior.
303 We then estimated the effects of these independent variants on CHD risk stratified by smoking
304 status and mutually adjusted the effects of these variants for each other.

305

306 **Analysis of eQTLs and regulatory features at the chr15q25.1 gene-smoking interaction locus**

307 eQTL analyses: We mined publicly available databases to identify genotype-related expression
308 differences (eQTLs) in *ADAMTS7* and the *CHRNA4-A3-A5* gene cluster in order to understand the
309 directionality of the association of expression of these genes with CHD and smoking behavior.
310 Specifically, we investigated data available from the GTEx consortium,²³ the HapMap consortium
311 (restricted to European populations), and the Multiple Tissue Human Expression Resource
312 (MuTHER).²⁴ We also analyzed expression data in 147 donor HAOEC lines.²⁵ We used a nominal P-
313 value of 0.002 to account for multiple testing involved in the eQTL analyses.

314 Regulatory features of the chr. 15q25.1 region: Data from ENCODE²⁶ were explored as described in
315 eMethods. CHIP-seq experiments were performed on confluent HCASMC (Cell Applications 350-05a
316 & Lonza CC-2583; cultured in SmGM-2 BulletKit media; Lonza) as described.²⁷ TCF21 (Abcam
317 ab49475), Jun (Santa Cruz Biotechnology sc-1694), JunD (Santa Cruz Biotechnology sc-74), and
318 CEBP (Santa Cruz Biotechnology sc-150) transcription factor binding was interrogated and H3K27ac
319 data were acquired using the same CHIP protocol with an anti-H3K27ac antibody (Abcam; ab4729).
320 Reads were aligned to the human genome (GRCh37p13) using STAR.²⁸

321

322 **Analyses of *ADAMTS7* and *CHRNA3-A5* gene expression in vascular cells and tissues**

323 *ADAMTS7* and *CHRNA3-A5* gene expression in vascular cells: *ADAMTS7* and *CHRNA3-A5*
324 mRNA levels were measured in cultured human coronary artery smooth muscle cells (HCASMC;
325 Lonza CC-2583, Lonza Walkersville, MD), human coronary artery endothelial cells (HCAEC, Lonza
326 CC-2585), human aortic smooth muscle cells (HAoSMC, Lonza CC-2571), human aortic endothelial
327 cells (HAoEC, Lonza CC-2535), human aortic adventitial fibroblasts (HAoAF, Lonza CC-7014), and
328 human acute monocytic leukemia cell line (THP-1, ATCC TIB-202). Further details are provided in
329 eMethods.

330 *ADAMTS7* and *CHRNA3-A5* gene expression in response to cigarette smoke extract: HCASMC
331 were grown to confluence and cigarette smoke extract experiments performed at passage-7.
332 Cigarette smoke extract was custom-prepared by Arista Laboratories (Richmond, VA). Briefly, the
333 condensate was generated by smoking Marlboro Red King Size Hard Pack cigarettes on an
334 analytical smoke machine under International Organization for Standardization smoking conditions.
335 The smoke condensate was collected on 92 mm filter pads and extracted from each pad in DMSO
336 by shaking to obtain a solution of ~20 mg/mL final concentration of the total particulate matter.
337 Serum starved (24 hrs) HCASMC were treated with 0.5% or 1.0% cigarette smoke extract (v/v) for 4,
338 12, and 24 hrs in serum reduced conditions (0.5% FBS in DMEM). Details on RNA preparation and
339 q-PCR are provided in **Supplementary Methods**.

340

341

342

343

344 RESULTS

345 Description of the participating studies

346 Of the 37 studies participating in the CARDIoGRAMplusC4D consortium, information on
347 “ever-smoking” was available in 30 studies, yielding a total sample size of 60,919 CHD cases and
348 80,243 controls. All studies recruited participants of European ancestry, except PROMIS (South
349 Asian),¹⁵ LOLIPOP (South Asian)²⁹ and FGENTCARD (Lebanese).³⁰ Number of CHD cases and
350 controls and percentages that were “ever-smokers” are provided in **Supplementary Table 1**. As
351 expected, in all the participating studies, association of “ever-smoking” status with CHD risk was
352 directionally consistent with an increased risk of CHD (**Supplementary Figure 2**).

353 New variants associated with CHD at established loci

354 **Supplementary Figure 3** and **Supplementary Table 2** present effect estimates for the
355 association with CHD for (i) the most significant variant that we identified at known CHD loci in the
356 current CARDIoGRAMplusC4D consortium analysis as well as for (ii) the top SNP previously
357 reported at each of these established CHD loci. Of the 45 established CHD loci, we identified 32 for
358 which we discovered a more statistically significant SNP in association with CHD risk in our dataset
359 than the prior reported top variant. All of these 32 SNPs were in moderate to high LD ($r^2 > 0.6$) with
360 the previously published variants.⁷⁻⁹ In our primary gene-smoking interaction analyses, at each of the
361 CHD loci, we, therefore, used the SNP with the most significant CHD association (**Supplementary**
362 **Figure 3** and **Supplementary Table 2**). Because the smoking behavior phenotype (captured as
363 cigarettes per day [CPD]) was not available in all CARDIoGRAMplusC4D studies, we used the top
364 variant previously reported for CPD¹⁰ at each locus (**Supplementary Figure 4**).

365 Analyses of the APOE locus.

366 The effect of rs6857, the lead CHD variant at the *APOE* locus, was similar in “ever-
367 smokers” compared to “never-smokers” (**Supplementary Table 3**). Specifically, the CHD OR for the
368 T allele at rs6857 was found to be 1.10 (P-value 7.93×10^{-4}) in “never-smokers” (12,159 CHD cases
369 and 22,932 controls) which was quantitatively similar to the CHD OR of 1.09 (P-value: 8.68×10^{-5})
370 observed in “ever-smokers” (23,753 CHD cases and 24,019 controls) (interaction P-value: 0.76)
371 (**Supplementary Figure 5a**). Investigation in the PROMIS study of the *APOE* epsilon genotypes
372 yielded consistent findings; the OR for CHD among $\epsilon 4$ carriers in “never-smokers” was 1.13
373 compared to the CHD OR of 1.07 observed in “ever-smokers” (interaction P-value: 0.82)
374 (**Supplementary Figure 5a**).

375 Novel gene-smoking interaction effects on CHD at chromosome 15q25.1

376 Of the 50 loci, we identified effect-modification by “ever-smoking” status on CHD risk for the
377 lead variants at two distinct loci, rs7178051, in proximity of *ADAMTS7* (an established CHD locus),
378 and rs1051730, in proximity of *CHRNA4-A3-A5* (an established smoking behavior locus)
379 (**Supplementary Table 3**). Although associated with different traits and located in distinct LD blocks,
380 these two variants reside ~224 KBs apart on chr.15q25.1 and are in weak linkage disequilibrium
381 (LD) ($r^2 = 0.22$), raising the question of whether these two chr.15q25.1 gene-smoking interactions on
382 CHD are independent of each other.

383 At the *ADAMTS7* CHD locus, the T allele at the rs7178051 variant was found to be more
384 strongly and inversely associated with CHD risk in never-smokers (OR: 0.88; P-value: 7.02×10^{-16})
385 compared to a much weaker effect in ever-smokers (OR: 0.95; P-value: 8.64×10^{-4}) (P-value of
386 interaction: 8.57×10^{-5}) (**Table 1**). Thus, the protective impact of the rs7178051 T allele observed in
387 never-smokers was halved in people who smoked (**Figure-1**). This difference is not related to power
388 differences within strata because for this variant, there were less data available in the never-smoking
389 group (21,232 CHD cases and 38,713 controls) compared to the ever-smoking group (39,585 CHD
390 cases and 40,749 controls). There was no substantial evidence of heterogeneity for the interaction
391 beta across the participating studies (Heterogeneity chi-squared = 36.23 (d.f. = 25); P-value for the
392 χ^2 test of heterogeneity = 0.06; $I^2 = 31.0\%$; tau-squared ($\tau^2 = 0$). We further conducted sensitivity
393 analyses using a random effect model; the results remained unchanged and the interaction beta
394 remained significant (**Supplementary Figure 5b**). Although the frequency of rs7178051 was 39% in
395 Europeans compared to 28% in South Asians, further analyses stratified by ancestry (i.e., European
396 versus non-Europeans) showed similar results (**Supplementary Figure 5c**). Other variants
397 discovered through prior CHD GWAS at the *ADAMTS7* locus (e.g., rs7173743, rs4380028,
398 rs3825807) were in moderate to high LD ($r^2 > 0.50$) with rs7178051 and were also found to display a
399 similar pattern of gene-smoking interaction effects (**Table 1**).

400 At the *CHRNA4-A3-A5* smoking locus, the A allele at the rs1051730 variant had an inverse
401 trend (not significant after adjustment) of association with CHD in never-smokers (OR: 0.96; P-value:
402 1.56×10^{-2}) and a positive trend (not significant after adjustment) in ever-smokers (OR: 1.03; P-value:
403 1.53×10^{-2}) (P-value of interaction: 2.37×10^{-4}) (**Table 1 and Supplementary Table 3**). For this
404 variant, data on 20,559 CHD cases and 38,198 controls were available in the never-smoking group
405 whereas 38,923 CHD cases and 40,371 controls were available in the ever-smoking group. Similar
406 gene-smoking interaction patterns were observed for other variants (e.g., rs2036527, rs8034191)
407 that have been previously reported for CPD behavior at the *CHRNA4-A3-A5* gene cluster (**Table 1**).

408 Further interrogation of the chr15q21.1 region encompassing rs7178051 and rs1051730
409 across three distinct LD blocks (**Figure 1**) revealed multiple additional variants for which we
410 observed gene-smoking interactions in CHD (**Table 1** and **Figure 1**). Indeed, several SNPs (e.g.,
411 rs7178051, rs10083696, rs7176187, rs6495335, rs4887077) had genome-wide significant
412 associations with CHD in “never-smokers” but had weaker and less significant associations with
413 CHD in “ever-smokers” (**Figure 1**). Alleles clustered specifically around *ADAMTS7* rather than at the
414 *CHRNA4-A3-A5* genes appear to be protective of CHD in “never-smokers” but have attenuated
415 protective effects in “ever-smokers” (**Figure 2**).

416 Conditional analyses

417 To investigate the possibility that the two chr.15q25.1 gene-smoking interactions might
418 represent a single interaction locus across the entire region we undertook an approximate
419 conditional and joint analyses¹⁷⁻²² using summary data derived from CARDIoGRAMplus4D for CHD
420 and from the TGC for smoking behavior. In-addition to rs7178051, we identified one other variant,
421 rs11072794 in low LD with rs7178051 ($r^2=0.20$) that was associated independently with CHD
422 (**Figure 3a**; red triangles) (**Figure 3b & Supplementary Figure 6b**; red triangles). We also
423 confirmed two variants (rs1051730 and rs684513) located in two different LD blocks that were
424 independently associated with smoking behavior in the TGC data¹⁰ (**Figure 3d & Supplementary**
425 **Figure 6b**; grey circles).

426 In analyses of the CHD variants, both rs7178051 and rs11072794 remained strongly
427 associated with CHD after adjusting for the top CPD variants (rs1051730 and rs684513) (**Figure 3d**,
428 red triangles) whereas their weak association with CPD was lost after adjusting for the top CPD
429 variants (**Figure 3d**; grey circles); e.g., the P-value for rs7178051 association with CPD was 1×10^{-5}
430 in unadjusted analyses but attenuated to 0.55 after adjusting for rs1051730 and rs684513. In
431 analyses of the CPD variants, both rs1051730 and rs684513 remained strongly associated with CPD
432 after adjusting for the top CHD variants (rs7178051 and rs11072794) (**Figure 3b**, grey circles)
433 whereas their weak association with CHD was lost after adjusting for the top CHD variants (**Figure**
434 **3b**, red triangles). As expected, conditional analyses that included all four of these variants resulted
435 in a null association of the region with both CHD and CPD (**Supplementary Figure 6b**). To
436 underscore the validity of the conditional approach using summary data, we used individual
437 participant data from an expanded PROMIS sample involving 9,025 MI cases and 8,506 controls.
438 We found that the OR conferred by allelic variation at rs7178051 remained associated with MI risk
439 independent of the two CPD variants (rs1051730 and rs684513) and rs11072794 (the second CHD

440 SNP) (**Supplementary Figure 6c**). Conversely, the apparent effect of allelic variation at rs1051730
441 (the top CPD variant) on CHD risk was lost when we adjusted for the other three variants,
442 rs7178051, rs11072794 and rs684513 (**Supplementary Figure 6c**).

443 Next, using summary level data we examined the association of each of these four variants
444 with CHD risk separately in “ever-smokers” and “never-smokers” while mutually adjusting for the
445 other three variants (**Figure 4 & Supplementary Figure 7**). In these analyses, only allelic variation
446 at rs7178051 was found to have independent genome-wide significant effects on CHD in never-
447 smokers. rs7178051 was also the only one of these four variants with significant differences in the
448 effect estimate for gene-CHD associations between the two smoking groups (P-value for the χ^2 test
449 of heterogeneity: 5.4×10^{-5}) after adjusting for the effects of other variants (rs11072794, rs1051730
450 and rs684513). These conditional analyses suggest that (a) variants located near the *ADAMTS7*
451 gene but not *CHRNA4-A3-A5* genes have independent effects on CHD, (b) a single independent
452 gene-smoking interaction signal for CHD exists on chr.15q.25.1 which is localized at the *ADAMTS7*
453 CHD locus (marked by rs7178051) and (c) an apparent interaction signal observed at the nearby
454 *CHRNA4-A3-A5* CPD locus (marked by rs1051730) is not independent of the *ADAMTS7*
455 (rs7178051) interaction signal.

456 To assess the robustness of conditional analyses methodology that uses summary level data
457 (i.e., GCTA)¹⁷⁻²², we conducted sensitivity analyses in the PROMIS dataset (9,025 MI cases and
458 8,506 controls). We assessed the association of rs7178051 (top CHD SNP) and rs1051730 (top
459 CPD SNP) after mutually adjusting for each other by conducting (i) standard logistic regression using
460 individual participant data and (ii) summary level data in PROMIS using the GCTA method
461 (**Supplementary Table 4**). The top CHD SNP was found associated with CHD risk in PROMIS
462 independent of the top CPD variant using both the methods, in-contrast the effect on CHD of the top
463 CPD SNP attenuated sharply when adjusted for the top CHD SNP – the effect estimates obtained
464 using the two methods were very similar (**Supplementary Table 4**).

465 Finally, to further demonstrate that the gene-smoking interaction effect in CHD at rs7178051 is
466 independent of the *CHRNA4-A3-A5* CPD locus, we conducted sensitivity analyses in the PROMIS
467 study by restricting our gene-environment interaction analysis to subjects who do not carry the minor
468 alleles of rs1051730 and rs684513 (the two SNPs associated with CPD) at the *CHRNA4-A3-A5*
469 locus. The T allele at the rs7178051 variant was associated with CHD only in never-smokers (OR:
470 0.88; P-value: 0.01) compared to a weaker and non-significant association in ever-smokers (OR:
471 0.94; P-value: 0.21) (**Supplementary Table 5**). The effect estimates obtained in each of the

472 categories defined by smoking status in PROMIS were similar to the effect estimates obtained in our
473 overall meta-analyses that utilized data in all participants (**Supplementary Table 5**).

474 Analysis of eQTLs and regulatory features at the chr15q25.1 gene-smoking interaction locus.

475 We mined publicly available eQTL data from the HapMap consortium,¹¹ GTEx consortium²³
476 and the MuTHER consortium²⁴ as well as data from 147 HAOEC lines²⁵ to examine the association
477 between mRNA expression of *ADAMTS7* and *CHRN* genes with CHD, CPD and gene-smoking
478 interaction SNPs at the chr15q25.1 locus. SNP-mRNA associations with p-values <0.002 (correction
479 for 20 tests) are presented (**Figure 5**). The top two CHD variants (rs7178051, rs11072794) are
480 associated with reduced *ADAMTS7* expression (e.g., rs11072794 $p=6.01 \times 10^{-21}$ in MuTHER LCL,
481 $n=850$; and rs7178051 $p=0.0029$ in HAOEC, $n=147$) but have no association with expression of
482 *CHRN* genes in any cell or tissue examined. In contrast, the top two CPD variants (rs1051730 and
483 rs684513) were associated with *CHRN* gene expression (e.g., rs1051730 $p=6.9 \times 10^{-7}$ for *CHRNA5* in
484 GTEx skeletal muscle and nerve tissue) but have no association with *ADAMTS7* in these cells or
485 tissues. These findings complement conditional analyses suggesting that gene-CHD and gene-
486 smoking interaction effects on CHD are likely mediated by *ADAMTS7* whereas the smoking behavior
487 effect appears to be mediated through the *CHRNA3-5* gene cluster.

488 In analysis of data from the ENCODE project²⁶ and for human aortic tissue in NIH
489 Roadmap Epigenomics project, *ADAMTS7* was associated with RNAseq reads and an active
490 transcription mark, H3K36me3, whereas *CHRN* genes had low/absent RNAseq reads and were
491 positive for repressive marks, H3K27me3 and H3K9me3 (**Supplementary Figure 8**). In HCASMC
492 ChIPseq data, rs7178051 the top CHD and gene-smoking CHD interacting SNP, is located in a
493 region with active regulatory marks H3K4me1 and H3K4me3 as well as transcription factor binding
494 site for TCF21, an important HCASMC transcription factor also associated with CAD. This ChIPseq
495 pattern was observed also in human aortic tissue (**Figure 6**). These regulatory data suggest active
496 transcription of *ADAMTS7*, but not *CHRN* genes, in vascular cells and aortic tissue and reveal that
497 rs7178051, the lead gene-smoking CHD interaction SNP, overlaps active transcription marks and
498 transcription factor binding regions in HCASMC.

499 *ADAMTS7* and *CHRNA3-5* expression in vascular cells and their response to cigarette smoke
500 extract

501 In order to explore which genes at the chr15q25.1 locus are expressed in CHD-relevant
502 vascular cells, we performed q-PCR of *ADAMTS7* and the *CHRNA3-5* genes in primary human

503 vascular cells and in the THP1 human monocyte cell line (**Supplementary Figure 9 & Figure 5**).
504 Whilst *ADAMTS7* mRNA was expressed abundantly in all vascular cell types, mRNA was below
505 detection or expressed at a very low level for each of the genes in the *CHRNA4-A3-A5* cluster in any
506 of these cell types (**Supplementary Figure 9**). Next, we explored the effect of cigarette smoke
507 extract on gene expression in HCASMC, a cell type of particular relevance to vascular responses to
508 cigarette smoke products^{31, 32} as well as to *ADAMTS7* vascular functions in atherosclerosis and
509 CHD.³³ In primary HCASMC, cigarette smoke extract exposure increased *ADAMTS7* mRNA levels
510 by over 2-fold (**Figure 5**) but did not affect expression of the *CHRN* genes (not shown). Thus, in
511 contrast to *CHRN* genes, *ADAMTS7* is both expressed and modulated by cigarette smoke extract in
512 CHD-relevant vascular cells providing biological support for *ADAMTS7*, but not *CHRN* genes, in the
513 gene-smoking interaction at chr15q25.1.

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529 **DISCUSSION**

530 We conducted a gene-environment interaction study at fifty loci associated with either CHD
531 or smoking behavior and found evidence of effect-modification of genotype-related CHD risk by
532 smoking-behavior at the chr.15q21.1 CHD locus. Specifically, we observed highly significant
533 attenuation of the cardio-protective effects associated with alleles at this locus in people who
534 smoked cigarettes. Conditional analyses identified an LD block located at the *ADAMTS7* gene that
535 accounted for both the main effect on CHD as well as the gene-smoking interactions in CHD. Data
536 from expression and cell studies support our genetic analysis, suggesting that the underlying
537 mechanism relates to genotype differences in the effect of smoking on expression of *ADAMTS7* in
538 vascular tissue.

539 Our findings have novel mechanistic and clinical implications. These human genomic data
540 provide new insights into the mechanism of CHD in cigarette smokers. Identification of gene-
541 smoking interaction at the chr15q21.1 locus suggests a specific role in smoking-related CHD for
542 *ADAMTS7* and its substrates, vascular matrix and vascular smooth muscle cell biology more
543 broadly. Such insights can help to prioritize translational strategies for smoking-related CHD and
544 present opportunities to study lifestyle interventions and pharmacological strategies to lower CHD in
545 individuals who smoke cigarettes. Thus, inhibition of *ADAMTS7* represents a novel potential
546 therapeutic strategy for CHD that may have particular benefits in individuals who smoke cigarettes.
547 All smokers should receive counseling for smoking cessation yet such broad public health strategies
548 have failed to reach or impact smoking behavior in a large portion of nicotine-addicted individuals.
549 Our data provides a human genomic context for consideration of targeting specific genetically at-risk
550 individuals via intensified preventive strategies and development of novel pharmacological
551 treatments.

552 Our study also represents a realistic strategy for performing gene-environment interaction
553 studies using contemporary genetic data. We illustrate that identifying joint effects of genetic and
554 lifestyle factors in CHD requires very large sample sizes, yet such analyses are biologically
555 informative when studies are adequately powered. In this context, an important observation in our
556 large sample is the lack of effect modification by smoking behavior on CHD at the *APOE* locus, a
557 previously reported smoking interaction locus.¹²⁻¹⁴ This finding is consistent with a recent meta-
558 analysis that found no evidence of effect modification by smoking for *APOE* genotypes and CHD
559 risk.³⁴ These studies raise concerns that most prior gene-environment interactions studies in CHD
560 have been prone to the same biases (i.e., limited statistical power and false positive associations) as

561 candidate gene studies investigating main effects in the pre-GWAS era. The present study differs
562 from previous studies by being much larger and, importantly, it includes genomic and functional
563 follow-up data supporting the plausibility of the observed gene-environment interaction.

564 *ADAMTS7* (or the A disintegrin and metalloproteinase with thrombospondin motifs-7) is a
565 member of the ADAMTS family of secreted zinc metalloproteases.^{35, 36} We previously discovered
566 and replicated genetic variation at the *ADAMTS7* locus in association with coronary atherosclerosis
567 and MI.⁷⁻⁹ Both *in vivo* and *in vitro* studies suggest that *ADAMTS7* modulates VSMC phenotype
568 switching and migration and that this may be mediated via cartilage oligomeric matrix protein
569 (COMP) or thrombospondin-1 (TSP-1),^{32,33} i.e. putative *ADAMTS7* substrates expressed in vascular
570 tissue. Genetic variation at *ADAMTS7*, however, has no relationship with traditional risk factors or
571 mechanistic biomarkers; hence the directional impact of *ADAMTS7* expression on CHD risk and the
572 underlying biological mechanisms have been unclear.³²

573 Our gene-smoking interaction analyses provide novel insights into the directional impact of
574 the *ADAMTS7* locus on CHD, the underlying mechanisms of CHD in smokers, and how such
575 findings ultimately might translate towards achieving health benefits in society. Our human eQTL
576 interrogations reveal that common alleles that relate to lower CHD risk at the *ADAMTS7* locus are
577 also associated with reduced *ADAMTS7* expression, implying an atherogenic role of the gene. This
578 is supported by our recent *in vivo* experimental studies; *Adamts7* deficiency protected against diet-
579 induced atherosclerosis in both the *Ldlr*^{-/-} and *ApoE*^{-/-} mouse models, reduced neointima formation
580 following arterial injury, and decreased VSMC migration *in vitro*.³³ In our smoking-stratified analyses,
581 we observed CHD protective effect which was attenuated in smokers. Thus, smoking exposure may
582 overcome the genetic effect of protective alleles that act by reducing *ADAMTS7* expression. Such a
583 possibility is supported by our HCASMC data that reveals increased *ADAMTS7* expression in
584 HCASMC exposed to cigarette smoke extract. These human genome-smoking studies are the first to
585 implicate a specific locus as causal in mediating increased risk of CHD in smokers. Additional
586 translational studies are needed to establish the precise mechanisms of atheroprotection for alleles
587 at the *ADAMTS7* locus, how cigarette smoking impacts these genetic effects, and whether deletion
588 or inhibition of *ADAMTS7* *in vivo* attenuates the specific acceleration of atherosclerosis conferred by
589 cigarette smoking.

590 Strengths and limitations of our study merit consideration. This is a large study that
591 conducted gene-smoking interaction analyses in CHD by using GWAS data. We observed
592 substantial heterogeneity across study samples in our initial quality control analyses of “ever-

593 smoking” status with CHD risk. This is similar, however, to the heterogeneity reported in a recent
594 meta-analysis that pooled risk ratios from all the past prospective studies with information on
595 association of “ever-smoking” with incident CHD events.⁵ We recognize that other smoking related
596 phenotypes are important e.g., “current smoking” may have a more pronounced role than “ever-
597 smoking” in plaque rupture and thrombosis in patients with MI. We were however unable to
598 distinguish between “former” versus “current” smokers within “Ever Smokers” in our current
599 analyses; furthermore we were not able to analyze graded exposure to cigarette smoking such as
600 “pack-years”. Given the use of multiple studies and meta-analyses of data, we used only one
601 analytical approach to investigate gene-smoking interactions. This approach, however, was feasible
602 and powerful in this large-scale consortium setting. While we used a fixed effects approach in our
603 meta-analyses, a random effects meta-analysis yielded qualitatively similar results (data not shown).
604 The lack of replication is partially offset by a large sample size, consistency across study cohorts
605 and racial groups and supplemental genomic and experimental evidence supporting biological
606 plausibility. This approach is also consistent with recent recommendations³⁷ which favor use of a
607 powerful discovery experiment using all data rather than reducing power by splitting available
608 dataset for discovery and validation. While our *in vitro* studies support a role for *ADAMTS7* in the
609 gene-smoking interaction, it will be important to confirm that *Adamts7* deficiency protect against
610 cigarette-smoke acceleration of atherosclerosis in rodent models.

611 Our interaction analyses, conditional analyses, eQTL interrogations and cell studies
612 suggest that *ADAMTS7*, but not the *CHRNA4-A3-A5* gene cluster, is likely causal at 15q21.1 for
613 gene-smoking interaction effects in CHD. Yet, analyses are not definitive. Although top interacting
614 SNPs and CHD SNPs (e.g., rs7178051) were associated with *ADAMTS7*, but not *CHRNA4-A3-A5*,
615 expression in LCLs, large-scale eQTL data and allele specific expression data (e.g., via RNA
616 sequencing) are not available for vascular tissues limiting causal inference. In our small HCAEC
617 datasets, we did however find that alleles at rs7178051 associate with *ADAMTS7* expression but not
618 with any *CHRNA4-A3-A5* genes suggesting, at least in one vascular cell type, that the gene-smoking
619 interaction is mediated via *ADAMTS7*.

620

621

622

623

624

625 **Conclusions**

626 We provide novel evidence for allelic variation exhibiting gene-smoking interaction in CHD
627 at the chr.15q21.1 locus. The protective effect conferred by variation at this locus in never-smokers
628 is markedly attenuated in people who are ever-smokers. Stepwise conditional analyses, gene
629 expression data in vascular cells, eQTL interrogation, and cigarette smoke extract exposure in
630 HCASMC suggest that *ADAMTS7* accounts for both the gene-smoking interaction in CHD and the
631 CHD main effect on chr.15q21.1. Our findings reveal interactions of genetic variants and key lifestyle
632 determinants in the etiology of CHD, provide new insights into the potential mechanisms of CHD in
633 cigarette smokers, and facilitate precision medicine advances in cigarette-smoking related CHD. Our
634 work motivates future large-scale studies investigating joint effects of genes and environment in
635 CHD using existing complex-disease consortia datasets and genome-wide discovery approaches.
636 This will provide opportunities to detect additional and novel loci displaying gene-environment
637 interactions revealing genetic contexts for targeting intensive lifestyle interventions and novel
638 therapeutics.

639

640

641

642

643

644

645

646

647

648

649

650

651

652

653

654

655

656 **Collaborators:**

657 Stanley L. Hazen MD,¹ W.H. Wilson Tang MD,¹ Perttu P Salo PhD,^{2,3} Marja-Liisa Lokki PhD,⁴
658 Markku S Nieminen PhD,⁵ Antti-Pekka Sarin MSc,^{2,6} Alun Evans MSc,⁷ Jean Ferrières MD,⁸ Jarmo
659 Virtamo PhD,² Frank Kee PhD,⁹ David-Alexandre Trégouët PhD,¹⁰ Dominique Arveiler PhD,¹¹
660 Philippe Amouyel PhD,¹² Paolo Brambilla PhD,¹³ Annette Peters PhD,¹⁴ Melanie Waldenberger
661 PhD,^{14,15} Giovanni Veronesi PhD,¹⁶ Giancarlo Cesana PhD,¹⁷ Satu Männistö PhD,² Pekka Jousilahti
662 PhD,² Antti M Jula PhD,² Kennet Harald PhD,² Albert Hofman PhD,¹⁸ Oscar H. Franco PhD,¹⁸ Andre
663 G. Uitterlinden PhD.¹⁹

664 ¹ Departments of Cardiovascular Medicine and Cellular and Molecular Medicine, Cleveland Clinic, Cleveland,
665 OH 44195

666 ² National Institute for Health and Welfare, Helsinki, Finland

667 ³ University of Helsinki, Institute for Molecular Medicine, Finland (FIMM)

668 ⁴ Transplantation Laboratory, Haartman Institute, Helsinki, Finland

669 ⁵ Heart and Lung Center HUH, Helsinki University Hospital and Helsinki University, Finland

670 ⁶ University of Helsinki, Institute for Molecular Medicine, Finland (FIMM)

671 ⁷ Centre for Public Health, The Queen's University of Belfast, Belfast, Northern Ireland

672 ⁸ INSERM UMR 1027, Department of Cardiology, Toulouse University School of Medicine, Rangueil Hospital,
673 31059 Toulouse, France

674 ⁹ UKCRC Centre of Excellence for Public Health (NI), Queens University of Belfast, Northern Ireland

675 ¹⁰ Institut National pour la Santé et la Recherche Médicale (INSERM), Unité Mixte de Recherche en Santé

676 (UMR_S) 1166, F-75013, Paris, France AND Sorbonne Universités, Université Pierre et Marie Curie, Paris,
677 France; and Institute for Cardiometabolism and Nutrition, Paris, France

678 ¹¹ Department of Epidemiology and Public Health, EA3430, University of Strasbourg, Faculty of Medicine,
679 Strasbourg, France

680 ¹² Institut Pasteur de Lille, INSERM U744, Université Lille Nord de France, F-59000 Lille, France

681 ¹³ Laboratory Medicine, Hospital of Desio, Department of Health Sciences, University of Milano, Bicocca, Italy

682 ¹⁴ Institute of Epidemiology II, Helmholtz Zentrum München, Neuherberg, Germany AND German Center for
683 Cardiovascular Disease Research (DZHK e.V.), Munich, Germany Melanie Waldenberger

684 ¹⁵ Research Unit of Molecular Epidemiology, Helmholtz Zentrum München, Neuherberg, Germany

685 ¹⁶ EPIMED Research Center, Department of Clinical and Sperimental Medicine, University of Insubria, Varese,
686 Italy

687 ¹⁷ University of Milano, Bicocca, Italy

688 ¹⁸ Department of Epidemiology, Erasmus University Medical center, Rotterdam, The Netherlands

689 ¹⁹ Department of Internal Medicine, Erasmus University Medical center, Rotterdam, The Netherlands

690

691

692

693

694

695

696

697

698

699

700

701

702 **Sources of Funding**

703 Dr. Saleheen has received funding from the National Institutes of Health, the Fogarty International,
704 the Wellcome Trust, the British Heart Foundation, Pfizer, Genentech, Regeneron and Eli Lilly
705 pharmaceuticals. This work was supported in part by R01-HL-111694 and K24-HL-107643 from the
706 National Institutes of Health to Dr. Reilly.

707
708 **PROMIS.** Genotyping in PROMIS was funded by the Wellcome Trust, UK and Pfizer. Fieldwork in
709 the PROMIS study has been supported through funds available to investigators at the Center for
710 Non-Communicable Diseases, Pakistan and the University of Cambridge, UK.

711
712 **EPIC-CVD Consortium.** CHD case ascertainment and validation, genotyping, and clinical chemistry
713 assays in EPIC-CVD were principally supported by grants awarded to the University of Cambridge
714 from the European Union (EU) Framework Programme 7 (HEALTH-F2-2012-279233), the United
715 Kingdom (UK) Medical Research Council (G0800270) and British Heart Foundation (SP/09/002), the
716 UK National Institute for Health Research Cambridge Biomedical Research Centre, and the
717 European Research Council (268834). Scientists at the EPIC-CVD Coordinating Centre have also
718 been supported by grants from the US National Institutes of Health, Merck, Novartis,
719 GlaxoSmithKline, and Pfizer.

720
721 **WTCCC.** Recruitment of the WTCCC CAD cases was funded by the British Heart Foundation.
722 Collection of controls and genotyping was funded by the Wellcome Trust. Chris Nelson and Nilesh
723 Samani are funded by the British Heart Foundation. Nilesh Samani is a National Institute for Health
724 Research (NIHR) Senior Investigator.

725
726 **SCARF-SHEEP.** The investigators would like to acknowledge the Swedish Heart-Lung Foundation,
727 the Swedish Research Council, the Strategic Cardiovascular Programme of Karolinska Institutet and
728 the Stockholm County Council, the Strategic support for epidemiological research at Karolinska
729 Institutet and the Stockholm County Council. Rona J Strawbridge is supported by Strategic Research
730 Support (SRP) Diabetes Program at Karolinska Institutet.

731 **CARDIOGENICS and THESIAs.** Professor Deloukas' work forms part of the research themes
732 contributing to the translational research portfolio of Barts Cardiovascular Biomedical Research Unit
733 which is supported and funded by the National Institute for Health Research. Analysis was supported
734 by British Heart Foundation (BHF) grant (Deloukas) RG/14/5/30893. Professor Schunkert was

735 supported by grants from the Fondation Leducq (*CADgenomics*: Understanding CAD Genes,
736 12CVD02), the German Federal Ministry of Education and Research (BMBF) within the framework of
737 the *e:AtheroSysMed* (e:Med) research and funding concept (grant 01ZX1313A-2014), and the
738 European Union Seventh Framework Programme FP7/2007-2013 under grant agreement n°
739 HEALTH-F2-2013-601456 (*CVgenes-at-target*). Further grants were received from the Deutsche
740 Forschungsgemeinschaft (DFG) as part of the Sonderforschungsbereich CRC 1123 (B2). Thorsten
741 Kessler MD was supported by a German Centre for Cardiovascular Research (DZHK) Rotation
742 Grant.

743 **Rotterdam Study.** The Rotterdam Study is supported by the Erasmus Medical Center and Erasmus
744 University Rotterdam; the Netherlands Organization for Scientific Research; the Netherlands
745 Organization for Health Research and Development (ZonMw); the Research Institute for Diseases in
746 the Elderly; the Ministry of Education, Culture and Science; the Ministry of Health, Welfare and
747 Sports; the European Commission; and the Municipality of Rotterdam. Support for genotyping was
748 provided by the Netherlands Organisation of Scientific Research (NOW) Investments (No.
749 175.010.2005.011, 911-03-012), the Research Institute for Diseases in the Elderly (014-93-015;
750 RIDE2), the Netherlands Genomics Initiative/Netherlands Consortium for Healthy Aging project No.
751 050-060-810. Abbas Dehghan is supported by an NWO grant (veni, 916.12.154) and the EUR
752 Fellowship. Oscar H. Franco works in ErasmusAGE, a center for aging research across the life
753 course funded by Nestlé Nutrition (Nestec Ltd.), Metagenics Inc., and AXA. Nestlé Nutrition (Nestec
754 Ltd.), Metagenics Inc., and AXA had no role in design and conduct of the study; collection,
755 management, analysis, and interpretation of the data; and preparation, review, or approval of the
756 manuscript.

757
758 **Expression studies.** These studies were supported in part by a Transatlantic Network of Excellence
759 grant 10CVD03 from the Fondation Leducq. Analysis of expression quantitative trait loci (eQTL) in
760 endothelial cells was supported by a Transatlantic Networks of Excellence Award (12CVD02) from
761 Fondation Leducq (to Dr. Jake Lusis and team)

762 **Diabetes Heart Study.** This study was supported in part by NIH R01 HL67348, NIH R01 HL092301,
763 and NIH R01 NS058700 to Donald W. Bowden.

764 **Cleveland Clinic Study.** This study was supported in part by NIH grants R01ES021801,
765 3R01ES021801-03S1, and R01ES025786.

766 **Family Heart Study (FamHS).** The FamHS is funded by NIH grant R01HL117078 grant.

767 **MORGAM.** This work has been sustained by the MORGAM Project's funding: European Union FP 7
768 projects ENGAGE (HEALTH-F4-2007-201413), CHANCES (HEALTH-F3-2010-242244) and
769 BiomarCaRE (278913). This funding has supported central coordination, workshops and part of the
770 activities of the MORGAM Data Centre, at THL in Helsinki, Finland. MORGAM Participating Centres
771 are funded by regional and national governments, research councils, charities, and other local
772 sources. The PRIME Study was supported by grants from Inserm, Merck Sharp and Dohme-Chibret
773 Laboratory, the French Research Agency and the Foundation Heart and Arteries. We also thank the
774 following organisations that allowed the recruitment of participants for the PRIME: the health
775 screening centres organised by the SocialSecurity of Lille (Institut Pasteur), Strasbourg, Toulouse,
776 and Tourcoing; the occupational medicine services of Haute-Garonne and of the Urban Community
777 of Strasbourg; the Association Inter-entreprises des Services Médicaux du Travail de Lille et
778 environs; the Comité pour le Développement de la Médecine du Travail; the Mutuelle Générale des
779 Postes, Télégraphes et Téléphones du Bas-Rhin; the Laboratoire d'Analyses de l'Institut de Chimie
780 Biologique de la Faculté de Médecine de Strasbourg. We also gratefully acknowledge the teams of
781 the Lille, Strasbourg and Toulouse centres for their dedicate work and relentness energy in following
782 up their cohorts; the contribution of the members of the event validation committees : L Guize; C
783 Morrison; M-T Guillanneuf; and M Giroud and the Alliance Partnership Programme for its financial
784 support. The KORA study was initiated and financed by the Helmholtz Zentrum München – German
785 Research Center for Environmental Health, which is funded by the German Federal Ministry of
786 Education and Research (BMBF) and by the State of Bavaria. KK was supported by the Orion-
787 Farnos Research Foundation and Academy of Finland (grant number 250207).

788 **DILGOM.** This work was enabled through a grant #139635 from the Academy of Finland and a grant
789 from the Finnish Foundation for Cardiovascular Research. The DILGOM project is also supported by
790 the Academy of Finland (grant numbers 136895 and 263836). We are grateful for the THL DNA
791 laboratory for its skillful work to produce the DNA samples used in this study. We also acknowledge
792 the Academy of Finland (136895 and 263836), Funding from Academy of Finland, grant number:
793 118065, and Juho Vainio Foundation.

794

795

796

797

798

799 **Acknowledgements**

800 We would like to thank the CARDIoGRAMplusC4D consortium, the EPIC-CVD and the PROMIS
801 study for contributing data.

802 **PROMIS.** We also acknowledge the contributions made by the following: Zeeshan Ozair, Usman
803 Ahmed, Abdul Hakeem, Hamza Khalid, Naeem Khan, Sadiq Khan, Ayaz Ali, Madad Ali, Saeed
804 Ahmed, Muhammad Waqar Khan, Muhammad Razaq Khan, Abdul Ghafoor, Mir Alam, Riazuddin,
805 Muhammad Irshad Javed, Abdul Ghaffar, Tanveer Baig Mirza, Muhammad Shahid, Jabir Furqan,
806 Muhammad Iqbal Abbasi, Tanveer Abbas, Rana Zulfiqar, Muhammad Wajid, Irfan Ali, Muhammad
807 Ikhtlaq, Danish Sheikh, Muhammad Imran, Nadeem Sarwar, Adam Butterworth, Matthew Walker and
808 Hannah Lombardi, Shahid Abbas, Faisal Majeed, Saba Akhtar, Nadeem Qamar, Khan Shah Zaman,
809 Zia Yaqoob, Tahir Saghir, Syed Nadeem Hasan Rizvi, Anis Memon, Nadeem Hayyat Mallick,
810 Mohammad Ishaq, Syed Zahed Rasheed, Fazal-ur-Rehman Memon, Khalid Mahmood, and
811 Naveeduddin Ahmed.

812 **EPIC-CVD.** We thank all EPIC participants and staff for their contribution to the study, the laboratory
813 teams at the Medical Research Council Epidemiology Unit for sample management and Cambridge
814 Genomic Services for genotyping, Sarah Spackman for data management, and the team at the
815 EPIC-CVD Coordinating Centre for study coordination and administration.

816 **Disclosures**

817 Dr. Saleheen has received funding from Pfizer, Genentech, Regeneron and Eli Lilly pharmaceuticals.

818

819

820

821

822

823

824

825

826

827

828

829

830

831 **References**

- 832 (1) Mangino M, Spector T. Understanding coronary artery disease using twin studies. *Heart*.
833 2013;99:373-375.
- 834 (2) Lusia AJ, Mar R, Pajukanta P. Genetics of atherosclerosis. *Annu Rev Genomics Hum Genet*.
835 2004;5:189-218.
- 836 (3) Teo KK, Ounpuu S, Hawken S, Pandey MR, Valentin V, Hunt D, Diaz R, Rashed W, Freeman R,
837 Jiang L, Zhang X, Yusuf S; INTERHEART Study Investigators. Tobacco use and risk of myocardial
838 infarction in 52 countries in the INTERHEART study: a case-control study. *Lancet*. 2006;368:647-
839 658.
- 840 (4) Doll R, Peto R, Boreham J, Sutherland I. Mortality in relation to smoking: 50 years' observations
841 on male British doctors. *BMJ*. 2004;328:1519.
- 842 (5) Huxley RR, Woodward M. Cigarette smoking as a risk factor for coronary heart disease in women
843 compared with men: a systematic review and meta-analysis of prospective cohort studies. *Lancet*.
844 2011;378:1297-1305.
- 845 (6) Ezzati M, Lopez AD. Estimates of global mortality attributable to smoking in 2000. *Lancet*.
846 2003;362:847-852.
- 847 (7) CARDIoGRAMplusC4D Consortium., Deloukas P, Kanoni S, Willenborg C, Farrall M, Assimes
848 TL, Thompson JR, Ingelsson E, Saleheen D, Erdmann J, Goldstein BA, Stirrups K, König IR, Cazier
849 JB, Johansson A, Hall AS, Lee JY, Willer CJ, Chambers JC, Esko T, Folkersen L, Goel A,
850 Grundberg E, Havulinna AS, Ho WK, Hopewell JC, Eriksson N, Kleber ME, Kristiansson K,
851 Lundmark P, Lyytikäinen LP, Rafelt S, Shungin D, Strawbridge RJ, Thorleifsson G, Tikkanen E, Van
852 Zuydam N, Voight BF, Waite LL, Zhang W, Ziegler A, Absher D, Altshuler D, Balmforth AJ, Barroso I,
853 Braund PS, Burgdorf C, Claudi-Boehm S, Cox D, Dimitriou M, Do R; DIAGRAM Consortium.;
854 CARDIOGENICS Consortium., Doney AS, El Mokhtari N, Eriksson P, Fischer K, Fontanillas P,
855 Franco-Cereceda A, Gigante B, Groop L, Gustafsson S, Hager J, Hallmans G, Han BG, Hunt SE,
856 Kang HM, Illig T, Kessler T, Knowles JW, Kolovou G, Kuusisto J, Langenberg C, Langford C,
857 Leander K, Lokki ML, Lundmark A, McCarthy MI, Meisinger C, Melander O, Mihailov E, Maouche S,
858 Morris AD, Müller-Nurasyid M; MuTHER Consortium., Nikus K, Peden JF, Rayner NW, Rasheed A,
859 Rosinger S, Rubin D, Rumpf MP, Schäfer A, Sivananthan M, Song C, Stewart AF, Tan ST,
860 Thorgeirsson G, van der Schoot CE, Wagner PJ; Wellcome Trust Case Control Consortium., Wells
861 GA, Wild PS, Yang TP, Amouyel P, Arveiler D, Basart H, Boehnke M, Boerwinkle E, Brambilla P,
862 Cambien F, Cupples AL, de Faire U, Dehghan A, Diemert P, Epstein SE, Evans A, Ferrario MM,
863 Ferrières J, Gauguier D, Go AS, Goodall AH, Gudnason V, Hazen SL, Holm H, Iribarren C, Jang Y,
864 Kähönen M, Kee F, Kim HS, Klopp N, Koenig W, Kratzer W, Kuulasmaa K, Laakso M, Laaksonen R,
865 Lee JY, Lind L, Ouwehand WH, Parish S, Park JE, Pedersen NL, Peters A, Quertermous T, Rader
866 DJ, Salomaa V, Schadt E, Shah SH, Sinisalo J, Stark K, Stefansson K, Trégouët DA, Virtamo J,
867 Wallentin L, Wareham N, Zimmermann ME, Nieminen MS, Hengstenberg C, Sandhu MS, Pastinen
868 T, Syvänen AC, Hovingh GK, Dedoussis G, Franks PW, Lehtimäki T, Metspalu A, Zalloua PA,
869 Siegbahn A, Schreiber S, Ripatti S, Blankenberg SS, Perola M, Clarke R, Boehm BO, O'Donnell C,
870 Reilly MP, März W, Collins R, Kathiresan S, Hamsten A, Kooner JS, Thorsteinsdottir U, Danesh J,
871 Palmer CN, Roberts R, Watkins H, Schunkert H, Samani NJ. Large-scale association analysis
872 identifies new risk loci for coronary artery disease. *Nat Genet*. 2013;45:25-33.

873 (8) Coronary Artery Disease (C4D) Genetics Consortium. A genome-wide association study in
874 Europeans and South Asians identifies five new loci for coronary artery disease. *Nat Genet.*
875 2011;43:339-344.

876 (9) Schunkert H, König IR, Kathiresan S, Reilly MP, Assimes TL, Holm H, Preuss M, Stewart AF,
877 Barbalic M, Gieger C, Absher D, Aherrahrou Z, Allayee H, Altshuler D, Anand SS, Andersen K,
878 Anderson JL, Ardissino D, Ball SG, Balmforth AJ, Barnes TA, Becker DM, Becker LC, Berger K, Bis
879 JC, Boekholdt SM, Boerwinkle E, Braund PS, Brown MJ, Burnett MS, Buysschaert I; Cardiogenics.,
880 Carlquist JF, Chen L, Cichon S, Codd V, Davies RW, Dedoussis G, Dehghan A, Demissie S,
881 Devaney JM, Diemert P, Do R, Doering A, Eifert S, Mokhtari NE, Ellis SG, Elosua R, Engert JC,
882 Epstein SE, de Faire U, Fischer M, Folsom AR, Freyer J, Gigante B, Girelli D, Gretarsdottir S,
883 Gudnason V, Gulcher JR, Halperin E, Hammond N, Hazen SL, Hofman A, Horne BD, Illig T,
884 Iribarren C, Jones GT, Jukema JW, Kaiser MA, Kaplan LM, Kastelein JJ, Khaw KT, Knowles JW,
885 Kolovou G, Kong A, Laaksonen R, Lambrechts D, Leander K, Lettre G, Li M, Lieb W, Loley C, Lotery
886 AJ, Mannucci PM, Maouche S, Martinelli N, McKeown PP, Meisinger C, Meitinger T, Melander O,
887 Merlini PA, Mooser V, Morgan T, Mühleisen TW, Muhlestein JB, Münzel T, Musunuru K, Nahrstaedt
888 J, Nelson CP, Nöthen MM, Olivieri O, Patel RS, Patterson CC, Peters A, Peyvandi F, Qu L, Quyyumi
889 AA, Rader DJ, Rallidis LS, Rice C, Rosendaal FR, Rubin D, Salomaa V, Sampietro ML, Sandhu MS,
890 Schadt E, Schäfer A, Schillert A, Schreiber S, Schrezenmeir J, Schwartz SM, Siscovick DS,
891 Sivananthan M, Sivapalaratnam S, Smith A, Smith TB, Snoep JD, Soranzo N, Spertus JA, Stark K,
892 Stirrups K, Stoll M, Tang WH, Tennstedt S, Thorgeirsson G, Thorleifsson G, Tomaszewski M,
893 Uitterlinden AG, van Rij AM, Voight BF, Wareham NJ, Wells GA, Wichmann HE, Wild PS, Willenborg
894 C, Wittman JC, Wright BJ, Ye S, Zeller T, Ziegler A, Cambien F, Goodall AH, Cupples LA,
895 Quertermous T, März W, Hengstenberg C, Blankenberg S, Ouwehand WH, Hall AS, Deloukas P,
896 Thompson JR, Stefansson K, Roberts R, Thorsteinsdottir U, O'Donnell CJ, McPherson R, Erdmann
897 J; CARDIoGRAM Consortium., Samani NJ. Large-scale association analysis identifies 13 new
898 susceptibility loci for coronary artery disease. *Nat Genet* 2011;43:333-338.

899 (10) Tobacco and Genetics Consortium. Genome-wide meta-analyses identify multiple loci
900 associated with smoking behavior. *Nat Genet.* 2010;42:441-447.

901 (11) International HapMap 3 Consortium., Altshuler DM, Gibbs RA, Peltonen L, Altshuler DM, Gibbs
902 RA, Peltonen L, Dermitzakis E, Schaffner SF, Yu F, Peltonen L, Dermitzakis E, Bonnen PE,
903 Altshuler DM, Gibbs RA, de Bakker PI, Deloukas P, Gabriel SB, Gwilliam R, Hunt S, Inouye M, Jia X,
904 Palotie A, Parkin M, Whittaker P, Yu F, Chang K, Hawes A, Lewis LR, Ren Y, Wheeler D, Gibbs RA,
905 Muzny DM, Barnes C, Darvishi K, Hurler M, Korn JM, Kristiansson K, Lee C, McCarroll SA, Nemesh
906 J, Dermitzakis E, Keinan A, Montgomery SB, Pollack S, Price AL, Soranzo N, Bonnen PE, Gibbs
907 RA, Gonzaga-Jauregui C, Keinan A, Price AL, Yu F, Anttila V, Brodeur W, Daly MJ, Leslie S,
908 McVean G, Moutsianas L, Nguyen H, Schaffner SF, Zhang Q, Ghorji MJ, McGinnis R, McLaren W,
909 Pollack S, Price AL, Schaffner SF, Takeuchi F, Grossman SR, Shlyakhter I, Hostetter EB, Sabeti
910 PC, Adebamowo CA, Foster MW, Gordon DR, Licinio J, Manca MC, Marshall PA, Matsuda I, Ngare
911 D, Wang VO, Reddy D, Rotimi CN, Royal CD, Sharp RR, Zeng C, Brooks LD, McEwen JE.
912 Integrating common and rare genetic variation in diverse human populations. *Nature.* 2010;467:52-
913 58.

914 (12) Humphries SE, Talmud PJ, Hawe E, Bolla M, Day IN, Miller GJ. Apolipoprotein E4 and coronary
915 heart disease in middle-aged men who smoke: a prospective study. *Lancet.* 2001;358:115-119.

916 (13) Grammer TB, Hoffmann MM, Scharnagl H, Kleber ME, Silbernagel G, Pilz S, Tomaschitz A,
917 Lerchbaum E, Siekmeier R, März W. Smoking, apolipoprotein E genotypes, and mortality (the
918 Ludwigshafen Risk and Cardiovascular Health study). *Eur Heart J.* 2013;34:1298-1305.

919 (14) Gustavsson J, Mehlig K, Leander K, Strandhagen E, Björck L, Thelle DS, Lissner L, Blennow K,
920 Zetterberg H, Nyberg F. Interaction of apolipoprotein E genotype with smoking and physical inactivity
921 on coronary heart disease risk in men and women. *Atherosclerosis*. 2012;220:486-492.

922 (15) Saleheen D, Zaidi M, Rasheed A, Ahmad U, Hakeem A, Murtaza M, Kayani W, Faruqui A,
923 Kundi A, Zaman KS, Yaqoob Z, Cheema LA, Samad A, Rasheed SZ, Mallick NH, Azhar M, Jooma
924 R, Gardezi AR, Memon N, Ghaffar A, Fazal-ur-Rehman, Khan N, Shah N, Ali Shah A, Samuel M,
925 Hanif F, Yameen M, Naz S, Sultana A, Nazir A, Raza S, Shazad M, Nasim S, Javed MA, Ali SS,
926 Jafree M, Nisar MI, Daood MS, Hussain A, Sarwar N, Kamal A, Deloukas P, Ishaq M, Frossard P,
927 Danesh J. The Pakistan Risk of Myocardial Infarction Study: a resource for the study of genetic,
928 lifestyle and other determinants of myocardial infarction in South Asia. *Eur J Epidemiol*.
929 2009;24:329-338.

930 (16) Teslovich TM, Musunuru K, Smith AV, Edmondson AC, Stylianou IM, Koseki M, Pirruccello JP,
931 Ripatti S, Chasman DI, Willer CJ, Johansen CT, Fouchier SW, Isaacs A, Peloso GM, Barbalic M,
932 Ricketts SL, Bis JC, Aulchenko YS, Thorleifsson G, Feitosa MF, Chambers J, Orho-Melander M,
933 Melander O, Johnson T, Li X, Guo X, Li M, Shin Cho Y, Jin Go M, Jin Kim Y, Lee JY, Park T, Kim K,
934 Sim X, Twee-Hee Ong R, Croteau-Chonka DC, Lange LA, Smith JD, Song K, Hua Zhao J, Yuan X,
935 Luan J, Lamina C, Ziegler A, Zhang W, Zee RY, Wright AF, Witteman JC, Wilson JF, Willemsen G,
936 Wichmann HE, Whitfield JB, Waterworth DM, Wareham NJ, Waeber G, Vollenweider P, Voight BF,
937 Vitart V, Uitterlinden AG, Uda M, Tuomilehto J, Thompson JR, Tanaka T, Surakka I, Stringham HM,
938 Spector TD, Soranzo N, Smit JH, Sinisalo J, Silander K, Sijbrands EJ, Scuteri A, Scott J,
939 Schlessinger D, Sanna S, Salomaa V, Saharinen J, Sabatti C, Ruukonen A, Rudan I, Rose LM,
940 Roberts R, Rieder M, Psaty BM, Pramstaller PP, Pichler I, Perola M, Penninx BW, Pedersen NL,
941 Pattaro C, Parker AN, Pare G, Oostra BA, O'Donnell CJ, Nieminen MS, Nickerson DA, Montgomery
942 GW, Meitinger T, McPherson R, McCarthy MI, McArdle W, Masson D, Martin NG, Marroni F,
943 Mangino M, Magnusson PK, Lucas G, Luben R, Loos RJ, Lokki ML, Lettre G, Langenberg C, Launer
944 LJ, Lakatta EG, Laaksonen R, Kyvik KO, Kronenberg F, König IR, Khaw KT, Kaprio J, Kaplan LM,
945 Johansson A, Jarvelin MR, Janssens AC, Ingelsson E, Igl W, Kees Hovingh G, Hottenga JJ, Hofman
946 A, Hicks AA, Hengstenberg C, Heid IM, Hayward C, Havulinna AS, Hastie ND, Harris TB,
947 Haritunians T, Hall AS, Gyllenstein U, Guiducci C, Groop LC, Gonzalez E, Gieger C, Freimer NB,
948 Ferrucci L, Erdmann J, Elliott P, Ejebe KG, Döring A, Dominiczak AF, Demissie S, Deloukas P, de
949 Geus EJ, de Faire U, Crawford G, Collins FS, Chen YD, Caulfield MJ, Campbell H, Burt NP,
950 Bonnycastle LL, Boomsma DI, Boekholdt SM, Bergman RN, Barroso I, Bandinelli S, Ballantyne CM,
951 Assimes TL, Quertermous T, Altshuler D, Seielstad M, Wong TY, Tai ES, Feranil AB, Kuzawa CW,
952 Adair LS, Taylor HA Jr, Borecki IB, Gabriel SB, Wilson JG, Holm H, Thorsteinsdottir U, Gudnason V,
953 Krauss RM, Mohlke KL, Ordovas JM, Munroe PB, Kooner JS, Tall AR, Hegele RA, Kastelein JJ,
954 Schadt EE, Rotter JI, Boerwinkle E, Strachan DP, Mooser V, Stefansson K, Reilly MP, Samani NJ,
955 Schunkert H, Cupples LA, Sandhu MS, Ridker PM, Rader DJ, van Duijn CM, Peltonen L, Abecasis
956 GR, Boehnke M, Kathiresan S. Biological, clinical and population relevance of 95 loci for blood lipids.
957 *Nature* 2010;466:707-713.

958 (17) Yang J, Ferreira T, Morris AP, Medland SE; Genetic Investigation of ANthropometric Traits
959 (GIANT) Consortium.; DIAbetes Genetics Replication And Meta-analysis (DIAGRAM) Consortium.,
960 Madden PA, Heath AC, Martin NG, Montgomery GW, Weedon MN, Loos RJ, Frayling TM, McCarthy
961 MI, Hirschhorn JN, Goddard ME, Visscher PM. Conditional and joint multiple-SNP analysis of GWAS
962 summary statistics identifies additional variants influencing complex traits. *Nat Genet*. 2012;44:369-
963 373.

964 (18) Yang J, Lee SH, Goddard ME, Visscher PM. Genome-wide complex trait analysis (GCTA): methods,
965 data analyses, and interpretations. *Methods Mol Biol*. 2013; 1019:215-236.

966 (19) Keller MF, Ferrucci L, Singleton AB. Genome-Wide Analysis of the Heritability of Amyotrophic Lateral
967 Sclerosis. *JAMA Neurol.* 2014;71:1123-1134.

968 (20) van Rheenen W, Shatunov A, Dekker AM, McLaughlin RL, Diekstra FP, Pulit SL, van der Spek RA,
969 Vösa U, de Jong S, Robinson MR, Yang J, Fogh I, van Doormaal PT, Tazelaar GH, Koppers M, Blokhuis
970 AM, Sproviero W, Jones AR, Kenna KP, van Eijk KR, Harschnitz O, Schellevis RD, Brands WJ, Medic J,
971 Menelaou A, Vajda A, Ticozzi N, Lin K, Rogelj B, Vrabec K, Ravnik-Glavač M, Koritnik B, Zidar J,
972 Leonardis L, Grošelj LD, Millicamps S, Salachas F, Meininger V, de Carvalho M, Pinto S, Mora JS,
973 Rojas-García R, Polak M, Chandran S, Colville S, Swingler R, Morrison KE, Shaw PJ, Hardy J, Orrell
974 RW, Pittman A, Sidle K, Fratta P, Malaspina A, Topp S, Petri S, Abdulla S, Drepper C, Sendtner M,
975 Meyer T, Ophoff RA, Staats KA, Wiedau-Pazos M, Lomen-Hoerth C, Van Deerlin VM, Trojanowski JQ,
976 Elman L, McCluskey L, Basak AN, Tunca C, Hamzeiy H, Parman Y, Meitinger T, Lichtner P, Radivojkov-
977 Blagojevic M, Andres CR, Maurel C, Bensimon G, Landwehrmeyer B, Brice A, Payan CA, Saker-Delye S,
978 Dürr A, Wood NW, Tittmann L, Lieb W, Franke A, Rietschel M, Cichon S, Nöthen MM, Amouyel P,
979 Tzourio C, Dartigues JF, Uitterlinden AG, Rivadeneira F, Estrada K, Hofman A, Curtis C, Blauw HM, van
980 der Kooij AJ, de Visser M, Goris A, Weber M, Shaw CE, Smith BN, Pansarasa O, Cereda C, Del Bo R,
981 Comi GP, D'Alfonso S, Bertolin C, Sorarù G, Mazzini L, Pensato V, Gellera C, Tiloca C, Ratti A, Calvo A,
982 Moglia C, Brunetti M, Arcuti S, Capozzo R, Zecca C, Lunetta C, Penco S, Riva N, Padovani A, Filosto M,
983 Muller B, Stuit RJ; PARALS Registry.; SLALOM Group.; SLAP Registry.; FALS Sequencing Consortium.;
984 SLAGEN Consortium.; NNIPPS Study Group., Blair I, Zhang K, McCann EP, Fifita JA, Nicholson GA,
985 Rowe DB, Pamphlett R, Kiernan MC, Grosskreutz J, Witte OW, Ringer T, Prell T, Stubendorff B, Kurth I,
986 Hübner CA, Leigh PN, Casale F, Chio A, Beghi E, Pupillo E, Tortelli R, Logroscino G, Powell J, Ludolph
987 AC, Weishaupt JH, Robberecht W, Van Damme P, Franke L, Pers TH, Brown RH, Glass JD, Landers JE,
988 Hardiman O, Andersen PM, Corcia P, Vourc'h P, Silani V, Wray NR, Visscher PM, de Bakker PI, van Es
989 MA, Pasterkamp RJ, Lewis CM, Breen G, Al-Chalabi A, van den Berg LH, Veldink JH. Genome-wide
990 association analyses identify new risk variants and the genetic architecture of amyotrophic lateral
991 sclerosis. *Nat Genet.* 2016;48:1043-1048.

992 (21) Torres JM, Gamazon ER, Parra EJ, Below JE, Valladares-Salgado A, Wachter N, Cruz M, Hanis CL,
993 Cox NJ. Cross-tissue and tissue-specific eQTLs: partitioning the heritability of a complex trait. *Am J Hum*
994 *Genet.* 2014;95:521-534.

995 (22) Bailey JN, Loomis SJ, Kang JH, Allingham RR, Gharahkhani P, Khor CC, Burdon KP, Aschard H,
996 Chasman DI, Igo RP Jr, Hysi PG, Glastonbury CA, Ashley-Koch A, Brilliant M, Brown AA, Budenz DL,
997 Buil A, Cheng CY, Choi H, Christen WG, Curhan G, De Vivo I, Fingert JH, Foster PJ, Fuchs C,
998 Gaasterland D, Gaasterland T, Hewitt AW, Hu F, Hunter DJ, Khawaja AP, Lee RK, Li Z, Lichter PR,
999 Mackey DA, McGuffin P, Mitchell P, Moroi SE, Perera SA, Pepper KW, Qi Q, Realini T, Richards JE,
1000 Ridker PM, Rimm E, Ritch R, Ritchie M, Schuman JS, Scott WK, Singh K, Sit AJ, Song YE, Tamimi RM,
1001 Topouzis F, Viswanathan AC, Verma SS, Vollrath D, Wang JJ, Weisschuh N, Wissinger B, Wollstein G,
1002 Wong TY, Yaspan BL, Zack DJ, Zhang K, Study EN; ANZRAG Consortium., Weinreb RN, Pericak-Vance
1003 MA, Small K, Hammond CJ, Aung T, Liu Y, Vithana EN, MacGregor S, Craig JE, Kraft P, Howell G,
1004 Hauser MA, Pasquale LR, Haines JL, Wiggs JL. Genome-wide association analysis identifies TXNRD2,
1005 ATXN2 and FOXC1 as susceptibility loci for primary open-angle glaucoma. *Nat Genet.* 2016;48:189-194

1006 (23) GTEx Consortium. The Genotype-Tissue Expression (GTEx) project. *Nat Genet.* 2013;45:580-5.

1007 (24) Grundberg E, Small KS, Hedman ÅK, Nica AC, Buil A, Keildson S, Bell JT, Yang TP, Meduri E,
1008 Barrett A, Nisbett J, Sekowska M, Wilk A, Shin SY, Glass D, Travers M, Min JL, Ring S, Ho K,
1009 Thorleifsson G, Kong A, Thorsteindottir U, Ainali C, Dimas AS, Hassanali N, Ingle C, Knowles D,
1010 Krestyaninova M, Lowe CE, Di Meglio P, Montgomery SB, Parts L, Potter S, Surdulescu G,
1011 Tzaprouni L, Tsoka S, Bataille V, Durbin R, Nestle FO, O'Rahilly S, Soranzo N, Lindgren CM,
1012 Zondervan KT, Ahmadi KR, Schadt EE, Stefansson K, Smith GD, McCarthy MI, Deloukas P,
1013 Dermitzakis ET, Spector TD; Multiple Tissue Human Expression Resource (MuTHER) Consortium..

- 1014 Mapping cis- and trans-regulatory effects across multiple tissues in twins. *Nat Genet.* 2012;44:1084-
1015 1089.
- 1016 (25) Erbilgin A, Civelek M, Romanoski CE, Pan C, Hagopian R, Berliner JA, Lusis AJ. Identification
1017 of CAD candidate genes in GWAS loci and their expression in vascular cells. *J Lipid Res.*
1018 2013;54:1894-1905.
- 1019 (26) ENCODE Project Consortium. A user's guide to the encyclopedia of DNA elements (ENCODE).
1020 *PLoS Biol.* 2011;9:e1001046.
- 1021 (27) Miller CL, Anderson DR, Kundu RK, Raiesdana A, Nürnberg ST, Diaz R, Cheng K, Leeper NJ,
1022 Chen CH, Chang IS, Schadt EE, Hsiung CA, Assimes TL, Quertermous T. Disease-related growth
1023 factor and embryonic signaling pathways modulate an enhancer of TCF21 expression at the 6q23.2
1024 coronary heart disease locus. *PLoS Genet.* 2013;9:e1003652.
- 1025 (28) Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras
1026 TR. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics.* 2013;29:15-21.
- 1027 (29) Kooner JS, Saleheen D, Sim X, Sehmi J, Zhang W, Frossard P, Been LF, Chia KS, Dimas AS,
1028 Hassanali N, Jafar T, Jowett JB, Li X, Radha V, Rees SD, Takeuchi F, Young R, Aung T, Basit A,
1029 Chidambaram M, Das D, Grundberg E, Hedman AK, Hydrie ZI, Islam M, Khor CC, Kowlessur S,
1030 Kristensen MM, Liju S, Lim WY, Matthews DR, Liu J, Morris AP, Nica AC, Pinidiyapathirage JM,
1031 Prokopenko I, Rasheed A, Samuel M, Shah N, Shera AS, Small KS, Suo C, Wickremasinghe AR,
1032 Wong TY, Yang M, Zhang F; DIAGRAM; MuTHER, Abecasis GR, Barnett AH, Caulfield M, Deloukas
1033 P, Frayling TM, Froguel P, Kato N, Katulanda P, Kelly MA, Liang J, Mohan V, Sanghera DK, Scott J,
1034 Seielstad M, Zimmet PZ, Elliott P, Teo YY, McCarthy MI, Danesh J, Tai ES, Chambers JC. Genome-
1035 wide association study in individuals of South Asian ancestry identifies six new type 2 diabetes
1036 susceptibility loci. *Nat Genet.* 2011;43:984-999.
- 1037 (30) Saade S, Cazier JB, Ghassibe-Sabbagh M, Youhanna S, Badro DA, Kamatani Y, Hager J,
1038 Yeretzian JS, El-Khazen G, Haber M, Salloum AK, Douaihy B, Othman R, Shasha N, Kabbani S,
1039 Bayeh HE, Chammas E, Farrall M, Gauguier D, Platt DE, Zalloua PA; FGENTCARD consortium.
1040 Large scale association analysis identifies three susceptibility loci for coronary artery disease. *PLoS*
1041 *One.* 2011;6:e29427.
- 1042 (31) Tsaprouni LG, Yang TP, Bell J, Dick KJ, Kanoni S, Nisbet J, Viñuela A, Grundberg E, Nelson
1043 CP, Meduri E, Buil A, Cambien F, Hengstenberg C, Erdmann J, Schunkert H, Goodall AH,
1044 Ouwehand WH, Dermizakis E, Spector TD, Samani NJ, Deloukas P. Cigarette smoking reduces
1045 DNA methylation levels at multiple genomic loci but the effect is partially reversible upon cessation.
1046 *Epigenetics.* 2014;9:1382-1396.
- 1047 (32) Starke RM, Ali MS, Jabbour PM, Tjoumakaris SI, Gonzalez F, Hasan DM, Rosenwasser RH,
1048 Owens GK, Koch WJ, Dumont AS. Cigarette smoke modulates vascular smooth muscle phenotype:
1049 implications for carotid and cerebrovascular disease. *PLoS One.* 2013;8:e71954.
- 1050 (33) Bauer RC, Tohyama J, Cui J, Cheng L, Yang J, Zhang X, Ou K, Paschos GK, Zheng XL,
1051 Parmacek MS, Rader DJ, Reilly MP. Knockout of Adamts7, a novel coronary artery disease locus in
1052 humans, reduces atherosclerosis in mice. *Circulation.* 2015;131:1202-1213.
- 1053 (34) Holmes MV, Frikke-Schmidt R, Melis D, Luben R, Asselbergs FW, Boer JM, Cooper J, Palmén
1054 J, Horvat P, Engmann J, Li KW, Onland-Moret NC, Hofker MH, Kumari M, Keating BJ, Hubacek JA,

1055 Adamkova V, Kubinova R, Bobak M, Khaw KT, Nordestgaard BG, Wareham N, Humphries SE,
1056 Langenberg C, Tybjaerg-Hansen A, Talmud PJ. A systematic review and meta-analysis of 130,000
1057 individuals shows smoking does not modify the association of APOE genotype on risk of coronary
1058 heart disease. *Atherosclerosis*. 2014;237:5-12.

1059 (35) Pu X, Xiao Q, Kiechl S, Chan K, Ng FL, Gor S, Poston RN, Fang C, Patel A, Senver EC, Shaw-
1060 Hawkins S, Willeit J, Liu C, Zhu J, Tucker AT, Xu Q, Caulfield MJ, Ye S. ADAMTS7 cleavage and
1061 vascular smooth muscle cell migration is affected by a coronary-artery-disease-associated variant.
1062 *Am J Hum Genet*. 2013;92:366-374.

1063 (36) Wang L, Zheng J, Bai X, Liu B, Liu CJ, Xu Q, Zhu Y, Wang N, Kong W, Wang X. ADAMTS-7
1064 mediates vascular smooth muscle cell migration and neointima formation in balloon-injured rat
1065 arteries. *Circ Res*. 2009;104:688-698.

1066
1067 (37) Hutter CM, Mechanic LE, Chatterjee N, Kraft P, Gillanders EM; NCI Gene-Environment Think
1068 Tank. Gene-environment interactions in cancer epidemiology: a National Cancer Institute Think Tank
1069 report. *Genet Epidemiol*. 2013;37:643-657.

1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097

1098 **Table-1.** Novel genotype-smoking interaction findings in coronary heart disease at the chromosome 15q25.1 locus

Variant	Association	allele	LD with rs7178051*	LD with rs1051730^	Never Smokers					Ever Smokers					
					N cases	N controls	N Total	Beta (SE)	P-value	N cases	N controls	N Total	Beta (SE)	P-value	P-value interaction
*rs7178051 ⁴	CHD (NPR)	T/C	-	0.22	21232	38713	59945	-0.13 (0.01)	1.30E-16	39585	40749	80334	-0.05 (.01)	2.49E-04	8.57E-05
†rs1051730 ¹⁶	SB (known)	A/G	0.22	-	20559	38198	58757	-0.04 (0.02)	0.02	38923	40371	79294	0.03 (0.01)	0.02	2.37E-04
Other variants on chr.15q25.1 with significant gene-smoking interactions on CHD															
rs7173743 ¹	CHD (Known)	C/T	0.61	0.18	21050	37955	59005	-0.11 (0.01)	2.73E-13	39044	39559	78603	-0.04 (0.01)	8.60E-04	9.29E-05
rs10083696 ²	CHD (Novel)	A/G	1.0	0.22	19721	36206	55927	-0.11 (0.02)	1.60E-12	38807	40018	78825	-0.05 (0.01)	2.72E-04	5.15E-05
rs7176187 ³	CHD (Novel)	T/C	1.0	0.24	21232	38713	59945	-0.12 (0.01)	7.02E-16	39585	40749	80334	-0.04 (0.01)	8.64E-04	6.93E-05
rs6495335 ⁵	CHD (Novel)	G/T	1.0	0.22	20144	37217	57361	-0.13 (0.02)	2.39E-15	36448	38203	74651	-0.04 (0.01)	1.69E-03	9.51E-04
rs4380028 ⁶	CHD (Known)	T/C	1	0.22	21232	38713	59945	-0.12 (0.01)	2.20E-15	39585	40749	80334	-0.04 (.01)	1.03E-03	5.44E-04
rs3825807 ⁷	CHD (Known)	G/A	0.52	0.43	17137	28633	45771	-0.09 (0.02)	2.82E-08	30071	29014	59086	-0.03 (0.01)	0.04	2.6E-03
rs3813565 ⁸	CHD (NPR)	T/G	0.43	0.56	19466	35830	55296	-0.08 (0.02)	5.08E-07	36642	37759	74401	-0.01 (0.01)	0.42	3.05E-04
rs11638490 ⁹	CHD (NPR)	T/C	0.44	0.51	20465	37897	58362	-0.08 (0.01)	6.90E-08	38533	39690	78223	-0.01 (0.01)	0.28	2.25E-04
rs11072791 ¹¹	CHD (NPR)	A/C	0.44	0.51	19289	35944	55233	-0.08 (0.02)	2.83E-07	35245	36635	71880	-.005 (0.01)	0.68	1.06E-04
rs922692 ¹²	CHD (NPR)	A/C	0.44	0.50	20559	38198	58757	-0.08 (0.01)	2.81E-07	38923	40371	79294	-0.01 (0.01)	0.29	2.75E-04
rs11638372 ¹³	CHD (NPR)	T/C	0.44	0.50	21232	38713	59945	-0.08 (0.01)	6.92E-08	39585	40749	80334	-0.01 (0.01)	0.23	3.16E-04
rs4887077 ¹⁴	CHD (NPR)	T/C	0.44	0.50	21232	38713	59945	-0.08 (0.01)	4.71E-08	39585	40749	80334	-0.02 (0.01)	0.20	3.92E-05
rs12899135 ¹⁵	CHD (NPR)	G/A	0.39	0.56	20377	37440	57817	-0.07 (0.02)	3.97E-06	38382	39181	77563	-0.01 (0.01)	0.58	4.54E-04
rs684513 ¹⁸	SB (Known)	C/G	0.01	0.10	12517	21054	33572	-0.01 (0.02)	0.67	24641	24487	49129	0.03 (0.02)	0.18	0.08
rs2036527 ¹⁹	SB (Known)	A/G	0.17	0.90	20559	38198	58757	-0.04 (0.02)	0.02	38923	40371	79294	0.03 (0.01)	0.02	2.14E-04
rs10519203 ²⁰	CHD (NPR)	G/A	0.19	0.93	21232	38713	59945	-0.04 (0.01)	5.93E-03	39585	40749	80334	0.03 (0.01)	0.03	1.27E-04
rs8034191 ²¹	SB (Known)	C/T	0.19	1.0	19251	32131	51382	-0.05 (0.02)	2.62E-03	34925	34047	68972	0.02 (0.01)	0.06	3.91E-05

1099 CHD = coronary heart disease; SB = smoking behavior; NPR: Not a previously reported variant with disease risk
 1100 *lead variant in association with CHD in our dataset; † lead variant in association with SB
 1101 ¹⁻²¹each number refers to the physical location of the variant in figure-

1102
 1103
 1104
 1105
 1106
 1107
 1108
 1109
 1110

1111 **Figure Legends**

1112

1113 **Figure 1.** (a) Regional association analyses at the chromosome 15q25.1 locus in association with
1114 CHD risk stratified by smoking status. Association P-values for genetic variants with CHD risk in
1115 “never-smokers” (green squares) and “ever-smokers” (red triangles). (b) Longitudinal bars represent
1116 gene-smoking CHD interaction P-values at the chromosome 15q25.1 locus; bars in blue are P-
1117 values for variants listed in Table-1 and each variant has been assigned a unique identification
1118 number based on its physical location; (c) LD-blocks at the 15q25.1 locus visualized through
1119 HAPLOVIEW using LD estimates in the HapMAP-2 CEU reference population.

1120

1121 **Figure 2.** Several variants at chromosome 15q21.1 have stronger effects on CHD risk in “never-
1122 smokers” compared to “ever-smokers”. Variants with the strongest interaction P-value are displayed.

1123

1124 **Figure 3.** Step-wise conditional analysis of genetic variation at the chromosome 15q21.1 locus with
1125 CHD (red triangles) and smoking behavior (cigarettes per day, CPD; grey circles). At the
1126 chromosome 15q21.1 locus, analyses adjusted for rs7178051 and rs11072794 completely
1127 attenuated the gene-CHD associations whereas gene-smoking remained unchanged. Analyses
1128 adjusted for rs1051730 and rs684513 completely attenuated the gene-smoking associations
1129 whereas gene-CHD effect remained unchanged.

1130 **Figure 4.** Analyses mutually adjusted for rs7178051, rs11072794, rs1051730 and rs684513 at
1131 15q21.1 on CHD and smoking behavior; gene-CHD interaction analyses were only found significant
1132 for rs7178051. Analyses on the left panel show associations of rs7178051, rs11072794, rs1051730
1133 and rs684513 with CHD risk mutually adjusted for each other. Analyses on the right panel show
1134 associations of rs7178051, rs11072794, rs1051730 and rs684513 with smoking behavior mutually
1135 adjusted for each other.

1136 **Figure 5.** (a) *ADAMTS7* and *CHRNA3-4-5* mRNA levels were measured in HCASMC. Cells were
1137 cultured to confluence, total RNA was extracted and cDNA generated. q-PCR was performed for
1138 *ACTB*, *GAPDH*, *TBP*, *ADAMTS7*, *CHRNA3*, *CHRNA4*, *CHRNA5* (95°C 15s, 60°C 1min). Delta Cts
1139 were calculated as follows: $(Ct_{ACTB} + Ct_{GAPDH} + Ct_{TBP})/3 - Ct_{TARGET\ GENE}$. Fold changes are derived
1140 from delta delta Cts based on formula $FC = 2^{-\Delta\Delta Ct}$. (b) Confluent HCASMC were exposed to cigarette
1141 smoke extract. Serum starved (x24 hrs.) confluent HCASMC were treated with 0.5% or 1.0%
1142 cigarette smoke extract (v/v) for 4, 12, and 24 hrs. in serum reduced conditions (0.5% FBS in
1143 DMEM). Total RNA was extracted, cDNA generated preparation and q-PCR performed for

1144 *ADAMTS7* by Taqman and normalized to *GAPDH*. The Average Ct for *ADAMTS7* at baseline was
1145 28.25. Results were presented as means \pm SEM, and data were analyzed using Student's t-Test. (c)
1146 expression and eQTL Data from the GTEx consortium, the HapMap consortium (restricted to
1147 European populations), the Multiple Tissue Human Expression Resource (MuTHER) and in 147
1148 donor HAoEC lines. Association of the independent lead variants identified in our conditional
1149 analyses with expression of *ADAMTS7* and genes in the *CHRNA4-A3-A5* cluster. A P-value
1150 threshold of 0.002 was set to account for multiple testing involved in the eQTL analyses.

1151 **Figure 6.** Genome browser view of regulatory features at rs7178051 on Chr15q21.1. ChIP-seq
1152 experiments were performed on confluent HCASMC for TCF21, Jun, JunD, CEBP and H3K4me1,
1153 H3K27me3, H3K27ac. DNaseI hypersensitivity data for human AoSMC were acquired from the
1154 ENCODE project. Human aortic tissue H3K4me1, H3K9me3, H3K27me3, and H3K36me3 ChIP-seq
1155 data were acquired from the NIH Roadmap Epigenomics Project. HCASMC = human coronary
1156 artery smooth muscle cells; AoSMC = human aortic smooth muscle cells.

1157

1158

1159

1160