

University of Dundee

Corrigendum

Munoz, Ivan M.; Jowsey, Paul A.; Toth, Rachel; Rouse, John

Published in:
Nucleic Acids Research

DOI:
[10.1093/nar/gkx1140](https://doi.org/10.1093/nar/gkx1140)

Publication date:
2017

Licence:
CC BY-NC

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Munoz, I. M., Jowsey, P. A., Toth, R., & Rouse, J. (2017). Corrigendum: Phospho-epitope binding by the BRCT domains of hPTIP controls multiple aspects of the cellular response to DNA damage. *Nucleic Acids Research*, 45(22), 13094. <https://doi.org/10.1093/nar/gkx1140>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Corrigendum

Phospho-epitope binding by the BRCT domains of hPTIP controls multiple aspects of the cellular response to DNA damage

Ivan M. Munoz, Paul A. Jowsey, Rachel Toth and John Rouse*

MRC Protein Phosphorylation Unit, Sir James Black Centre, University of Dundee, Dundee DD1 5EH, Scotland, UK

Nucleic Acids Research (2007) 35(16) 5312–5322, <https://doi.org/10.1093/nar/gkm493>

The authors wish to make the following correction to their article:

Duplicated loading control lanes were noticed in Figure 1C. A new figure to correct the mistake in Figure 1C is presented here to maintain the integrity of the research record. This correction does not alter the conclusions of the article.




Figure 1C. HEK293 cells were co-transfected with FLAG-hPTIP and full-length HA-53BP1 bearing the indicated mutations. Cells were exposed to IR (0 Gy or 20 Gy) and after cell lysis anti-FLAG immunoprecipitates were subjected to SDS-PAGE and western blotting with the indicated antibodies. The lowest panels show HA-53BP1 and FLAG-hPTIP levels in cell extracts.

*To whom correspondence should be addressed. Tel: +44 1382 385490; Fax: +44 1382 223778; Email: j.rouse@dundee.ac.uk
Present address: Paul A. Jowsey, NIHR Health Protection Research Unit, Wolfson Building, Newcastle University, Newcastle upon Tyne, NE2 4AA, United Kingdom.