

University of Dundee

Striate palmoplantar keratoderma resulting from a missense mutation in DSG1

Vodo, D.; O'Toole, E. A.; Malchin, N.; Lahav, A.; Adir, N.; Sarig, O.

Published in:
British Journal of Dermatology

DOI:
[10.1111/bjd.16320](https://doi.org/10.1111/bjd.16320)

Publication date:
2018

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Vodo, D., O'Toole, E. A., Malchin, N., Lahav, A., Adir, N., Sarig, O., Green, K., Smith, F. J. D., & Sprecher, E. (2018). Striate palmoplantar keratoderma resulting from a missense mutation in DSG1. *British Journal of Dermatology*, 179(3), 755-757. Advance online publication. <https://doi.org/10.1111/bjd.16320>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

PROFESSOR ELI SPRECHER (Orcid ID : 0000-0002-4093-1032)

Article type : Research Letter

Striate palmoplantar keratoderma resulting from a missense mutation in *DSG1*

D. Vodo^{1,2}, E.A. O'Toole³, N. Malchin¹, A. Lahav⁴, N. Adir⁴, O. Sarig¹, K. Green⁵, F. J.D. Smith⁶, E. Sprecher^{1,2}

¹Department of Dermatology, Tel Aviv Sourasky Medical Center, Tel Aviv, Israel;

²Department of Human Molecular Genetics and Biochemistry, Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel; ³Centre for Cell Biology and

Cutaneous Research, The Blizard Institute, Barts and The London School of

Medicine and Dentistry, Queen Mary University of London, London, UK; ⁴Schulich

Faculty of chemistry, Technion, Haifa, Israel; ⁵Departments of Pathology and

Dermatology, Northwestern University Feinberg School of Medicine, Chicago, Illinois,

USA; ⁶Pachyonychia Congenita Project, School of Life Sciences, University of Dundee, UK

Running title: SPPK resulting from missense mutations in DSG1

Corresponding Author: Eli Sprecher, MD, PhD

Department of Dermatology

Tel Aviv Sourasky Medical Center

6 Weizmann Street, Tel Aviv 64239, Israel

E-mail: elisp@tlvmc.gov.il

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/bjd.16320

This article is protected by copyright. All rights reserved.

Accepted Article

Funding resources: This work has been supported in part by the Pachyonychia Congenita Project.

Conflict of interests: The authors have declared no conflicting interests.

Dear Editor, palmoplantar keratodermas (PPKs, OMIM #144200) refer to a large phenotypically and genetically heterogeneous group of keratinisation disorders characterised by marked hyperkeratosis on the surface of the palms and soles.¹ Striate palmoplantar keratoderma (SPPK) features linear hyperkeratosis of the volar aspects of the fingers, extending onto the palms, as well as focal plantar keratoderma. The disease is most commonly caused by heterozygous mutations in *DSG1* encoding desmoglein 1 (*DSG1*)² (SPPK1, OMIM #148700) but can also arise from mutations in *DSP* encoding desmoplakin (*DSP*)³ (SPPK2, OMIM #612908) or *KRT1* encoding keratin 1 (*K1*)⁴ (SPPK3, OMIM #607654). To date, more than 25 mutations in *DSG1* have been identified in patients with SPPK1. These mutations were uniformly found to result in a premature termination codon (PTC).^{5,6} In this study, we present, to our knowledge, the first case of SPPK caused by a missense (rather than PTC-causing) mutation affecting a residue predicted to be of critical importance for *DSG1*-mediated adhesion.

The patient was a 27-year-old female of Jewish origin. She reported thickening of the skin of palms and soles since 1 year of age, resulting in pain while walking. She denied any additional systemic or dermatologic manifestations apart from yellow discolouration of all toenails. Her father and her sister reportedly displayed similar clinical features. On examination, the patient exhibited foci of plantar yellowish fissured hyperkeratotic skin involving weight bearing areas as well as subtle linear

Accepted Article

hyperkeratosis of the volar aspects of the fingers (Fig. 1a). Haematoxylin and eosin staining of a skin biopsy showed hypergranulosis, marked orthohyperkeratosis and widening of the intercellular spaces between keratinocytes. (Fig. 1b).

To identify the causative mutation underlying SPPK in the patient, after obtaining ethical approval and informed consent, we used direct sequencing of gDNA to fully sequence all coding exons of *DSG1*, *KRT1*, *DSP* and *KRT16*. We discovered a heterozygous A to G transition at position 254 (c.254A>G) within exon 4 (Fig. 1c) of the *DSG1* gene (reference sequence Ensembl accession number ENST00000257192.4), resulting in a p.Y85C substitution. The sequence variation was not detected in 629 Jewish healthy controls and was not found in the following genomic databases: NCBI, UCSC, HGMD, ESP, ExAC, 1000 genomes and ENSEMBL, comprising a total of more than 58,000 individual sequences. The p.Y85C variation is predicted to be deleterious by two prediction tools (Polyphen2 score 1, range 0–1; SIFT scores 0, range 1–0) and was found to affect a highly conserved residue (Conseq score = 9, range 1-9). No mutations were identified in *KRT1*, *DSP* or *KRT16*.

To assess the consequences of the mutation, we modeled the wildtype and the mutant DSG1 proteins using three prediction tools (Fig. 1d): SWISS PROT⁷ and LOMETS⁸ predictions were based on the crystal structure previously determined for the human DSG2 ectodomain (PDB code 5ERD, chain A, 36% identity) while PHYRE2⁹ was used based on the human DSG3 ectodomain (PDB code 5EQX, chain A, 55% identity). Similarity between the model structures was assessed using PyMOL alignment (The PyMOL Molecular Graphics System, Version 1.8

Schrödinger, LLC).

Like other members of the cadherin superfamily, desmosomal cadherins are thought to form trans-adhesive dimers, which rely on the interaction between the conserved tryptophan (W2) of one molecule and a specific hydrophobic pocket, serving as an acceptor site, on the interacting molecule from the opposing cell.¹⁰⁻¹³ This was recently shown to depend predominantly on a heterotypic interaction between the donor site of a desmocollin molecule and the acceptor pocket of a DSG molecule,¹⁴ which is lined by the side chains of specific conserved residues including Y85 (numbered Y36 in the final DSG chain).¹²⁻¹⁴

Accordingly, we found that the affected amino acid, Y85, resides within the first extracellular domain (EC1) of the protein (Fig. 1e), inside a hydrophobic-positively charged area (Fig. 1f). The p.Y85C mutation could affect the binding capability of the hydrophobic pocket in several ways. First, while the hydroxyl side chain of Y85 was found to reach the inner part of this pocket, the C85 mutation is lacking the length to do so (Fig. 1f). Additionally, the mutant C85 residue is predicted to aberrantly interact with C127, a part of the adhesive interface of the pocket,¹³ and could potentially form a disulfide bond either through an intramolecular cis-interaction, due to proximity of the two amino acids, or through a trans-interaction with another bound desmosomal cadherin. When we used the same modeling tools to examine the effect of p.V83D and p.R86C, both highly rare heterozygous non-pathogenic variants in close proximity to p.Y85C, we discovered that although residing close to the hydrophobic pocket, they are not predicted to disturb its binding capability or to have an effect on DSG1 dimerisation (Fig. 1g).

To conclude, we describe here the first case of SPPK resulting from a heterozygous missense (as opposed to nonsense or frameshift) mutation in the *DSG1* gene to our knowledge. Protein modeling predicts this mutation to abolish DSG1-mediated adhesive function, thus resulting in what could be manifested as functional haploinsufficiency. Of interest, DSG1 has been shown to downregulate the Ras/MAPK pathway by interacting with Erbin; elevated Ras activity, resulting from DSG1 deficiency, is therefore thought to trigger cellular mechanisms leading to hyperkeratosis in SPPK.¹⁵ Whether missense mutations in the extracellular domain affect this regulatory pathway remains to be determined. In addition to the relevance of these data to PPK diagnostics, the putative mode of action of the mutation identified in the present study emphasises the importance of the role of DSG1-mediated adhesive function in palmoplantar skin.

Acknowledgements

The authors wish to thank the patient and her family for their collaboration. This work complies with the Committee on Publication Ethics (COPE) guidelines on good publication.

References

- 1 Itin PH, Fistarol SK. Palmoplantar keratodermas. *Clin Dermatol* 2005; **23**: 15-22.
- 2 Rickman L, Simrak D, Stevens HP *et al.* N-terminal deletion in a desmosomal cadherin causes the autosomal dominant skin disease striate palmoplantar keratoderma. *Hum Mol Genet* 1999; **8**: 971-6.

- 3 Armstrong DK, McKenna KE, Purkis PE *et al.* Haploinsufficiency of
desmoplakin causes a striate subtype of palmoplantar keratoderma. *Hum Mol*
Genet 1999; **8**: 143-8.
- 4 Whittock NV, Smith FJ, Wan H *et al.* Frameshift mutation in the V2 domain of
human keratin 1 results in striate palmoplantar keratoderma. *J Invest Dermatol*
2002; **118**: 838-44.
- 5 Hershkovitz D, Lugassy J, Indelman M *et al.* Novel mutations in DSG1 causing
striate palmoplantar keratoderma. *Clin Exp Dermatol* 2009; **34**: 224-8.
- 6 Lovgren ML, McAleer MA, Irvine AD *et al.* Mutations in desmoglein-1 cause
diverse inherited palmoplantar keratoderma phenotypes: Implications for
genetic screening. *Br J Dermatol* 2016.
- 7 Boeckmann B, Bairoch A, Apweiler R *et al.* The SWISS-PROT protein
knowledgebase and its supplement TrEMBL in 2003. *Nucleic Acids Res* 2003;
31: 365-70.
- 8 Wu S, Zhang Y. LOMETS: a local meta-threading-server for protein structure
prediction. *Nucleic Acids Res* 2007; **35**: 3375-82.
- 9 Kelley LA, Mezulis S, Yates CM *et al.* The Phyre2 web portal for protein
modeling, prediction and analysis. *Nat Protoc* 2015; **10**: 845-58.
- 10 Al-Amoudi A, Diez DC, Betts MJ *et al.* The molecular architecture of cadherins
in native epidermal desmosomes. *Nature* 2007; **450**: 832-7.
- 11 Boggon TJ, Murray J, Chappuis-Flament S *et al.* C-cadherin ectodomain
structure and implications for cell adhesion mechanisms. *Science* 2002; **296**:
1308-13.
- 12 Shapiro L, Fannon AM, Kwong PD *et al.* Structural basis of cell-cell adhesion
by cadherins. *Nature* 1995; **374**: 327-37.

- 13 Patel SD, Chen CP, Bahna F *et al.* Cadherin-mediated cell-cell adhesion: sticking together as a family. *Curr Opin Struct Biol* 2003; **13**: 690-8.
- 14 Harrison OJ, Brasch J, Lasso G *et al.* Structural basis of adhesive binding by desmocollins and desmogleins. *Proc Natl Acad Sci U S A* 2016; **113**: 7160-5.
- 15 Harmon RM, Simpson CL, Johnson JL *et al.* Desmoglein-1/Erbin interaction suppresses ERK activation to support epidermal differentiation. *J Clin Invest* 2013; **123**: 1556-70.

Legends to figures

Figure 1. Clinical features, mutation analysis and protein modeling. (a) The patient presented with foci of plantar hyperkeratosis involving weight bearing areas with multiple skin fissures (left panel). Linear hyperkeratosis of the volar aspects of the fingers was also observed (right panel); (b) Hematoxylin and eosin staining of a skin biopsy obtained from the skin of the patient demonstrates hypergranulosis, marked orthohyperkeratosis and widening (arrows) of the intercellular spaces between keratinocytes, compatible with keratoderma (scale bar = 100 μ m); (c) Direct sequencing of gDNA revealed a A>G transition at position c.254 within exon 4 of *DSG1* (upper panel), compared to wild-type sequence (lower panel). (d) Comparison between the molecular model of the human DSG1 protein based on three prediction tools: LOMETS (green), SWISS PROT (magenta) and PHYRE2 (cyan) shows that all the structures are similar (RMSD<4 with PyMOL alignment) and that the Y85 amino acid is located in the first domain (in red, highlighted by black oval). (e) A schematic representation of the DSG1 molecule shows that the p.Y85C mutation is located

inside the EC1 domain of the protein. EC, extracellular domain; EA, extracellular anchoring domain; P, propeptide region; SP, signal peptide; TM, transmembrane domain (protein scheme modified from Hershkovitz et al⁵); (f) Mutation modeling was performed based on the above models. The protein molecules are overlaid with semi-transparent calculated electrostatic potential (blue-positive, red-negative) created using PyMOL. Y85 (in red) is located in a hydrophobic-positively charged pocket-shaped surface, predicted to serve as a pathway for DSG1 binding. While the hydroxyl of the tyrosine lies within the inner part of the pocket (left panel), the mutant C85 substitution is lacking the length to reach its inner rim (right panel). (g) Modeling of two rare heterozygous variants in close proximity to Y85 (in red) was performed based on the above models. The p.V83D variant (in purple, right panel) resides further away from the hydrophobic pocket and will lead to a negatively charged glutamic acid that is not predicted to alter the negatively charged area of the wild type p.V83 (in purple, left panel). p.R86 (in green, left panel) is located in proximity of the hydrophobic pocket and of p.C127 (in black) but faces the opposite direction. The p.R86C variant (in green, right panel) is not predicted to affect the binding capability of the hydrophobic pocket or to form a disulphide bond with p.C127.

