



**University of Dundee**

**A Posteriori Error Analysis for Evolution Nonlinear Schrodinger Equations Up to the Critical Exponent**

Katsaounis, Theodoros ; Kyza, Irene

*Published in:*  
SIAM Journal on Numerical Analysis

*DOI:*  
[10.1137/16M1108029](https://doi.org/10.1137/16M1108029)

*Publication date:*  
2018

*Document Version*  
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

*Citation for published version (APA):*  
Katsaounis, T., & Kyza, I. (2018). A Posteriori Error Analysis for Evolution Nonlinear Schrodinger Equations Up to the Critical Exponent. *SIAM Journal on Numerical Analysis*, 56(3), 1405-1434.  
<https://doi.org/10.1137/16M1108029>

**General rights**

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## A POSTERIORI ERROR ANALYSIS FOR EVOLUTION NONLINEAR SCHRÖDINGER EQUATIONS UP TO THE CRITICAL EXPONENT

THEODOROS KATSAOUNIS AND IRENE KYZA

**ABSTRACT.** We provide a posteriori error estimates in the  $L^\infty([0, T]; L^2(\Omega))$ -norm for relaxation time discrete and fully discrete schemes for a class of evolution nonlinear Schrödinger equations up to the critical exponent. In particular for the discretization in time we use the relaxation Crank-Nicolson-type scheme introduced by Besse in [9]. The space discretization consists of finite element spaces that are allowed to change between time steps. The estimates are obtained using the reconstruction technique. Through this technique the problem is converted to a perturbation of the original partial differential equation and this makes it possible to use nonlinear stability arguments as in the continuous problem. Our analysis includes as special cases the cubic and quintic nonlinear Schrödinger equations in one spatial dimension and the cubic nonlinear Schrödinger equation in two spatial dimensions. Numerical results illustrate that the estimates are indeed of optimal order of convergence.

### 1. INTRODUCTION

In this paper we provide error control for a class of evolution nonlinear Schrödinger (NLS) equations, up to the critical exponent through *rigorous a posteriori error analysis*. *To the best of our knowledge, this is the first time that rigorous a posteriori error estimates are proven for NLS equations*. More specifically, we consider the initial and boundary value problem

$$(1.1) \quad \begin{cases} \partial_t u - i\alpha \Delta u = i\lambda f(u) & \text{in } \Omega \times (0, T], \\ u = 0 & \text{on } \partial\Omega \times (0, T], \\ u(\cdot, 0) = u_0 & \text{in } \Omega, \end{cases}$$

where  $\Omega \subset \mathbb{R}^d$ ,  $d = 1, 2$ , is a bounded, convex, polygonal domain for  $d = 2$  and a finite interval for  $d = 1$ ,  $T < \infty$ ,  $\alpha > 0$ ,  $\lambda \in \mathbb{R}$ , and where  $f(u)$  denotes the nonlinear term. In particular, the nonlinear term has the form

$$f(z) := |z|^{2p} z, \quad \frac{1}{2} \leq p \leq p^* \quad \text{with} \quad p^* := \frac{2}{d}.$$

The exponent  $p^* = \frac{2}{d}$  in  $d$  dimensions is *critical* in the following sense: For  $u_0 \in H_0^1(\Omega) \cap H^2(\Omega)$  and  $\frac{1}{2} \leq p < p^*$  problem (1.1) admits a unique global solution  $u \in C([0, T]; H_0^1(\Omega) \cap H^2(\Omega)) \cap C^1([0, T]; L^2(\Omega))$ , for *any*  $T > 0$  cf., e.g., [4, 10, 13]. In contrast, for the critical exponent  $p^* = \frac{2}{d}$ , global existence of a solution  $u$  of (1.1) is guaranteed *only* under a smallness condition, elaborated in equation (1.3). If this smallness condition does not hold, blowup solutions of problem (1.1) exist [40, 41, 42].

In particular the range of exponents  $p$  with  $\frac{1}{2} \leq p \leq p^*$  includes the cubic and quintic NLS in one spatial dimension and the cubic NLS in two spatial dimensions. Models of the form (1.1) are widely used in many areas of applied sciences. For example they appear in nonlinear optics and lasers [23], water waves [21], quantum hydrodynamics [25] and Bose-Einstein condensates [48]. More applications are discussed in [49].

There is still a large activity on NLS equations in the area of partial differential equations (PDEs) and analysis community, cf. e.g., [4, 10, 17, 40, 41, 42, 44, 50] and the references therein. Moreover

---

*Date:* March 6, 2018.

*Key words and phrases.* Evolution NLS, power nonlinearities, a posteriori error control, reconstruction technique, relaxation Crank-Nicolson-type scheme, finite elements.

Work partially supported by Excellence Award 1456 of the Greek Ministry of Research and Education.

this activity includes qualitative and asymptotic questions, cf. e.g., [8, 16, 37, 53] and the references therein. A particular example is the semiclassical behavior of NLS equations, i.e. the regime where  $0 < \alpha \ll 1$ ,  $\lambda \sim \frac{1}{\alpha}$ .

Those are the main reasons that (1.1) has attracted the interest of the numerical analysis community as well. Several papers exist in the literature dealing with discretization methods for (1.1) their stability and convergence properties through a priori error analysis; here we mention a few: [1, 5, 6, 7, 9, 19, 27, 28, 51, 54, 56]. Popular methods for the discretization in time of (1.1) are Crank-Nicolson-type and time-splitting-type methods, while for the spatial discretization spectral or finite element methods are usually used.

However, there is a very limited literature on the a posteriori error control for the evolution Schrödinger equations. For linear evolution Schrödinger equations a posteriori error estimates for Crank-Nicolson finite element schemes can be found in [22, 29, 31], whilst for time-splitting spectral methods can be found in [33]. To the best of our knowledge *no a posteriori error bounds exist in the literature for evolution NLS equations*. Nevertheless, developing such estimates is important. Not only they will provide mathematical guarantees on how accurate the numerical approximation is, but they will also highlight qualitative characteristics of the exact solution of (1.1) not known before, via rigorous error control. Additionally, the a posteriori error bounds can lead to the development of an adaptive algorithm that will significantly reduce the computational cost. That was the case for example in [29] with the a posteriori estimator of linear evolution Schrödinger equations. Adaptive algorithms based on *heuristic* mesh selection criteria exist in the literature for various cases of NLS equations (1.1), cf. [2, 24, 45, 52, 55]. Usually the criteria used for the construction of adaptive algorithms in these cases are based on structural properties known for the exact solution.

*Our main contribution in this paper is the rigorous proof of optimal order a posteriori error bounds in the  $L^\infty([0, T]; L^2(\Omega))$ -norm for the NLS equation (1.1) up to the critical exponent*, when it is discretized by a relaxation Crank-Nicolson finite element scheme. With the term *optimal order* we mean that the a posteriori estimator reduces with the same order as the exact error. The relaxation Crank-Nicolson scheme we use for the discretization in time is a generalization to variable time steps of the relaxation scheme introduced earlier by Besse in [9] for the time discretization of (1.1) for constant time steps. The reason we also use the relaxation scheme, rather than the standard Crank-Nicolson scheme, is because that way the nonlinear term is computed explicitly. Thus we avoid solving a nonlinear equation, that would have added an error, difficult to handle a posteriori. Moreover, the relaxation scheme exhibits mass conservation, same as the standard Crank-Nicolson scheme, thus reflecting the mass conservation of the continuous problem (1.1), cf. (1.4) below.

The a posteriori error estimates will be obtained using the reconstruction technique proposed by Akrivis, Makridakis & Nocketto, [3, 38]. Through this technique we will be able to obtain an error equation of a similar form to the NLS equation in (1.1).

The derivation of the estimates is then based on energy techniques and on nonlinear stability arguments for (1.1). It is interesting to note that the PDE satisfied by the error, poses new mathematical challenges when aiming at meaningful error control. For this reason the handling of the nonlinearity is delicate and technical.

The reconstruction technique has been used successfully in proving a posteriori error bounds for linear problems, [3, 14, 29]. In this work, using the reconstruction technique, we introduce *novel time-space reconstructions* and prove rigorous a posteriori error bounds for (1.1). To the best of our knowledge this is the first time in the literature where the reconstruction technique has been used successfully for a highly nonlinear problem such as (1.1).

We next mention some of the main tools used in the subsequent analysis. For this, we need to introduce some notation. We denote by  $\|\cdot\|$  the  $L^2$ -norm in  $\Omega$ , while for  $1 \leq q \leq \infty$ ,  $q \neq 2$ , we denote by  $\|\cdot\|_{L^q}$  the  $L^q$ -norm in  $\Omega$ . We equip  $H_0^1(\Omega)$  with the norm  $\|\nabla v\|$  and we denote by  $H^{-1}(\Omega)$  the dual of  $H_0^1(\Omega)$  under that norm; we denote by  $\|\cdot\|_{H^{-1}}$  the norm in  $H^{-1}(\Omega)$ . By  $\langle \cdot, \cdot \rangle$  we indicate both the  $L^2$ -inner product, or the  $H^{-1} - H_0^1$  duality pairing in  $\Omega$ , depending on the context. In what follows, global constants or functionals depending on the initial condition  $u_0$  that are introduced in the paper may also depend on the dimension  $d$ , the exponent  $p$  and the parameters  $\alpha$  and  $\lambda$ . For simplicity, where there is no confusion, we avoid writing those dependences on the

definitions, but we mention them precisely each time we define such a quantity (e.g., (1.2) and (1.3) below).

A key role in the derivation of the a posteriori estimates of the paper will be played by the Gagliardo-Nirenberg inequality, [12]

$$(1.2) \quad \|v\|_{L^{2p+2}} \leq \beta \|\nabla v\|^\zeta \|v\|^{1-\zeta}, \quad \forall v \in H_0^1(\Omega),$$

where  $\zeta := \frac{pd}{2(p+1)}$  and  $\beta$  is an absolute constant depending on  $\Omega$ ,  $d$  and  $p$ . For the cases of critical exponent  $p = p^* (= \frac{2}{d})$ , (1.1) admits a unique global solution  $u \in C([0, T]; H_0^1(\Omega) \cap H^2(\Omega)) \cap C^1([0, T]; L^2(\Omega))$  for every  $T > 0$ , if  $u_0 \in H_0^1(\Omega) \cap H^2(\Omega)$  and

$$(1.3) \quad \Gamma(u_0) := \frac{\beta^{2p+2}\lambda}{\alpha(p+1)} \|u_0\|^{2p} < 1.$$

For  $\lambda \leq 0$ , inequality (1.3) is automatically satisfied; problem (1.1) is then known as the defocusing NLS equation. If  $\lambda > 0$ , problem (1.1) is called the focusing NLS equation. In this case, if (1.3) is not satisfied, i.e., if  $\Gamma(u_0) \geq 1$ , the solution  $u$  of (1.1) may blow up in the  $H^1$ -norm in some finite time  $T^* < \infty$  (cf., e.g., [40, 41, 42] and the references therein). The focusing cases we consider in this paper are those with  $\Gamma(u_0) < 1$ , thus a global solution of (1.1) exists. A posteriori error analysis for controlling the error close to blowup for focusing cases with  $\Gamma(u_0) \geq 1$  is a very interesting question and currently under investigation.

Problem (1.1) satisfies two conservation laws that will be instrumental in the subsequent analysis. In particular, for  $t \geq 0$ , we have

$$(1.4) \quad \|u(t)\| = \|u_0\|, \quad \text{mass conservation,}$$

$$(1.5) \quad \|\nabla u(t)\|^2 - \frac{\lambda}{\alpha(p+1)} \|u(t)\|_{L^{2p+2}}^{2p+2} = \|\nabla u_0\|^2 - \frac{\lambda}{\alpha(p+1)} \|u_0\|_{L^{2p+2}}^{2p+2}, \quad \text{energy conservation.}$$

Using the Gagliardo-Nirenberg inequality (1.2) and the two conservation laws (1.4) and (1.5) we prove a novel a posteriori estimate for the gradient of the solution:

$$\|\nabla u(t)\| \leq G(u_0),$$

where  $G(u_0)$  is a computable quantity depending only on  $u_0$  and known constants, see (3.11), (3.12). This estimate is an essential ingredient of the proof of the final a posteriori error estimates. Similar estimates are used in the a priori analysis for establishing existence of solutions for (1.1), cf. [26, 41].

The final estimates include an exponential term of the  $L^{2p}([0, T]; L^\infty(\Omega))$ -norm of the approximation; this is due to Gronwall's inequality. This is an improvement compared to the existing results on the a priori error analysis, where the exponential of the  $L^\infty([0, T]; L^\infty(\Omega))$ -norm of the approximation appears. Although any exponential term may seem pessimistic, it actually reflects the nonlinear nature of the problem. A similar term appears in the a posteriori error analysis for semilinear parabolic equations with possible blowup in finite time; cf. [15, 32]. In fact, as illustrated in [15, 34], this term enables the proposition of an efficient time-space adaptive algorithm leading to blowup detection and accurate numerical approximation of blowup times. In the cases of NLS equations (1.1) we expect that the exponential term will also be proven beneficial towards the development of an efficient time-space adaptive algorithm in the spirit of [15]. This is the subject of a forthcoming paper. In this paper we investigate the behavior of the exponential term numerically in the last section of the paper.

The paper is organized as follows. In Sections 2, 3 we consider only time discretization using a relaxation Crank-Nicolson-type scheme. In particular, in Section 2 we introduce some additional notation and generalize the relaxation scheme of [9] to variable time steps. We propose a novel time reconstruction and study its properties. Section 3 is devoted to the proof of optimal order a posteriori error estimates in the  $L^\infty([0, T]; L^2(\Omega))$ -norm for the time discrete scheme. The analysis requires a careful use of the Gagliardo-Nirenberg inequality; the two conservation laws are then used for the boundedness of the solution in the  $H^1(\Omega)$ -norm. The fully discrete relaxation Crank-Nicolson finite element scheme is introduced in Section 4. In this case, we use finite element spaces that are allowed to change from one time step to another. With the help of the elliptic

reconstruction of [38] and the time reconstruction of Section 2 we define an appropriate time-space reconstruction and study its properties. In Section 5 we provide a posteriori error estimates for the fully discrete scheme. To obtain the estimates we use the machinery developed in Section 3 for the time discrete scheme as well as residual-type estimators to control the terms coming from the use of the elliptic reconstruction. The analysis is quite technical, but eventually leads to a posteriori error bounds in the  $L^\infty([0, T]; L^2(\Omega))$ -norm that are expected to be of optimal order of accuracy. Finally, in Section 6 we present numerical experiments, using uniform partitions in space and time. More specifically, we verify that the obtained estimators are indeed of optimal order of accuracy and we study the behavior of the exponential term.

## 2. TIME DISCRETE SCHEMES

As already mentioned in the Introduction, we first consider time discrete schemes, in an attempt to present clearly the main ideas of the technical analysis caused by the nonlinear nature of problem (1.1).

**2.1. A relaxation Crank-Nicolson-type method.** We consider a partition  $0 =: t_0 < t_1 < \dots < t_N := T$  of  $[0, T]$  and let  $k_n := t_{n+1} - t_n$  and  $I_n := (t_n, t_{n+1}]$ ,  $0 \leq n \leq N - 1$ , denote the variable time steps and subintervals of  $[0, T]$ , respectively. Let also  $k := \max_{1 \leq n \leq N} k_n$ . We further assume that there exists an absolute constant  $c \in \mathbb{R}^+$  such that

$$(2.1) \quad k_n \leq ck_{n-1}, \quad 1 \leq n \leq N - 1.$$

The mild constraint (2.1) between consecutive time steps appears in the analysis of other time-stepping methods with variable time steps. For example, it appears in [39], in the a posteriori error analysis for discontinuous Galerkin in time methods.

The relaxation Crank-Nicolson-type scheme for (1.1) is defined as follows: We seek approximations  $U^n \in H_0^1(\Omega)$  to  $u(t_n)$ ,  $0 \leq n \leq N$ , such that

$$(2.2) \quad \begin{cases} \frac{k_{n-1}}{k_n + k_{n-1}} \Phi^{n+\frac{1}{2}} + \frac{k_n}{k_n + k_{n-1}} \Phi^{n-\frac{1}{2}} = |U^n|^{2p}, & 0 \leq n \leq N - 1, \\ \bar{\partial}U^n - i\alpha\Delta U^{n+\frac{1}{2}} = i\lambda\Phi^{n+\frac{1}{2}}U^{n+\frac{1}{2}}, & 0 \leq n \leq N - 1, \end{cases}$$

with  $k_{-1} := k_0$ ,  $\Phi^{-\frac{1}{2}} = |u_0|^{2p}$  and  $U^0 = u_0$ . In (2.2) we also used the notation

$$(2.3) \quad \bar{\partial}U^n := \frac{U^{n+1} - U^n}{k_n} \quad \text{and} \quad U^{n+\frac{1}{2}} := \frac{U^{n+1} + U^n}{2}.$$

The relaxation scheme (2.2) was introduced for constant time steps by Besse in [9] for the numerical solution of (1.1). Here we generalize Besse's method to the case of variable time steps.

We next present briefly how method (2.2) can be obtained to make it clear to the reader that the generalisation of Besse's scheme in [9] indeed leads to (2.2) for variable time steps. First we rewrite the NLS equation in (1.1) equivalently as the system of the following two equations:

$$(2.4) \quad \begin{cases} \phi = |u|^{2p} & \text{in } \Omega \times (0, T], \\ \partial_t u - i\alpha\Delta u = i\lambda\phi u & \text{in } \Omega \times (0, T]. \end{cases}$$

Recall that the Crank-Nicolson method for (1.1) reads as: for  $0 \leq n \leq N$ , find  $U^n \in H_0^1(\Omega)$  such that

$$(2.5) \quad \bar{\partial}U^n - i\alpha\Delta U^{n+\frac{1}{2}} = i\lambda|U^{n+\frac{1}{2}}|^{2p}U^{n+\frac{1}{2}}, \quad 0 \leq n \leq N - 1,$$

with  $U^0 = u_0$ . Note that using the equivalent system-form (2.4),  $|U^{n+\frac{1}{2}}|^{2p}$  is an approximation of

$$\phi(t_{n+\frac{1}{2}}) = |u(t_{n+\frac{1}{2}})|^{2p} \quad \text{with} \quad t_{n+\frac{1}{2}} := \frac{t_{n+1} + t_n}{2}.$$

The idea now is to replace  $|U^{n+\frac{1}{2}}|^{2p}$  in (2.5) by an appropriate approximation of  $\phi(t_{n+\frac{1}{2}})$ , in an attempt to avoid the costly numerical treatment of the nonlinearity. This is where the first equation

of (2.2) is involved. In particular, it is conjectured that if,  $\Phi^{-\frac{1}{2}} = |u_0|^{2p}$  and provided regularity on the exact solution  $u$  of (1.1), then

$$(2.6) \quad |u(t_{n+\frac{1}{2}})|^{2p} = \phi(t_{n+\frac{1}{2}}) = \Phi^{n-\frac{1}{2}} + \frac{k_n + k_{n-1}}{k_{n-1}} \left( |u(t_n)|^{2p} - \Phi^{n-\frac{1}{2}} \right) + \mathcal{O}(k^2), \quad 0 \leq n \leq N-1.$$

Neglecting  $\mathcal{O}(k^2)$  in (2.6) and replacing  $\phi(t_{n+\frac{1}{2}})$  by  $\Phi^{n+\frac{1}{2}}$  and  $u(t_n)$  by  $U^n$ , we obtain the first equation in (2.2). We prove conjecture (2.6) formally:

$$(2.7) \quad \phi(t_{n+\frac{1}{2}}) = \phi(t_{n-\frac{1}{2}}) + \frac{k_n + k_{n-1}}{2} \partial_t \phi(t)|_{t=t_{n-\frac{1}{2}}} + \mathcal{O}\left(\frac{1}{2}(k_n^2 + k_{n-1}^2)\right),$$

where

$$(2.8) \quad \partial_t \phi(t)|_{t=t_{n-\frac{1}{2}}} = \frac{2}{k_{n-1}} \left( \phi(t_n) - \phi(t_{n-\frac{1}{2}}) \right) + \mathcal{O}(k_{n-1}).$$

Invoking (2.8) in (2.7) we deduce

$$\phi(t_{n+\frac{1}{2}}) = \phi(t_{n-\frac{1}{2}}) + \frac{k_n + k_{n-1}}{k_{n-1}} \left( \phi(t_n) - \phi(t_{n-\frac{1}{2}}) \right) + \mathcal{O}(k^2),$$

since  $k = \max_{0 \leq n \leq N-1} k_n$ . An inductive argument and (2.1) lead to (2.6) (recall that  $\phi(t_n) = |u(t_n)|^{2p}$ ).

Note that with scheme (2.2), the nonlinear term at each time  $t_n$  is computed explicitly, avoiding the solution of a nonlinear equation, as in the case of the Crank-Nicolson method (2.5). At the same time, the term involving the laplacian ( $\Delta u$ ) is discretized in time implicitly, preserving good stability properties of the numerical scheme (2.2). Actually, the second equation in (2.2) produces approximations that coincide with the Crank-Nicolson approximations of the linear Schrödinger equation

$$(2.9) \quad \partial_t \tilde{u} - i\alpha \Delta \tilde{u} = i\lambda V(x, t) \tilde{u} \quad \text{in } \Omega \times (0, T], \quad V(x, t) := \Phi^{n+\frac{1}{2}}(x), \quad \text{in } \Omega \times I_n,$$

with potential  $V$  is a piecewise constant in time function defined through  $\Phi^{n+\frac{1}{2}}$ ,  $0 \leq n \leq N-1$ . From this point of view, the relaxation scheme (2.2) may also be regarded as a linearised Crank-Nicolson method for the NLS problem (1.1).

Next we show formally that scheme (2.2) is expected to produce second order approximations to (1.1). To this end, combining (2.2), (2.6) and (2.1), we expect that

$$(2.10) \quad \Phi^{n+\frac{1}{2}} = |u(t_{n+\frac{1}{2}})|^{2p} + \mathcal{O}(k^2), \quad 0 \leq n \leq N-1,$$

as  $|z|^{2p}$  is a locally Lipschitz continuous function for  $p \geq \frac{1}{2}$ . Since  $U^n$ ,  $0 \leq n \leq N$ , are the Crank-Nicolson approximations to (2.9), we expect that  $\tilde{u}(t_n) - U^n = \mathcal{O}(k^2)$ ,  $0 \leq n \leq N$ . Inserting (2.10) to (2.9) gives

$$\partial_t \tilde{u} - i\alpha \Delta \tilde{u} = i\lambda \left( |u(t_{n+\frac{1}{2}})|^{2p} + \mathcal{O}(k^2) \right) \tilde{u} \quad \text{in } \Omega \times I_n,$$

for  $0 \leq n \leq N-1$ . Using once more that  $|z|^{2p}$  is a locally Lipschitz continuous function for  $p \geq \frac{1}{2}$  and (1.1) we obtain, for  $0 \leq n \leq N-1$ ,

$$\partial_t (u - \tilde{u}) - i\alpha \Delta (u - \tilde{u}) = i\lambda |u(t)|^{2p} (u - \tilde{u}) + i\lambda \mathcal{O}(k^2) \tilde{u} \quad \text{in } \Omega \times I_n.$$

Therefore from the stability of the NLS (1.1), the continuity of  $u, \tilde{u}$  in  $[0, T]$ , and the fact that  $\tilde{u}$  is a bounded function in time and space, we arrive at  $u(t) - \tilde{u}(t) = \mathcal{O}(k^2)$ ,  $t \in [0, T]$ . Thus, by writing  $u(t_n) - U^n = (u(t_n) - \tilde{u}(t_n)) + (\tilde{u}(t_n) - U^n)$ , we immediately obtain  $u(t_n) - U^n = \mathcal{O}(k^2)$ ,  $0 \leq n \leq N-1$ . For constant time steps, stability and convergence results for scheme (2.2) can be found in [9].

Moreover, the relaxation scheme (2.2) satisfies  $\|U^n\| = \|U^0\|$ ,  $0 \leq n \leq N$ , which is the discrete analogue of the mass conservation (1.4). This can easily be proven by taking the  $L^2$ -inner product with  $U^{n+\frac{1}{2}}$  in the second equation of (2.2) and then real parts and by noting that  $\Phi^{n+\frac{1}{2}}$ ,  $0 \leq n \leq N-1$ , is always real. Also, as it has been proven in [9], for  $p = 1$  and uniform time steps, the numerical scheme (2.2) also satisfies a discrete analogue of the energy conservation (1.5).

Our goal is to derive optimal order a posteriori error bounds for (1.1) when it is discretized in time by (2.2). In order to achieve this, we first define the continuous in time approximation  $U(t)$  to  $u(t)$ ,  $0 \leq t \leq T$ , by linearly interpolating between the nodal values  $U^n$  and  $U^{n+1}$ , i.e., let  $U : [0, T] \rightarrow H_0^1(\Omega)$  be defined as

$$(2.11) \quad U(t) := \ell_0^n(t)U^n + \ell_1^n(t)U^{n+1} = U^{n+\frac{1}{2}} + (t - t_{n+\frac{1}{2}})\bar{\partial}U^n, \quad t \in I_n,$$

with  $\ell_0^n(t) := \frac{t_{n+1}-t}{k_n}$  and  $\ell_1^n(t) := \frac{t-t_n}{k_n}$ ,  $0 \leq n \leq N-1$ .

It is clear that if  $u(t_n) - U^n = \mathcal{O}(k^2)$ ,  $0 \leq n \leq N$ , then for  $t \in [0, T]$ , we have  $u(t) - U(t) = \mathcal{O}(k^2)$ , as well. However, it is well known, cf., e.g., [3, 22, 36], that using  $U(t)$  in the a posteriori error analysis leads to estimates of first instead of second order of accuracy, which is the expected order of accuracy of method (2.2). For this reason we shall use a reconstruction  $\hat{U}$  of  $U$ , [3]. An alternative reconstruction for  $U$  and the scheme (2.2) can be proposed by following similar arguments as in [36].

**2.2. Time-reconstruction and its properties.** We introduce now a reconstruction  $\hat{U}$  of  $U$ . To this end,  $\hat{U}(t)$ ,  $t \in [0, T]$ , is defined to be the piecewise quadratic polynomial in  $t$ ,

$$(2.12) \quad \hat{U}(t) := U^n + i\alpha \int_{t_n}^t \Delta U(s) ds + i\lambda \int_{t_n}^t \Phi^{n+\frac{1}{2}} U(s) ds, \quad t \in I_n.$$

As expected  $\hat{U}$  coincides with the Crank-Nicolson reconstruction of [3] (see also [31]) for the linear Schrödinger equation (2.9). It is also worth mentioning that  $\hat{U}(t) \in H_0^1(\Omega)$ ,  $t \in [0, T]$ , provided compatibility conditions on the initial data, cf., [30, 31]. However, as we will see on Section 4, where fully discrete schemes are studied,  $\hat{U}(t) \in H_0^1(\Omega)$ ,  $t \in [0, T]$ , automatically, without further assumptions on the initial data. Therefore, from now on we assume that  $\hat{U}$  belongs to  $H_0^1(\Omega)$ .

**Proposition 2.1** (properties of  $\hat{U}$ ). *The reconstruction  $\hat{U}$  defined in (2.12) is equivalently written as*

$$(2.13) \quad \hat{U}(t) = U^n + \frac{i}{2}(t - t_n)(\alpha\Delta + \lambda\Phi^{n+\frac{1}{2}})(U(t) + U^n), \quad t \in I_n.$$

Moreover, it satisfies

$$(2.14) \quad \partial_t \hat{U} - i\alpha\Delta U = i\lambda\Phi^{n+\frac{1}{2}}U \quad \text{in } I_n$$

and  $\hat{U}(t_n^+) = U^n$ ,  $\hat{U}(t_{n+1}) = U^{n+1}$ ,  $0 \leq n \leq N-1$ ; in particular,  $\hat{U}$  is a time-continuous function.

*Proof.* First we note that  $\int_{t_n}^t U(s) ds = \frac{1}{2}(t - t_n)(U(t) + U^n)$ , and therefore, (2.13) is obtained from (2.12), whilst (2.14) is directly obtained by differentiation in time of (2.12). Finally,  $U(t_n^+) = U^n$  is obvious once more from (2.12), while (2.13) gives, for  $t = t_{n+1}$ ,

$$\hat{U}(t_{n+1}) = U^n + i\alpha k_n \Delta U^{n+\frac{1}{2}} + i\lambda k_n \Phi^{n+\frac{1}{2}} U^{n+\frac{1}{2}},$$

whence  $\hat{U}(t_{n+1}) = U^{n+1}$  follows from (2.2).  $\square$

The a posteriori quantity  $\hat{r}$ , by which  $\hat{U}$  misses satisfying the NLS equation in (1.1), is the residual and it is defined as

$$(2.15) \quad \hat{r} := \partial_t \hat{U} - i\alpha\Delta \hat{U} - i\lambda f(\hat{U}) \quad \text{in } I_n.$$

By virtue of (2.14), we have that  $\hat{r}$  is written as

$$(2.16) \quad \hat{r} = -i\alpha\Delta(\hat{U} - U) - i\lambda(f(\hat{U}) - f(U)) - i\lambda(|U|^{2p} - \Phi^{n+\frac{1}{2}})U \quad \text{in } I_n.$$

We would like  $\hat{r}$  to decrease as fast as the order of the method, i.e., we would like  $\hat{r}$  to be of second order of accuracy, provided that  $u(t_n) - U^n = \mathcal{O}(k^2)$ ,  $0 \leq n \leq N$ . Towards that direction, in the next proposition we compute the difference  $\hat{U} - U$ .

**Proposition 2.2** (the difference  $\hat{U} - U$ ). *The difference  $\hat{U} - U$  can be expressed as*

$$(2.17) \quad (\hat{U} - U)(t) = -\frac{i}{2}(t - t_n)(t_{n+1} - t)(\alpha\Delta + \lambda\Phi^{n+\frac{1}{2}})\bar{\partial}U^n, \quad t \in I_n.$$

*Proof.* We compute  $\hat{U} - U$  by subtracting (2.11) from (2.13) and we use (2.2). The expression (2.17) follows then by basic algebraic manipulations.  $\square$

Given that the relaxation scheme (2.2) produces second order approximations to the exact solution at the nodes  $t_n$ ,  $0 \leq n \leq N$ , we conclude that  $\hat{U} - U$  is expected to be of second order of accuracy.

Using this, (2.10), and the fact that  $f(z) = |z|^{2p}z$  is also a locally Lipschitz function for  $p \geq \frac{1}{2}$ , and resorting back to (2.16), we deduce that  $\hat{r}$  is expected to be second order accurate.

**2.3. Variational formulation & Error equation.** Problem (1.1) is equivalently written, in variational form, for  $t \in [0, T]$ , as

$$(2.18) \quad \begin{cases} \langle \partial_t u(t), v \rangle + i\alpha \langle \nabla u(t), \nabla v \rangle = i\lambda \langle f(u(t)), v \rangle, & \forall v \in H_0^1(\Omega), \\ u(\cdot, 0) = u_0 & \text{in } \Omega. \end{cases}$$

Similarly, the reconstruction  $\hat{U}$  defined in (2.12) satisfies, for  $t \in I_n$ , the problem

$$(2.19) \quad \begin{cases} \langle \partial_t \hat{U}(t), v \rangle + i\alpha \langle \nabla \hat{U}(t), \nabla v \rangle = i\lambda \langle f(\hat{U}(t)), v \rangle + \langle \hat{r}(t), v \rangle, & \forall v \in H_0^1(\Omega), \\ \hat{U}(\cdot, 0) = u_0 & \text{in } \Omega. \end{cases}$$

We denote by  $\hat{e} := u - \hat{U}$  the error between the exact solution  $u$  and the reconstruction  $\hat{U}$ . Then subtracting the first equation of (2.19) from the first equation of (2.18) we derive the basic error equation

$$(2.20) \quad \langle \partial_t \hat{e}(t), v \rangle + i\alpha \langle \nabla \hat{e}(t), \nabla v \rangle = i\lambda \langle f(u(t)) - f(\hat{U}(t)), v \rangle - \langle \hat{r}(t), v \rangle, \quad \forall v \in H_0^1(\Omega), \quad t \in I_n,$$

with  $\hat{e}(0) = 0$ . Taking  $v = \hat{e}$  in the error equation (2.20) and then real parts reveals

$$(2.21) \quad \frac{1}{2} \frac{d}{dt} \|\hat{e}(t)\|^2 = \lambda \operatorname{Re} i \langle f(u) - f(\hat{U}), \hat{e} \rangle(t) - \operatorname{Re} \langle \hat{r}, \hat{e} \rangle(t), \quad t \in I_n.$$

In Section 3 we analyse how we treat the term  $\operatorname{Re} i \langle f(u) - f(\hat{U}), \hat{e} \rangle$  on the right-hand side of (2.21) to derive an a posteriori error bound for (1.1).

### 3. A POSTERIORI ERROR CONTROL FOR TIME DISCRETE SCHEMES

In view of (2.21) it is evident that we have to handle the term  $\operatorname{Re} i \langle f(u) - f(\hat{U}), \hat{e} \rangle$ , which arises due to the nonlinear nature of the problem. To this direction we will use the Gagliardo-Nirenberg inequality (1.2), the conservation laws (1.4) and (1.5) and some other ingredients presented in the next subsection.

**3.1. Main Ingredients.** First we write

$$f(u) - f(\hat{U}) = |u|^{2p}u - |\hat{U}|^{2p}\hat{U} = |u|^{2p}(u - \hat{U}) + \hat{U}(|u|^{2p} - |\hat{U}|^{2p}) = |u|^{2p}\hat{e} + \hat{U}(|u|^{2p} - |\hat{U}|^{2p}).$$

Thus

$$(3.1) \quad \begin{aligned} \operatorname{Re} i \langle f(u) - f(\hat{U}), \hat{e} \rangle(t) &= \operatorname{Re} i \langle \hat{U} (|u|^{2p} - |\hat{U}|^{2p}), \hat{e} \rangle(t) \\ &= \left[ \operatorname{Re} i \int_{\Omega} \frac{\hat{U} (|u|^{2p} - |\hat{U}|^{2p})}{\hat{e}} \hat{e} \bar{\hat{e}} dx \right] (t) \\ &= \left[ \int_{\Omega} \operatorname{Re} i \left( \frac{\hat{U} (|u|^{2p} - |\hat{U}|^{2p})}{\hat{e}} \right) |e|^2 dx \right] (t). \end{aligned}$$

Applying the Mean Value Theorem to the real-valued function  $y^{2p}$  we have that, for every  $(x, t) \in \Omega \times [0, T]$ , there exists  $s(x, t) \in [0, 1]$  such that

$$(3.2) \quad \begin{aligned} |u|^{2p} - |\hat{U}|^{2p} &= 2p \left( s|u| + (1-s)|\hat{U}| \right)^{2p-1} (|u| - |\hat{U}|) \\ &= 2p \left( s|\hat{U} + \hat{e}| + (1-s)|\hat{U}| \right)^{2p-1} (|u| - |\hat{U}|). \end{aligned}$$



Invoking (3.2) to(3.1) we obtain

$$\begin{aligned} |\operatorname{Rei}\langle f(u) - f(\hat{U}), \hat{e} \rangle(t)| &= 2p \left| \left[ \int_{\Omega} \operatorname{Rei} \left( \frac{\hat{U} (s|\hat{U} + \hat{e}| + (1-s)|\hat{U}|)^{2p-1} (|u| - |\hat{U}|)}{\hat{e}} \right) |e|^2 dx \right] (t) \right| \\ &\leq \left[ \int_{\Omega} 2p(|\hat{U}| + s|\hat{e}|)^{2p-1} |\hat{U}| |\hat{e}|^2 dx \right] (t), \end{aligned}$$

where in the last inequality we used that  $2p - 1 \geq 0$  (as  $p \geq \frac{1}{2}$ ) and  $||u| - |\hat{U}|| \leq |u - \hat{U}| = |\hat{e}|$ . Therefore

$$|\operatorname{Rei}\langle f(u) - f(\hat{U}), \hat{e} \rangle(t)| \leq 2p \langle (|\hat{U}| + s|\hat{e}|)^{2p-1} |\hat{U}| \hat{e}, \hat{e} \rangle(t),$$

and since  $|\hat{U}| \leq |\hat{U}| + s|\hat{e}|$  (recall that  $s(x, t) \in [0, 1]$  for all  $(x, t) \in \Omega \times [0, T]$ ), we conclude that

$$(3.3) \quad |\operatorname{Rei}\langle f(u) - f(\hat{U}), \hat{e} \rangle(t)| \leq 2p \langle (|\hat{U}| + s|\hat{e}|)^{2p} \hat{e}, \hat{e} \rangle(t).$$

Next to handle the term  $(|\hat{U}| + s|\hat{e}|)^{2p}$  in (3.3) we use the standard inequality

$$(3.4) \quad (a + b)^q \leq \gamma(q)(a^q + b^q) \quad \text{with} \quad \gamma(q) := \begin{cases} 2^{q-1}, & q \geq 1 \\ 1, & 0 \leq q \leq 1, \end{cases}$$

valid for any  $a, b \geq 0$  and  $q \geq 0$ . Applying (3.4) to  $(|\hat{U}| + s|\hat{e}|)^{2p}$  we get  $(|\hat{U}| + s|\hat{e}|)^{2p} \leq 2^{2p-1}(|\hat{e}|^{2p} + |\hat{U}|^{2p})$ ; recall that  $0 \leq s \leq 1$  and  $p \geq \frac{1}{2}$ . Hence (3.3) becomes

$$(3.5) \quad |\operatorname{Rei}\langle f(u) - f(\hat{U}), \hat{e} \rangle(t)| \leq 2^{2p} p \left( \|\hat{e}(t)\|_{L^{2p+2}}^{2p+2} + \|\hat{U}(t)\|_{L^\infty}^{2p} \|\hat{e}(t)\|^2 \right).$$

To estimate  $\|\hat{e}(t)\|_{L^{2p+2}}^{2p+2}$  we refer to the Gagliardo-Nirenberg inequality (1.2):

$$(3.6) \quad \|\hat{e}(t)\|_{L^{2p+2}}^{2p+2} \leq B \|\nabla \hat{e}(t)\|^{pd} \|\hat{e}(t)\|^{p(2-d)+2} = B \|\nabla \hat{e}(t)\|^{pd} \|\hat{e}(t)\|^{p(2-d)} \|\hat{e}(t)\|^2,$$

with  $B := \beta^{2p+2}$ . Since  $d \leq 2$  and  $\frac{1}{2} \leq p \leq \frac{2}{d}$ , we have that  $p(2-d) \geq 0$  and thus by (3.4)

$$(3.7) \quad \begin{aligned} \|\hat{e}(t)\|^{p(2-d)} &\leq \gamma(p(2-d)) \left( \|u(t)\|^{p(2-d)} + \|\hat{U}(t)\|^{p(2-d)} \right) \\ &= \gamma(p(2-d)) \left( \|u_0\|^{p(2-d)} + \|\hat{U}(t)\|^{p(2-d)} \right), \end{aligned}$$

where the last equality holds because of (1.4). Set  $A := |\lambda| 2^{2p} p \max\{1, \gamma(p(2-d)) B\}$ . Then using (3.6) and (3.7) in (3.5) we deduce, for  $\lambda \neq 0$ <sup>1</sup> and  $t \in [0, T]$ ,

$$(3.8) \quad |\operatorname{Rei}\langle f(u) - f(\hat{U}), \hat{e} \rangle(t)| \leq \frac{A}{|\lambda|} \left( \left( \|u_0\|^{p(2-d)} + \|\hat{U}(t)\|^{p(2-d)} \right) \|\nabla \hat{e}(t)\|^{pd} + \|\hat{U}(t)\|_{L^\infty}^{2p} \right) \|\hat{e}(t)\|^2,$$

whereas returning back to the error equation (2.21) we see that it can be written, for  $t \in I_n$ , as

$$(3.9) \quad \frac{d}{dt} \|\hat{e}(t)\| \leq A \left( \left( \|u_0\|^{p(2-d)} + \|\hat{U}(t)\|^{p(2-d)} \right) \|\nabla \hat{e}(t)\|^{pd} + \|\hat{U}(t)\|_{L^\infty}^{2p} \right) \|\hat{e}(t)\| + \|\hat{r}(t)\|.$$

From (3.9) it is clear that we can bound the  $L^\infty([0, T]; L^2(\Omega))$ -norm of  $\hat{e}$ , using Gronwall's inequality, as long as we have an estimation for  $\|\nabla \hat{e}(t)\|$ . We do this in the next subsection.

**Remark 3.1** (the  $3d$  case). If  $d = 3$ , we have that  $p(2-d) < 0$  and (3.7) fails. This is the reason why the three dimensional spatial case cannot be included in the analysis of this paper.

<sup>1</sup>if  $\lambda = 0$  then  $A=0$ , i.e., the problem is reduced to the linear one.

**3.2. A posteriori error bound in the  $L^\infty([0, T]; L^2(\Omega))$ -norm.** In order to state the main theorem of this section, we first estimate a posteriori the gradient term in (3.9). Since  $pd \geq \frac{1}{2} > 0$ , (3.4) implies

$$(3.10) \quad \|\nabla \hat{e}(t)\|^{pd} \leq \gamma(pd) \left( \|\nabla \hat{U}(t)\|^{pd} + \|\nabla u(t)\|^{pd} \right).$$

In the next lemma we bound a posteriori the term  $\|\nabla u(t)\|$ .

**Lemma 3.1** (a posteriori bound for  $\|\nabla u(t)\|$ ). *For  $t \in [0, T]$ , the following estimate holds true*

$$(3.11) \quad \|\nabla u(t)\| \leq G(u_0),$$

with

$$(3.12) \quad G(u_0) := \begin{cases} \left( \|\nabla u_0\|^2 - \frac{\lambda}{\alpha(p+1)} \|u_0\|_{L^{2p+2}}^{2p+2} \right)^{1/2}, & \lambda \leq 0, \\ \left( \|\nabla u_0\|^{2-pd} + \frac{\lambda}{\alpha(p+1)} \left( \beta^{2p+2} \|u_0\|^{p(2-d)+2} - \frac{\|u_0\|_{L^{2p+2}}^{2p+2}}{\|\nabla u_0\|^{pd}} \right) \right)^{1/(2-pd)}, & \lambda > 0, \frac{1}{2} \leq p < \frac{2}{d}, \\ \frac{1}{(1-\Gamma(u_0))^{1/2}} \left( \|\nabla u_0\|^2 - \frac{\lambda}{\alpha(p+1)} \|u_0\|_{L^{2p+2}}^{2p+2} \right)^{1/2}, & pd = 2, \lambda > 0, \Gamma(u_0) < 1, \end{cases}$$

where  $\beta$  denotes the constant in the Gagliardo-Nirenberg inequality (1.2) and  $\Gamma(u_0)$  is given in (1.3).

*Proof.* We divide the proof into three parts. First we assume that  $\lambda \leq 0$ . Then from the energy conservation (1.5) we readily obtain

$$\|\nabla u(t)\|^2 \leq \|\nabla u_0\|^2 - \frac{\lambda}{\alpha(p+1)} \|u_0\|_{L^{2p+2}}^{2p+2}.$$

Next we assume that  $\lambda > 0$  and  $\frac{1}{2} \leq p < \frac{2}{d}$ . Using the conservation laws (1.4)-(1.5) and invoking the Gagliardo-Nirenberg inequality (1.2) to the term  $\|u(t)\|_{L^{2p+2}}^{2p+2}$  leads to

$$(3.13) \quad \begin{aligned} \|\nabla u(t)\|^{pd} & \left( \|\nabla u(t)\|^{2-pd} - \frac{B\lambda}{\alpha(p+1)} \|u_0\|^{p(2-d)+2} \right) \\ & \leq \|\nabla u_0\|^{pd} \left( \|\nabla u_0\|^{2-pd} - \frac{\lambda}{\alpha(p+1)} \frac{\|u_0\|_{L^{2p+2}}^{2p+2}}{\|\nabla u_0\|^{pd}} \right). \end{aligned}$$

Hence if  $\|\nabla u(t)\| > \|\nabla u_0\|$  we must have, from (3.13), that

$$(3.14) \quad \|\nabla u(t)\| \leq \left( \|\nabla u_0\|^{2-pd} - \frac{\lambda}{\alpha(p+1)} \frac{\|u_0\|_{L^{2p+2}}^{2p+2}}{\|\nabla u_0\|^{pd}} + \frac{B\lambda}{\alpha(p+1)} \|u_0\|^{p(2-d)+2} \right)^{1/(2-pd)},$$

because  $2 - pd > 0$ . Note now that (1.2) implies  $\frac{\|u_0\|_{L^{2p+2}}^{2p+2}}{\|\nabla u_0\|^{pd}} \leq B \|u_0\|^{p(2-d)+2}$ . This observation in combination with (3.14) give (3.11) for  $\lambda > 0$  and  $\frac{1}{2} \leq p < \frac{2}{d}$ .

Finally we consider the case  $\lambda > 0$ ,  $\Gamma(u_0) < 1$  and  $pd = 2$ , i.e., the critical exponent. Using the same argumentation as for the derivation of (3.13) we obtain

$$(1 - \Gamma(u_0)) \|\nabla u(t)\|^2 \leq \|\nabla u_0\|^2 - \frac{\lambda}{\alpha(p+1)} \|u_0\|_{L^{2p+2}}^{2p+2},$$

which implies (3.11) for this case as well, and the proof is complete.  $\square$

**Remark 3.2** (the critical exponent). In view of (3.12) it is obvious that  $G(u_0)$  blows up when  $\Gamma(u_0) \rightarrow 1$ . This is reasonable as estimate (3.11) is proven uniformly for all  $t$ , and for  $\Gamma(u_0) \geq 1$  the  $H^1$ -norm of  $u$  may blow up in finite time.

**Remark 3.3** (computational cost of  $G(u_0)$ ). Although  $G(u_0)$  in (3.12) seems complicated, it is a global quantity and it is computed only once numerically.

If we define

$$(3.15) \quad \mathcal{H}(\hat{U}, u_0; t) := A \left( \gamma(pd) \left( \|u_0\|^{p(2-d)} + \|\hat{U}(t)\|^{p(2-d)} \right) \left( G(u_0)^{pd} + \|\nabla \hat{U}(t)\|^{pd} \right) + \|\hat{U}(t)\|_{L^\infty}^{2p} \right),$$

we derive in light of (3.9), (3.10) and (3.11)

$$(3.16) \quad \frac{d}{dt} \|\hat{e}(t)\| \leq \mathcal{H}(\hat{U}, u_0; t) \|\hat{e}(t)\| + \|\hat{r}(t)\|, \quad t \in I_n.$$

Thus, using Gronwall's inequality and keeping in mind that  $\hat{e} = u - \hat{U}$  is a time-continuous function with  $\hat{e}(0) = 0$ , we arrive at the main theorem of the section.

**Theorem 3.1** (a posteriori error estimate in  $L^\infty(0, T; L^2(\Omega))$ ). *The following local and global a posteriori error estimates in the  $L^\infty([0, T]; L^2(\Omega))$ -norm are valid:*

(3.17)

$$\sup_{t \in I_n} \|(u - \hat{U})(t)\| \leq \exp \left( \int_{I_n} \mathcal{H}(\hat{U}, u_0; t) dt \right) \left( \|(u - \hat{U})(t_n)\| + \int_{I_n} \|\hat{r}(t)\| dt \right), \quad 0 \leq n \leq N - 1,$$

and

$$(3.18) \quad \|(u - \hat{U})(t)\| \leq \exp \left( \int_0^t \mathcal{H}(\hat{U}, u_0; s) ds \right) \int_0^t \|\hat{r}(s)\| ds, \quad t \in [0, T],$$

where  $\mathcal{H}(\hat{U}, u_0; t)$  is given in (3.15),  $u$  is the solution of (1.1),  $\hat{U}$  is the reconstruction (2.13) for the relaxation Crank-Nicolson-type scheme (2.2) and  $\hat{r}$  is the residual given in (2.16).  $\square$

**Remark 3.4** (focusing NLS). Estimates (3.17) and (3.18) blow up exponentially when  $\Gamma(u_0) \rightarrow 1$ , because  $G(u_0)$  (and thus  $\mathcal{H}(\hat{U}, u_0; t)$ ) blows up in this case (cf., (3.12)). Thus the methodology of this section cannot be generalised to the focusing NLS equation with the aim to control the error close to the blowup time. In contrast to the corresponding parabolic equation with blowup, cf., [15], energy methods are not appropriate for the focusing NLS with blowup and other techniques should be applied for the a posteriori error analysis of these equations.

**Remark 3.5** (local estimate (3.17)). Estimate (3.17) may be used for the proposition of an efficient adaptive algorithm as in [15, 34]. This estimate is more appropriate for adaptivity due its local nature. Note that  $\|(u - \hat{U})(t_n)\|$  comes from the previous time step, so if we have  $\max_{t \in I_{n-1}} \|(u - \hat{U})(t)\| \leq$

TOL, where TOL is a given tolerance, then TOL can replace  $\|(u - \hat{U})(t_n)\|$  in (3.17).

**3.3. An improved estimate for the one-dimensional case.** In this subsection we consider the one spatial dimension,  $d = 1$ . Let  $\eta(t)$  denote the right-hand side in (3.18) to the power  $p$ . Then in view of (3.18), (3.6) takes the form

$$\|\hat{e}(t)\|_{L^{2p+2}}^{2p+2} \leq B \|\nabla \hat{e}(t)\|^p \eta(t) \|\hat{e}(t)\|^2.$$

Plugging the above estimate in (3.5) and using (3.10) and (3.11) we deduce

$$|\operatorname{Re} \langle f(u) - f(\hat{U}), \hat{e} \rangle(t)| \leq \frac{A}{|\lambda|} \left( \eta(t) \left( G(u_0)^p + \|\nabla \hat{U}(t)\|^p \right) + \|\hat{U}(t)\|_{L^\infty}^{2p} \right) \|\hat{e}(t)\| + \|\hat{r}(t)\|, \quad t \in I_n.$$

Hence the error equation (2.21) takes now the form

$$\frac{d}{dt} \|\hat{e}(t)\| \leq \mathcal{K}(\hat{U}, u_0; t) \|\hat{e}(t)\| + \|\hat{r}(t)\|, \quad t \in I_n,$$

with

$$(3.19) \quad \mathcal{K}(\hat{U}, u_0; t) := A \left( \eta(t) \left( G(u_0)^p + \|\nabla \hat{U}(t)\|^p \right) + \|\hat{U}(t)\|_{L^\infty}^{2p} \right).$$

Thus Gronwall's inequality leads to:

**Theorem 3.2** (improved a posteriori error estimate in  $L^\infty(0, T; L^2(\Omega))$  for  $d = 1$ ). *In the case  $d = 1$ , the following estimates hold:*

(3.20)

$$\sup_{t \in I_n} \|(u - \hat{U})(t)\| \leq \exp\left(\int_{I_n} \mathcal{K}(\hat{U}, u_0; t) dt\right) \left(\|(u - \hat{U})(t_n)\| + \int_{I_n} \|\hat{r}(t)\| dt\right), \quad 0 \leq n \leq N-1,$$

(3.21)

$$\|(u - \hat{U})(t)\| \leq \exp\left(\int_0^t \mathcal{K}(\hat{U}, u_0; s) ds\right) \int_0^t \|\hat{r}(s)\| ds,$$

where  $u$  is the solution of (1.1),  $\hat{U}$  is the reconstruction (2.13) for the relaxation Crank-Nicolson-type scheme (2.2),  $\hat{r}$  denotes the residual (2.16) and  $\mathcal{K}(\hat{U}, u_0; s)$  is given by (3.19).  $\square$

**Remark 3.6** (why is (3.21) an improved estimate?). Note that since  $\eta(t) = \mathcal{O}(k^{2p})$ ,  $t \in [0, T]$ , we have that

$$(3.22) \quad \int_0^t \mathcal{K}(\hat{U}, u_0; s) ds = \mathcal{O}(k^{2p}) + A \int_0^t \|\hat{U}(s)\|_{L^\infty}^{2p} ds.$$

From this point of view, (3.21) is an improved estimate compared to (3.18).

**Remark 3.7** (the triangle inequality (3.10)). Note that for the derivation of the a posteriori error estimates in Theorems 3.1 and 3.2 we use the triangle inequality (3.10) and then we control  $\|\nabla u(t)\|$  from the data of the NLS problem (1.1). Since  $\int_0^T \|\nabla \hat{e}(s)\|^{p^d} ds$  will appear in the exponential term in the estimates, using (3.10) to estimate this term could make the estimates unpractical if  $\|\nabla u(t)\|$  or  $\int_0^T \|\nabla \hat{U}(s)\|^{p^d} ds$  (or both) are large.

This issue is substantially improved in the one-dimensional case, by using the improved estimates (3.20), (3.21) of Theorem 3.2. In that case, the terms that are coming from the approximation of  $\int_0^T \|\nabla \hat{e}(s)\|^{p^d} ds$  are multiplied by  $\eta(t) = \mathcal{O}(k^{2p})$ . In particular, in view of (3.22) (see Remark 3.6 above) the dominant term in the exponential is  $\int_0^T \|\hat{U}(s)\|_{L^\infty}^{2p} ds$ . This term appears due to the nonlinear nature of the problem (cf. (3.3) and (3.5)). If this term is large, again the estimates will be unpractical for uniform partitions. However, as in the case of [15, 34], the proposition of an efficient time-space adaptive algorithm (using the local estimate (3.20)) could provide a way to control the error.

For the two-dimensional case, where the improved estimates of Theorem 3.2 do not hold, the issue of unpractical estimates for uniform partitions may be raised when  $\|\nabla u(t)\|$  or  $\int_0^T \|\nabla \hat{U}(s)\|^{p^d} ds$  or  $\int_0^T \|\hat{U}(s)\|^{2p} ds$  (or all of them) are large. Again, the idea there is to use the local estimate (3.17) and propose an efficient time-space adaptive algorithm in order to control the error. This issue will be further investigated in a forthcoming paper.

#### 4. FULLY DISCRETE SCHEMES

In this section we study fully discrete schemes. A Galerkin-type finite element method is used for the spatial discretization while for the discretization in time we use the relaxation Crank-Nicolson-type method (2.2). We begin the section with some notation and the introduction of the fully discrete scheme.

**4.1. Notation & The method.** We keep the same notation as in time discretization section for the partition of  $[0, T]$ , i.e., we denote by  $k_n$  and  $I_n$ ,  $0 \leq n \leq N-1$ , the variable time steps and subintervals, respectively. For the spatial discretization, we follow the same notation as in [29]. More precisely, we consider a family of conforming, shape regular triangulations  $\{\mathcal{T}_n\}_{n=0}^N$  of  $\Omega$  (for  $d = 1$ , the elements of  $\mathcal{T}_n$  are just finite intervals). We additionally assume that each triangulation  $\mathcal{T}_n$ ,  $1 \leq n \leq N$ , is a refinement of a macro-triangulation of  $\Omega$  and that every two consecutive triangulations  $\mathcal{T}_n$  and  $\mathcal{T}_{n+1}$ ,  $0 \leq n \leq N-1$ , are compatible. We refer to [20, 35] for precise definitions on these properties.

For an element  $K \in \mathcal{T}_n$ , we denote its boundary by  $\partial K$  and by  $h_K$  its diameter. Let also  $\underline{h}_n := \min_{K \in \mathcal{T}_n} h_K$ . By  $h_n$  we denote the local mesh-size function on  $\mathcal{T}_n$ , defined as

$$h_n(x) := h_K \quad \text{for } K \in \mathcal{T}_n \text{ and } x \in K.$$

Let  $\Sigma_n(K)$  be the set of internal sides of  $K \in \mathcal{T}_n$  (points for  $d = 1$ , edges for  $d = 2$ ) and define  $\Sigma_n := \bigcup_{K \in \mathcal{T}_n} \Sigma_n(K)$ . To any side  $e \in \Sigma_n$ , we associate a unit vector  $\mathbf{n}_e$  on  $e$  and for  $x \in e$  and a function  $v$ , we define

$$J[\nabla v](x) := \lim_{\delta \rightarrow 0} \left[ \nabla v(x + \delta \mathbf{n}_e) - \nabla v(x - \delta \mathbf{n}_e) \right] \cdot \mathbf{n}_e.$$

To each triangulation  $\mathcal{T}_n$ , we associate the finite element space  $\mathbb{V}^n$ ,

$$\mathbb{V}^n := \{V_n \in H_0^1(\Omega) : \forall K \in \mathcal{T}_n, V_n|_K \in \mathbb{P}^r\},$$

where  $\mathbb{P}^r$  denotes the space of polynomials in  $d$  variables of degree at most  $r$ .

With  $\widehat{\mathcal{T}}_{n+1} := \mathcal{T}_{n+1} \wedge \mathcal{T}_n$  we denote the finest common coarsening triangulation of  $\mathcal{T}_{n+1}$  and  $\mathcal{T}_n$ , by  $\widehat{h}_{n+1}$  its local mesh-size function and by  $\widehat{\mathbb{V}}^{n+1} := \mathbb{V}^{n+1} \cap \mathbb{V}^n$  its corresponding finite element space. Finally, let  $\check{\Sigma}_{n+1} := \Sigma_{n+1} \cup \Sigma_n$ , and for  $K \in \widehat{\mathcal{T}}_{n+1}$ , let  $\check{\Sigma}_K^{n+1} := \check{\Sigma}_{n+1} \cap K$ , where the element  $K \in \widehat{\mathcal{T}}_{n+1}$  is taken to be closed.

To introduce a fully discrete method, we will also need the definitions of the  $L^2$ -projection and of the discrete laplacian onto  $\mathbb{V}^n$ . To this end, the  $L^2$ -projection  $\mathcal{P}^n : L^2 \rightarrow \mathbb{V}^n$  is defined as

$$\langle \mathcal{P}^n v, V_n \rangle = \langle v, V_n \rangle, \quad \forall V_n \in \mathbb{V}^n,$$

and every  $v \in L^2(\Omega)$ . Moreover, the discrete laplacian  $-\Delta^n : H_0^1(\Omega) \rightarrow \mathbb{V}^n$  is defined as

$$(4.1) \quad \langle -\Delta^n v, V_n \rangle = \langle \nabla v, \nabla V_n \rangle, \quad \forall V_n \in \mathbb{V}^n,$$

and every  $v \in H_0^1(\Omega)$ .

With the notation and definitions so far, we can now define the modified relaxation Crank-Nicolson-Galerkin-type fully discrete scheme. For  $0 \leq n \leq N$ , we seek approximations  $U^n \in \mathbb{V}^n$  to  $u(t_n)$  such that, for  $0 \leq n \leq N-1$ ,

$$(4.2) \quad \begin{cases} \frac{k_{n-1}}{k_n + k_{n-1}} \Phi^{n+\frac{1}{2}} + \frac{k_n}{k_n + k_{n-1}} \mathcal{P}^{n+1} \Phi^{n-\frac{1}{2}} = \mathcal{P}^{n+1} (|U^n|^{2p}), \\ \frac{U^{n+1} - \mathcal{P}^{n+1} U^n}{k_n} - i\alpha \frac{\Delta^{n+1} U^{n+1} + \mathcal{P}^{n+1} \Delta^n U^n}{2} = i\lambda \mathcal{P}^{n+1} \left( \Phi^{n+\frac{1}{2}} U^{n+\frac{1}{2}} \right), \end{cases}$$

with  $\Phi^{-\frac{1}{2}} = \mathcal{P}^0 (|u_0|^{2p})$ ,  $U^0 = \mathcal{P}^0 u_0$ . Scheme (4.2) is not the standard finite element scheme for (2.2). The above modified scheme was introduced by Bänsch, Karakatsani & Makridakis in [14] for the a posteriori analysis of the heat equation with mesh change. Its main advantage is that it avoids the existence of the term  $\|(\Delta^{n+1} - \Delta^n)U^n\|$  in the a posteriori error analysis, which oscillates when there is mesh change. This schemes was also used recently in [29] for linear Schrödinger equations.

**4.2. Space reconstruction.** The main tool leading to a posteriori error estimates of optimal order in space in the  $L^\infty([0, T]; L^2(\Omega))$ -norm via energy techniques is the elliptic reconstruction. It was originally introduced by Makridakis & Nochetto in [38] for finite element semidiscrete schemes for the heat equation.

For  $V_n \in \mathbb{V}^n$ , the *elliptic reconstruction*  $\mathcal{R}^n V_n \in H_0^1(\Omega)$  of  $V_n$  is defined to be the unique solution of the elliptic equation

$$(4.3) \quad \langle \nabla \mathcal{R}^n V_n, \nabla v \rangle = \langle -\Delta^n V_n, v \rangle, \quad \forall v \in H_0^1(\Omega).$$

Using the elliptic reconstruction, we define the *space reconstruction*  $\omega : [0, T] \in H_0^1(\Omega)$  of the piecewise linear interpolated  $U$  (see (2.11)) as

$$(4.4) \quad \omega(t) := \ell_0^n(t) \mathcal{R}^n U^n + \ell_1^n(t) \mathcal{R}^{n+1} U^{n+1}, \quad t \in I_n.$$

The space reconstruction (4.4) will allow us to handle efficiently a posteriori the spatial error using the elliptic theory. More precisely, using  $\omega$  in the analysis below, terms of the form  $\|(\mathcal{R}^n - \mathbf{I})V_n\|$ ,  $\|(\mathcal{R}^n - \mathbf{I})V_n\|_{L^\infty}$  and  $\|(\mathcal{R}^{n+1} - \mathbf{I})V_{n+1} - (\mathcal{R}^n - \mathbf{I})V_n\|$  will appear. To estimate these terms, we will use residual-type elliptic error estimators.

To this end, for a given  $V_n \in \mathbb{V}^n$ ,  $0 \leq n \leq N$ , we define the following  $L^2$ - and  $L^\infty$ -residual-type estimators:

$$(4.5) \quad \eta_{2, \mathbb{V}^n}(V_n) := \left\{ \sum_{K \in \mathcal{T}_n} \left( \|h_K^2(\Delta - \Delta^n)V_n\|_{L^2(K)}^2 + \|h_K^{\frac{3}{2}}J[\nabla V_n]\|_{L^2(\partial K)}^2 \right) \right\}^{\frac{1}{2}},$$

$$(4.6) \quad \eta_{\infty, \mathbb{V}^n}(V_n) := \max_{K \in \mathcal{T}_n} \left\{ \|h_K^2(\Delta - \Delta^n)V_n\|_{L^\infty(K)} + \|h_K J[\nabla V_n]\|_{L^\infty(\partial K)} \right\},$$

where, for  $p = 1, \infty$ ,  $\|\cdot\|_{L^p(K)}$  and  $\|\cdot\|_{L^p(\partial K)}$  denote the  $L^p$ -norm in  $K$  and on  $\partial K$ , respectively. In the one-dimensional case,  $d = 1$ , the term with the discontinuities in (4.5) and (4.6) vanishes. For  $V_{n+1} \in \mathbb{V}^{n+1}$  and  $V_n \in \mathbb{V}^n$ ,  $0 \leq n \leq N$ , we also define

$$(4.7) \quad \eta_{2, \widehat{\mathbb{V}}^{n+1}}(V_{n+1}, V_n) := \left\{ \sum_{K \in \widehat{\mathcal{T}}_n} \left( \|h_K^2[(\Delta - \Delta^{n+1})V_{n+1} - (\Delta - \Delta^n)V_n]\|_{L^2(K)}^2 + \|h_K^{\frac{3}{2}}J[\nabla V_{n+1} - \nabla V_n]\|_{L^2(\widehat{\Sigma}_K^{n+1})}^2 \right) \right\}^{\frac{1}{2}}.$$

In view of the definitions (4.5) and (4.7), the next lemma is standard. For its proof we refer, for example, in [38, 35] (for the case  $d = 1$ , see [30]).

**Lemma 4.1.** *For all  $V_n \in \mathbb{V}^n$ ,  $0 \leq n \leq N$ , we have*

$$(4.8) \quad \|(\mathcal{R}^n - \mathbb{I})V_n\| \leq C_2 \eta_{2, \mathbb{V}^n}(V_n),$$

where  $C_2$  depends only on  $\Omega$  and the shape regularity of the family of triangulations  $\{\mathcal{T}_n\}_{n=0}^N$ . Furthermore, for all  $V_{n+1} \in \mathbb{V}^{n+1}$  and  $V_n \in \mathbb{V}^n$ ,  $0 \leq n \leq N - 1$ , it holds

$$(4.9) \quad \|(\mathcal{R}^{n+1} - \mathbb{I})V_{n+1} - (\mathcal{R}^n - \mathbb{I})V_n\| \leq \widehat{C}_2 \eta_{2, \widehat{\mathbb{V}}^{n+1}}(V_{n+1}, V_n),$$

where  $\widehat{C}_2$  depends only on  $\Omega$ , the shape regularity of the triangulations, and the number of refinement steps necessary to pass from  $\mathcal{T}_n$  to  $\mathcal{T}_{n+1}$ .  $\square$

A similar estimate to (4.8) holds for  $\|(\mathcal{R}^n - \mathbb{I})V_n\|_{L^\infty}$ ; for its proof we refer to [43] (for  $d = 1$ , see [30]).

**Lemma 4.2.** *For every  $V_n \in \mathbb{V}^n$ ,  $0 \leq n \leq N$ , the following estimate is valid:*

$$(4.10) \quad \|(\mathcal{R}^n - \mathbb{I})V_n\|_{L^\infty} \leq C_\infty (\ln \underline{h}_n)^2 \eta_{\infty, \mathbb{V}^n}(V_n),$$

where  $C_\infty$  depends only on  $\Omega$  and the shape regularity of the family of triangulations.  $\square$

**4.3. Time-space reconstruction and its properties.** As already discussed in Section 2 to handle efficiently the time-error due to the discretization with the relaxation Crank-Nicolson scheme, we need a reconstruction in time. Similarly, for the fully discrete scheme (4.2) we will use a time reconstruction of  $\omega$ . Thus, we end up with a time-space reconstruction  $\widehat{U} : [0, T] \rightarrow H_0^1(\Omega)$  which is defined as follows:

$$(4.11) \quad \begin{aligned} \widehat{U}(t) := & \mathcal{R}^n U^n + \frac{t - t_n}{k_n} (\mathcal{R}^{n+1} \mathcal{P}^{n+1} U^n - \mathcal{R}^n U^n) + i\alpha \int_{t_n}^t \mathcal{R}^{n+1} \Theta(s) ds \\ & + i\lambda \int_{t_n}^t \mathcal{R}^{n+1} \mathcal{P}^{n+1} \left( \Phi^{n+\frac{1}{2}} U(s) \right) ds, \quad t \in I_n, \end{aligned}$$

$$(4.12) \quad \Theta(t) := \ell_0^n(t) \mathcal{P}^{n+1} \Delta^n U^n + \ell_1^n(t) \Delta^{n+1} U^{n+1}, \quad t \in I_n;$$

compare with (2.12). We choose to use the same notation  $\widehat{U}$  for the time-space reconstruction as for the time-reconstruction in time discrete schemes (cf., Sections 2, 3). This is done in an attempt to simplify the notation as well as for a direct comparison with the time discrete schemes. Note that  $\widehat{U}$  in (4.11) coincides with the time-space reconstruction of [29] for the linear Schrödinger equation (2.9). Denoting by

$$(4.13) \quad W(t) := i\alpha \Theta(t) + i\lambda \mathcal{P}^{n+1} \left( \Phi^{n+\frac{1}{2}} U(t) \right), \quad t \in I_n,$$

we have that  $\hat{U}$  is equivalently written as

$$(4.14) \quad \hat{U}(t) = \mathcal{R}^n U^n + \frac{t - t_n}{k_n} (\mathcal{R}^{n+1} \mathcal{P}^{n+1} U^n - \mathcal{R}^n U^n) + \int_{t_n}^t \mathcal{R}^{n+1} W(s) ds, \quad t \in I_n,$$

whilst the second equation in method (4.2) is written as

$$(4.15) \quad \frac{U^{n+1} - \mathcal{P}^{n+1} U^n}{k_n} - W(t_{n+\frac{1}{2}}) = 0, \quad 0 \leq n \leq N-1.$$

For  $0 \leq n \leq N-1$ ,  $t \in I_n$ , we further define

$$(4.16) \quad \tilde{\partial} \Theta^{n+1} := \partial_t \Theta(t) = \frac{\Delta^{n+1} U^{n+1} - \mathcal{P}^{n+1} \Delta^n U^n}{k_n}$$

and similarly,

$$(4.17) \quad \tilde{\partial} W^{n+1} := \partial_t W(t) = i\alpha \tilde{\partial} \Theta^n + i\lambda \mathcal{P}^{n+1} \left( \Phi^{n+\frac{1}{2}} \bar{\partial} U^n \right).$$

Using (4.14), (4.15) and notations (4.13), (4.16), (4.17) we can prove the next propositions. We state them here without any proofs and we refer to [29] (Proposition 2.1 and Lemma 2.5) for further details (compare also with Propositions 2.1 and 2.2 of this paper).

**Proposition 4.1** (properties of  $\hat{U}$ ). *The reconstruction  $\hat{U}$  in (4.11) satisfies  $\hat{U}(t_n^+) = \mathcal{R}^n U^n$  and  $\hat{U}(t_{n+1}) = \mathcal{R}^{n+1} U^{n+1}$ ,  $0 \leq n \leq N-1$ . In particular,  $\hat{U}$  is continuous in time on  $[0, T]$ . Furthermore it satisfies*

$$(4.18) \quad \partial_t \hat{U} - i\alpha \mathcal{R}^{n+1} \Theta = i\lambda \mathcal{R}^{n+1} \mathcal{P}^{n+1} \left( \Phi^{n+\frac{1}{2}} U \right) + \frac{\mathcal{R}^{n+1} \mathcal{P}^{n+1} U^n - \mathcal{R}^n U^n}{k_n} \quad \text{in } I_n. \quad \square$$

**Proposition 4.2** (the difference  $\hat{U} - \omega$ ). *The difference  $\hat{U} - \omega$  can be expressed as*

$$(4.19) \quad \begin{aligned} (\hat{U} - \omega)(t) &= -\frac{1}{2} (t_{n+1} - t)(t - t_n) \mathcal{R}^{n+1} \tilde{\partial} W^{n+1} \\ &= -\frac{i}{2} (t_{n+1} - t)(t - t_n) \mathcal{R}^{n+1} \left( \alpha \tilde{\partial} \Theta^{n+1} + \lambda \mathcal{P}^{n+1} \left( \Phi^{n+\frac{1}{2}} \bar{\partial} U^n \right) \right) \quad t \in I_n. \quad \square \end{aligned}$$

## 5. A POSTERIORI ERROR CONTROL FOR FULLY DISCRETE SCHEMES

We are now ready to prove the main a posteriori error bound for the fully discrete scheme (4.2). We split the error  $e := u - U$  as

$$e = u - U := \hat{\rho} + \sigma + \epsilon \quad \text{with} \quad \hat{\rho} := u - \hat{U}, \quad \sigma := \hat{U} - \omega \quad \text{and} \quad \epsilon := \omega - U,$$

and we refer to  $\hat{\rho}$  as the main error, to  $\sigma$  as the time-reconstruction error, and to  $\epsilon$  as the elliptic error.

**5.1. Estimation of  $\sigma$  &  $\epsilon$ .** In this subsection we give two simple propositions for the a posteriori estimation of the time-reconstruction and elliptic errors.

**Proposition 5.1** (estimation of  $\sigma$ ). *For  $0 \leq n \leq N-1$ , the following estimates hold:*

$$(5.1) \quad \max_{t_n \leq t \leq t_{n+1}} \|\sigma(t)\| \leq \varepsilon_{n+1}^{\text{T},0} \quad \text{with} \quad \varepsilon_{n+1}^{\text{T},0} := \frac{k_n^2}{8} \left[ \|\tilde{\partial} W^{n+1}\| + C_2 \eta_{2, \mathbb{V}^{n+1}}(\tilde{\partial} W^{n+1}) \right] \quad \text{and}$$

$$(5.2) \quad \max_{t_n \leq t \leq t_{n+1}} \|\sigma(t)\|_{L^\infty} \leq \varepsilon_{n+1}^{\text{T},\infty} \quad \text{with} \quad \varepsilon_{n+1}^{\text{T},\infty} := \frac{k_n^2}{8} \left[ \|\tilde{\partial} W^{n+1}\|_{L^\infty} + C_\infty (\ln \underline{h}_{n+1})^2 \eta_{\infty, \mathbb{V}^{n+1}}(\tilde{\partial} W^{n+1}) \right].$$

In particular, for  $1 \leq m \leq N$ , we have that

$$(5.3) \quad \max_{0 \leq t \leq t_m} \|\sigma(t)\| \leq \mathcal{E}_m^{\text{T},0} \quad \text{with} \quad \mathcal{E}_m^{\text{T},0} := \max_{1 \leq n \leq m} \varepsilon_n^{\text{T},0} \quad \text{and}$$

$$(5.4) \quad \max_{0 \leq t \leq t_m} \|\sigma(t)\|_{L^\infty} \leq \mathcal{E}_m^{\text{T},\infty} \quad \text{with} \quad \mathcal{E}_m^{\text{T},\infty} := \max_{1 \leq n \leq m} \varepsilon_n^{\text{T},\infty}.$$

*Proof.* We write  $\mathcal{R}^{n+1}\tilde{\partial}W^{n+1} = \tilde{\partial}W^{n+1} + (\mathcal{R}^{n+1} - \mathbf{I})\tilde{\partial}W^{n+1}$  and the estimates follow directly from (4.19) and the elliptic properties (4.5), (4.6).  $\square$

**Proposition 5.2** (estimation of  $\epsilon$ ). *The next local estimates are valid, for  $0 \leq n \leq N - 1$ ,*

$$(5.5) \quad \max_{t_n \leq t \leq t_{n+1}} \|\epsilon(t)\| \leq C_2 \varepsilon_{n+1}^{S,0} \quad \text{with} \quad \varepsilon_{n+1}^{S,0} := \max \{ \eta_{2, \mathbb{V}^{n+1}}(U^{n+1}), \eta_{2, \mathbb{V}^n}(U^n) \} \quad \text{and}$$

$$(5.6) \quad \max_{t_n \leq t \leq t_{n+1}} \|\epsilon(t)\|_{L^\infty} \leq C_\infty \varepsilon_{n+1}^{S,\infty} \quad \text{with} \quad \varepsilon_{n+1}^{S,\infty} := \max \left\{ (\ln \underline{h}_{n+1})^2 \eta_{\infty, \mathbb{V}^{n+1}}(U^{n+1}), (\ln \underline{h}_n)^2 \eta_{\infty, \mathbb{V}^n}(U^n) \right\}.$$

*In particular, for  $1 \leq m \leq N$ , the following global estimates hold true:*

$$(5.7) \quad \max_{0 \leq t \leq t_m} \|\epsilon(t)\| \leq C_2 \mathcal{E}_m^{S,0} \quad \text{with} \quad \mathcal{E}_m^{S,0} := \max_{0 \leq n \leq m} \eta_{2, \mathbb{V}^n}(U^n),$$

$$(5.8) \quad \max_{0 \leq t \leq t_m} \|\epsilon(t)\|_{L^\infty} \leq C_\infty \mathcal{E}_m^{S,\infty} \quad \text{with} \quad \mathcal{E}_m^{S,\infty} := \max_{0 \leq n \leq m} (\ln \underline{h}_n)^2 \eta_{\infty, \mathbb{V}^n}(U^n).$$

*Proof.* For  $t \in I_n$ , we write  $\epsilon(t) = \ell_0^n(t)(\mathcal{R}^n - \mathbf{I})U^n + \ell_1^{n+1}(t)(\mathcal{R}^{n+1} - \mathbf{I})U^{n+1}$  and the estimates follow from (4.5) and (4.6).  $\square$

**5.2. Estimation of  $\hat{\rho}$ .** In this subsection we estimate a posteriori the main error  $\hat{\rho}$  using energy techniques. The estimation of  $\hat{\rho}$  is based on the analysis of Section 3. We first have from (4.18) that  $\hat{U}$  satisfies, in  $I_n$ ,  $0 \leq n \leq N - 1$ , the equation

$$(5.9) \quad \langle \partial_t \hat{U}, v \rangle + i\alpha \langle \nabla \hat{U}, \nabla v \rangle = i\lambda \langle f(\hat{U}), v \rangle + \langle R, v \rangle, \quad \forall v \in H_0^1(\Omega)$$

with

$$R(t) := \mathcal{R}^{n+1}W(t) + \frac{\mathcal{R}^{n+1}\mathcal{P}^{n+1}U^n - \mathcal{R}^n U^n}{k_n} - i\lambda f(\hat{U}(t)) - i\alpha \Delta(\omega + \sigma)(t), \quad t \in I_n.$$

Furthermore, the definition of the elliptic reconstruction (4.3) and (4.4) lead to

$$\langle \Delta \omega, v \rangle = \ell_0^n(t) \langle \Delta^n U^n, v \rangle + \ell_1^n(t) \langle \Delta^{n+1} U^{n+1}, v \rangle, \quad \forall v \in H_0^1(\Omega)$$

and by (4.19)

$$\langle \Delta \sigma, v \rangle = -\frac{1}{2}(t_{n+1} - t)(t - t_n) \langle \Delta^{n+1} \tilde{\partial}W^{n+1}, v \rangle, \quad \forall v \in H_0^1(\Omega).$$

Therefore, subtracting (5.9) from the first equation of (2.18) and using the above, leads to the next proposition:

**Proposition 5.3** (error equation for  $\hat{\rho}$ ). *The main error  $\hat{\rho}$  satisfies the following equation in  $I_n$ ,  $0 \leq n \leq N - 1$ ,*

$$(5.10) \quad \langle \partial_t \hat{\rho}, v \rangle + i\alpha \langle \nabla \hat{\rho}, \nabla v \rangle = i\lambda \langle f(u) - f(\hat{U}), v \rangle + \sum_{j=1}^3 \langle R_j, v \rangle, \quad \forall v \in H_0^1(\Omega),$$

where the residuals  $R_j$ ,  $1 \leq j \leq 3$ , are given by

$$(5.11) \quad R_1(t) := (\mathbf{I} - \mathcal{R}^{n+1})W(t) - \frac{\mathcal{R}^{n+1}\mathcal{P}^{n+1}U^n - \mathcal{R}^n U^n}{k_n} + i\alpha \ell_0^n(t)(\mathbf{I} - \mathcal{P}^{n+1})\Delta^n U^n,$$

$$(5.12) \quad R_2(t) := -\frac{i\alpha}{2}(t_{n+1} - t)(t - t_n)\Delta^{n+1}\tilde{\partial}W^{n+1},$$

$$(5.13) \quad R_3(t) := i\lambda \left( f(\hat{U}) - \mathcal{P}^{n+1} \left( \Phi^{n+\frac{1}{2}} U \right) \right) (t). \quad \square$$

The next lemma is taken from [29] (Lemma 3.1) and we refer there for its proof.



**Lemma 5.1** (the residual  $R_1$ ). *For  $t \in I_n$ , the residual  $R_1$  can be rewritten equivalently as*

$$(5.14) \quad R_1(t) = (t - t_{n+\frac{1}{2}})(\mathbf{I} - \mathcal{R}^{n+1})\tilde{\delta}W^{n+1} - \frac{(\mathcal{R}^{n+1} - \mathbf{I})U^{n+1} - (\mathcal{R}^n - \mathbf{I})U^n}{k_n} \\ + (\mathbf{I} - \mathcal{P}^{n+1}) \left( i\alpha \ell_0^n(t) \Delta^n U^n + \frac{U^n}{k_n} \right). \quad \square$$

Equation (5.10) is of the same form as (2.20). Therefore, to estimate a posteriori  $\hat{\rho}$  in the  $L^\infty([0, T]; L^2(\Omega))$ -norm we will follow similar arguments as for the time discrete schemes. However note that in the case of fully discrete schemes the reconstruction  $\hat{U}$  (as well as  $\omega$ ) are no longer computable quantities. Thus before proceeding as in the time discrete case we need two auxiliary lemmata that will allow us to handle this issue.

The first one gives a corresponding to (3.10) estimate for  $\|\hat{\rho}(t)\|^{pd}$ . In the proof of this lemma we use, for  $d = 2$ , the Clément interpolant ([11, 18, 47]) or, for  $d = 1$ , the piecewise linear interpolant  $\hat{\mathcal{I}}_{n+1}z \in \hat{\mathbb{V}}^{n+1} := \mathbb{V}^{n+1} \cap \mathbb{V}^n$  of  $z \in H_0^1(\Omega)$ . We summarise next some of the properties of  $\hat{\mathcal{I}}_{n+1}z$  [11, 18, 47, 30]. The following estimates hold:

$$(5.15) \quad \|\nabla(z - \hat{\mathcal{I}}_{n+1}z)\| \leq D_{2,0}\|\nabla z\| \quad \text{and}$$

$$(5.16) \quad \sum_{K \in \hat{\mathcal{T}}_n} \|h_K^{-1}(z - \hat{\mathcal{I}}_{n+1}z)\|_{L^2(K)}^2 \leq D_{2,1}^2\|\nabla z\|^2,$$

where  $D_{2,0}, D_{2,1}$  are absolute constants depending only on the shape regularity of the family of triangulations and on the number of bisections necessary to pass from  $\mathcal{T}_n$  to  $\mathcal{T}_{n+1}$ .

**Lemma 5.2** (estimation of  $\|\nabla\hat{\rho}(t)\|^{pd}$ ). *For  $t \in I_n$ , we define*

$$(5.17) \quad M(t) := \Delta^n U^n + \frac{t - t_n}{k_n} (\Delta^{n+1} \mathcal{P}^{n+1} U^n - \Delta^n U^n) + \frac{t - t_n}{2} \Delta^{n+1} (W(t) + W(t_n)) \quad \text{and}$$

$$(5.18) \quad Z(t) := U^n + \frac{t - t_n}{k_n} (\mathcal{P}^{n+1} U^n - U^n) + \frac{t - t_n}{2} (W(t) + W(t_n)).$$

*Then the following estimate holds for  $t \in I_n$ ,  $0 \leq n \leq N - 1$ ,*

$$(5.19) \quad \|\nabla\hat{\rho}(t)\| \leq \|\nabla u(t)\| + \max\{D_{2,0}, D_{2,1}\} \left( \|\nabla Z(t)\| + \|\hat{h}_{n+1}M(t)\| \right).$$

*In particular, due to (3.4) we have that*

$$(5.20) \quad \|\nabla\hat{\rho}(t)\|^{pd} \leq \gamma(pd) \left( \|\nabla u(t)\|^{pd} + \left[ \max\{D_{2,0}, D_{2,1}\} \left( \|\nabla Z(t)\| + \|\hat{h}_{n+1}M(t)\| \right) \right]^{pd} \right).$$

*Proof.* Since  $\nabla\hat{\rho} = \nabla u - \nabla\hat{U}$ , we write

$$(5.21) \quad \|\nabla\hat{\rho}(t)\|^2 = \langle \nabla(u - \hat{U}), \nabla\hat{\rho}(t) \rangle = \langle \nabla u, \nabla\hat{\rho}(t) \rangle - \langle \nabla\hat{U}, \nabla\hat{\rho}(t) \rangle.$$

Further, since  $\hat{\rho}(\cdot, t) \in H_0^1(\Omega)$  for all  $t \in [0, T]$ , from the definition of the elliptic reconstruction (4.3) we obtain

$$-\langle \nabla \mathcal{R}^n U^n, \nabla\hat{\rho}(t) \rangle = \langle \Delta^n U^n, \hat{\rho}(t) \rangle, \quad -\langle \mathcal{R}^{n+1} \mathcal{P}^{n+1} U^n - \mathcal{R}^n U^n, \nabla\hat{\rho}(t) \rangle = \langle \Delta^{n+1} \mathcal{P}^{n+1} U^n - \Delta^n U^n, \hat{\rho}(t) \rangle,$$

and

$$-\langle \int_{t_n}^t \nabla \mathcal{R}^{n+1} W(s) ds, \nabla\hat{\rho}(t) \rangle = \langle \int_{t_n}^t \Delta^{n+1} W(s) ds, \hat{\rho}(t) \rangle = \frac{t - t_n}{2} \langle \Delta^{n+1} (W(t) + W(t_n)), \hat{\rho}(t) \rangle;$$

cf. (4.13) for the last equality. Using the relations above and the definition (4.14) of  $\hat{U}$ , we deduce

$$-\langle \nabla\hat{U}, \nabla\hat{\rho}(t) \rangle = \langle M, \hat{\rho}(t) \rangle \leq \|M(t)\|_{H^{-1}} \|\nabla\hat{\rho}(t)\|.$$

Therefore, (5.21) takes the form

$$(5.22) \quad \|\nabla\hat{\rho}(t)\|^2 \leq (\|\nabla u(t)\| + \|M(t)\|_{H^{-1}}) \|\nabla\hat{\rho}(t)\|, \quad \text{or} \quad \|\nabla\hat{\rho}(t)\| \leq \|\nabla u(t)\| + \|M(t)\|_{H^{-1}}.$$

For the estimation of  $\|M(t)\|_{H^{-1}}$  we use the definition of the discrete laplacian (4.1) to obtain

$$(5.23) \quad \begin{aligned} \|M(t)\|_{H^{-1}} &= \sup_{\substack{z \in H_0^1(\Omega) \\ z \neq 0}} \frac{\langle M(t), z \rangle}{\|\nabla z\|} = \sup_{\substack{z \in H_0^1(\Omega) \\ z \neq 0}} \left\{ \frac{\langle M(t), \widehat{\mathcal{I}}_{n+1} z \rangle}{\|\nabla z\|} + \frac{\langle M(t), z - \widehat{\mathcal{I}}_{n+1} z \rangle}{\|\nabla z\|} \right\}, \\ &= \sup_{\substack{z \in H_0^1(\Omega) \\ z \neq 0}} \left\{ \frac{\langle \nabla Z(t), \nabla \widehat{\mathcal{I}}_{n+1} z \rangle}{\|\nabla z\|} + \frac{\langle \widehat{h}_{n+1} M(t), \widehat{h}_{n+1}^{-1}(z - \widehat{\mathcal{I}}_{n+1} z) \rangle}{\|\nabla z\|} \right\} \end{aligned}$$

where, recall that  $\widehat{\mathcal{I}}_{n+1} z \in \widehat{\mathbb{V}}^{n+1}$  denotes, for  $d = 2$ , the Clément interpolant and, for  $d = 1$ , the piecewise linear interpolant of  $z \in H_0^1(\Omega)$ . The desired result (5.19) is then obtained by applying (5.15), (5.16) in the last equality of (5.23) and inserting the result back to (5.22).  $\square$

Next we would like to deduce an estimate for  $R_3$ . For this we write  $R_3 := i\lambda(R_{31} + R_{32} + R_{33})$  with

$$(5.24) \quad R_{31} := f(\widehat{U}) - f(\omega), \quad R_{32} := f(\omega) - f(U), \quad \text{and} \quad R_{33} := f(U) - \mathcal{P}^{n+1} \left( \Phi^{n+\frac{1}{2}} U \right).$$

The term  $R_{33}$  is already an a posteriori quantity, while we estimate  $R_{31}$  and  $R_{32}$  in the next lemma.

**Lemma 5.3** (estimation of  $R_{31}$  and  $R_{32}$ ). *For  $t \in I_n$ ,  $0 \leq n \leq N - 1$ , we have that*

$$(5.25) \quad \|R_{31}(t)\| \leq \mathcal{L}_{31}^{n+1} (t_{n+1} - t)(t - t_n) \quad \text{with}$$

$$(5.26)$$

$$\mathcal{L}_{31}^{n+1} := \left( p + \frac{1}{2} \right) \left( \varepsilon_{n+1}^{\text{T},\infty} + \varepsilon_{n+1}^{\text{S},\infty} + \max \{ \|U^n\|_{L^\infty}, \|U^{n+1}\|_{L^\infty} \} \right)^{2p} \left( \|\tilde{\delta} W^{n+1}\| + C_2 \eta_{2,\mathbb{V}^{n+1}}(\tilde{\delta} W^{n+1}) \right)$$

and

$$(5.27) \quad \|R_{32}(t)\| \leq C_2 \mathcal{L}_{32}^{n+1} \max \{ \eta_{2,\mathbb{V}^{n+1}}(U^{n+1}), \eta_{2,\mathbb{V}^n}(U^n) \} \quad \text{with}$$

$$(5.28) \quad \mathcal{L}_{32}^{n+1} := (2p + 1) \left( C_\infty \varepsilon_{n+1}^{\text{S},\infty} + \max \{ \|U^n\|_{L^\infty}, \|U^{n+1}\|_{L^\infty} \} \right)^{2p},$$

where  $C_2, C_\infty$  are the constants in (4.8) and (4.10), respectively, and  $\varepsilon_{n+1}^{\text{T},\infty}, \varepsilon_{n+1}^{\text{S},\infty}$  are the estimators in (5.2) and (5.6), respectively.

*Proof.* Using (??) with  $z_1 = \widehat{U}$  and  $z_2 = \omega$  and  $v = \sigma$  we estimate

$$|R_{31}| \leq (2p + 1) |s\widehat{U} + (1 - s)\omega|^{2p} |\sigma| = (2p + 1) |s\sigma + \epsilon + U|^{2p} |\sigma|,$$

where recall that  $s = s(t) \in [0, 1]$ . Furthermore, since  $2p \geq 1$ , we have that

$$\|R_{31}(t)\| \leq (2p + 1) (\|\sigma(t)\|_{L^\infty} + \|\epsilon\|_{L^\infty} + \|U(t)\|_{L^\infty})^{2p} \|\sigma(t)\|, \quad t \in I_n.$$

Hence, estimate (5.25) is then obtained using (5.2), (5.6), the definition of  $U$  and (4.19). Similarly we deduce estimate (5.27), but now instead of using (5.2) and (4.19), we use (5.5).  $\square$

We are now ready to state and prove the main theorem of the subsection.

**Theorem 5.1** (estimation of  $\hat{\rho}$ ). *For  $t \in I_n$ , let*

$$\begin{aligned} \mathcal{M}(U, u_0; t) &:= A \left( \gamma(pd) \left( \|u_0\|^{p(2-d)} + \left( \varepsilon_{n+1}^{\text{T},0} + C_2 \varepsilon_{n+1}^{\text{S},0} + \|U(t)\| \right)^{p(2-d)} \right) \right. \\ &\quad \left. \times \left( G(u_0)^{pd} + \left[ \max \{ D_{2,0}, D_{2,1} \} \left( \|\nabla Z(t)\| + \|\widehat{h}_{n+1} M(t)\| \right) \right]^{pd} \right) + \left( \varepsilon_{n+1}^{\text{T},\infty} + C_\infty \varepsilon_{n+1}^{\text{S},\infty} + \|U(t)\|_{L^\infty} \right)^{2p} \right), \end{aligned}$$

where  $A$  is the constant in (3.8),  $\gamma(pd)$  is as in (3.4),  $C_2, C_\infty$  are the constants in (4.8), (4.9),  $D_{2,0}, D_{2,1}$  are the constants in (5.15), (5.16),  $\varepsilon_{n+1}^{\text{T},0}, \varepsilon_{n+1}^{\text{T},\infty}, \varepsilon_{n+1}^{\text{S},0}, \varepsilon_{n+1}^{\text{S},\infty}$  are as in (5.1), (5.2), (5.5), (5.6), respectively.  $M(t), Z(t)$  are given by (5.17), (5.18),  $G(u_0)$  is given by (3.12).

Then, for  $0 \leq n \leq N-1$ , the following local estimate holds true for the main error  $\hat{\rho} := u - \hat{U}$ :

$$(5.29) \quad \sup_{t \in I_n} \|\hat{\rho}(t)\| \leq \exp \left( \int_{I_n} \mathcal{M}(U, u_0; t) dt \right) \\ \times \left( \|\hat{\rho}(t_n)\| + \varepsilon_{n+1}^{\text{T},1} + \varepsilon_{n+1}^{\text{T},2} + C_2(\varepsilon_{n+1}^{\text{S},1} + \varepsilon_{n+1}^{\text{S},2}) + \widehat{C}_2 \varepsilon_{n+1}^{\text{S},3} + \varepsilon_{n+1}^{\text{C}} + \varepsilon_{n+1}^{\text{D}} \right)$$

with

$$\varepsilon_{n+1}^{\text{T},1} := \frac{\alpha k_n^3}{12} \|\Delta^{n+1} \tilde{\partial} W^{n+1}\| \quad \text{and} \quad \varepsilon_{n+1}^{\text{T},2} := \frac{k_n^3}{6} \mathcal{L}_{31}^{n+1}, \\ \varepsilon_{n+1}^{\text{S},1} := \frac{k_n^2}{4} \eta_{2, \mathbb{V}^{n+1}}(\tilde{\partial} W^{n+1}), \quad \varepsilon_{n+1}^{\text{S},2} := \mathcal{L}_{32}^{n+1} \max \{ \eta_{2, \mathbb{V}^{n+1}}(U^{n+1}), \eta_{2, \mathbb{V}^n}(U^n) \}, \\ \text{and} \quad \varepsilon_{n+1}^{\text{S},3} := k_n \eta_{2, \widehat{\mathbb{V}}^{n+1}} \left( \frac{U_{n+1}}{k_n}, \frac{U_n}{k_n} \right), \\ \varepsilon_{n+1}^{\text{C}} := \int_{I_n} \|(I - \mathcal{P}^{n+1}) \left( \frac{U^n}{k_n} + i\alpha \ell_0^n(t) \Delta^n U^n \right)\| dt, \\ \text{and} \quad \varepsilon_{n+1}^{\text{D}} := \int_{I_n} \|(f(U) - \mathcal{P}^{n+1}(\Phi^{n+\frac{1}{2}} U))(t)\| dt,$$

with  $\widehat{C}_2$  the constant in (4.10),  $\mathcal{L}_{31}^{n+1}$  and  $\mathcal{L}_{32}^{n+1}$  as in (5.26) and (5.28), respectively, and where  $\|\hat{\rho}(0)\|$  is estimated as  $\|\hat{\rho}(0)\| \leq \|u_0 - U^0\| + C_2 \eta_{2, \mathbb{V}^0}(U^0)$ . In particular, for  $1 \leq m \leq N$ , the following global estimate is valid for  $\hat{\rho}$ :

$$(5.30) \quad \max_{0 \leq t \leq t_m} \|\hat{\rho}(t)\| \leq \exp \left( \sum_{n=0}^{m-1} \int_{I_n} \mathcal{M}(U, u_0; t) dt \right) \times \left( \|u_0 - U^0\| + C_2 \eta_{2, \mathbb{V}^0}(U^0) \right. \\ \left. + \mathcal{E}_m^{\text{T},1} + \mathcal{E}_m^{\text{T},2} + C_2(\mathcal{E}_m^{\text{S},1} + \mathcal{E}_m^{\text{S},2}) + \widehat{C}_2 \mathcal{E}_m^{\text{S},3} + \mathcal{E}_m^{\text{C}} + \mathcal{E}_m^{\text{D}} \right),$$

with  $\mathcal{E}_m^{\text{T},j} := \sum_{n=1}^m \varepsilon_n^{\text{T},j}$ ,  $j = 1, 2$ ,  $\mathcal{E}_m^{\text{S},j} := \sum_{n=1}^m \varepsilon_n^{\text{S},j}$ ,  $j = 1, 2, 3$ ,  $\mathcal{E}_m^{\text{C}} := \sum_{n=1}^m \varepsilon_n^{\text{C}}$ , and  $\mathcal{E}_m^{\text{D}} := \sum_{n=1}^m \varepsilon_n^{\text{D}}$ .

*Proof.* Equation (5.10) is of the same form as (2.20). Therefore, using (5.20) instead of (3.10) and proceeding similarly to the proof of Theorem 3.1 leads to

$$(5.31) \quad \sup_{t \in I_n} \|\hat{\rho}(t)\| \leq \exp \left( \int_{I_n} \mathcal{H}(\hat{U}, u_0; t) dt \right) \times \left( \|\hat{\rho}(t_n)\| + \sum_{j=1}^3 \int_{I_n} \|R_j(t)\| dt \right), \quad 0 \leq n \leq N-1,$$

where  $\mathcal{H}(\hat{U}, u_0; t)$  is given, in this case, by

$$\mathcal{H}(\hat{U}, u_0; t) := A \left( \gamma(pd) \left( \|u_0\|^{p(2-d)} + \|\hat{U}(t)\|^{p(2-d)} \right) \times \right. \\ \left. \left( G(u_0)^{pd} + \left[ \max\{D_{2,0}, D_{2,1}\} \|\nabla Z(t)\| + \widehat{h}_{n+1} M(t) \right]^{pd} \right) + \|\hat{U}(t)\|_{L^\infty}^{2p} \right);$$

compare with (3.15).

Moreover, from (5.1) and (5.5) we get

$$\|\hat{U}(t)\| \leq \|\sigma(t)\| + \|\epsilon(t)\| + \|U(t)\| \leq \varepsilon_{n+1}^{\text{T},0} + C_2 \varepsilon_{n+1}^{\text{S},0} + \|U(t)\|,$$

while similarly, from (5.2) and (5.6) we deduce

$$\|\hat{U}(t)\| \leq \varepsilon_{n+1}^{\text{T},\infty} + C_\infty \varepsilon_{n+1}^{\text{S},\infty} + \|U(t)\|.$$

Hence (5.31) becomes

$$(5.32) \quad \sup_{t \in I_n} \|\hat{\rho}(t)\| \leq \exp \left( \int_{I_n} \mathcal{M}(U, u_0; t) dt \right) \times \left( \|\hat{\rho}(t_n)\| + \sum_{j=1}^3 \int_{I_n} \|R_j(t)\| dt \right), \quad 0 \leq n \leq N-1.$$

On the other hand, due to (5.14), (4.8) and (4.10), it is easily seen that

$$(5.33) \quad \int_{I_n} \|R_1(t)\| dt \leq C_2 \varepsilon_{n+1}^{\text{S},1} + \widehat{C}_2 \varepsilon_{n+1}^{\text{S},3} + \varepsilon_{n+1}^{\text{C}}.$$

Additionally, from (5.12), (5.13), (5.24), (5.25), and (5.27), we deduce

$$(5.34) \quad \int_{I_n} \|R_2(t)\| dt \leq \varepsilon_{n+1}^{\text{T},1} \quad \text{and} \quad \int_{I_n} \|R_3(t)\| dt \leq \varepsilon_{n+1}^{\text{T},2} + C_2 \varepsilon_{n+1}^{\text{S},2} + \varepsilon_{n+1}^{\text{D}}.$$

Plugging (5.33) and (5.34) into (5.32) we readily obtain (5.29). Estimate (5.30) follows from (5.29) by summing over  $n$ .  $\square$

Similarly to Theorem 3.2, we can also derive an improved estimate for  $\|\hat{\rho}(t)\|$  in the one-dimensional case.

**Theorem 5.2** (improved estimate for  $\hat{\rho}$  in  $d = 1$ ). *With the notation of Theorem 5.1 we further denote by  $\eta(t)$  the right-hand side in (5.30) to the power  $p$ . Let*

$$(5.35) \quad \begin{aligned} \mathcal{N}(U, u_0; t) := & A \left( \eta(t) \left( G(u_0)^p + \left[ \max\{D_{2,0}, D_{2,1}\} \left( \|\nabla Z(t)\| + \|\hat{h}_{n+1} M(t)\| \right) \right]^p \right) \right. \\ & \left. + \left( \varepsilon_{n+1}^{\text{T},\infty} + C_\infty \varepsilon_{n+1}^{\text{S},\infty} + \|U(t)\|_{L^\infty} \right)^{2p} \right). \end{aligned}$$

Then, if  $d = 1$ , the following local and global estimates are valid:

$$(5.36) \quad \begin{aligned} \sup_{t \in I_n} \|\hat{\rho}(t)\| \leq & \exp \left( \int_{I_n} \mathcal{N}(U, u_0; t) dt \right) \times \left( \|\hat{\rho}(t_n)\| + \varepsilon_{n+1}^{\text{T},1} + \varepsilon_{n+1}^{\text{T},2} \right. \\ & \left. + C_2 (\varepsilon_{n+1}^{\text{S},1} + \varepsilon_{n+1}^{\text{S},2}) + \widehat{C}_2 \varepsilon_{n+1}^{\text{S},3} + \varepsilon_{n+1}^{\text{C}} + \varepsilon_{n+1}^{\text{D}} \right), \quad 0 \leq n \leq N-1, \end{aligned}$$

$$(5.37) \quad \begin{aligned} \max_{0 \leq t \leq t_m} \|\hat{\rho}(t)\| \leq & \exp \left( \sum_{n=0}^{m-1} \int_{I_n} \mathcal{N}(U, u_0; t) dt \right) \times \left( \|u_0 - U^0\| + C_2 \eta_{2,\nabla^0}(U^0) \right. \\ & \left. + \mathcal{E}_m^{\text{T},1} + \mathcal{E}_m^{\text{T},2} + C_2 (\mathcal{E}_m^{\text{S},1} + \mathcal{E}_m^{\text{S},2}) + \widehat{C}_2 \mathcal{E}_m^{\text{S},3} + \mathcal{E}_m^{\text{C}} + \mathcal{E}_m^{\text{D}} \right), \quad 1 \leq m \leq N. \quad \square \end{aligned}$$

### 5.3. A posteriori error bounds in the $L^\infty([0, T]; L^2(\Omega))$ -norm for fully discrete schemes.

We conclude the section by presenting the theorems with the a posteriori error estimates in the  $L^\infty([0, T]; L^2(\Omega))$ -norm for the error  $e = u - U$ .

**Theorem 5.3** (a posteriori error estimate in the  $L^\infty(0, T; L^2(\Omega))$ ). *With the notation of Theorem 5.1 we further define*

$$\begin{aligned} \varepsilon_{n+1}^{\text{LOC}} := & \exp \left( \int_{I_n} \mathcal{M}(U, u_0; t) dt \right) \times \left( \varepsilon_n^{\text{LOC}} + \varepsilon_{n+1}^{\text{T},1} + \varepsilon_{n+1}^{\text{T},2} + C_2 (\varepsilon_{n+1}^{\text{S},1} + \varepsilon_{n+1}^{\text{S},2}) + \widehat{C}_2 \varepsilon_{n+1}^{\text{S},3} \right. \\ & \left. + \varepsilon_{n+1}^{\text{C}} + \varepsilon_{n+1}^{\text{D}} \right), \quad 0 \leq n \leq N-1, \end{aligned}$$

with  $\varepsilon_0^{\text{LOC}} := \|u_0 - U^0\| + C_2 \eta_{2,\nabla^0}(U^0)$ . Let also  $\mathcal{E}_m^{\text{GLOB}}$ ,  $1 \leq m \leq N$ , be the right-hand side in (5.30). Then the following estimates are valid:

$$(5.38) \quad \sup_{t \in I_n} \|(u - U)(t)\| \leq C_2 \varepsilon_{n+1}^{\text{S},0} + \varepsilon_{n+1}^{\text{T},0} + \varepsilon_{n+1}^{\text{LOC}}, \quad 0 \leq n \leq N-1,$$

$$(5.39) \quad \max_{0 \leq t \leq t_m} \|(u - U)(t)\| \leq C_2 \mathcal{E}_m^{\text{S},0} + \mathcal{E}_m^{\text{T},0} + \mathcal{E}_m^{\text{GLOB}}, \quad 1 \leq m \leq N,$$

where  $u$  is the exact solution of (1.1),  $U$  denotes the continuous, linear, piecewise interpolant  $U(t) = \ell_0^n(t)U^n + \ell_1^n(t)U^{n+1}$ ,  $t \in I_n$ , and  $\{U^n\}_{n=0}^N$  are the modified relaxation Crank-Nicolson-Galerkin approximations, obtained by (4.2).

*Proof.* We just note that

$$\|(u - U)(t)\| = \|e(t)\| \leq \|\epsilon(t)\| + \|\sigma(t)\| + \|\hat{\rho}(t)\|$$

and we apply (5.7), (5.1), and (5.29) to obtain (5.38), while for (5.39) we use (5.7), (5.3) and (5.30).  $\square$

The last theorem of the section is the improved a posteriori error estimate for  $u - U$  in the  $L^\infty([0, T]; L^2(\Omega))$ -norm for the one-dimensional case.

**Theorem 5.4** (improved a posteriori error estimate in the  $L^\infty(0, T; L^2(\Omega))$  for  $d = 1$ ). *With the notation of Theorems 3.2 and 5.3 we further define  $\tilde{\varepsilon}_n^{\text{LOC}}$ ,  $0 \leq n \leq N$ , and  $\tilde{\mathcal{E}}_m^{\text{GLOB}}$ ,  $1 \leq m \leq N$ , as in Theorem 5.3, but with  $\mathcal{M}$  replaced by  $\mathcal{N}$ . Then, for  $d = 1$ , the following local and global estimates hold:*

$$(5.40) \quad \begin{aligned} \sup_{t \in I_n} \|(u - U)(t)\| &\leq C_2 \varepsilon_{n+1}^{\text{S},0} + \varepsilon_{n+1}^{\text{T},0} + \tilde{\varepsilon}_{n+1}^{\text{LOC}}, \quad 0 \leq n \leq N - 1, \\ \max_{0 \leq t \leq t_m} \|(u - U)(t)\| &\leq C_2 \mathcal{E}_m^{\text{S},0} + \mathcal{E}_m^{\text{T},0} + \tilde{\mathcal{E}}_m^{\text{GLOB}}, \quad 1 \leq m \leq N. \quad \square \end{aligned}$$

## 6. NUMERICAL EXPERIMENTS

In this section we report a series of numerical experiments which verify the theoretical results obtained in the previous sections. The modified relaxation Crank-Nicolson-Galerkin scheme (4.2) and the corresponding a posteriori space and time estimators with homogeneous Dirichlet boundary conditions were implemented in a double precision C-code using B-splines of degree  $r$ ,  $r \in \mathbb{N}$ , as a basis for the finite element space  $\mathbb{V}^n$ ,  $0 \leq n \leq N$ . The numerical results are for the one-dimensional case  $d = 1$ ; in this case  $\Omega = [a, b]$ . For the implementations we use uniform partitions in space and time and we set equal to one the absolute constants  $A, \beta, C_2, \hat{C}_2, C_\infty, D_{2,0}, D_{2,1}$  appearing in the final a posteriori error estimators.

Our goals in this section are : (a) to asses the magnitude of the constants  $\mathcal{L}_{31}^n$ ,  $\mathcal{L}_{32}^n$ ,  $\mathcal{M}(U, u_0; t)$  and  $\mathcal{N}(U, u_0; t)$  involved in the estimators, (b) to verify the correct order of convergence for the space and time estimators, and (c) to verify the accuracy of the Crank-Nicolson relaxation scheme and of its a posteriori time estimator for variable time steps.

To facilitate the process we choose to work with a soliton type exact solution of (1.1). In particular we choose, [9]

$$(6.1) \quad u(x, t) = i \operatorname{sech}(x - x_0 - 4\omega t) e^{i(2\omega x + (1 - 4\omega^2)t)},$$

which is an exact solution of (1.1) with  $p = 1$ ,  $\alpha = 1$ ,  $\lambda = 2$  and  $u_0 = u(\cdot, 0)$ . The function  $u$  represents a solitary wave, initially located around  $x = x_0$ , which travels to the right with speed  $4\omega$ . All computations in Subsections 6.1 and 6.2 are performed in  $[a, b] \times [0, T] = [-30, 30] \times [0, 1]$  with  $x_0 = 0$  and  $\omega = 0.3$ .

**Remark 6.1.** The development of a time-space adaptive algorithm based on the a posteriori error estimators of the paper and its numerical validation in terms of accuracy, effectiveness and robustness for the approximation of the solution  $u$  of (1.1) will be the subject of a forthcoming paper. The discussion of Subsection 6.3 is a first indication that time adaptivity is possible. In this forthcoming work we will also study further cases including for example nonlinearities to the critical exponent  $p^* = \frac{2}{d}$ , as well as particular examples for (1.1) in the semiclassical regime, cf. [53].

In the next subsections we consider a series of different runs for the NLS problem (1.1) with exact solution (6.1). Let  $\ell \in \mathbb{N}$  counts the different realizations (runs),  $h(\ell)$  the corresponding meshsize and  $M(\ell) = 1 + \lfloor \frac{b-a}{h(\ell)} \rfloor \in \mathbb{N}$  where  $\lfloor \cdot \rfloor$  denotes the integral part of a real number. When there is no danger of confusion we drop the dependence on  $\ell$  and we write just  $h$  and  $M$ . Since we discretize in space by B-splines of degree  $r$ , the expected order of convergence for the modified relaxation Crank-Nicolson finite element scheme is  $r + 1$  in space, whilst in time is 2. This motivates the relation between the mesh size  $h$  and the time step  $k$ . More specifically in all the computations of the next two subsections we choose

$$(6.2) \quad k \sim h^{\frac{r+1}{2}}.$$

**6.1. Behavior of  $\mathcal{L}_{31}^n$ ,  $\mathcal{L}_{32}^n$  and  $\mathcal{M}, \mathcal{N}$ .** First, we quantify the quantities  $\mathcal{L}_{31}^n$ ,  $\mathcal{L}_{32}^n$  defined in (5.26), (5.28), respectively. For  $0 \leq n \leq N-1$ ,  $\mathcal{L}_{31}^n$  appears in the local time estimator  $\varepsilon_{n+1}^{T,2}$ , cf. (5.29), and thus in the global time estimator  $\mathcal{E}_N^{T,2}$ , cf. (5.30). Similarly, for  $0 \leq n \leq N-1$ ,  $\mathcal{L}_{32}^n$  is in the local space estimator  $\varepsilon_{n+1}^{S,2}$ , cf. (5.29), and thus in the global space estimator  $\mathcal{E}_N^{S,2}$ , cf. (5.30). Estimators  $\mathcal{E}_N^{T,2}$  and  $\mathcal{E}_N^{S,2}$  are expected to be of optimal order of accuracy as long as the following global quantities

$$\mathcal{L}_{31} := \max_{0 \leq n \leq N-1} \mathcal{L}_{31}^n \quad \text{and} \quad \mathcal{L}_{32} := \max_{0 \leq n \leq N-1} \mathcal{L}_{32}^n,$$

tend to a constant value for different realizations. In Table 1 we compute the values of  $\mathcal{L}_{31}, \mathcal{L}_{32}$  for linear ( $r = 1$ ), quadratic ( $r = 2$ ) and cubic ( $r = 3$ ) B-splines and a series of different space discretizations  $h$ ; the time step is chosen according to (6.2).

We are also interested in studying the magnitude of the quantity  $\mathcal{E}_N^{\mathcal{M}} := \sum_{n=0}^{N-1} \int_{I_n} \mathcal{M}(U, u_0; t) dt$ , which appears in the exponential of (5.30) and thus on the final global a posteriori error estimator (5.39). Our findings are reported again in Table 1. Since the considered example is in the one spatial dimension, the improved a posteriori error estimate (5.40) holds (see also (5.37)). The difference of this estimate compared to (5.39) is the existence of  $\mathcal{E}_N^{\mathcal{N}} := \sum_{n=0}^{N-1} \int_{I_n} \mathcal{N}(U, u_0; t) dt$  instead of  $\mathcal{E}_N^{\mathcal{M}} = \sum_{n=0}^{N-1} \int_{I_n} \mathcal{M}(U, u_0; t) dt$  in the exponential term. In Table 1 we confirm that indeed  $\mathcal{E}_N^{\mathcal{N}}$  is smaller than  $\mathcal{E}_N^{\mathcal{M}}$ . More precisely, in view of (5.35), we expect that, as the temporal and spatial mesh sizes tend to zero,  $\mathcal{E}_N^{\mathcal{N}}$  will tend to  $\int_0^T \|u(t)\|_{L^\infty}^{2p} dt$ , which is equal to 1 for this particular example. This is verified in Table 1 for quadratic and cubic B-splines. For linear B-splines, we need to consider smaller mesh sizes; however, it is clear that  $\mathcal{E}_N^{\mathcal{N}}$  reduces as  $h$  reduces in this case as well. Although at first glance the improvement on  $\mathcal{E}_N^{\mathcal{N}}$  compared to  $\mathcal{E}_N^{\mathcal{M}}$  seems minor, we should keep in mind that these quantities appear in the exponential, therefore even minor improvements have a significant impact on the size of the total a posteriori error estimator.

	$r = 1$				$r = 2$				$r = 3$			
$M$	$\mathcal{L}_{31}$	$\mathcal{L}_{32}$	$\mathcal{E}_N^{\mathcal{M}}$	$\mathcal{E}_N^{\mathcal{N}}$	$\mathcal{L}_{31}$	$\mathcal{L}_{32}$	$\mathcal{E}_N^{\mathcal{M}}$	$\mathcal{E}_N^{\mathcal{N}}$	$\mathcal{L}_{31}$	$\mathcal{L}_{32}$	$\mathcal{E}_N^{\mathcal{M}}$	$\mathcal{E}_N^{\mathcal{N}}$
2400	6.639	3.070	7.847	17.423	6.359	3.001	7.820	2.233	6.321	3.000	7.820	1.164
3600	6.520	3.038	7.831	9.757	6.338	3.000	7.817	1.613	6.317	3.000	7.817	1.071
4800	6.464	3.025	7.825	6.737	6.329	3.000	7.816	1.382	6.315	3.000	7.816	1.040
6000	6.432	3.017	7.821	5.187	6.325	3.000	7.815	1.267	6.315	3.000	7.815	1.025
7200	6.411	3.013	7.820	4.263	6.322	3.000	7.814	1.201	6.314	3.000	7.815	1.018
8400	6.397	3.010	7.818	3.658	6.320	3.000	7.813	1.158	6.314	3.000	7.815	1.013
9600	6.386	3.008	7.818	3.234	6.319	3.000	7.813	1.128	6.314	3.000	7.815	1.010

TABLE 1. The computed quantities  $\mathcal{L}_{31}$ ,  $\mathcal{L}_{32}$ ,  $\mathcal{E}_N^{\mathcal{M}}$  and  $\mathcal{E}_N^{\mathcal{N}}$ .

**6.2. EOC of the estimators.** Our purpose in this subsection is to compute the *experimental order of convergence (EOC)* of the space and time estimators at the final time  $T = 1$  for the NLS problem (1.1) with exact solution (6.1). For this, we choose quadratic B-splines ( $r = 2$ ). Hence the expected order of convergent of the modified relaxation Crank-Nicolson finite element scheme is in this case 3 in space and 2 in time; thus we take  $k \sim h^{3/2}$ , cf. (6.2).

The EOC for each space estimator  $\mathcal{E}_N^{S,j}$ ,  $0 \leq j \leq 3$ , is computed as follows

$$(6.3) \quad \text{EOC}_{S,j} := \frac{\log(\mathcal{E}_N^{S,j}(\ell)/\mathcal{E}_N^{S,j}(\ell+1))}{\log(M(\ell+1)/M(\ell))},$$

where  $\mathcal{E}_N^{S,j}(\ell)$  and  $\mathcal{E}_N^{S,j}(\ell+1)$  denote the value of the estimators in two consecutive implementations with mesh sizes  $h(\ell)$  and  $h(\ell+1)$  respectively. Similarly for the time estimators  $\mathcal{E}_N^{T,j}$ ,  $0 \leq j \leq 2$ , the EOC is computed as

$$(6.4) \quad \text{EOC}_{T,j} := \frac{\log(\mathcal{E}_N^{T,j}(\ell)/\mathcal{E}_N^{T,j}(\ell+1))}{\log(k(\ell+1)/k(\ell))},$$

where  $k(\ell)$ ,  $k(\ell+1)$  are the time steps of consecutive realizations.

In Table 2 the values of the space estimators  $\mathcal{E}_N^{S,j}$  along with the corresponding  $\text{EOC}_{S,j}$ ,  $0 \leq j \leq 3$ , are presented. Note that the estimator  $\mathcal{E}_N^{S,1}$  is expected to be of optimal third order in space and of first order in time (cf. Theorem 5.1), i.e., it is a superconvergent term. Due to the choice  $k \sim h^{3/2}$  we have  $k \times h^3 = h^{3/2} \times h^3 = h^{9/2}$ , and the EOC we expect to observe is 4.5. This EOC is indeed observed in Table 2 for  $\mathcal{E}_N^{S,1}$ , while, as it is shown in the same table, the other three space estimators exhibit the correct order. Similarly, in Table 3 the values of the time estimators  $\mathcal{E}_N^{T,j}$  along with the  $\text{EOC}_{T,j}$ ,  $0 \leq j \leq 2$ , are presented. We note that all time estimators exhibit the correct second order. We are also interested in computing the *effectivity index* which is the ratio between an a

$M$	$\mathcal{E}_N^{S,0}$	$\text{EOC}_{S,0}$	$\mathcal{E}_N^{S,1}$	$\text{EOC}_{S,1}$	$\mathcal{E}_N^{S,2}$	$\text{EOC}_{S,2}$	$\mathcal{E}_N^{S,3}$	$\text{EOC}_{S,3}$
2400	1.3817E-05	–	2.64149E-07	–	4.1466E-05	–	2.5055E-05	–
3600	4.0936E-06	3.0001	4.25720E-08	4.5018	1.2283E-05	3.0007	7.4230E-06	3.0003
4800	1.7270E-06	3.0000	1.16949E-08	4.4912	5.1813E-06	3.0003	3.1315E-06	3.0001
6000	8.8421E-07	3.0000	4.28570E-09	4.4988	2.6527E-06	3.0002	1.6033E-06	3.0001
7200	5.1169E-07	3.0000	1.88822E-09	4.4956	1.5351E-06	3.0001	9.2783E-07	3.0000
8400	3.2223E-07	3.0000	9.44191E-10	4.4960	9.6671E-07	3.0001	5.8429E-07	3.0000
9600	2.1587E-07	3.0000	5.20390E-10	4.4615	6.4762E-07	3.0000	3.9142E-07	3.0000

TABLE 2. Space a posteriori estimators  $\mathcal{E}_N^{S,j}$  and corresponding  $\text{EOC}_{S,j}$ ,  $0 \leq j \leq 3$ .

$k^{-1}$	$\mathcal{E}_N^{T,0}$	$\text{EOC}_{T,0}$	$\mathcal{E}_N^{T,1}$	$\text{EOC}_{T,1}$	$\mathcal{E}_N^{T,2}$	$\text{EOC}_{T,2}$
252	8.2786E-06	–	3.1156E-04	–	1.6442E-05	–
464	2.4426E-06	2.0055	9.2190E-05	2.0007	4.8667E-06	2.0002
715	1.0288E-06	2.0032	3.8879E-05	2.0003	2.0526E-06	2.0001
1000	5.2600E-07	2.0021	1.9891E-05	2.0000	1.0501E-06	2.0000
1314	3.0466E-07	2.0016	1.1528E-05	1.9993	6.0850E-07	2.0000
1656	1.9182E-07	2.0012	7.2678E-06	1.9957	3.8324E-07	2.0000
2023	1.2854E-07	2.0010	4.8767E-06	1.9943	2.5686E-07	2.0000

TABLE 3. Time a posteriori estimators  $\mathcal{E}_N^{T,j}$  and corresponding  $\text{EOC}_{T,j}$ ,  $0 \leq j \leq 2$ .

posteriori error estimator and the exact error. The effectivity index is a tool providing information on the quality of the estimator. To that end, we denote by  $E_{\text{exact}} := \max_{0 \leq n \leq N} \|u(t_n) - U^n\|$  the exact error, by  $\mathcal{E}_N$  the sum of all time and space estimators and we compute the effectivity index  $ei$  as  $ei := \mathcal{E}_N/E_{\text{exact}}$ . The exact error  $E_{\text{exact}}$ , the estimator  $\mathcal{E}_N$  and the effectivity index  $ei$  are presented in Table 4. We observe that the computed effectivity index  $ei$  stabilizes to a fixed value, indicating in this case that the sum of all time and space estimators  $\mathcal{E}_N$  is about 26 times larger than the actual error. Note that the exponential term  $e^{\mathcal{E}_N^M}$ , or  $e^{\mathcal{E}_N^N}$  for the case of the improved estimate (5.40), is not included in the definition of  $\mathcal{E}_N$ . In order to get an indication of how bigger is the actual total a posteriori error estimator with respect to the actual error we should multiply the third column with the effectivity index in Table 4 by  $e^{\mathcal{E}_N^M}$ , or  $e^{\mathcal{E}_N^N}$  for the case of the improved estimate (5.40); cf.

Table 1 for the corresponding values of  $\mathcal{E}_N^{\mathcal{M}}$  and  $\mathcal{E}_N^{\mathcal{N}}$ . At this point, the importance of the improved estimate (5.40) for the one-dimensional case becomes apparent. Finally we would like to note that

$\mathcal{E}_N$	$E_{\text{exact}}$	$ei$
4.1688E-04	1.6587E-05	25.13
1.2334E-04	4.9118E-06	25.11
5.2012E-05	2.0186E-06	25.77
2.6612E-05	1.0202E-06	26.09
1.5418E-05	5.8838E-07	26.20
9.7171E-06	3.7056E-07	26.22
6.5176E-06	2.4818E-07	26.26

TABLE 4. A posteriori estimator  $\mathcal{E}_N$ , exact error  $E_{\text{exact}}$  and effectivity index  $ei$ .

the data estimator  $\mathcal{E}_N^{\text{D}}$  is, for this example, much smaller than the time and space estimators, whilst the coarsening estimator  $\mathcal{E}_N^{\text{C}}$  is zero in the case of uniform partitions. This is the reason we don't include these estimators in  $\mathcal{E}_N$ .

**6.3. Accuracy and variable time steps.** We investigate briefly the accuracy of the time discretization in the case of variable time steps. The expected second order accuracy is illustrated through a numerical experiment. We consider again (6.1) as the exact solution of (1.1) and we compute its approximation by the numerical method. The parameters  $x_0$ ,  $\omega$  and  $[a, b]$  are as in the previous experiments, while the final time is  $T = 5$  for this experiment. To isolate the temporal error, we take a sufficiently fine spatial mesh consisting of  $M = 3000$  points and quintic B-splines, thus the spatial discretization error is almost negligible. We split the time interval  $[0, T] = \cup_{j=0}^4 [T_j, T_{j+1}]$  with  $T_j = j$ ,  $0 \leq j \leq 4$ . In each subinterval  $[T_j, T_{j+1}]$ , we use a different time step  $k_j = 0.05/2^{5-j}$ ,  $1 \leq j \leq 5$  and we compute the exact error  $E_{\text{exact}}^j := \|u(T_j) - U^{T_j}\|$  at the endpoint  $T_j$ ,  $1 \leq j \leq 5$ , where  $U^{T_j}$  denotes the corresponding approximation at  $T_j$ . We also compute the total a posteriori time estimator  $\mathcal{E}_{T_j}^{\text{T}} := \mathcal{E}_{T_j}^{\text{T},0} + \mathcal{E}_{T_j}^{\text{T},1} + \mathcal{E}_{T_j}^{\text{T},2}$  in the interval  $[0, T_j]$ ,  $1 \leq j \leq 5$ . Our findings are reported in Table 6.3.

$j$	$k_j$	$E_{\text{exact}}^j$	$\text{EOC}_E$	$\mathcal{E}_{T_j}^{\text{T}}$	$\text{EOC}_{\text{T}}$
1	3.125E-03	1.1601E-05		2.0110E-05	
2	6.250E-03	4.6623E-05	2.0068	9.8040E-05	2.2854
3	1.250E-02	1.8770E-04	2.0093	4.1007E-04	2.0644
4	2.500E-02	7.5547E-04	2.0090	1.6622E-03	2.0191
5	5.000E-02	3.0345E-03	2.0060	6.7331E-03	2.0182

TABLE 5. Exact error  $E_{\text{exact}}^n$  and total time estimator  $\mathcal{E}_n^{\text{T}}$  with their corresponding EOC.

#### ACKNOWLEDGMENTS

I.K. is grateful to Prof. Charalambos Makridakis for suggesting the problem and for fruitful discussions. The authors would like to thank Prof. Georgios Akrivis and the anonymous reviewers for their valuable comments and suggestions.



## REFERENCES

- [1] G. Akrivis, V. Dougalis, O. Karakashian, *On fully discrete Galerkin methods of second-order temporal accuracy for the nonlinear Schrödinger equation*, Numer. Math. **59** (1991) 31–53.
- [2] G.D. Akrivis, V.A. Dougalis, O.A. Karakashian, W.R. McKinney, *Numerical approximation of blow-up of radially symmetric solutions of the nonlinear Schrödinger equation*, SIAM J. Sci. Comput. **25** (2003) 186–212.
- [3] G. Akrivis, Ch. Makridakis, R.H. Nochetto, *A posteriori error estimates for the Crank-Nicolson method for parabolic equations*, Math. Comp. **75** (2005) 511–531.
- [4] R. Anton, *Strichartz inequalities for Lipschitz metrics on manifolds and nonlinear Schrödinger equation on domains*, Bull. Soc. Math. France **136** (2008) 27–65.
- [5] P. Antonelli, A. Athanassoulis, Z. Huang, P.A. Markowich, *Numerical Simulations of X-Ray Free Electron Lasers (XFEL)*, Multiscale Model. Sim. **12** (2014) 1607–1621.
- [6] W. Bao, Y. Cai, *Uniform and optimal error estimates of an exponential wave integrator sine pseudospectral method for the nonlinear Schrödinger equation with wave operator*, SIAM J. Numer. Anal. **52** (2014) 1103–1127.
- [7] W. Bao, S. Jin, P.A. Markowich, *Numerical study of time-splitting spectral discretizations of nonlinear Schrödinger equations in the semiclassical regimes*, SIAM J. Sci. Comput. **25** (2003) 27–64.
- [8] M. Bertola, A. Tovbis, *Universality for the focusing nonlinear Schrödinger equation at the gradient catastrophe point: rational breathers and poles of the Triconquée solution to Painlevé-I*, Comm. Pure Appl. Math. **66** (2013) 678–752.
- [9] Ch. Besse, *A relaxation scheme for the nonlinear Schrödinger equation*, SIAM J. Numer. Anal. **42** (2004) 934–952.
- [10] J. Bourgain, *Global solutions of nonlinear Schrödinger equations*, American Mathematical Society, Providence, RI, 1999.
- [11] S.C. Brenner, L.R. Scott, *The Mathematical Theory of Finite Element Methods*, 2<sup>nd</sup> edition, Springer, New York, 2002.
- [12] H. Brezis, *Functional analysis, Sobolev spaces and partial differential equations*, University text, Springer, New York, 2011.
- [13] H. Brezis, T. Gallouet, *Nonlinear Schrödinger evolution equations*, Nonlinear Anal. **4** (1980), 677–681.
- [14] E. Bänsch, F. Karakatsani, Ch. Makridakis, *A posteriori error control for fully discrete Crank-Nicolson schemes*, SIAM J. Numer. Anal. **50** (2012) 2845–2872.
- [15] A. Cangiani, E.H. Georgoulis, I. Kyza, S. Metcalfe, *Adaptivity and blow-up detection for nonlinear evolution problems*, SIAM J. Sci. Comput. **38**, pp. A3833–A3856.
- [16] R. Carles, *WKB analysis for nonlinear Schrödinger equations with potential*, Comm. Math. Phys. **269** (2007) 195–221.
- [17] Th. Cazenave, *Semilinear Schrödinger equations*, American Mathematical Society, Providence, RI, 2003.
- [18] Ch. Clément, *Approximation by finite element functions using local regularization*, Analyse Numérique **9** (1975) 77–84.
- [19] M. Delfour, M. Fortin, G. Payre, *Finite-difference solutions of a non-linear Schrödinger equation*, J. Comput. Phys. **44** (1981) 277–288.
- [20] A. Demlow, O. Lakkis, Ch. Makridakis, *A posteriori error estimates in the maximum norm for parabolic problems*, SIAM J. Numer. Anal. **47** (2009) 2157–2176.
- [21] K.B. Dysthe, *Note on a modification to the nonlinear Schrödinger equation for application to deep water waves*, Proc. R. Soc. London Ser. A **369** (1979) 105–114.
- [22] W. Dörfler, *A time-and space-adaptive algorithm for the linear time-dependent Schrödinger equation*, Numer. Math. **73** (1996) 419–448.
- [23] A. Hasegawa, Y. Kodama, *Solitons in optical communications*, Oxford University Press, USA, 1995.
- [24] S. Jimenez, I.M. Llorente, A.M. Mancho, V.M. Perez-Garcia, L. Vazquez, *A numerical scheme for the simulation of blow-up in the nonlinear Schrödinger equation*, Appl. Math. Comput. **134** (2003), 271–291.
- [25] A. Jüngel, M.C. Mariani, D. Rial, *Local existence of solutions to the transient quantum hydrodynamic equations*, Math. Models Methods Appl. Sci. **12** (2002) 485–495.
- [26] A. Jüngel, R.-M. Weishäupl, *Blow-up in two component nonlinear Schrödinger systems with an external driven field*, Math. Models Methods Appl. Sci. **23** (2013) 1699–1727.
- [27] O. Karakashian, Ch. Makridakis, *A space-time finite element method for the nonlinear Schrödinger equation: the discontinuous Galerkin method*, Math. Comp. **67** (1998) 479–499.
- [28] O. Karakashian, Ch. Makridakis, *A space-time finite element method for the nonlinear Schrödinger equation: the continuous Galerkin method*, SIAM J. Numer. Anal. **36** (1999) 1779–1807.
- [29] Th. Katsaounis, I. Kyza, *A posteriori error control and adaptivity for Crank-Nicolson finite element approximations for the linear Schrödinger equation*, Numer. Math. **129** (2015) 55–90.
- [30] I. Kyza, *A posteriori error estimates for approximations of semilinear parabolic and Schrödinger-type equations*, PhD Thesis, University of Crete, 2009.
- [31] I. Kyza, *A posteriori error analysis for the Crank-Nicolson method for linear Schrödinger equations*, ESAIM Math. Model. Numer. Anal. **45** (2011) 761–778.
- [32] I. Kyza, Ch. Makridakis, *Analysis for time discrete approximations of blow-up solutions of semilinear parabolic equations*, SIAM J. Numer. Anal. **49** (2011) 405–426.

- [33] I. Kyza, Ch. Makridakis, M. Plexousakis, *Error control for time-splitting spectral approximations of the semi-classical Schrödinger equation*, IMA J. Numer. Anal. **31** (2011) 416–441.
- [34] I. Kyza, S. Metcalfe, T.P. Wihler, *hp-Adaptive Galerkin time stepping methods for nonlinear initial value problems*, J. Sci. Comput. **75** (2018), 111–127.
- [35] O. Lakkis, Ch. Makridakis, *Elliptic reconstruction and a posteriori error estimates for fully discrete linear parabolic problems*, Math. Comp. **75** (2006) 1627–1658.
- [36] A. Lozinski, M. Picasso, V. Prachittham, *An anisotropic error estimator for the Crank-Nicolson method: Application to a parabolic problem*, SIAM J. Sci. Comput. **31** (2009) 2757–2783.
- [37] G. Lyng, P.D. Miller, *The  $N$ -soliton of the focusing nonlinear Schrödinger equation for  $N$  large*, Comm. Pure Appl. Math. **60** (2007) 951–1026.
- [38] Ch. Makridakis, R.H. Nochetto, *Elliptic reconstruction and a posteriori error estimates for parabolic problems*, SIAM J. Numer. Anal. **41** (2003) 1585–1594.
- [39] Ch. Makridakis, R.H. Nochetto, *A posteriori error analysis for higher order dissipative methods for evolution problems*, Numer. Math. **104** (2006) 489–514.
- [40] F. Merle, P. Raphael, *On a sharp lower bound on the blow up rate for critical nonlinear Schrödinger equation*, J. Amer. Math. Soc. **19** (2006) 37–90.
- [41] F. Merle, P. Raphael, *The blow-up dynamic and upper bound on the blow-up rate for critical nonlinear Schrödinger equation*, Ann. of Math. **161** (2005) 157–222.
- [42] F. Merle, P. Raphael, *On universality of blow up profile for  $L^2$  critical nonlinear Schrödinger equation*, Invent. Math. **156** (2004) 565–672.
- [43] R.H. Nochetto, A. Schmidt, K.G. Siebert, A. Veiser, *Pointwise a posteriori error estimates for monotone semilinear problems*, Numer. Math. **104** (2006) 515–538.
- [44] T. Ozawa, Y. Tsutsumi, *The nonlinear Schrödinger limit and the initial layer of the Zakharov equations*, Differential Integral Equations, **5** (1992) 721–745.
- [45] M. Plexousakis, *An adaptive nonconforming finite element method for the nonlinear Schrödinger equation*, PhD Thesis, University of Tennessee, 1996.
- [46] W. Rudin, *Real & Complex Analysis*, 2<sup>nd</sup> edition, Tata McGraw-Hill, New York, 1974.
- [47] L.R. Scott, S. Zhang, *Finite element interpolation of non-smooth functions satisfying boundary conditions*, Math. Comp. **54** (1990) 483–493.
- [48] K.E. Strecker, G.B. Partridge, A.G. Truscott, R.G. Hulet, *Formation and propagation of matter-wave soliton trains*, Nature **417** (2002) 150–153.
- [49] C. Sulem, P.-L. Sulem, *The nonlinear Schrödinger equation: self-focusing and wave collapse*, Springer Science & Business Media, 1999.
- [50] T. Tao, *Nonlinear dispersive equations: local and global analysis*, American Mathematical Society, Providence, RI, 2006.
- [51] M. Thalhammer, *Convergence analysis of high-order time-splitting pseudo-spectral methods for nonlinear Schrödinger equations*, SIAM J. Numer. Anal. **50** (2012) 3231–3258.
- [52] M. Thalhammer, J. Abhau, *A numerical study of adaptive space and time discretizations for Gross-Pitaevskii equations*, J. Comput. Phys. **231** (2012) 6665–6681.
- [53] A. Tovbis, S. Venakides, X. Zhou, *On semiclassical (zero dispersion limit) solutions of the focusing nonlinear Schrödinger equation*, Comm. Pure Appl. Math., **57** (2004) 877–985.
- [54] H. Wang, *Numerical studies on the split-step finite difference method for nonlinear Schrödinger equations*, Appl. Math. Comp. **170** (2005) 17–35.
- [55] R. Zhang, K. Zhang, Y.S. Zhou, *Numerical study of time-splitting, space-time adaptive wavelet scheme for Schrödinger equations*, J. Comput. Appl. Math. **195** (2006) 263–273.
- [56] G. Zouraris, *On the convergence of a linear two-step finite element method for the nonlinear Schrödinger equation*, M2AN Math. Model. Numer. Anal. **35** (2001) 389–405.

(Theodoros Katsaounis) COMPUTER, ELECTRICAL AND MATHEMATICAL SCIENCES & ENGINEERING, KAUST, SAUDI ARABIA, & IACM-FORTH, HERAKLION, GREECE, & DEPT. OF MATHEMATICS AND APPLIED MATHEMATICS, UNIV. OF CRETE, GREECE

*E-mail address:* theodoros.katsaounis@kaust.edu.sa

(Irene Kyza) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DUNDEE, DUNDEE DD1 4HN, SCOTLAND, UK

*E-mail address:* ikyza@dundee.ac.uk