



University of Dundee

An Expanded Mixed Finite Element Method for Two-dimensional Sobolev Equations

Li, Na; Lin, Ping; Gao, Fuzheng

Published in:
Journal of Computational and Applied Mathematics

DOI:
[10.1016/j.cam.2018.08.041](https://doi.org/10.1016/j.cam.2018.08.041)

Publication date:
2019

Licence:
CC BY-NC-ND

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Li, N., Lin, P., & Gao, F. (2019). An Expanded Mixed Finite Element Method for Two-dimensional Sobolev Equations. *Journal of Computational and Applied Mathematics*, 348, 342-355.
<https://doi.org/10.1016/j.cam.2018.08.041>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

An Expanded Mixed Finite Element Method for Two-dimensional Sobolev Equations[★]

Na Li^{a,b}, Ping Lin^{c,*}, Fuzheng Gao^d

^a*School of Mathematics and Physics, University of Science and Technology Beijing, Beijing 100083, China*

^b*School of Data and Computer Science, Shandong Women's University, Jinan 250300, China*

^c*Department of Mathematics, University of Dundee, Dundee DD1 4HN, UK*

^d*School of Mathematics, Shandong University, Jinan 250100, China*

Abstract

In this paper we provide an expanded mixed finite element method for a class of two-dimensional Sobolev equations. The optimal error estimates for a semi-discrete scheme and a fully discrete scheme are obtained. Also numerical examples are stated to verify our theoretical results.

Keywords: Sobolev Equation; Expanded Mixed Finite Element; Error Estimate; Numerical Experiment

1 Introduction

Let $\Omega \subset R^2$ be a bounded domain with piecewise smooth boundary $\partial\Omega$. For fixed $0 < T < \infty$, we consider the following initial and boundary value problem:

$$\begin{cases} u_t - \mu\Delta u_t - \gamma\Delta u = f, & (x, y, t) \in \Omega \times (0, T], \\ u(x, y, t) = 0, & (x, y, t) \in \partial\Omega \times (0, T], \\ u(x, y, 0) = \varphi_0(x, y), & (x, y) \in \Omega, \end{cases} \quad (1.1)$$

where u_t denotes the time derivative of the function u , μ and γ are two positive constants, the source term $f(x, y, t)$ and the initial value function $\varphi_0(x, y)$ are sufficiently smooth.

The above equation is characterized by the occurrence of mixed time and space derivatives appearing in the highest-order term, which we call Sobolev equation. As a class of important Sobolev equations, Eq.(1.1) possesses the important physical background. In 1960, Eq.(1.1) appeared in the theory developed by Barenblatt, Zheltov and Kochia for flow through fissured

*This work was supported by the National Natural Science Foundation of China [grant number 11771040]; the Project of Shandong Province Higher Educational Science and Technology Program (grant no. J16LI53).

*Corresponding author.

Email addresses: dajiena2002@163.com (Na Li), p.lin@dundee.ac.uk (Ping Lin).

rock [1], where u represents the average pressure of the liquid in the fissures in the neighborhood of the given point. Its one-dimensional counterpart $u_t = u_{xx} + au_{txx}$ was derived by Coleman and Noll as governing the simple shearing motion of a fluid of a second grade[2]. Subsequently, when Chen and Gurtin studied the heat conduction problems in different viscous media, they came to the conclusion that Eq.(1.1) related to be the simplified energy equation, and u denotes the conductive temperature[3]. Based on that concept, Ting performed further mathematical studies of the theory by considering a simple cooling process and thereby derived physically significant results[4]. The more and more extensive applications of this kind of equation in mathematics and physics have attracted the attention of many scholars. So far, a few different schemes have been proposed for the numerical solution of this type of equations([5, 6, 7, 8]).

The mixed finite element method, basically a finite element method [9] with constrained conditions, plays an important role in studying the numerical solution of higher order partial differential equations(PDEs) or PDEs including two (or more) unknown functions. Its general theory was proposed by Babuska [10] and Brezzi [11] in the early 1970s, and later on improved and incorporated with adaptivity by Falk and Osborn[12]. The mixed finite element method has now been widely used for solving fluid flow and transport equations[13, 14], for example, when the governing equations describing two-phase flow within a petroleum reservoir are expressed as a fractional flow formulation (i.e., in terms of a global pressure and a saturation), mixed methods can be very efficient and accurate in solving the pressure equation. However, mixed finite element methods have not yet applied to groundwater hydrology, for underground reservoirs often need to specify complex boundary conditions involving combinations of individual fluid fluxes and pressures, and it is sometimes impractical to express them in terms of the total quantifies. Consequently, two-pressure formulations are commonly used by hydrologists, since the complex individual boundary conditions can easily be handled. However, the coefficient in the two-pressure formulation may tend to zero because of low permeability, so that its reciprocal is not readily usable as in standard mixed finite element methods. Therefore, it is not feasible to apply mixed methods to a two-pressure formulation.

The expanded mixed finite element(EMFE) method[15, 16, 17], a new formulation expanding the standard mixed formulation, introduces three (or more) auxiliary variables for practical problems. This method was developed and analyzed by Arbogast, Wheeler and Yotov ([18]). Chen([19, 20]) detailed the EMFE method for linear and quasi-linear second-order elliptic problems. Woodward and Dawson([21]) proposed an analysis of EMFE method applied to Richards' equation, which simulates the flow of the water into a variably saturated porous medium. EMFE method can be applied to the above two-pressure formulation, so that individual boundary conditions can be handled. In addition, this method works for small diffusion or low permeability fluid problems. Using this method, we can get optimal order error estimates for certain nonlinear problems, while standard mixed formulation sometimes gives only suboptimal error estimates([22]). While the papers mentioned above are the introduction of two variables, in this paper we will introduce three parameter variables to get better approximation of more physical quantities.

The purpose of this paper is to present the expanded mixed finite element method for the 2D Sobolev equation. We show the existence and uniqueness of the solution of the mixed finite element method, and obtain its optimal error estimates. The outline of this paper is as follows. In Sect.2, we describe a semi-discrete formulation for Eq. (1.1) and provide the error estimates for the solution of the formulation. The optimal error estimate is obtained. In Sect.3, we present a fully discrete formulation from the semi-discrete formulation and analyze its error. In Sect.4, we provide two numerical examples to verify that numerical results are consistent with theoretical

conclusions.

Throughout this paper we use C to denote a generic positive constant independent of the discretization parameters h and t unless otherwise stated, which has different values in different appearances. We also adopt the standard definitions and notations of Sobolev spaces and their norms in [9] and [23].

2 Semi-discrete Scheme Based on EMFE Method

2.1 Semi-discrete Scheme and Existence and Uniqueness of Solution

By introducing three auxiliary variables

$$\mathbf{q} = \nabla u, \quad w = u_t, \quad \mathbf{p} = \nabla w,$$

the problem (1.1) can be rewritten into the following equivalent first-order differential system with regard to the solution $\{u, \mathbf{q}, w, \mathbf{p}\}$. It reads that, $\forall (x, y, t) \in \Omega \times (0, T]$

$$u_t = w, \quad \mathbf{q}_t = \mathbf{p}, \quad w - \mu \operatorname{div}(\mathbf{p}) = f + \gamma \operatorname{div}(\mathbf{q}), \quad \mathbf{p} - \nabla w = 0, \quad (2.1)$$

with boundary value

$$u(x, y, t) = 0, \quad (x, y, t) \in \partial\Omega \times (0, T],$$

and initial value

$$u(x, y, 0) = \varphi_0(x, y), \quad \mathbf{q}(x, y, 0) = \nabla \varphi_0(x, y), \quad (x, y) \in \Omega.$$

Let

$$W = H(\operatorname{div}, \Omega) = \{\tau \mid \tau \in (L^2(\Omega))^2, \operatorname{div} \tau \in L^2(\Omega)\}$$

normed by $\|\tau\|_W = \|\tau\| + \|\operatorname{div} \tau\|$, and let $V = L^2(\Omega)$.

Note that $u(x, y, t)|_{\partial\Omega} = 0$ implies $w(x, y, t)|_{\partial\Omega} = u_t(x, y, t)|_{\partial\Omega} = 0$ and integrating by parts, we obtain the weak form of (2.1): find a map $\{u, \mathbf{q}, w, \mathbf{p}\} : [0, T] \rightarrow V \times W \times V \times W$ such that

$$\left\{ \begin{array}{ll} (u_t, v) = (w, v), & \forall v \in V, 0 < t \leq T, \\ (\mathbf{q}_t, \tau) = (\mathbf{p}, \tau), & \forall \tau \in W, 0 < t \leq T, \\ (w, v) - \mu(\operatorname{div} \mathbf{p}, v) = (f, v) + \gamma(\operatorname{div} \mathbf{q}, v), & \forall v \in V, 0 < t \leq T, \\ (\mathbf{p}, \tau) + (w, \operatorname{div} \tau) = 0, & \forall \tau \in W, 0 < t \leq T. \end{array} \right. \quad (2.2)$$

where v, τ can take different values in different equations.

In order to clarify a proper finite element approximation procedure for $\{u, \mathbf{p}, w, \mathbf{q}\}$, we consider the finite-dimensional subspace $V_h \times W_h \times V_h \times W_h$ of $V \times W \times V \times W$ associated with a quasi-uniform partition τ_h of Ω into triangles, where the diameter of τ_h is not greater than h ($0 < h < 1$), and every angle of each triangle is bounded below by a positive constant. The boundary elements of τ_h are allowed to have one curvilinear edge. We choose $V_h \times W_h \times V_h \times W_h$ as the Raviart-Thomas-Nedelec space [24, 25, 26] of index $k \geq 0$ and introduce the L^2 -projection $R_h : V \rightarrow V_h$, and the Raviart-Thomas projection [26] $\pi_h : H^1(\Omega)^2 \rightarrow W_h$, which have the following useful commuting property:

$$\operatorname{div} \circ \pi_h = R_h \circ \operatorname{div} : H^1(\Omega)^2 \rightarrow V_h.$$

These projections have the following approximation properties([27, 28]):

$$\|v - R_h v\|_{-s} \leq Ch^{l+s} \|v\|_l, \quad 0 \leq l, s \leq k+1, \quad (2.3)$$

$$\|v - R_h v\|_{0,q} \leq Ch^l \|v\|_{l,q}, \quad 0 \leq l \leq k+1, 1 \leq q \leq \infty, \quad (2.4)$$

$$\|\tau - \pi_h \tau\|_{0,q} \leq Ch^l \|\tau\|_{l,q}, \quad \frac{1}{q} < l \leq k+1, 1 \leq q \leq \infty, \quad (2.5)$$

$$\|\operatorname{div}(\tau - \pi_h \tau)\| \leq Ch^l \|\operatorname{div} \tau\|_l, \quad 0 \leq l \leq k+1. \quad (2.6)$$

Our semi-discrete scheme, that is, the continuous-in-time mixed finite element approximation to (2.2) is defined by determining $\{u_h, \mathbf{q}_h, w_h, \mathbf{p}_h\} : [0, T] \rightarrow V_h \times W_h \times V_h \times W_h$ such that

$$\begin{cases} (u_{h,t}, v_h) = (w_h, v_h), & \forall v_h \in V_h, 0 < t \leq T, \\ (\mathbf{q}_{h,t}, \tau_h) = (\mathbf{p}_h, \tau_h), & \forall \tau_h \in W_h, 0 < t \leq T, \\ (w_h, v_h) - \mu(\operatorname{div} \mathbf{p}_h, v_h) = (f, v_h) + \gamma(\operatorname{div} \mathbf{q}_h, v_h), & \forall v_h \in V_h, 0 < t \leq T, \\ (\mathbf{p}_h, \tau_h) + (w_h, \operatorname{div} \tau_h) = 0, & \forall \tau_h \in W_h, 0 < t \leq T. \\ (u_h(0), v_h) = (\varphi_0, v_h), & \forall v_h \in V_h, \\ (\mathbf{q}_h(0), \tau_h) = (\nabla \varphi_0, \tau_h), & \forall \tau_h \in W_h. \end{cases} \quad (2.7)$$

Theorem 1 *The problem (2.7) has the unique solution.*

Proof. In fact, since (2.7) is linear, it suffices to show that the associated homogeneous system

$$(u_{h,t}, v_h) = (w_h, v_h), \quad \forall v_h \in V_h, 0 < t \leq T, \quad (2.8)$$

$$(\mathbf{q}_{h,t}, \tau_h) = (\mathbf{p}_h, \tau_h), \quad \forall \tau_h \in W_h, 0 < t \leq T, \quad (2.9)$$

$$(w_h, v_h) - \mu(\operatorname{div} \mathbf{p}_h, v_h) = \gamma(\operatorname{div} \mathbf{q}_h, v_h), \quad \forall v_h \in V_h, 0 < t \leq T, \quad (2.10)$$

$$(\mathbf{p}_h, \tau_h) + (w_h, \operatorname{div} \tau_h) = 0, \quad \forall \tau_h \in W_h, 0 < t \leq T, \quad (2.11)$$

$$(u_h(0), v_h) = 0, \quad \forall v_h \in V_h, \quad (2.12)$$

$$(\mathbf{q}_h(0), \tau_h) = 0, \quad \forall \tau_h \in W_h, \quad (2.13)$$

has only the trivial solution.

Let $v_h = w_h$ in (2.10), we can get that

$$\|w_h\|^2 - \mu(\operatorname{div} \mathbf{p}_h, w_h) = \gamma(\operatorname{div} \mathbf{q}_h, w_h). \quad (2.14)$$

Choosing $\tau_h = \mathbf{p}_h$ and $\tau_h = \mathbf{q}_h$ in (2.11) respectively, then we have

$$\|\mathbf{p}_h\|^2 + (w_h, \operatorname{div} \mathbf{p}_h) = 0, \quad (2.15)$$

$$(\mathbf{p}_h, \mathbf{q}_h) + (w_h, \operatorname{div} \mathbf{q}_h) = 0. \quad (2.16)$$

From (2.14), (2.15) and (2.16), using the ε -inequality, we obtain

$$\|w_h\|^2 + \mu \|\mathbf{p}_h\|^2 = -\gamma(\mathbf{p}_h, \mathbf{q}_h) \leq \gamma \|\mathbf{p}_h\| \|\mathbf{q}_h\| \leq \varepsilon \|\mathbf{p}_h\|^2 + C \|\mathbf{q}_h\|^2. \quad (2.17)$$

From (2.17) and taking $\varepsilon < \mu$, we get

$$\|w_h\|^2 + \|\mathbf{p}_h\|^2 \leq C \|\mathbf{q}_h\|^2. \quad (2.18)$$

From (2.9) we know that if we choose $\tau_h = \mathbf{q}_{h,t}$ and $\tau_h = \mathbf{p}_h$ respectively, then we have

$$\|\mathbf{q}_{h,t}\| = \|\mathbf{p}_h\|. \quad (2.19)$$

Combining (2.13),(2.18) with (2.19) yields

$$\|\mathbf{q}_h\| = \left\| \int_0^t \mathbf{q}_{h,t} dt \right\| \leq \int_0^t \|\mathbf{q}_{h,t}\| dt \leq C \int_0^t \|\mathbf{q}_h\| dt. \quad (2.20)$$

Using Gronwall's lemma, we have $\|\mathbf{q}_h\| = 0$, and from (2.18), $\|w_h\| = \|\mathbf{p}_h\| = 0$. Hence $\mathbf{q}_h \equiv \mathbf{0}$, $w_h \equiv 0$ and $\mathbf{p}_h \equiv \mathbf{0}$. By taking $v_h = u_{h,t}$ and $v_h = w_h$ in (2.8), we obtain $\|u_{h,t}\| = \|w_h\|$, and then $u_{h,t} = 0$, considering (2.12), we get $u_h \equiv 0$. \square

Therefore the solution $\{u_h, \mathbf{q}_h, w_h, \mathbf{p}_h\}$ of (2.7) is well defined.

2.2 Error Estimate

In study of mixed methods for parabolic problems, we usually introduce a mixed elliptic projection related to our equations. According to our Sobolev equations, we modify this idea and define a map $\{\bar{u}, \bar{\mathbf{q}}, \bar{w}, \bar{\mathbf{p}}\} : [0, T] \rightarrow V_h \times W_h \times V_h \times W_h$ such that

$$(u - \bar{u}, v_h) = 0, \quad \forall v_h \in V_h, 0 < t \leq T, \quad (2.21)$$

$$(\mathbf{q} - \bar{\mathbf{q}}, \tau_h) = 0, \quad \forall \tau_h \in W_h, 0 < t \leq T, \quad (2.22)$$

$$(w - \bar{w}, v_h) - \mu(\operatorname{div}(\mathbf{p} - \bar{\mathbf{p}}), v_h) = \gamma(\operatorname{div}(\mathbf{q} - \bar{\mathbf{q}}), v_h), \quad \forall v_h \in V_h, 0 < t \leq T, \quad (2.23)$$

$$(\mathbf{p} - \bar{\mathbf{p}}, \tau_h) + (w - \bar{w}, \operatorname{div}\tau_h) = 0, \quad \forall \tau_h \in W_h, 0 < t \leq T, \quad (2.24)$$

To begin with, let us demonstrate the existence and uniqueness of the solution of (2.21)-(2.24). Similar to the proof of Theorem 1, since (2.21)-(2.24) is linear, we only need to prove that there is only the trivial solution to the associated homogeneous system

$$(\bar{u}, v_h) = 0, \quad \forall v_h \in V_h, 0 < t \leq T, \quad (2.25)$$

$$(\bar{\mathbf{q}}, \tau_h) = 0, \quad \forall \tau_h \in W_h, 0 < t \leq T, \quad (2.26)$$

$$(\bar{w}, v_h) - \mu(\operatorname{div}\bar{\mathbf{p}}, v_h) = \gamma(\operatorname{div}\bar{\mathbf{q}}, v_h), \quad \forall v_h \in V_h, 0 < t \leq T, \quad (2.27)$$

$$(\bar{\mathbf{p}}, \tau_h) + (\bar{w}, \operatorname{div}\tau_h) = 0, \quad \forall \tau_h \in W_h, 0 < t \leq T. \quad (2.28)$$

Let $v_h = \bar{u}$ in (2.25) and $\tau_h = \bar{\mathbf{q}}$ in (2.26), we can easily get $\bar{u} = 0, \bar{\mathbf{q}} = \mathbf{0}$.

By taking $v_h = \bar{w}$ in (2.27) and $\tau_h = \bar{\mathbf{p}}$ in (2.28), we obtain

$$\|\bar{w}\|^2 + \mu\|\bar{\mathbf{p}}\|^2 = \gamma(\operatorname{div}\bar{\mathbf{q}}, \bar{w}) = 0,$$

which implies that $\bar{w} = 0$ and $\bar{\mathbf{p}} = \mathbf{0}$.

So, the existence and uniqueness of the solution of (2.21)-(2.24) have been demonstrated and $\{\bar{u}, \bar{\mathbf{q}}, \bar{w}, \bar{\mathbf{p}}\}$ in (2.21)-(2.24) is well defined. Next we give some error estimates of $\{\bar{u}, \bar{\mathbf{q}}, \bar{w}, \bar{\mathbf{p}}\}$.

Lemma 1 *Let $\{u, \mathbf{q}, w, \mathbf{p}\}$ and $\{\bar{u}, \bar{\mathbf{q}}, \bar{w}, \bar{\mathbf{p}}\}$ satisfy the relation (2.2) and (2.21)-(2.24), respectively. Assume that $\{u, \mathbf{q}, w, \mathbf{p}\}$ are sufficiently smooth and that Ω is 2-regular (for the definition of*

2-regularity see [24]). Then for all $0 < t \leq T$, there exists a constant $C > 0$ independent of h and t , such that

$$\begin{aligned} \|u - \bar{u}\| &\leq Ch^l \|u\|_l, \\ \|\mathbf{q} - \bar{\mathbf{q}}\| &\leq Ch^l \|\mathbf{q}\|_l, \\ \|w - \bar{w}\| &\leq Ch^l (\|w\|_l + \|\mathbf{q}\|_l + \|\mathbf{p}\|_l + \|\operatorname{div}\mathbf{q}\|_l + \|\operatorname{div}\mathbf{p}\|_l), \\ \|\mathbf{p} - \bar{\mathbf{p}}\| + \|\operatorname{div}(\mathbf{p} - \bar{\mathbf{p}})\| + \|\operatorname{div}(\mathbf{q} - \bar{\mathbf{q}})\| &\leq Ch^l (\|\mathbf{q}\|_l + \|\mathbf{p}\|_l + \|\operatorname{div}\mathbf{q}\|_l + \|\operatorname{div}\mathbf{p}\|_l), \end{aligned}$$

where $0 < l \leq k + 1$.

Proof. Let $v_h = R_h u - \bar{u}$ in (2.21), we get

$$\|R_h u - \bar{u}\|^2 = -(u - R_h u, R_h u - \bar{u}) \leq \|u - R_h u\| \|R_h u - \bar{u}\|,$$

which implies that $\|R_h u - \bar{u}\| \leq \|u - R_h u\| \leq Ch^l \|u\|_l$. Noting (2.3), we obtain

$$\|u - \bar{u}\| \leq \|u - R_h u\| + \|R_h u - \bar{u}\| \leq Ch^l \|u\|_l.$$

Similarly, let $\tau_h = \pi_h \mathbf{q} - \bar{\mathbf{q}}$ in (2.22), we get $\|\pi_h \mathbf{q} - \bar{\mathbf{q}}\| \leq Ch^l \|\mathbf{q}\|_l$, then $\|\mathbf{q} - \bar{\mathbf{q}}\| \leq Ch^l \|\mathbf{q}\|_l$.

Let $v_h = \operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})$ and $v_h = \operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})$ in (2.23) respectively, we have

$$\begin{aligned} (R_h w - \bar{w}, \operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})) - \mu \|\operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})\|^2 - \gamma (\operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}}), \operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})) \\ = -(w - R_h w, \operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})) + \mu (\operatorname{div}(\mathbf{p} - \pi_h \mathbf{p}), \operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})) + \\ \gamma (\operatorname{div}(\mathbf{q} - \pi_h \mathbf{q}), \operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})), \end{aligned} \quad (2.29)$$

$$\begin{aligned} (R_h w - \bar{w}, \operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})) - \mu (\operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}}), \operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})) - \gamma \|\operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})\|^2 \\ = -(w - R_h w, \operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})) + \mu (\operatorname{div}(\mathbf{p} - \pi_h \mathbf{p}), \operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})) + \\ \gamma (\operatorname{div}(\mathbf{q} - \pi_h \mathbf{q}), \operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})). \end{aligned} \quad (2.30)$$

Taking $\tau_h = \pi_h \mathbf{p} - \bar{\mathbf{p}}$ and $\tau_h = \pi_h \mathbf{p} - \bar{\mathbf{p}}$ in (2.24) respectively,

$$\|\pi_h \mathbf{p} - \bar{\mathbf{p}}\|^2 + (R_h w - \bar{w}, \operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})) = -(\mathbf{p} - \pi_h \mathbf{p}, \pi_h \mathbf{p} - \bar{\mathbf{p}}) - (w - R_h w, \operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})), \quad (2.31)$$

$$(\pi_h \mathbf{p} - \bar{\mathbf{p}}, \pi_h \mathbf{q} - \bar{\mathbf{q}}) + (R_h w - \bar{w}, \operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})) = -(\mathbf{p} - \pi_h \mathbf{p}, \pi_h \mathbf{q} - \bar{\mathbf{q}}) - (w - R_h w, \operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})). \quad (2.32)$$

Let $\tau_h = \pi_h \mathbf{p} - \bar{\mathbf{p}}$ in (2.22), we have

$$(\pi_h \mathbf{q} - \bar{\mathbf{q}}, \pi_h \mathbf{p} - \bar{\mathbf{p}}) = -(\mathbf{q} - \pi_h \mathbf{q}, \pi_h \mathbf{p} - \bar{\mathbf{p}}). \quad (2.33)$$

Combining (2.29)-(2.33), we obtain

$$\begin{aligned} &\mu \|\pi_h \mathbf{p} - \bar{\mathbf{p}}\|^2 + \gamma^2 \|\operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})\|^2 + \mu^2 \|\operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})\|^2 \\ &= -\mu^2 (\operatorname{div}(\mathbf{p} - \pi_h \mathbf{p}), \operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})) - \gamma^2 (\operatorname{div}(\mathbf{q} - \pi_h \mathbf{q}), \operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})) - \\ &\quad \mu (\mathbf{p} - \pi_h \mathbf{p}, \pi_h \mathbf{p} - \bar{\mathbf{p}}) - \gamma (\mathbf{p} - \pi_h \mathbf{p}, \pi_h \mathbf{q} - \bar{\mathbf{q}}) + \gamma (\mathbf{q} - \pi_h \mathbf{q}, \pi_h \mathbf{p} - \bar{\mathbf{p}}) - \\ &\quad \mu \gamma (\operatorname{div}(\mathbf{p} - \pi_h \mathbf{p}), \operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})) - \mu \gamma (\operatorname{div}(\mathbf{q} - \pi_h \mathbf{q}), \operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})) \\ &\leq \mu^2 \|\operatorname{div}(\mathbf{p} - \pi_h \mathbf{p})\| \|\operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})\| + \gamma^2 \|\operatorname{div}(\mathbf{q} - \pi_h \mathbf{q})\| \|\operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})\| + \\ &\quad \mu \|\mathbf{p} - \pi_h \mathbf{p}\| \|\pi_h \mathbf{p} - \bar{\mathbf{p}}\| + \gamma \|\mathbf{p} - \pi_h \mathbf{p}\| \|\pi_h \mathbf{q} - \bar{\mathbf{q}}\| + \gamma \|\mathbf{q} - \pi_h \mathbf{q}\| \|\pi_h \mathbf{p} - \bar{\mathbf{p}}\| + \\ &\quad \mu \gamma \|\operatorname{div}(\mathbf{p} - \pi_h \mathbf{p})\| \|\operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})\| + \mu \gamma \|\operatorname{div}(\mathbf{q} - \pi_h \mathbf{q})\| \|\operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})\|. \end{aligned}$$

Using ε -inequality, there is a sufficiently small ε such that

$$\begin{aligned}
& \mu \|\pi_h \mathbf{p} - \bar{\mathbf{p}}\|^2 + \gamma^2 \|\operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})\|^2 + \mu^2 \|\operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})\|^2 \\
& \leq C(\|\mathbf{q} - \pi_h \mathbf{q}\|^2 + \|\mathbf{p} - \pi_h \mathbf{p}\|^2 + \|\operatorname{div}(\mathbf{q} - \pi_h \mathbf{q})\|^2 + \|\operatorname{div}(\mathbf{p} - \pi_h \mathbf{p})\|^2) + \\
& \quad \varepsilon(\|\pi_h \mathbf{p} - \bar{\mathbf{p}}\|^2 + \|\operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})\|^2 + \|\operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})\|^2) \\
& \leq Ch^{2l}(\|\mathbf{q}\|_l^2 + \|\mathbf{p}\|_l^2 + \|\operatorname{div} \mathbf{q}\|_l^2 + \|\operatorname{div} \mathbf{p}\|_l^2) + \varepsilon(\|\pi_h \mathbf{p} - \bar{\mathbf{p}}\|^2 + \\
& \quad \|\operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})\|^2 + \|\operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})\|^2),
\end{aligned}$$

with that,

$$\|\pi_h \mathbf{p} - \bar{\mathbf{p}}\|^2 + \|\operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})\|^2 + \|\operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})\|^2 \leq Ch^{2l}(\|\mathbf{q}\|_l^2 + \|\mathbf{p}\|_l^2 + \|\operatorname{div} \mathbf{q}\|_l^2 + \|\operatorname{div} \mathbf{p}\|_l^2),$$

then,

$$\begin{aligned}
& \|\mathbf{p} - \bar{\mathbf{p}}\| + \|\operatorname{div}(\mathbf{q} - \bar{\mathbf{q}})\| + \|\operatorname{div}(\mathbf{p} - \bar{\mathbf{p}})\| \\
& \leq \|\mathbf{p} - \pi_h \mathbf{p}\| + \|\pi_h \mathbf{p} - \bar{\mathbf{p}}\| + \|\operatorname{div}(\mathbf{q} - \pi_h \mathbf{q})\| + \|\operatorname{div}(\pi_h \mathbf{q} - \bar{\mathbf{q}})\| + \\
& \quad \|\operatorname{div}(\mathbf{p} - \pi_h \mathbf{p})\| + \|\operatorname{div}(\pi_h \mathbf{p} - \bar{\mathbf{p}})\| \\
& \leq Ch^l(\|\mathbf{q}\|_l + \|\mathbf{p}\|_l + \|\operatorname{div} \mathbf{q}\|_l + \|\operatorname{div} \mathbf{p}\|_l).
\end{aligned}$$

Let $v_h = R_h w - \bar{w}$ in (2.23), we have

$$\begin{aligned}
\|R_h w - \bar{w}\|^2 &= - (w - R_h w, R_h w - \bar{w}) + \mu(\operatorname{div}(\mathbf{p} - \bar{\mathbf{p}}), R_h w - \bar{w}) + \gamma(\operatorname{div}(\mathbf{q} - \bar{\mathbf{q}}), R_h w - \bar{w}) \\
&\leq (\|w - R_h w\| + \mu\|\operatorname{div}(\mathbf{p} - \bar{\mathbf{p}})\| + \gamma\|\operatorname{div}(\mathbf{q} - \bar{\mathbf{q}})\|)\|R_h w - \bar{w}\|,
\end{aligned}$$

which yields

$$\begin{aligned}
\|R_h w - \bar{w}\| &\leq \|w - R_h w\| + \gamma\|\operatorname{div}(\mathbf{q} - \bar{\mathbf{q}})\| + \mu\|\operatorname{div}(\mathbf{p} - \bar{\mathbf{p}})\| \\
&\leq Ch^l(\|w\|_l + \|\mathbf{q}\|_l + \|\mathbf{p}\|_l + \|\operatorname{div} \mathbf{q}\|_l + \|\operatorname{div} \mathbf{p}\|_l),
\end{aligned}$$

thus, we conclude

$$\begin{aligned}
\|w - \bar{w}\| &\leq \|w - R_h w\| + \|R_h w - \bar{w}\| \\
&\leq Ch^l(\|w\|_l + \|\mathbf{q}\|_l + \|\mathbf{p}\|_l + \|\operatorname{div} \mathbf{q}\|_l + \|\operatorname{div} \mathbf{p}\|_l). \quad \square
\end{aligned}$$

Subtracting (2.7) from (2.2), we get the error equations

$$\begin{cases}
((u - u_h)_t, v_h) = (w - w_h, v_h), & \forall v_h \in V_h, 0 < t \leq T, \\
((\mathbf{q} - \mathbf{q}_h)_t, \tau_h) = (\mathbf{p} - \mathbf{p}_h, \tau_h), & \forall \tau_h \in W_h, 0 < t \leq T, \\
(w - w_h, v_h) - \mu(\operatorname{div}(\mathbf{p} - \mathbf{p}_h), v_h) = \gamma(\operatorname{div}(\mathbf{q} - \mathbf{q}_h), v_h), & \forall v_h \in V_h, 0 < t \leq T, \\
(\mathbf{p} - \mathbf{p}_h, \tau_h) + (w - w_h, \operatorname{div} \tau_h) = 0, & \forall \tau_h \in W_h, 0 < t \leq T.
\end{cases} \quad (2.34)$$

Using (2.21)-(2.24), (2.34) can be written as

$$((\bar{u} - u_h)_t, v_h) = (w - w_h, v_h), \quad \forall v_h \in V_h, 0 < t \leq T, \quad (2.35)$$

$$((\bar{\mathbf{q}} - \mathbf{q}_h)_t, \tau_h) = (\mathbf{p} - \mathbf{p}_h, \tau_h), \quad \forall \tau_h \in W_h, 0 < t \leq T, \quad (2.36)$$

$$(\bar{w} - w_h, v_h) - \mu(\operatorname{div}(\bar{\mathbf{p}} - \mathbf{p}_h), v_h) = \gamma(\operatorname{div}(\bar{\mathbf{q}} - \mathbf{q}_h), v_h), \quad \forall v_h \in V_h, 0 < t \leq T, \quad (2.37)$$

$$(\bar{\mathbf{p}} - \mathbf{p}_h, \tau_h) + (\bar{w} - w_h, \operatorname{div} \tau_h) = 0, \quad \forall \tau_h \in W_h, 0 < t \leq T. \quad (2.38)$$

Theorem 2 Let $\{u, \mathbf{q}, w, \mathbf{p}\}$ be the solution of (2.2) and $\{u_h, \mathbf{q}_h, w_h, \mathbf{p}_h\}$ be that of (2.7). Assume $\{u, \mathbf{q}, w, \mathbf{p}\}$ are sufficiently smooth and that Ω is 2-regular. Then there exists a constant $C > 0$ independent of h and t , such that

$$\begin{aligned} \|(u - u_h)(t)\|^2 + \|(\mathbf{q} - \mathbf{q}_h)(t)\|^2 &\leq Ch^{2l}[\|u\|_l^2 + \|\mathbf{q}\|_l^2 + \int_0^t (\|w\|_l^2 + \|\mathbf{q}\|_l^2 + \|\mathbf{p}\|_l^2 + \|\operatorname{div}\mathbf{q}\|_l^2 \\ &\quad + \|\operatorname{div}\mathbf{p}\|_l^2)dt], \\ \|(w - w_h)(t)\|^2 + \|(\mathbf{p} - \mathbf{p}_h)(t)\|^2 &\leq Ch^{2l}[\|u\|_l^2 + \|w\|_l^2 + \|\mathbf{q}\|_l^2 + \|\mathbf{p}\|_l^2 + \|\operatorname{div}\mathbf{q}\|_l^2 + \|\operatorname{div}\mathbf{p}\|_l^2 \\ &\quad + \int_0^t (\|w\|_l^2 + \|\mathbf{q}\|_l^2 + \|\mathbf{p}\|_l^2 + \|\operatorname{div}\mathbf{q}\|_l^2 + \|\operatorname{div}\mathbf{p}\|_l^2)dt], \end{aligned}$$

where $0 < t \leq T, 0 < l \leq k + 1$.

Proof. Let $v_h = \bar{w} - w_h$ in (2.37), we get

$$\|\bar{w} - w_h\|^2 - \mu(\operatorname{div}(\bar{\mathbf{p}} - \mathbf{p}_h), \bar{w} - w_h) = \gamma(\operatorname{div}(\bar{\mathbf{q}} - \mathbf{q}_h), \bar{w} - w_h). \quad (2.39)$$

Taking $\tau_h = \bar{\mathbf{p}} - \mathbf{p}_h$ and $\tau_h = \bar{\mathbf{q}} - \mathbf{q}_h$ in (2.38) respectively, we obtain

$$\|\bar{\mathbf{p}} - \mathbf{p}_h\|^2 + (\bar{w} - w_h, \operatorname{div}(\bar{\mathbf{p}} - \mathbf{p}_h)) = 0, \quad (2.40)$$

$$(\bar{\mathbf{p}} - \mathbf{p}_h, \bar{\mathbf{q}} - \mathbf{q}_h) + (\bar{w} - w_h, \operatorname{div}(\bar{\mathbf{q}} - \mathbf{q}_h)) = 0. \quad (2.41)$$

From (2.39), (2.40) and (2.41), using ε - inequality, for a sufficiently small ε that satisfy

$$\|\bar{w} - w_h\|^2 + \mu\|\bar{\mathbf{p}} - \mathbf{p}_h\|^2 = -\gamma(\bar{\mathbf{p}} - \mathbf{p}_h, \bar{\mathbf{q}} - \mathbf{q}_h) \leq \varepsilon\|\bar{\mathbf{p}} - \mathbf{p}_h\|^2 + C\|\bar{\mathbf{q}} - \mathbf{q}_h\|^2,$$

obviously,

$$\|\bar{w} - w_h\|^2 + \|\bar{\mathbf{p}} - \mathbf{p}_h\|^2 \leq C\|\bar{\mathbf{q}} - \mathbf{q}_h\|^2. \quad (2.42)$$

Let $v_h = \bar{u} - u_h$ in (2.35), and $\tau_h = \bar{\mathbf{q}} - \mathbf{q}_h$ in (2.36), we get

$$\begin{aligned} \frac{d}{dt}\|\bar{u} - u_h\|^2 &= 2(w - w_h, \bar{u} - u_h) \\ &\leq 2(\|w - \bar{w}\|^2 + \|\bar{w} - w_h\|^2 + \|\bar{u} - u_h\|^2) \\ &\leq C[h^{2l}(\|w\|_l^2 + \|\mathbf{q}\|_l^2 + \|\mathbf{p}\|_l^2 + \|\operatorname{div}\mathbf{q}\|_l^2 + \|\operatorname{div}\mathbf{p}\|_l^2) + \|\bar{\mathbf{q}} - \mathbf{q}_h\|^2 + \|\bar{u} - u_h\|^2], \end{aligned}$$

$$\begin{aligned} \frac{d}{dt}\|\bar{\mathbf{q}} - \mathbf{q}_h\|^2 &= 2(\mathbf{p} - \mathbf{p}_h, \bar{\mathbf{q}} - \mathbf{q}_h) \\ &\leq 2(\|\mathbf{p} - \bar{\mathbf{p}}\|^2 + \|\bar{\mathbf{p}} - \mathbf{p}_h\|^2 + \|\bar{\mathbf{q}} - \mathbf{q}_h\|^2) \\ &\leq C[h^{2l}(\|\mathbf{q}\|_l^2 + \|\mathbf{p}\|_l^2 + \|\operatorname{div}\mathbf{q}\|_l^2 + \|\operatorname{div}\mathbf{p}\|_l^2) + \|\bar{\mathbf{q}} - \mathbf{q}_h\|^2], \end{aligned}$$

obviously,

$$\begin{aligned} \frac{d}{dt}(\|\bar{u} - u_h\|^2 + \|\bar{\mathbf{q}} - \mathbf{q}_h\|^2) &\leq C[h^{2l}(\|w\|_l^2 + \|\mathbf{q}\|_l^2 + \|\mathbf{p}\|_l^2 + \|\operatorname{div}\mathbf{q}\|_l^2 + \|\operatorname{div}\mathbf{p}\|_l^2) + \\ &\quad \|\bar{u} - u_h\|^2 + \|\bar{\mathbf{q}} - \mathbf{q}_h\|^2], \end{aligned}$$

with that, for $0 < t \leq T$, we have

$$\begin{aligned} \|(\bar{u} - u_h)(t)\|^2 + \|(\bar{\mathbf{q}} - \mathbf{q}_h)(t)\|^2 &\leq C[h^{2l} \int_0^t (\|w\|_l^2 + \|\mathbf{p}\|_l^2 + \|\mathbf{q}\|_l^2 + \|\operatorname{div}\mathbf{p}\|_l^2 + \|\operatorname{div}\mathbf{q}\|_l^2) dt \\ &\quad + \int_0^t (\|\bar{u} - u_h\|^2 + \|\bar{\mathbf{q}} - \mathbf{q}_h\|^2) dt]. \end{aligned}$$

Using Gronwall's lemma, we have

$$\|(\bar{u} - u_h)(t)\|^2 + \|(\bar{\mathbf{q}} - \mathbf{q}_h)(t)\|^2 \leq Ch^{2l} \int_0^t (\|w\|_l^2 + \|\mathbf{q}\|_l^2 + \|\mathbf{p}\|_l^2 + \|\operatorname{div}\mathbf{q}\|_l^2 + \|\operatorname{div}\mathbf{p}\|_l^2) dt,$$

noticing (2.42),

$$\|(\bar{w} - w_h)(t)\|^2 + \|(\bar{\mathbf{p}} - \mathbf{p}_h)(t)\|^2 \leq Ch^{2l} \int_0^t (\|w\|_l^2 + \|\mathbf{q}\|_l^2 + \|\mathbf{p}\|_l^2 + \|\operatorname{div}\mathbf{q}\|_l^2 + \|\operatorname{div}\mathbf{p}\|_l^2) dt.$$

Finally, we get

$$\begin{aligned} \|(u - u_h)(t)\|^2 + \|(\mathbf{q} - \mathbf{q}_h)(t)\|^2 &\leq Ch^{2l} [\|u\|_l^2 + \|\mathbf{q}\|_l^2 + \int_0^t (\|w\|_l^2 + \|\mathbf{q}\|_l^2 + \|\mathbf{p}\|_l^2 + \|\operatorname{div}\mathbf{q}\|_l^2 \\ &\quad + \|\operatorname{div}\mathbf{p}\|_l^2) dt], \\ \|(w - w_h)(t)\|^2 + \|(\mathbf{p} - \mathbf{p}_h)(t)\|^2 &\leq Ch^{2l} [\|u\|_l^2 + \|w\|_l^2 + \|\mathbf{q}\|_l^2 + \|\mathbf{p}\|_l^2 + \|\operatorname{div}\mathbf{q}\|_l^2 + \|\operatorname{div}\mathbf{p}\|_l^2 \\ &\quad + \int_0^t (\|w\|_l^2 + \|\mathbf{q}\|_l^2 + \|\mathbf{p}\|_l^2 + \|\operatorname{div}\mathbf{q}\|_l^2 + \|\operatorname{div}\mathbf{p}\|_l^2) dt]. \quad \square \end{aligned}$$

3 Fully Discrete Scheme Based on EMFE Method

Let $0 = t_0 < t_1 < \dots < t_N = T$ be a partition of the domain $[0, T]$, $\Delta t_n = t_n - t_{n-1}$ ($n = 1, 2, \dots, N$), $(u_h^n, \mathbf{q}_h^n, w_h^n, \mathbf{p}_h^n)$ is the approximation of $(u, \mathbf{q}, w, \mathbf{p})$ at t_n . In order to make the scheme more stable, the Euler backward difference scheme with first order accuracy is adopted to discretize the time variables t . If the two order accuracy is to be reached, the Crank-Nicolson format can be considered.

Introduce the mark: $\partial_t u^n = \frac{u^n - u^{n-1}}{\Delta t}$, $\partial_t \mathbf{q}^n = \frac{\mathbf{q}^n - \mathbf{q}^{n-1}}{\Delta t}$. Then, we get the fully discrete scheme: find $(u_h^n, \mathbf{q}_h^n, w_h^n, \mathbf{p}_h^n)$ such that

$$\begin{cases} (\partial_t u_h^n, v_h) = (w_h^n, v_h), & \forall v_h \in V_h, \\ (\partial_t \mathbf{q}_h^n, \tau_h) = (\mathbf{p}_h^n, \tau_h), & \forall \tau_h \in W_h, \\ (w_h^n, v_h) - \mu(\operatorname{div}\mathbf{p}_h^n, v_h) = (f^n, v_h) + \gamma(\operatorname{div}\mathbf{q}_h^n, v_h), & \forall v_h \in V_h, \\ (\mathbf{p}_h^n, \tau_h) + (w_h^n, \operatorname{div}\tau_h) = 0, & \forall \tau_h \in W_h, \\ u_h^0 = \varphi_0(x, y), \quad \mathbf{q}_h^0 = \nabla\varphi_0(x, y). \end{cases} \quad (3.1)$$

Theorem 3 *The solution of (3.1) exists uniquely.*

Proof. Since (3.1) is linear, it suffices to show that the associated homogeneous system has only the trivial solution, that is, let $u_h^{n-1} = 0$, $\mathbf{q}_h^{n-1} = \mathbf{0}$, $f^n = 0$, we get the associated homogeneous system:

$$(u_h^n, v_h) = \Delta t(w_h^n, v_h), \quad \forall v_h \in V_h, \quad (3.2)$$

$$(\mathbf{q}_h^n, \tau_h) = \Delta t(\mathbf{p}_h^n, \tau_h), \quad \forall \tau_h \in W_h, \quad (3.3)$$

$$(w_h^n, v_h) - \mu(\operatorname{div}(\mathbf{p}_h^n), v_h) = \gamma(\operatorname{div}(\mathbf{q}_h^n), v_h), \quad \forall v_h \in V_h, \quad (3.4)$$

$$(\mathbf{p}_h^n, \tau_h) + (w_h^n, \operatorname{div}\tau_h) = 0, \quad \forall \tau_h \in W_h. \quad (3.5)$$

Taking $v_h = w_h^n$ in (3.4) and $\tau_h = \mathbf{p}_h^n$, $\tau_h = \mathbf{q}_h^n$ in (3.5) respectively, we have

$$\|w_h^n\|^2 - \mu(\operatorname{div}\mathbf{p}_h^n, w_h^n) = \gamma(\operatorname{div}\mathbf{q}_h^n, w_h^n), \quad (3.6)$$

$$\|\mathbf{p}_h^n\|^2 + (w_h^n, \operatorname{div}\mathbf{p}_h^n) = 0, \quad (3.7)$$

$$(\mathbf{p}_h^n, \mathbf{q}_h^n) + (w_h^n, \operatorname{div}\mathbf{q}_h^n) = 0. \quad (3.8)$$

Combine (3.6), (3.7) with (3.8), using ε -inequality, we obtain

$$\|w_h^n\|^2 + \mu\|\mathbf{p}_h^n\|^2 = \gamma(\operatorname{div}\mathbf{q}_h^n, w_h^n) = -\gamma(\mathbf{p}_h^n, \mathbf{q}_h^n) \leq \varepsilon\|\mathbf{p}_h^n\|^2 + C\|\mathbf{q}_h^n\|^2,$$

with that

$$\|w_h^n\|^2 + \|\mathbf{p}_h^n\|^2 \leq C\|\mathbf{q}_h^n\|^2. \quad (3.9)$$

Let $v_h = u_h^n$ and $v_h = w_h^n$ in (3.2) respectively, we have

$$\|u_h^n\|^2 = \Delta t(w_h^n, u_h^n) = (\Delta t)^2\|w_h^n\|^2. \quad (3.10)$$

Similarly, take $\tau_h = \mathbf{q}_h^n$ and $\tau_h = \mathbf{p}_h^n$ in (3.3) respectively, we have

$$\|\mathbf{q}_h^n\|^2 = (\Delta t)^2\|\mathbf{p}_h^n\|^2. \quad (3.11)$$

It thus follows that

$$\|u_h^n\|^2 + \|\mathbf{q}_h^n\|^2 \leq (\Delta t)^2(\|w_h^n\|^2 + \|\mathbf{p}_h^n\|^2) \leq C(\Delta t)^2\|\mathbf{q}_h^n\|^2. \quad (3.12)$$

If Δt is sufficiently small such that $C(\Delta t)^2 < 1$, we then get $\|u_h^n\| = 0$, $\|\mathbf{q}_h^n\| = 0$, noting (3.10) and (3.11) yields $\|w_h^n\| = 0$, $\|\mathbf{p}_h^n\| = 0$. Hence we can obtain

$$u_h^n = 0, \mathbf{q}_h^n = \mathbf{0}, w_h^n = 0, \mathbf{p}_h^n = \mathbf{0}. \quad \square$$

Next, we analyze the error estimate of the format (3.1). Subtracting (3.1) from (2.2), we get the error equations

$$\begin{cases} (u_t^n - \partial_t u_h^n, v_h) = (w^n - w_h^n, v_h), & \forall v_h \in V_h, \\ (\mathbf{q}_t^n - \partial_t \mathbf{q}_h^n, \tau_h) = (\mathbf{p}^n - \mathbf{p}_h^n, \tau_h), & \forall \tau_h \in W_h, \\ (w^n - w_h^n, v_h) - \mu(\operatorname{div}(\mathbf{p}^n - \mathbf{p}_h^n), v_h) = \gamma(\operatorname{div}(\mathbf{q}^n - \mathbf{q}_h^n), v_h), & \forall v_h \in V_h, \\ (\mathbf{p}^n - \mathbf{p}_h^n, \tau_h) + (w^n - w_h^n, \operatorname{div}\tau_h) = 0, & \forall \tau_h \in W_h. \end{cases} \quad (3.13)$$

Firstly, we introduce the marks:

$$\begin{aligned} u^n - u_h^n &= u^n - \bar{u}^n + \bar{u}^n - u_h^n = \rho_u^n + \theta_u^n, \\ \mathbf{q}^n - \mathbf{q}_h^n &= \mathbf{q}^n - \bar{\mathbf{q}}^n + \bar{\mathbf{q}}^n - \mathbf{q}_h^n = \rho_{\mathbf{q}}^n + \theta_{\mathbf{q}}^n, \\ w^n - w_h^n &= w^n - \bar{w}^n + \bar{w}^n - w_h^n = \rho_w^n + \theta_w^n, \\ \mathbf{p}^n - \mathbf{p}_h^n &= \mathbf{p}^n - \bar{\mathbf{p}}^n + \bar{\mathbf{p}}^n - \mathbf{p}_h^n = \rho_{\mathbf{p}}^n + \theta_{\mathbf{p}}^n. \end{aligned}$$

using (2.21)-(2.24), the error equations can be rewritten as

$$(\theta_u^n - \theta_u^{n-1}, v_h) = \Delta t(\rho_w^n, v_h) + \Delta t(\theta_w^n, v_h) - \Delta t(u_t^n - \partial_t u^n, v_h), \quad \forall v_h \in V_h, \quad (3.14)$$

$$(\theta_{\mathbf{q}}^n - \theta_{\mathbf{q}}^{n-1}, \tau_h) = \Delta t(\rho_{\mathbf{p}}^n, \tau_h) + \Delta t(\theta_{\mathbf{p}}^n, \tau_h) - \Delta t(\mathbf{q}_t^n - \partial_t \mathbf{q}^n, \tau_h), \quad \forall \tau_h \in W_h, \quad (3.15)$$

$$(\theta_w^n, v_h) - \mu(\operatorname{div} \theta_{\mathbf{p}}^n, v_h) = \gamma(\operatorname{div} \theta_{\mathbf{q}}^n, v_h), \quad \forall v_h \in V_h, \quad (3.16)$$

$$(\theta_{\mathbf{p}}^n, \tau_h) + (\theta_w^n, \operatorname{div} \tau_h) = 0, \quad \forall \tau_h \in W_h, \quad (3.17)$$

$$\theta_u^0 = 0, \quad \theta_{\mathbf{q}}^0 = \mathbf{0}. \quad (3.18)$$

By taking $v_h = \theta_w^n$ in (3.16) and $\tau_h = \theta_{\mathbf{p}}^n$, $\tau_h = \theta_{\mathbf{q}}^n$ in (3.17) respectively, we have

$$\begin{aligned} \|\theta_w^n\|^2 - \mu(\operatorname{div} \theta_{\mathbf{p}}^n, \theta_w^n) &= \gamma(\operatorname{div} \theta_{\mathbf{q}}^n, \theta_w^n), \\ \|\theta_{\mathbf{p}}^n\|^2 + (\theta_w^n, \operatorname{div} \theta_{\mathbf{p}}^n) &= 0, \\ (\theta_{\mathbf{p}}^n, \theta_{\mathbf{q}}^n) + (\theta_w^n, \operatorname{div} \theta_{\mathbf{q}}^n) &= 0. \end{aligned}$$

Applying ε -inequality, it thus follows that

$$\|\theta_w^n\|^2 + \mu\|\theta_{\mathbf{p}}^n\|^2 = \gamma(\operatorname{div} \theta_{\mathbf{q}}^n, \theta_w^n) = -\gamma(\theta_{\mathbf{p}}^n, \theta_{\mathbf{q}}^n) \leq \varepsilon\|\theta_{\mathbf{p}}^n\|^2 + C\|\theta_{\mathbf{q}}^n\|^2,$$

and then

$$\|\theta_w^n\|^2 + \|\theta_{\mathbf{p}}^n\|^2 \leq C\|\theta_{\mathbf{q}}^n\|^2. \quad (3.19)$$

Considering the following inequality:

$$\begin{aligned} \|u_t^n - \partial_t u^n\| &\leq C \int_{t_{n-1}}^{t_n} \|u_{tt}\| dt, \\ \|\mathbf{q}_t^n - \partial_t \mathbf{q}^n\| &\leq C \int_{t_{n-1}}^{t_n} \|\mathbf{q}_{tt}\| dt, \end{aligned}$$

taking $v_h = \theta_u^n$ in (3.14) yields

$$\begin{aligned} \|\theta_u^n\| &\leq \|\theta_u^{n-1}\| + \Delta t(\|\rho_w^n\| + \|\theta_w^n\|) + \Delta t\|u_t^n - \partial_t u^n\| \\ &\leq \|\theta_u^{n-1}\| + \Delta t \cdot h^l(\|w\|_l + \|\mathbf{p}\|_l + \|\mathbf{q}\|_l + \|\operatorname{div} \mathbf{p}\|_l + \|\operatorname{div} \mathbf{q}\|_l) + \\ &\quad C\Delta t\|\theta_{\mathbf{q}}^n\| + C\Delta t \int_{t_{n-1}}^{t_n} \|u_{tt}\| dt. \end{aligned}$$

then, take $\tau_h = \theta_{\mathbf{q}}^n$ in (3.15), we get

$$\begin{aligned} \|\theta_{\mathbf{q}}^n\| &\leq \|\theta_{\mathbf{q}}^{n-1}\| + \Delta t(\|\rho_{\mathbf{p}}^n\| + \|\theta_{\mathbf{p}}^n\|) + \Delta t\|\mathbf{q}_t^n - \partial_t \mathbf{q}^n\| \\ &\leq \|\theta_{\mathbf{q}}^{n-1}\| + \Delta t \cdot h^l(\|\mathbf{p}\|_l + \|\mathbf{q}\|_l + \|\operatorname{div} \mathbf{p}\|_l + \|\operatorname{div} \mathbf{q}\|_l) + C\Delta t\|\theta_{\mathbf{q}}^n\| + C\Delta t \int_{t_{n-1}}^{t_n} \|\mathbf{q}_{tt}\| dt. \end{aligned}$$

Using the two inequality above, we get

$$\begin{aligned} \|\theta_u^n\| + \|\theta_{\mathbf{q}}^n\| &\leq \|\theta_u^{n-1}\| + \|\theta_{\mathbf{q}}^{n-1}\| + 2\Delta t \cdot h^l (\|w\|_l + \|\mathbf{p}\|_l + \|\mathbf{q}\|_l + \|\operatorname{div}\mathbf{p}\|_l + \|\operatorname{div}\mathbf{q}\|_l) \\ &\quad + C\Delta t \|\theta_{\mathbf{q}}^n\| + C\Delta t \left(\int_{t_{n-1}}^{t_n} \|u_{tt}\| dt + \int_{t_{n-1}}^{t_n} \|\mathbf{q}_{tt}\| dt \right). \end{aligned}$$

If Δt is sufficiently small such that $C\Delta t < 1$, we obtain

$$\begin{aligned} \|\theta_u^n\| + \|\theta_{\mathbf{q}}^n\| &\leq C[\|\theta_u^{n-1}\| + \|\theta_{\mathbf{q}}^{n-1}\| + \Delta t \cdot h^l (\|w\|_l + \|\mathbf{p}\|_l + \|\mathbf{q}\|_l + \|\operatorname{div}\mathbf{p}\|_l + \|\operatorname{div}\mathbf{q}\|_l) + \\ &\quad \Delta t \left(\int_{t_{n-1}}^{t_n} \|u_{tt}\| dt + \int_{t_{n-1}}^{t_n} \|\mathbf{q}_{tt}\| dt \right)] \\ &\leq C[\|\theta_u^0\| + \|\theta_{\mathbf{q}}^0\| + n\Delta t \cdot h^l (\|w\|_l + \|\mathbf{p}\|_l + \|\mathbf{q}\|_l + \|\operatorname{div}\mathbf{p}\|_l + \|\operatorname{div}\mathbf{q}\|_l) + \\ &\quad \Delta t \left(\int_{t_0}^{t_n} \|u_{tt}\| dt + \int_{t_0}^{t_n} \|\mathbf{q}_{tt}\| dt \right)] \\ &\leq C[h^l (\|w\|_l + \|\mathbf{p}\|_l + \|\mathbf{q}\|_l + \|\operatorname{div}\mathbf{p}\|_l + \|\operatorname{div}\mathbf{q}\|_l) + \Delta t \left(\int_{t_0}^{t_n} \|u_{tt}\| dt + \int_{t_0}^{t_n} \|\mathbf{q}_{tt}\| dt \right)], \end{aligned}$$

using Theorem 2,

$$\begin{aligned} &\|u^n - u_h^n\| + \|\mathbf{q}^n - \mathbf{q}_h^n\| \\ &\leq \|\rho_u^n\| + \|\theta_u^n\| + \|\rho_{\mathbf{q}}^n\| + \|\theta_{\mathbf{q}}^n\| \\ &\leq C[h^l (\|w\|_l + \|\mathbf{p}\|_l + \|\mathbf{q}\|_l + \|\operatorname{div}\mathbf{p}\|_l + \|\operatorname{div}\mathbf{q}\|_l) + \Delta t \left(\int_{t_0}^{t_n} \|u_{tt}\| dt + \int_{t_0}^{t_n} \|\mathbf{q}_{tt}\| dt \right)]. \end{aligned}$$

Due to (3.19), we obtain

$$\begin{aligned} \|\theta_w^n\| + \|\theta_{\mathbf{p}}^n\| &\leq C[h^l (\|w\|_l + \|\mathbf{p}\|_l + \|\mathbf{q}\|_l + \|\operatorname{div}\mathbf{p}\|_l + \|\operatorname{div}\mathbf{q}\|_l) + \\ &\quad \Delta t \left(\int_{t_0}^{t_n} \|u_{tt}\| dt + \int_{t_0}^{t_n} \|\mathbf{q}_{tt}\| dt \right)], \end{aligned}$$

and

$$\begin{aligned} \|w^n - w_h^n\| + \|\mathbf{p}^n - \mathbf{p}_h^n\| &\leq C[h^l (\|w\|_l + \|\mathbf{p}\|_l + \|\mathbf{q}\|_l + \|\operatorname{div}\mathbf{p}\|_l + \|\operatorname{div}\mathbf{q}\|_l) + \\ &\quad \Delta t \left(\int_{t_0}^{t_n} \|u_{tt}\| dt + \int_{t_0}^{t_n} \|\mathbf{q}_{tt}\| dt \right)]. \end{aligned}$$

We summarize the results of the analysis into the following theorem.

Theorem 4 *Under the assumptions of Theorem 2, and $\Delta t \leq \delta$ (δ sufficiently small and independent of h) the fully discrete scheme has prior error estimate as following*

$$\begin{aligned} &\max_{1 \leq n \leq N} \|u^n - u_h^n\| + \max_{1 \leq n \leq N} \|\mathbf{q}^n - \mathbf{q}_h^n\| + \max_{1 \leq n \leq N} \|w^n - w_h^n\| + \max_{1 \leq n \leq N} \|\mathbf{p}^n - \mathbf{p}_h^n\| \\ &\leq C[h^l (\|w\|_{L^\infty(0,T;H^1(\Omega))} + \|\mathbf{q}\|_{L^\infty(0,T;H^1(\Omega))} + \|\mathbf{p}\|_{L^\infty(0,T;H^1(\Omega))} + \|\operatorname{div}\mathbf{q}\|_{L^\infty(0,T;H^1(\Omega))} + \\ &\quad \|\operatorname{div}\mathbf{p}\|_{L^\infty(0,T;H^1(\Omega))}) + \Delta t (\|u_{tt}\|_{L^1(0,T;H^0(\Omega))} + \|\mathbf{q}_{tt}\|_{L^1(0,T;H^0(\Omega))})], \end{aligned}$$

where $0 < t \leq T, 0 < l \leq k + 1$, C is a positive constant independent of h and Δt .

4 Numerical Experiment

In this section, two numerical examples are solved by the EMFE method in two cases ($k = 0$ and $k = 1$). All computations are performed under the P_k (piecewise polynomial of degree k) finite element space for u and w , and RT_k (Raviart-Thomas-Nedelec of index k) finite element space for \mathbf{q} and \mathbf{p} . FreeFem++ platform and Matlab are used to calculate and show the results.

Example 1. We choose the computational domain as $\bar{\Omega} = [0, 1] \times [0, 1]$, $a = 1$, $b = 1$, $c = 1$, the initial value function $u_0(x, y) = \sin(\pi x)\sin(\pi y)$, and the source term $f(x, y, t) = (1 + 4\pi^2)e^t \sin(\pi x)\sin(\pi y)$. It is easy to verify that the exact solution is $u = e^t \sin(\pi x)\sin(\pi y)$, we calculate the equation till the final time $T = 1$.

- Let $k = 0$, that is to say, $V_h = P_0$, $W_h = RT_0$. Table 1 lists the L_2 norms of errors between u_h , \mathbf{q}_h , w_h , \mathbf{p}_h and u , \mathbf{q} , w , \mathbf{p} with different subsections at $t = 0.2, 0.4, 0.6, 0.8, 1$, and Table 2 lists the corresponding convergence orders, from which we can see that the errors are quite small and the convergence orders are all about 1, the approximation effect is very good.
- When $k = 1$, i.e. $V_h = P_1$, $W_h = RT_1$. We plot the images of the analytical solution u and the numerical solution u_h at $t = 1$ as shown in Fig.1, and we can intuitively realize that the error between u_h , \mathbf{q}_h , w_h , \mathbf{p}_h and u , \mathbf{q} , w , \mathbf{p} are quite small, and the thinner the split, the smaller the error, until an almost complete coincidence. Similarly, we list L_2 error norms between u_h , \mathbf{q}_h , w_h , \mathbf{p}_h and u , \mathbf{q} , w , \mathbf{p} in Table 3, it can be seen that the approximation speed in this case is faster than that of $k = 0$. Table 4 lists the corresponding convergence orders, and the results show that the accuracy is in accordance with the theoretical analysis.

Example 2. We compute the same example given in [29]. The computational domain is $\bar{\Omega} = [0, 1] \times [0, 1]$, $a = 1$, $b = 1$, $c = 1$, the initial value function $u_0(x, y) = xy(1 - x)(1 - y)$, and the source term $f(x, y, t) = e^t[xy(1 - x)(1 - y) + 4x(1 - x) + 4y(1 - y)]$. The analytical solution is $u = e^t xy(1 - x)(1 - y)$.

In this example, we also show the long time stability of the method, thus we choose $T = 10$. To further show the good stability, we use a large time step. We notice that the analytical solution u contains the factor e^t , its value increases rapidly with time, we thus calculate the relative errors. It needs to be noted that the conclusions obtained in our theoretical analysis are also true for the relative errors.


- For $k = 0$, Table 5 lists the L_2 norms of relative errors between u_h , \mathbf{q}_h , w_h , \mathbf{p}_h and u , \mathbf{q} , w , \mathbf{p} at $t = 2, 4, 6, 8, 10$. The errors show similar rule with time and subdivision as in Example 1. The corresponding convergence orders are listed in Table 6, which is in accordance with the theoretical analysis.
- For $k = 1$, we also list the L_2 norms of relative errors of four unknown variables and corresponding convergence orders in Table 7 and Table 8, respectively, both of them illustrate the effectiveness of the method. Compared with [29], in which different mixed finite element methods are used to solve this problem, we get not only more variables, but also smaller errors under the same partition.

Table 1: L_2 norms of errors for u , \mathbf{q} , w , and \mathbf{p}


		$t = 0.2$	$t = 0.4$	$t = 0.6$	$t = 0.8$	$t = 1.0$
$\ u - u_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{5}$	9.63793e-3	1.97476e-2	3.06969e-2	4.28932e-2	5.67993e-2
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{10}$	4.94052e-3	1.01264e-2	1.57514e-2	2.20292e-2	2.92016e-2
	$h_3 = \frac{1}{32}, \Delta t_3 = \frac{1}{20}$	2.50091e-3	5.12698e-3	7.97771e-3	1.11626e-2	1.48055e-2
	$h_4 = \frac{1}{64}, \Delta t_4 = \frac{1}{40}$	1.25809e-3	2.57936e-3	4.01427e-3	5.61828e-3	7.45391e-3
$\ \mathbf{q} - \mathbf{q}_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{5}$	4.57217e-2	9.42955e-2	1.47518e-1	2.07395e-1	2.76223e-1
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{10}$	2.26422e-2	4.65397e-2	7.25957e-2	1.01809e-1	1.35313e-1
	$h_3 = \frac{1}{32}, \Delta t_3 = \frac{1}{20}$	1.12733e-2	2.31304e-2	3.60253e-2	5.04574e-2	6.69906e-2
	$h_4 = \frac{1}{64}, \Delta t_4 = \frac{1}{40}$	5.62575e-3	1.15323e-2	1.79473e-2	2.51207e-2	3.33338e-2
$\ w - w_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{5}$	9.63163e-3	1.98354e-2	3.11424e-2	4.38879e-2	5.85412e-2
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{10}$	4.73150e-3	9.80166e-3	1.53485e-2	2.15724e-2	2.87115e-2
	$h_4 = \frac{1}{32}, \Delta t_3 = \frac{1}{20}$	2.36306e-3	4.88055e-3	7.62302e-3	1.06946e-2	1.42146e-2
	$h_3 = \frac{1}{64}, \Delta t_4 = \frac{1}{40}$	1.18348e-3	2.43667e-3	3.62159e-3	5.12362e-3	7.07249e-3
$\ \mathbf{p} - \mathbf{p}_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{5}$	2.78376e-2	6.48858e-2	1.08826e-1	1.57067e-1	2.12917e-1
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{10}$	1.64798e-2	3.75976e-2	6.07238e-2	8.65551e-2	1.16061e-1
	$h_3 = \frac{1}{32}, \Delta t_3 = \frac{1}{20}$	9.37106e-3	2.03038e-2	3.21754e-2	4.54276e-2	6.05747e-2
	$h_4 = \frac{1}{64}, \Delta t_4 = \frac{1}{40}$	5.00782e-3	1.05482e-2	1.65567e-2	2.32655e-2	3.09375e-2

Table 2: Convergence orders of u , \mathbf{q} , w , and \mathbf{p}


		$t = 0.2$	$t = 0.4$	$t = 0.6$	$t = 0.8$	$t = 1.0$
$\ u - u_h\ $	h_1/h_2	0.964060	0.963556	0.962613	0.961332	0.959826
	h_2/h_3	0.982210	0.981940	0.981433	0.980744	0.979914
	h_3/h_4	0.991218	0.991096	0.990837	0.990480	0.990064
$\ \mathbf{q} - \mathbf{q}_h\ $	h_1/h_2	1.013865	1.018727	1.022935	1.026516	1.029533
	h_2/h_3	1.006104	1.008672	1.010874	1.012727	1.014270
	h_3/h_4	1.002793	1.004110	1.005244	1.006189	1.006973
$\ w - w_h\ $	h_1/h_2	1.025482	1.016979	1.020782	1.024637	1.027824
	h_2/h_3	1.001641	1.005982	1.009663	1.012304	1.014255
	h_3/h_4	0.997621	1.002133	1.073739	1.061647	1.007083
$\ \mathbf{p} - \mathbf{p}_h\ $	h_1/h_2	0.756336	0.787262	0.841689	0.859689	0.875408
	h_2/h_3	0.814415	0.888891	0.916304	0.930050	0.938096
	h_3/h_4	0.904030	0.944753	0.958543	0.965377	0.969359




(a) u, u_h at $h = \frac{1}{8}$ and $\Delta t = \frac{1}{5}$



(b) u, u_h at $h = \frac{1}{16}$ and $\Delta t = \frac{1}{20}$



(c) u, u_h at $h = \frac{1}{32}$ and $\Delta t = \frac{1}{80}$



(d) u, u_h at $h = \frac{1}{64}$ and $\Delta t = \frac{1}{320}$

Fig. 1: Comparison of numerical solution u_h with exact solution u at $t = 1$

Table 3: L_2 norms of errors for u , \mathbf{q} , w , and \mathbf{p}

		$t = 0.2$	$t = 0.4$	$t = 0.6$	$t = 0.8$	$t = 1.0$
$\ u - u_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{5}$	9.67088e-3	1.99375e-2	3.11787e-2	4.38175e-2	5.83378e-2
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{20}$	2.50417e-3	5.13787e-3	8.00183e-3	1.12069e-2	1.48783e-2
	$h_3 = \frac{1}{32}, \Delta t_3 = \frac{1}{80}$	6.31323e-4	1.29356e-3	2.01232e-3	2.81565e-3	3.73515e-3
	$h_4 = \frac{1}{64}, \Delta t_4 = \frac{1}{320}$	1.58101e-4	3.23839e-4	5.03615e-4	7.04489e-4	9.34362e-4
$\ \mathbf{q} - \mathbf{q}_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{5}$	4.25240e-2	8.76765e-2	1.37125e-1	1.92731e-1	2.56625e-1
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{20}$	1.10826e-2	2.27408e-2	3.54210e-2	4.96141e-2	6.58747e-2
	$h_3 = \frac{1}{32}, \Delta t_3 = \frac{1}{80}$	2.80302e-3	5.74391e-3	8.93647e-3	1.25053e-2	1.63174e-2
	$h_4 = \frac{1}{64}, \Delta t_4 = \frac{1}{320}$	7.03103e-4	1.44028e-3	2.24014e-3	3.13397e-3	4.15698e-3
$\ w - w_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{5}$	9.12603e-3	1.86851e-2	2.92441e-2	4.11360e-2	5.48032e-2
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{20}$	2.38605e-3	4.85550e-3	7.55967e-3	1.05903e-2	1.40633e-2
	$h_3 = \frac{1}{32}, \Delta t_3 = \frac{1}{80}$	6.05522e-4	1.22851e-3	1.90900e-3	2.67063e-3	3.54280e-3
	$h_4 = \frac{1}{64}, \Delta t_4 = \frac{1}{320}$	1.52210e-4	3.08387e-4	4.78838e-4	6.69553e-4	8.87910e-4
$\ \mathbf{p} - \mathbf{p}_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{5}$	4.53065e-2	8.62345e-2	1.32585e-1	1.85173e-1	2.45813e-1
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{20}$	1.15993e-2	2.22267e-2	3.41311e-2	4.75588e-2	6.29896e-2
	$h_3 = \frac{1}{32}, \Delta t_3 = \frac{1}{80}$	2.91513e-3	5.60208e-3	8.60146e-3	1.19788e-2	1.58561e-2
	$h_4 = \frac{1}{64}, \Delta t_4 = \frac{1}{320}$	7.29291e-4	1.40352e-3	2.15526e-3	3.00126e-3	3.97223e-3

Table 4: Convergence orders of u , \mathbf{q} , w , and \mathbf{p}

		$t = 0.2$	$t = 0.4$	$t = 0.6$	$t = 0.8$	$t = 1.0$
$\ u - u_h\ $	h_1/h_2	1.949315	1.956242	1.962159	1.967120	1.971221
	h_2/h_3	1.987882	1.989823	1.991470	1.992847	1.993972
	h_3/h_4	1.997532	1.998038	1.998467	1.998819	1.999113
$\ \mathbf{q} - \mathbf{q}_h\ $	h_1/h_2	1.939981	1.946907	1.952815	1.957767	1.961865
	h_2/h_3	1.983242	1.990210	1.986828	1.988211	2.013313
	h_3/h_4	1.995174	1.995684	1.996116	1.996476	1.972803
$\ w - w_h\ $	h_1/h_2	1.935363	1.944197	1.951750	1.957658	1.962325
	h_2/h_3	1.978373	1.982710	1.985506	1.987491	1.988973
	h_3/h_4	1.992116	1.994096	1.995208	1.995910	1.996405
$\ \mathbf{p} - \mathbf{p}_h\ $	h_1/h_2	1.965680	1.955971	1.957759	1.961090	1.964376
	h_2/h_3	1.992406	1.988259	1.988433	1.989229	1.990076
	h_3/h_4	1.998994	1.996913	1.996720	1.996843	1.997017

Table 5: L_2 norms of relative errors for u , \mathbf{q} , w , and \mathbf{p}

		$t = 0.2$	$t = 0.4$	$t = 0.6$	$t = 0.8$	$t = 1.0$
$\ u - u_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{2}$	1.18648e-1	1.21844e-1	1.21902e-1	1.21899e-1	1.21898e-1
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{4}$	6.13049e-2	6.27734e-2	6.28006e-2	6.27999e-2	6.27997e-2
	$h_3 = \frac{1}{32}, \Delta t_3 = \frac{1}{8}$	3.11322e-2	3.18297e-2	3.18431e-2	3.18431e-2	3.18431e-2
	$h_4 = \frac{1}{64}, \Delta t_4 = \frac{1}{16}$	1.56832e-2	1.60221e-2	1.60289e-2	1.60289e-2	1.60289e-2
$\ \mathbf{q} - \mathbf{q}_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{2}$	1.18571e-1	1.21997e-1	1.22100e-1	1.22104e-1	1.22104e-1
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{4}$	6.09759e-2	6.24793e-2	6.25171e-2	6.25181e-2	6.25181e-2
	$h_3 = \frac{1}{32}, \Delta t_3 = \frac{1}{8}$	3.09221e-2	3.16161e-2	3.16318e-2	3.16322e-2	3.16322e-2
	$h_4 = \frac{1}{64}, \Delta t_4 = \frac{1}{16}$	1.55709e-2	1.59028e-2	1.59099e-2	1.59101e-2	1.59101e-2
$\ w - w_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{2}$	1.14025e-1	1.17325e-1	1.17421e-1	1.17424e-1	1.17424e-1
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{4}$	5.81838e-2	5.96423e-2	5.96790e-2	5.96799e-2	5.96800e-2
	$h_4 = \frac{1}{32}, \Delta t_3 = \frac{1}{8}$	2.93591e-2	3.00335e-2	3.00490e-2	3.00494e-2	3.00494e-2
	$h_3 = \frac{1}{64}, \Delta t_4 = \frac{1}{16}$	1.47430e-2	1.50656e-2	1.50727e-2	1.50729e-2	1.50729e-2
$\ \mathbf{p} - \mathbf{p}_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{2}$	1.08438e-1	1.11635e-1	1.11726e-1	1.11729e-1	1.11729e-1
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{4}$	5.69473e-2	5.83682e-2	5.84031e-2	5.84040e-2	5.84040e-2
	$h_3 = \frac{1}{32}, \Delta t_3 = \frac{1}{8}$	2.91739e-2	2.98324e-2	2.98471e-2	2.98475e-2	2.98475e-2
	$h_4 = \frac{1}{64}, \Delta t_4 = \frac{1}{16}$	1.47639e-2	1.50791e-2	1.50859e-2	1.50860e-2	1.50860e-2

Table 6: Convergence orders of u , \mathbf{q} , w , and \mathbf{p}

		$t = 0.2$	$t = 0.4$	$t = 0.6$	$t = 0.8$	$t = 1.0$
$\ u - u_h\ $	h_1/h_2	0.952613	0.956810	0.956872	0.956852	0.956845
	h_2/h_3	0.977595	0.979780	0.979798	0.979781	0.979777
	h_3/h_4	0.989188	0.990310	0.990305	0.990305	0.990305
$\ \mathbf{q} - \mathbf{q}_h\ $	h_1/h_2	0.959440	0.965395	0.965740	0.965765	0.965765
	h_2/h_3	0.979601	0.982719	0.982875	0.982880	0.982880
	h_3/h_4	0.989786	0.991379	0.991451	0.991451	0.991451
$\ w - w_h\ $	h_1/h_2	0.970661	0.976103	0.976395	0.976410	0.976408
	h_2/h_3	0.986810	0.989763	0.989906	0.989909	0.989911
	h_3/h_4	0.993778	0.995314	0.995379	0.995379	0.995379
$\ \mathbf{p} - \mathbf{p}_h\ $	h_1/h_2	0.929171	0.935535	0.935848	0.935865	0.935865
	h_2/h_3	0.994949	0.968303	0.968454	0.968457	0.968457
	h_3/h_4	0.982604	0.984330	0.984390	0.984400	0.984400

Table 7: L_2 norms of relative errors for u , \mathbf{q} , w , and \mathbf{p}

		$t = 0.2$	$t = 0.4$	$t = 0.6$	$t = 0.8$	$t = 1.0$
$\ u - u_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{2}$	1.19091e-1	1.22462e-1	1.22553e-1	1.22555e-1	1.22555e-1
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{4}$	3.09925e-2	3.16823e-2	3.16971e-2	3.16973e-2	3.16973e-2
	$h_3 = \frac{1}{32}, \Delta t_3 = \frac{1}{8}$	7.82004e-3	7.98155e-3	7.98479e-3	7.98485e-3	7.98485e-3
	$h_4 = \frac{1}{64}, \Delta t_4 = \frac{1}{16}$	1.95945e-3	1.99916e-3	1.99962e-3	1.99963e-3	1.99963e-3
$\ \mathbf{q} - \mathbf{q}_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{2}$	1.18045e-1	1.21453e-1	1.21556e-1	1.21559e-1	1.21560e-1
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{4}$	3.08857e-2	3.15891e-2	3.16065e-2	3.16071e-2	3.16072e-2
	$h_3 = \frac{1}{32}, \Delta t_3 = \frac{1}{8}$	7.81346e-3	7.97852e-3	7.98240e-3	7.98254e-3	7.98255e-3
	$h_4 = \frac{1}{64}, \Delta t_4 = \frac{1}{16}$	1.96033e-3	2.00093e-3	2.00167e-3	2.00169e-3	2.00169e-3
$\ w - w_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{2}$	1.12330e-1	1.15519e-1	1.15610e-1	1.15612e-1	1.15612e-1
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{4}$	2.95040e-2	3.01662e-2	3.01811e-2	3.01814e-2	3.01814e-2
	$h_4 = \frac{1}{32}, \Delta t_3 = \frac{1}{8}$	7.44492e-3	7.59944e-3	7.60264e-3	7.60271e-3	7.60271e-3
	$h_3 = \frac{1}{64}, \Delta t_4 = \frac{1}{16}$	1.86538e-3	1.90328e-3	1.90394e-3	1.90395e-3	1.90395e-3
$\ \mathbf{p} - \mathbf{p}_h\ $	$h_1 = \frac{1}{8}, \Delta t_1 = \frac{1}{2}$	1.13293e-1	1.16536e-1	1.16629e-1	1.16631e-1	1.16631e-1
	$h_2 = \frac{1}{16}, \Delta t_2 = \frac{1}{4}$	2.94193e-2	3.00745e-2	3.00892e-2	3.00895e-2	3.00895e-2
	$h_3 = \frac{1}{32}, \Delta t_3 = \frac{1}{8}$	7.44548e-3	7.59886e-3	7.60204e-3	7.60210e-3	7.60210e-3
	$h_4 = \frac{1}{64}, \Delta t_4 = \frac{1}{16}$	1.86834e-3	1.90603e-3	1.90671e-3	1.90672e-3	1.90672e-3

Table 8: Convergence orders of u , \mathbf{q} , w , and \mathbf{p}

		$t = 0.2$	$t = 0.4$	$t = 0.6$	$t = 0.8$	$t = 1.0$
$\ u - u_h\ $	h_1/h_2	1.942073	1.950585	1.950983	1.950997	1.950997
	h_2/h_3	1.986670	1.988936	1.989024	1.989023	1.989023
	h_3/h_4	1.996727	1.997275	1.997529	1.997532	1.997532
$\ \mathbf{q} - \mathbf{q}_h\ $	h_1/h_2	1.934326	1.942899	1.943328	1.943336	1.943343
	h_2/h_3	1.982906	1.985234	1.985327	1.985329	1.985332
	h_3/h_4	1.994865	1.995450	1.995618	1.995629	1.995631
$\ w - w_h\ $	h_1/h_2	1.928761	1.937125	1.937549	1.937560	1.937560
	h_2/h_3	1.986582	1.988968	1.989073	1.989074	1.989074
	h_3/h_4	1.996787	1.997405	1.997512	1.997518	1.997518
$\ \mathbf{p} - \mathbf{p}_h\ $	h_1/h_2	1.945224	1.954163	1.954609	1.954619	1.954619
	h_2/h_3	1.982326	1.984686	1.984787	1.984790	1.984790
	h_3/h_4	1.994608	1.995212	1.995301	1.995305	1.995305

5 Conclusion and Perspective

In this article, we propose a EMFE method to solve a class of 2D Sobolev equation(1.1) by introducing three auxiliary variables. Optimal error estimates are obtained for both the semi-discrete and fully discrete schemes. Finally, two numerical examples are given to verify the optimal order of the proposed scheme. In future work, we will discuss how to extend this method to the analysis of high-dimensional nonlinear Sobolev equations.

References

- [1] G. I. Barenblatt, Iu. P. Zheltov, I. N. Kochina, Basic concepts in the theory of seepage of homogeneous liquids in fissured rocks[J], *Journal of Applied Mathematics and Mechanics*, 24(5), 1960, 1286-1303
- [2] B. D. Coleman, W. Noll, An Approximation Theorem for Functionals, with Applications in Continuum Mechanics[J], *Archive for Rational Mechanics and Analysis*, 6(1), 1960, 355-370
- [3] P. J. Chen, M. E. Gurtin, On a theory of heat conduction involving two temperatures[J], *Zeitschrift für Angewandte Mathematik und Physik ZAMP*, 19(4), 1968, 614-627
- [4] T. W. Ting, A cooling process according to two-temperature theory of heat conduction[J], *Journal of Mathematical Analysis and Applications*, 45(1), 1974, 23-31
- [5] N. Li, F. Gao, T. Zhang, An expanded mixed finite element method for Sobolev equation[J], *Journal of Computational Analysis and Applications*, 15(3), 2013, 535-543
- [6] T. Sun, A Godunov-mixed finite element method on changing meshes for the nonlinear Sobolev equations[J], *Abstract and Applied Analysis*, 2012, 2012, Article ID 413718, 19 pages
- [7] R. E. Ewing, Numerical solution of Sobolev partial differential equations[J], *SIAM Journal on Numerical Analysis*, 12, 1975, 345-363
- [8] D. Shi, Y. Zhang, High accuracy analysis of a new nonconforming mixed finite element scheme for Sobolev equations[J], *Applied Mathematics and Computation*, 218(7), 2011, 3176-3186
- [9] P. G. Ciarlet, *The Finite Element Method for Elliptic Problems*, Amsterdam, New York, 1978.
- [10] I. Babuska, Error-bounds for finite element method[J], *Numerische Mathematik*, 16(4), 1970, 322-333
- [11] F. Brezzi, On the existence, uniqueness and approximation of saddle-point problems arising from Lagrangian multipliers[J], *RAIRO Mathematical Modelling and Numerical Analysis*, 8(2), 1974, 129-151
- [12] R. S. Falk, J. E. Osborn, Error estimates for mixed methods[J], *RAIRO Analyse Numrique*, 14(3), 1980, 249-277
- [13] F. Brezzi, M. Fortin, *Mixed and Hybrid Finite Element Methods*[J], Springer Series in Computational Mathematics, Springer, Berlin, Germany, 15, 1991
- [14] J. E. Roberts, J. M. Thomas, *Mixed and hybrid methods*[J], *Handbook of Numerical Analysis*, 2, 1991, 523-639
- [15] L. Guo and H. Z. Chen, An expanded characteristic-mixed finite element method for a convection-dominated transport problem[J], *Journal of Computational Mathematics*, 23(5), 2005, 479-490
- [16] W. Liu, H. Rui, H. Guo, A two-grid method with expanded mixed element for nonlinear reaction-diffusion equations[J], *Acta Mathematicae Applicatae Sinica(English Series)*, 27(3), 2011, 495-502

- [17] H. Che, Z. Zhou, Z. Jiang, H1-Galerkin expanded mixed finite element methods for nonlinear pseudo-parabolic integrodifferential equations[J], Numerical Methods for Partial Differential Equations, 29(3), 2013, 799-817
- [18] T. Arbogast, M. F. Wheeler, I. Yotov, Mixed finite element methods for elliptic problems with tensor coefficients as cell-centered finite differences[J], SIAM Journal on Numerical Analysis, 34, 1997, 828-852
- [19] Z. Chen, Expanded mixed finite element methods for linear second-order elliptic problems(I)[J], RAIRO Mathematical Modelling and Numerical Analysis, 32(4), 1998, 479-499
- [20] Z. Chen, Expanded mixed finite element methods for quasilinear second-order elliptic problems[J], RAIRO Mathematical Modelling and Numerical Analysis, 32(4), 1998, 500-520
- [21] C. S. Woodward, C. N. Dawson, Analysis of expanded mixed finite element methods for a nonlinear parabolic equation modelling flow into variably saturated porous media[J], SIAM Journal on Numerical Analysis, 37, 2000, 701-724
- [22] Z. Chen, BDM mixed methods for a nonlinear elliptic problem[J], Computational and Applied Mathematics, 53, 1994, 207-223
- [23] R. A. Adams, Sobolev Spaces, Academic Press, New York, 1975
- [24] J. Jr. Douglas, J. E. Roberts, Global estimates for mixed methods for second order elliptic equations[J], Mathematical of computation, 44(169), 1985, 39-52
- [25] E. J. Park, Mixed finite element methods for nonlinear second-order elliptic problems[J], SIAM Journal on Numerical Analysis, 32, 1995, 865-885
- [26] P. A. Raviart, J. M. Thomas, A mixed finite element method for 2-nd order elliptic problems[J], Mathematical Aspects of the Finite Element Method, Lecture Notes in Math., Vol. 606, Springer-Verlag, Berlin, 1977, 292-315
- [27] R. Durán, Error analysis in L^p , $1 \leq p \leq \infty$, for mixed finite element methods for linear and quasilinear elliptic problems[J], Mathematical Modelling and Numerical Analysis, 22, 1988, 371-387
- [28] F. A. Milner, Mixed finite element methods for quasilinear second order elliptic problems[J], Mathematics of Computation, 44, 1985, 303-320
- [29] Y. H. Shi, D. Y. Shi, High accuracy analysis of a new mixed finite element method for Sobolev equations[J], Journal of Systems Science and Mathematical Sciences, 34(4), 2014, 452-463