



University of Dundee

Clinical correlates and outcome associated with changes in 6-Minute Walking Distance in Patients with Heart Failure

Ferreira, João Pedro; Metra, Marco; Anker, Stefan D.; Dickstein, Kenneth; Lang, Chim; Ng, Leong Loke

Published in:
European Journal of Heart Failure

DOI:
[10.1002/ejhf.1380](https://doi.org/10.1002/ejhf.1380)

Publication date:
2019

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Ferreira, J. P., Metra, M., Anker, S. D., Dickstein, K., Lang, C., Ng, L. L., Samani, N. J., Cleland, J. G. F., van Veldhuisen, D. J., Voors, A. A., & Zannad, F. (2019). Clinical correlates and outcome associated with changes in 6-Minute Walking Distance in Patients with Heart Failure: findings from the BIOSTAT-CHF study. *European Journal of Heart Failure*, 21(2), 218-226. <https://doi.org/10.1002/ejhf.1380>


General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Figure 1. Spline for the association between baseline 6MWT and the composite outcome of heart failure hospitalisation or death




Legend: 6MWT, 6-minute walking test distance; X-axis, distance in meters x 50 (1 =50m; 5 =250m;10 =500m).

Figure 2. Spline for the association between the change in 6MWT between baseline and 9 months and the subsequent risk of heart failure hospitalisation and death



Legend: 6MWT, 6-minute walking test distance; X-axis, distance in meters x 50 (5 =250m;10 =500m).

Figure 3. Forest plot showing hazard ratios for the primary outcome (heart failure hospitalisation or death) for patients subgrouped by 6MWT compared to other clinical features associated with an adverse prognosis in HF



Legend: 6MWT, 6-minute walking test; Afib, atrial fibrillation; COPD, chronic obstructive pulmonary disease; HF, heart failure; eGFR, estimated glomerular filtration rate.