



University of Dundee

Interventions for preventing oral mucositis in patients with cancer receiving treatment

Riley, Philip; Glenny, Anne Marie; Worthington, Helen V.; Littlewood, Anne; Clarkson, Jan E.; McCabe, Martin G.

Published in:
Cochrane Database of Systematic Reviews

DOI:
[10.1002/14651858.CD011552](https://doi.org/10.1002/14651858.CD011552)

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Riley, P., Glenny, A. M., Worthington, H. V., Littlewood, A., Clarkson, J. E., & McCabe, M. G. (2015). Interventions for preventing oral mucositis in patients with cancer receiving treatment: Oral cryotherapy. *Cochrane Database of Systematic Reviews*, 2015(2), 1-9. [CD011552]. <https://doi.org/10.1002/14651858.CD011552>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Interventions for preventing oral mucositis in patients with cancer receiving treatment: oral cryotherapy (Protocol)

Riley P, Glenny AM, Worthington HV, Littlewood A, Clarkson JE, McCabe MG



**THE COCHRANE
COLLABORATION®**

This is a reprint of a Cochrane protocol, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2015, Issue 2

<http://www.thecochranelibrary.com>

WILEY

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
BACKGROUND	1
OBJECTIVES	3
METHODS	3
ACKNOWLEDGEMENTS	6
REFERENCES	6
APPENDICES	7
CONTRIBUTIONS OF AUTHORS	8
DECLARATIONS OF INTEREST	8
SOURCES OF SUPPORT	8

[Intervention Protocol]

Interventions for preventing oral mucositis in patients with cancer receiving treatment: oral cryotherapy

Philip Riley¹, Anne-Marie Glenny¹, Helen V Worthington¹, Anne Littlewood¹, Jan E Clarkson², Martin G McCabe³

¹Cochrane Oral Health Group, School of Dentistry, The University of Manchester, Manchester, UK. ²Division of Oral Health Sciences, University of Dundee, Dundee, UK. ³Manchester Academic Health Science Centre, School of Cancer and Enabling Sciences, The University of Manchester, Manchester, UK

Contact address: Philip Riley, Cochrane Oral Health Group, School of Dentistry, The University of Manchester, JR Moore Building, Oxford Road, Manchester, M13 9PL, UK. philip.riley@manchester.ac.uk

Editorial group: Cochrane Oral Health Group.

Publication status and date: New, published in Issue 2, 2015.

Citation: Riley P, Glenny AM, Worthington HV, Littlewood A, Clarkson JE, McCabe MG. Interventions for preventing oral mucositis in patients with cancer receiving treatment: oral cryotherapy. *Cochrane Database of Systematic Reviews* 2015, Issue 2. Art. No.: CD011552. DOI: 10.1002/14651858.CD011552.

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

This is the protocol for a review and there is no abstract. The objectives are as follows:

To assess the effects of oral cryotherapy for preventing oral mucositis in patients with cancer who are receiving treatment.

BACKGROUND

Description of the condition

Treating cancer with chemotherapy, radiotherapy of the head and neck, or targeted therapy can cause toxic oral side effects (Al-Dasooqi 2013; Scully 2006; Sonis 2004). Perhaps the most widely researched of these side effects is oral mucositis (Al-Dasooqi 2013), which affects at least 75% of high risk patients (those receiving head and neck radiotherapy or high-dose chemotherapy) (Scully 2006). Oral mucositis may be under-reported in lower risk groups for various reasons: their tendency to be outpatients with less observation; less reporting of moderate mucositis; or patients and clinicians wishing to avoid any disruption to optimal cancer treatment (Scully 2006).

Simply put, oral mucositis affects the oral mucosa (the mucous membrane of moist tissue lining the oral cavity) and can lead to the

development of lesions (ulcers). However, the process that leads to oral mucositis is complex and multifactorial, with Sonis' five phase model being the currently accepted explanation for the sequence of events underlying the condition (Sonis 2004; Sonis 2009):

1. Initiation: DNA damage caused by chemotherapy or radiotherapy results in the loss of ability to proliferate in the basal cells of the epithelium (the external layers of cells lining the oral mucosa). This produces reactive oxygen species (ROS);
2. Primary damage response: Radiotherapy, chemotherapy, ROS, and DNA strand breaks all contribute to the activation of transcription factors such as nuclear factor kappa beta (NF- κ B), and sphingomyelinases. All this leads to the upregulation of pro-inflammatory cytokines (e.g. tumour necrosis factor alpha - TNF- α), nitric oxide, ceramide, and matrix metalloproteinases, resulting in the thinning of the epithelium through tissue injury and cell death, culminating with the destruction of the oral mucosa;

3. Signal amplification: Some of the molecules in the previous phase can lead to the exacerbation and prolonging of tissue injury through positive or negative feedback (e.g. TNF- α can positively feedback on NF- κ B thus inducing more pro-inflammatory cytokine production);

4. Ulceration: Bacteria colonise ulcers and their cell wall products infiltrate the submucosa (the connective tissues beneath the oral mucosa), activating tissue macrophages (white blood cells that respond to infection or damaged/dead cells), which results in further production of pro-inflammatory cytokines, inflammation, and pain;

5. Healing: Signalling from the extracellular matrix of the submucosa results in epithelial proliferation and differentiation, and thus a thickening of the epithelium. The local oral flora are reinstated.

Understanding of the pathobiology leading to mucosal toxicity as a result of targeted therapies (e.g. mammalian target of rapamycin (mTOR) inhibitor-associated stomatitis - mIAS) is currently limited, but it is thought to differ from chemotherapy- and radiotherapy-induced mucositis, and the clinical presentation of the ulcers is more similar to aphthous stomatitis (Al-Dasooqi 2013; Boers-Doets 2013; Peterson 2013a).

Chemotherapy-induced oral mucositis is regarded as an acute condition, with ulceration normally occurring one week after treatment, and resolving within three weeks of treatment (Sonis 2009). Radiotherapy-induced oral mucositis is chronic in nature, with ulceration normally occurring around two weeks into a seven-week treatment cycle, and resolving three to four weeks after treatment has ended (Sonis 2009).

Ulceration is the most significant phase as it leads to pain of varying severity, and difficulties with eating, swallowing, and talking (Scully 2006). This in turn leads to the consumption of pain relief medication, nutritional support (e.g. a feeding tube), treatment of the oral mucositis, specialist oral hygiene care, increased medical appointments and use of staff and resources, and, in some instances, hospitalisation (Jensen 2013; Miller 2001; Trotti 2003). Thus the negative impact on the quality of life of cancer patients, when they are already suffering, is severe (Elting 2008; Epstein 1999). Further problems can occur in immunosuppressed patients if whole bacteria on the ulcer surface cross into the underlying submucosa, potentially leading to bacteraemia and sepsis, which require antibiotics and hospitalisation, and can cause death (Jensen 2013; Peterson 2013a; Scully 2006).

Therefore, oral mucositis can be a dose-limiting condition, disrupting a patient's optimal cancer treatment plan and consequentially decreasing their chances of survival (Jensen 2013; Peterson 2013a; Sonis 2004). The additional costs associated with oral mucositis are significant, with one study reporting a median incremental cost of 18,515 US dollars per patient (Nonzee 2008).

Description of the intervention

Oral cryotherapy typically involves placing ice chips in the mouth five minutes prior to chemotherapy and continuing for 30 minutes (Lalla 2008). The ice chips are typically rounded to avoid any sharp edges or corners that may cause irritation to the patient, and also so that they can be easily moved around in the mouth (Karagöz lu 2005).

The advantages of using cryotherapy over other interventions are its availability, cost-effectiveness, ease of administration, and safety (in terms of lack of side-effects), and that it is well tolerated by patients (Peterson 2013b).

How the intervention might work

The use of ice chips in the mouth cools the oral tissues and causes the blood vessels to narrow (vasoconstriction), thus reducing blood flow to the area and therefore also restricting the amounts of the chemotherapy drugs delivered to the tissues (Lalla 2008; Peterson 2013b; Scully 2006). Cryotherapy may only be effective in the prevention of oral mucositis in patients receiving chemotherapy drugs that have a short half-life, such as bolus 5-fluorouracil (5-FU), bolus edatrexate, and high-dose melphalan (Lalla 2008; Peterson 2013b; Scully 2006). Considering the mechanism by which cryotherapy can prevent oral mucositis caused by chemotherapy, it is unclear whether or not it could have any effect on oral mucositis caused by radiotherapy (Lalla 2008). It is also unclear whether or not cryotherapy could have any role in the prevention of targeted therapy-induced stomatitis.

Why it is important to do this review

This review is the first of a series that will replace the previously published Cochrane review covering all interventions for the prevention of oral mucositis in patients with cancer receiving treatment (Worthington 2011). The Mucositis Study Group (MSG) of the Multinational Association of Supportive Care in Cancer/International Society of Oral Oncology (MASCC/ISOO) is a group that was set up in 1998 for the purpose of producing international evidence-based clinical practice guidelines for managing mucositis (both oral and gastrointestinal), which they first published in 2004, with the latest update published in 2014 (Lalla 2014). In order to facilitate the future updating of Cochrane reviews on this topic, and also to make them more usable to clinicians, guideline developers, and consumers, we have decided to divide the original Cochrane review into the same intervention categories as those used by MASCC/ISOO, which are as follows:

- Basic oral care/good clinical practice;
- Growth factors and cytokines;
- Anti-inflammatory agents;
- Antimicrobials, mucosal coating agents, anaesthetics, and analgesics;
- Laser and other light therapy;

- Cryotherapy;
- Natural and miscellaneous agents.

We believe that running in tandem with the MASCC/ISOO categories will enable the Cochrane reviews to more easily feed into such guidelines. We will also be able to be more thorough and rigorous in our assessment and summarising of the evidence in each of the categories, which was not feasible in a single Cochrane review approaching 150 included studies.

It is also important to do this review as it is consistently shown to be the most used review produced by the Cochrane Oral Health Group (in terms of full-text downloads). It was also ranked by an expert panel of oral medicine specialists as being the most important topic in the field of oral medicine in an international prioritisation exercise carried out by the Cochrane Oral Health Group in 2014 (<http://ohg.cochrane.org/priority-reviews>).

OBJECTIVES

To assess the effects of oral cryotherapy for preventing oral mucositis in patients with cancer who are receiving treatment.

METHODS

Criteria for considering studies for this review

Types of studies

We will include all randomised controlled trials (RCTs) of parallel design. It is possible to conduct cross-over studies in this area as patients may receive several chemotherapy sessions, with any mucositis completely healing in the periods between the sessions. However, we will exclude cross-over studies as we cannot discount any period effects, with mucositis risk increasing as patients receive further cycles of treatment (Scully 2006; Sonis 2009).

Types of participants

We will include all patients with cancer who are receiving treatment.

Types of interventions

We will include studies comparing oral cryotherapy for the prevention of oral mucositis (inclusive of targeted therapy-induced stomatitis) against placebo, no treatment, or any other treatment to prevent oral mucositis.

We will exclude studies assessing different cancer treatments where the primary outcome is survival/cure, with mucositis as a toxicity.

Types of outcome measures

We are in agreement with Williamson 2012 that, if clinical trials and systematic reviews are to be utilised, the outcomes assessed should be those considered important to patients, healthcare professionals, and other key stakeholders. If outcomes and outcome measures are inconsistent across studies, it will not be possible to compare and summarise research, and there is potential for outcome reporting bias, with the selective reporting of results based on statistical significance and favourability (Clarke 2007; Dwan 2008; Williamson 2005). This can lead to exaggerated estimates of effect in systematic reviews of interventions, leading to an incorrect belief that an intervention is more beneficial than it truly is (Clarke 2007). It is thought that the way to address this problem is to develop disease- or condition-specific core outcome sets to be used as a minimum when conducting and reporting clinical trials (Clarke 2007; Williamson 2012).

Therefore we will use the core outcome set produced by Bellm 2002, which is registered on the COMET (Core Outcome Measures in Effectiveness Trials) Initiative's website (www.comet-initiative.org), and is the only core outcome set for oral mucositis known to us.

Primary outcomes

Mucositis (at all levels of severity) using an appropriate, objective scale. We will use mucositis measured on a 0 to 4 point scale (none to severe) and dichotomise it as any mucositis (0 versus 1+), moderate to severe mucositis (0 to 1 versus 2+), and severe mucositis (0 to 2 versus 3+).

Some studies measure mucositis using a composite scale. If it is possible to extract the 'mucositis only' data from the total score, we will include the data in the analyses. If it is not possible, we will record the composite data in an additional table.

Secondary outcomes

- Interruptions to cancer treatment;
- Oral pain;
- Quality of life;
- Normalcy of diet (including use of percutaneous endoscopic gastrostomy (PEG) feeding tubes or total parenteral nutrition (TPN));
- Adverse events;
- Number of days in hospital;
- Number of days of treatment with opioid analgesics;
- Number of days unable to take medicine orally.

Search methods for identification of studies

For the identification of studies included or considered for this review, we will develop detailed search strategies for each database

searched. These will be based on the search strategy we have developed for MEDLINE (OVID) (Appendix 1), which we will revise appropriately for each database.

Electronic searches

We will search the following electronic databases:

- Cochrane Oral Health Group Trials Register (whole database);
- Cochrane Central Register of Controlled Trials (CENTRAL) (The Cochrane Library, current issue);
- MEDLINE via OVID (1946 to present) ([Appendix 1](#));
- EMBASE via OVID (1980 to present);
- CANCELIT via PubMed (1950 to present);
- CINAHL via EBSCO (1937 to present).

No restrictions will be placed on the language or date of publication when searching the electronic databases.

Searching other resources

We will search the following databases for ongoing trials:

- US National Institutes of Health Trials Register (<http://clinicaltrials.gov>);
- World Health Organization (WHO) International Clinical Trials Registry Platform (<http://apps.who.int/trialsearch/default.aspx>).

We will only include handsearching done as part of the Cochrane Worldwide Handsearching Programme and uploaded to CENTRAL (see the Cochrane Masterlist (<https://us.cochrane.org/master-list>) for details of journal issues searched to date).

Data collection and analysis

Selection of studies

Two review authors will independently screen the titles and abstracts retrieved from the electronic searches. We will obtain full text copies of all studies that appear to meet the inclusion criteria of the review, or where there is insufficient information in the title or abstract to make a clear judgement. Two review authors will independently assess the full text copies for eligibility and resolve any disagreements through discussion. We will consult a third review author if we cannot resolve disagreements.

On assessing the full text article, we will discard any studies that clearly do not meet the inclusion criteria. We will record all other studies that do not meet the inclusion criteria, along with reasons for exclusion, in the 'Characteristics of excluded studies' table.

Data extraction and management

Two review authors will independently extract the data from each included study using a specially designed data extraction form, which we will first pilot on a small sample of studies. We will contact study authors for clarification or missing data where necessary and feasible. We will resolve any disagreements through discussion, consulting a third review author to achieve consensus when necessary.

We will record the following data for each included study in the 'Characteristics of included studies' table.

- Trial design, location, number of centres, recruitment period;
- Inclusion/exclusion criteria, age and gender of participants, number randomised/analysed, any other potentially important prognostic factors (e.g. cancer type, cancer treatment, etc.);
- Detailed description of the intervention and comparator, including timing and duration. Information on compliance with the cryotherapy regimen;
- Details of the outcomes reported, including method of assessment and time(s) assessed;
- Details of sample size calculations, adverse effects, funding sources, declarations/conflicts of interest.

Assessment of risk of bias in included studies

Two review authors will independently assess the risk of bias of each included study using the Cochrane domain-based, two-part tool as described in Chapter 8 of the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). We will contact study authors for clarification or missing information where necessary and feasible. We will resolve any disagreements through discussion, consulting a third review author to achieve consensus when necessary.

We will complete a 'Risk of bias' table for each included study. For each domain of risk of bias, we will first describe what is reported to have happened in the study. This will provide the rationale for our judgement of whether that domain is at low, high, or unclear risk of bias.

We will assess the following domains:

1. Sequence generation (selection bias);
2. Allocation concealment (selection bias);
3. Blinding of outcome assessment (detection bias);
4. Incomplete outcome data (attrition bias);
5. Selective outcome reporting (reporting bias);
6. Other bias.

We will not include the risk of bias domain 'blinding of participants and personnel (performance bias)' as it will not be possible to blind participants or personnel. This could lead to differences in treatment for the patients (e.g. nurses could give people in the placebo group extra care), which could have an effect on the outcomes. Therefore we will consider all the studies to have a risk of performance bias.

Based on the remaining domains, we will categorise the overall risk of bias of individual studies. Studies will be categorised as being at low, high, or unclear risk of bias according to the following criteria:

- Low risk of bias (plausible bias unlikely to seriously alter the results) if all domains are at low risk of bias;
- High risk of bias (plausible bias that seriously weakens confidence in the results) if one or more domains are at high risk of bias; or
- Unclear risk of bias (plausible bias that raises some doubt about the results) if one or more domains are at unclear risk of bias.

We will also present the 'Risk of bias' summary graphically.

Measures of treatment effect

For continuous outcomes (e.g. oral pain on a visual analogue scale) where studies use the same scale, we will use the mean values and standard deviations (SDs) reported in the studies in order to express the estimate of effect as mean difference (MD) with 95% confidence interval (CI). Where different scales are used, we will express the treatment effect as standardised mean difference (SMD) with 95% CI.

For dichotomous outcomes (e.g. mucositis of any severity/no mucositis), we will express the estimate of effect as a risk ratio (RR) with 95% CI.

Unit of analysis issues

The participant will be the unit of analysis.

Dealing with missing data

Where feasible, we will attempt to contact the author(s) of included studies for clarification or missing data. We will use the methods described in Section 7.7.3 of the *Cochrane Handbook for Systematic Reviews of Interventions* to estimate missing SDs (Higgins 2011). We will not use any other statistical methods or perform any further imputation to account for missing data.

Assessment of heterogeneity

If a sufficient number of studies are included in any meta-analyses, we will assess clinical heterogeneity by examining the characteristics of the studies, the similarity between the types of participants, the interventions, and the outcomes. We will also assess heterogeneity statistically using a Chi² test, where a P value < 0.1 will indicate statistically significant heterogeneity. We will quantify heterogeneity using the I² statistic. A guide to interpretation of the I² statistic given in Section 9.5.2 of the *Cochrane Handbook for Systematic Reviews of Interventions* is as follows (Higgins 2011):

- 0% to 40%: Might not be important;
- 30% to 60%: May represent moderate heterogeneity;

- 50% to 90%: May represent substantial heterogeneity;
- 75% to 100%: Considerable heterogeneity.

Assessment of reporting biases

If at least 10 studies are included in a meta-analysis, we will assess publication bias according to the recommendations on testing for funnel plot asymmetry (Egger 1997), as described in Section 10.4 of the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). If asymmetry is identified, we will examine possible causes.

Data synthesis

We will only carry out meta-analyses where there are studies of similar comparisons reporting the same outcomes. We will combine MDs (SMDs if studies use different scales) for continuous data, and RRs for dichotomous data. Our general approach will be to use a random-effects model. With this approach, the CIs for the average intervention effect will be wider than those obtained using a fixed-effect approach, leading to a more conservative interpretation.

We will use an additional table to report the results from studies not suitable for inclusion in a meta-analysis.

Subgroup analysis and investigation of heterogeneity

Where there are sufficient studies, we intend to carry out the following subgroup analyses as mucositis incidence and severity may vary according to these factors:

- Cancer type;
- Cancer treatment;
- Age group (children versus adults).

Sensitivity analysis

If a sufficient number of studies are included in any meta-analyses, we will undertake sensitivity analyses to assess the robustness of the results by excluding studies with unclear or high overall risk of bias (whilst acknowledging that all studies are at high risk of performance bias because blinding of participants and personnel is not possible).

In meta-analyses that include several small studies and a single very large study, we will undertake a sensitivity analysis comparing the effect estimates from both random-effects and fixed-effect models. If these are different we will report on both analyses as part of the results section, and we will consider possible interpretation.

Presentation of main results

We will produce a 'Summary of findings' table for each comparison and for the main outcomes (listed below) using GRADE methods (GRADE 2004), and GRADEpro 2014 software. We will assess the

quality of the body of evidence for each comparison and outcome by considering the overall risk of bias of the included studies, the directness of the evidence, the inconsistency of the results, the precision of the estimates, and the risk of publication bias. We will categorise the quality of each body of evidence as high, moderate, low, or very low.

Main outcomes:

- Mucositis incidence;
- Interruptions to cancer treatment;
- Oral pain;
- Quality of life;

- Normalcy of diet;
- Adverse events;
- Number of days in hospital.

ACKNOWLEDGEMENTS

We would like to thank the Cochrane Oral Health Group editorial team for their help in preparing this systematic review protocol, and Professor Douglas E Peterson for his helpful feedback.

REFERENCES

Additional references

Al-Dasooqi 2013

Al-Dasooqi N, Sonis ST, Bowen JM, Bateman E, Blijlevens N, Gibson RJ, et al. Emerging evidence on the pathobiology of mucositis. *Supportive Care in Cancer* 2013;**21**(11):3233–41.

Bellm 2002

Bellm LA, Cunningham G, Durnell L, Eilers J, Epstein JB, Fleming T, et al. Defining clinically meaningful outcomes in the evaluation of new treatments for oral mucositis: oral mucositis patient provider advisory board. *Cancer Investigation* 2002;**20**(5-6):793–800.

Boers-Doets 2013

Boers-Doets CB, Raber-Durlacher JE, Treister NS, Epstein JB, Arends AB, Wiersma DR, et al. Mammalian target of rapamycin inhibitor-associated stomatitis. *Future Oncology* 2013;**9**(12):1883–92.

Clarke 2007

Clarke M. Standardising outcomes for clinical trials and systematic reviews. *Trials* 2007;**8**:39.

Dwan 2008

Dwan K, Altman DG, Arnaiz JA, Bloom J, Chan AW, Cronin E, et al. Systematic review of the empirical evidence of study publication bias and outcome reporting bias. *PLoS One* 2008;**3**(8):e3081.

Egger 1997

Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;**315**(7109):629–34.

Elting 2008

Elting LS, Keefe DM, Sonis ST, Garden AS, Spijkervet FK, Barasch A, et al. Patient-reported measurements of oral mucositis in head and neck cancer patients treated with radiotherapy with or without chemotherapy: demonstration of increased frequency, severity, resistance to palliation, and impact on quality of life. *Cancer* 2008;**113**(10):2704–13.

Epstein 1999

Epstein JB, Emerton S, Kolbinson DA, Le ND, Phillips N, Stevenson-Moore P, et al. Quality of life and oral function

following radiotherapy for head and neck cancer. *Head & Neck* 1999;**21**(1):1–11.

GRADE 2004

GRADE Working Group. Grading quality of evidence and strength of recommendations. *BMJ* 2004;**328**(7454):1490.

GRADEpro 2014

McMaster University. GRADEpro. McMaster University, 2014.

Higgins 2011

Higgins JPT, Green S (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Jensen 2013

Jensen SB, Peterson DE. Oral mucosal injury caused by cancer therapies: current management and new frontiers in research. *Journal of Oral Pathology & Medicine* 2014;**43**(2):81–90.

Karagözoğ lu 2005

Karagözoğ lu S, Filiz Ulusoy M. Chemotherapy: the effect of oral cryotherapy on the development of mucositis. *Journal of Clinical Nursing* 2005;**14**(6):754–65.

Lalla 2008

Lalla RV, Sonis ST, Peterson DE. Management of oral mucositis in patients who have cancer. *Dental Clinics of North America* 2008;**52**(1):61–77.

Lalla 2014

Lalla RV, Bowen J, Barasch A, Elting L, Epstein J, Keefe DM, et al. MASCC/ISOO clinical practice guidelines for the management of mucositis secondary to cancer therapy. *Cancer* 2014;**120**(10):1453–61.

Miller 2001

Miller M, Kearney N. Oral care for patients with cancer: a review of the literature. *Cancer Nursing* 2001;**24**(4):241–54.

Nonzee 2008

Nonzee NJ, Dandade NA, Patel U, Markossian T, Agulnik M, Argiris A, et al. Evaluating the supportive care costs

of severe radiochemotherapy-induced mucositis and pharyngitis. *Cancer* 2008;**113**(6):1446–52.

Peterson 2013a

Peterson D, Srivastava R, Lalla R. Oral mucosal injury in oncology patients: perspectives on maturation of a field. *Oral Diseases* 2013 Jul 19 [Epub ahead of print].

Peterson 2013b

Peterson DE, Ohrn K, Bowen J, Flidner M, Lees J, Loprinzi C, et al. Systematic review of oral cryotherapy for management of oral mucositis caused by cancer therapy. *Supportive Care in Cancer* 2013;**21**(1):327–32.

Scully 2006

Scully C, Sonis S, Diz PD. Oral mucositis. *Oral Diseases* 2006;**12**(3):229–41.

Sonis 2004

Sonis ST, Elting LS, Keefe D, Peterson DE, Schubert M, Hauer-Jensen M, et al. Perspectives on cancer therapy-induced mucosal injury: pathogenesis, measurement, epidemiology, and consequences for patients. *Cancer* 2004;**100**(9 Suppl):1995–2025.

Sonis 2009

Sonis ST. Mucositis: The impact, biology and therapeutic opportunities of oral mucositis. *Oral Oncology* 2009;**45**(12):1015–20.

Trotti 2003

Trotti A, Bellm LA, Epstein JB, Frame D, Fuchs HJ, Gwede CK, et al. Mucositis incidence, severity and associated outcomes in patients with head and neck cancer receiving radiotherapy with or without chemotherapy: a systematic literature review. *Radiotherapy and Oncology* 2003;**66**(3):253–62.

Williamson 2005

Williamson PR, Gamble C, Altman DG, Hutton JL. Outcome selection bias in meta-analysis. *Statistical Methods in Medical Research* 2005;**14**(5):515–24.

Williamson 2012

Williamson PR, Altman DG, Blazeby JM, Clarke M, Devane D, Gargon E, et al. Developing core outcome sets for clinical trials: issues to consider. *Trials* 2012;**13**:132.

References to other published versions of this review

Worthington 2011

Worthington HV, Clarkson JE, Bryan G, Furness S, Glennly AM, Littlewood A, et al. Interventions for preventing oral mucositis for patients with cancer receiving treatment. *Cochrane Database of Systematic Reviews* 2011, Issue 4. [DOI: 10.1002/14651858.CD000978.pub5]

* Indicates the major publication for the study

APPENDICES

Appendix I. MEDLINE (OVID) search strategy

1. exp NEOPLASMS/
2. exp LEUKEMIA/
3. exp LYMPHOMA/
4. exp RADIOTHERAPY/
5. exp Antineoplastic agents/
6. Bone Marrow Transplantation/
7. neoplasm\$.mp.
8. cancer\$.mp.
9. (leukaemi\$ or leukemi\$).mp.
10. (tumour\$ or tumor\$).mp.
11. malignan\$.mp.
12. neutropeni\$.mp.
13. carcino\$.mp.
14. adenocarcinoma\$.mp.
15. lymphoma\$.mp.
16. (radioth\$ or radiat\$ or irradiat\$).mp.
17. (bone adj marrow adj5 transplant\$).mp.
18. chemo\$.mp.
19. or/1-18
20. exp STOMATITIS/

21. Candidiasis, Oral/
22. stomatitis.mp.
23. mucositis.mp.
24. (oral adj6 mucos\$).mp.
25. (mycosis or mycotic).mp.
26. mIAS.ti,ab.
27. or/20-26
28. Cryotherapy/
29. cryotherap\$.mp.
30. (cold or freez\$ or ice).mp.
31. or/28-30
32. 19 and 27 and 31

CONTRIBUTIONS OF AUTHORS

Philip Riley: writing the Background and Methods sections.

Anne-Marie Glenny: writing the Methods section.

Helen V Worthington: writing the Methods section.

Anne Littlewood: writing the Methods section.

Jan E Clarkson: providing a clinical perspective.

Martin G McCabe: providing a clinical perspective.

DECLARATIONS OF INTEREST

Philip Riley: none known.

Anne-Marie Glenny: none known.

Helen V Worthington: none known.

Anne Littlewood: none known.

Jan E Clarkson: none known.

Martin G McCabe: none known.

SOURCES OF SUPPORT

Internal sources

- School of Dentistry, The University of Manchester, UK.
- Manchester Academic Health Sciences Centre (MAHSC) and the NIHR Manchester Biomedical Research Centre, UK.

External sources

- National Institute for Health Research (NIHR), UK.

The NIHR is the largest single funder of the Cochrane Oral Health Group.

Disclaimer:

The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the NIHR, NHS or the Department of Health.

- Cochrane Oral Health Group Global Alliance, Other.

The production of all our reviews is assisted by funding from our Global Alliance partners (<http://ohg.cochrane.org/>): British Association for the Study of Community Dentistry, UK; British Association of Oral Surgeons, UK; British Orthodontic Society, UK; British Society of Paediatric Dentistry, UK; British Society of Periodontology, UK; Canadian Dental Hygienists Association, Canada; Mayo Clinic, USA; National Center for Dental Hygiene Research & Practice, USA; New York University College of Dentistry, USA; and Royal College of Surgeons of Edinburgh, UK