

University of Dundee

Discovery of super soft-drug modulators of sphingosine-1-phosphate receptor 1

Bell, Mark; Foley, David; Naylor, Claire; Robinson, Colin; Riley, Jennifer; Epemolu, Rafiu

Published in:
Bioorganic & Medicinal Chemistry Letters

DOI:
[10.1016/j.bmcl.2018.07.044](https://doi.org/10.1016/j.bmcl.2018.07.044)

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Bell, M., Foley, D., Naylor, C., Robinson, C., Riley, J., Epemolu, R., ... Woodland, C. (2018). Discovery of super soft-drug modulators of sphingosine-1-phosphate receptor 1. *Bioorganic & Medicinal Chemistry Letters*, 28(19), 3255-3259. <https://doi.org/10.1016/j.bmcl.2018.07.044>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Discovery of super soft-drug modulators of sphingosine-1-phosphate receptor 1

Mark Bell^{a,*}, David Foley^a, Claire Naylor^a, Colin Robinson^a, Jennifer Riley^a, Ola Epemolu^a, Paul Scullion^a, Yoko Shishikura^a, Elad Katz^a, W.H. Irwin McLean^b, Paul Wyatt^a, Kevin D. Read^a, Andrew Woodland^{a,*}

^a The Drug Discovery Unit, Biological Chemistry and Drug Discovery, University of Dundee, Dundee DD1 4HN, UK

^b Dermatology and Genetic Medicine, Biological Chemistry and Drug Discovery, University of Dundee, Dundee DD1 5EH, UK

ARTICLE INFO

Keywords:

S1PR1
Soft-drug
Plasma stability
Psoriasis
Topical

ABSTRACT

The oral S1PR1 agonist ponesimod demonstrated substantial efficacy in a phase II clinical trial of psoriasis. Unfortunately, systemic side effects were observed, which included lymphopenia and transient bradycardia. We sought to develop a topical soft-drug S1PR1 agonist with an improved therapeutic index. By modifying ponesimod, we discovered an ester series of S1PR agonists. To increase metabolic instability in plasma we synthesised esters described as specific substrates for paraoxonase and butyrylcholinesterases, esterases present in human plasma.

Introduction

Psoriasis is a common chronic inflammatory skin disease that affects 2% of the population.¹ 52.3% of patients were dissatisfied with current treatments in a recent survey from the National Psoriasis Foundation in the US.¹ Recently approved biological drugs targeting disease relevant receptors, such as secukinumab² or ixekizumab³ for interleukin (IL)-17 and ustekinumab for IL-12/23⁴ have brought great benefit to patients with severe symptoms of the disease. However, a need remains for safe, convenient, efficacious therapies for mild and moderate psoriasis.

Sphingosine-1-phosphate receptor (S1PR) agonists are of interest to the pharmaceutical industry, due to their potential to treat diseases of the immune system such as psoriasis and multiple sclerosis as well as cancer.^{5,6} S1PR agonists, such as fingolimod and ponesimod (Fig. 1), initially activate sphingosine-1-phosphate receptors, but subsequently trigger receptor internalisation. This shuts down the sphingosine 1-phosphate signalling pathway, which then prevents the maturation and migration of lymphocytes.⁷ In 2010 fingolimod was approved for the treatment of relapsing/remitting multiple sclerosis and is the only S1PR1 agonist approved to date.⁸

In a phase 2 study in plaque psoriasis, a 40 mg oral dose of ponesimod, led to a 75% reduction in psoriasis area and severity index (PASI) score in 77% of the patients.⁹ This level of efficacy is competitive with other small molecules in development to treat psoriasis (apremilast¹⁰, tofacitinib¹¹ and sotrastaurin¹²). Ponesimod was not however

progressed into phase 3, possibly due to safety concerns. Oral use of pan-S1PR and S1PR1 selective modulators is associated with several severe side effects such as lymphopenia, bradycardia and dyspnoea, which limits their utility as therapies for non-life threatening chronic diseases.⁵

In mice, topical application of the S1PR agonist fingolimod led to a reduction in the number of Langerhans cells migrating to the lymph node.¹³ Work by Griffiths and co-workers has shown that Langerhans cell migration is higher in involved areas of patient's skin than in non-involved areas of skin.^{14,15} We hypothesised that local inhibition of Langerhans cell migration, through the effects of topical S1PR agonists, could restore a normal skin phenotype to patients with psoriasis.

Due to the potential of S1PR agonists to act as efficacious therapies, for the treatment of psoriasis, we decided to embark on developing a topical S1PR1 modulator using a fast-follower approach, inspired by ponesimod. Topical therapies apply drugs directly to the site of action and can lead to a reduction in the systemic exposure of a drug when compared to oral dosing. However, due to the chronic nature of psoriasis and the need to be able to treat large body surface areas (> 10%), there is a danger that topical application could lead to biologically relevant systemic drug concentrations. To maximise the therapeutic index we decided to develop a soft drug.^{16,17} Soft drugs are locally active, in this case in the skin, but then undergo rapid systemic metabolism, to metabolites, which are either inactive or rapidly cleared from the systemic circulation.¹⁸

* Corresponding authors.

E-mail addresses: m.u.bell@dundee.ac.uk (M. Bell), awoodland@dundee.ac.uk (A. Woodland).

<https://doi.org/10.1016/j.bmcl.2018.07.044>

Received 4 May 2018; Received in revised form 19 July 2018; Accepted 29 July 2018

Available online 30 July 2018

0960-894X/ © 2018 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).




Fig. 1. Selected S1PR1 modulators.

Esters can be metabolised in liver by esterases and cytochrome P450 enzymes.¹⁹ However, most tissues are capable of metabolising an ester. Importantly, human blood contains a wide range of esterases and the rate of ester hydrolysis in blood can lead to ultra-rapid clearance,²⁰ what we term a super-soft drug. Because S1PR agonists induce transient bradycardia, and as blood returning to circulation from skin passes through the heart before reaching the liver, we felt that a very rapid blood clearance mechanism could potentially discharge the cardiotoxicity risk. To select compounds which would be likely to demonstrate the desired pharmacokinetic profile, we targeted a human plasma stability half-life of < 5 min. We chose to first develop benzoate ester analogues of ponesimod to determine if they could demonstrate potential as super-soft drugs. The desired compound would have S1PR pXC₅₀ of > 7 and a logP in the range of 2–4 which is believed to be optimal for skin penetration.²¹

Compounds **4a** and **5a–d** were synthesised as shown in Scheme 1. The (*Z*)-2-(propylimino)-3-(*o*-tolyl)thiazolidin-4-one core **2** was synthesised utilising a one-pot, two step reaction. Only the *Z* isomer was observed. *N*-propylamine was reacted with 1-isothiocyanato-2-methylbenzene **1** to give the resulting thiourea, which was condensed with 2-bromoacetyl bromide followed by addition of pyridine to furnish the desired thiazolidin-4-one **2**. This core was then condensed with the appropriately substituted benzaldehyde **3a–b** to furnish **4a** and **5a**. Finally, esterification or amidation of **4a** gave the desired products. Based on the configurational analysis by X-ray, which characterised crystalline analogues of ponesimod as *Z,Z*²², all structurally related compounds **4a**, **5a–d**, **11**, **12a–f** and **15a, b** were assigned to the *Z,Z*-isomer.

We selected human plasma stability as a model for human blood metabolism. Esters **5a–c** and amide **5d** demonstrated half-lives of > 180 min in human plasma, far greater than the < 5 min we were aiming for. S1PR1 activity was measured using a PathHunter β-Arrestin recruitment assay.²³ All of the ester and amide compounds in Table 1 were active and had pIC₅₀ values ranging from 6.5 to 7.0, with the exception of compound **5a** where a chlorine ortho to the ester provided 100-fold improvement in activity compared to **5b**. Gratifyingly, the parent carboxylic acid **4a**, which is the desired metabolite of our esters/amides was less active, giving a pIC₅₀ of < 6 and meeting our targeted criteria for producing inactive metabolites.

Aqueous solubility is an important parameter for topically applied drugs. Increasing aqueous solubility, should directly increase skin penetration rates.²⁴ Increasing aqueous solubility also increases the range of formulation options that can be used in pre-clinical development. We

Table 1
Effect of R² substitution on S1PR1 activity, kinetic aqueous solubility and human plasma stability.

Compound	R ¹	R ²	Kinetic solubility (μM) ^a	CHI logD ^b	S1PR1 pIC ₅₀ ^c	H Plasma Stability (half-life min) ^d
Ponesimod	–	–	110	3.1	7.9 (0)	> 180
5a	Cl	OMe	39	> 4.4	8.6 (0.1)	> 180
5b	H	OMe	28	> 4.4	6.5 (0)	> 180
4a	H	OH	163	1.0	< 6 (0)	–
5c	H	OCH ₂ CH ₂ OH	110	3.4	6.9 (0)	> 180
5d	H	NHCH ₂ CH ₂ OH	> 250	2.3	7.0 (0)	> 180

^a The aqueous solubility of the test compounds was measured using laser nephelometry.

^b Reverse-phase HPLC method to determine the chromatographic hydrophobicity index (CHI).

^c All pIC₅₀s reported in this table correspond to n of 2, reported as their geometric mean. The range of pIC₅₀ values is provided in brackets.


^d Incubated in human plasma at 37 °C.

were pleased to see that introduction of a hydrogen bond donor at R² in **5c** led to a 4-fold increase in solubility relative to **5b**. Changing the ester linker in **5c** to an amide in **5d** further improved solubility to > 250 μM, probably due to improvements in lipophilicity (3.4 vs 2.3 respectively).

Encouraged by the solubility and activity data, we focused our efforts on optimising the metabolic profile of the esters. At this time, we identified that switching from the *o*-tolyl group used in ponesimod to an *o*-phenol **6** increased aqueous solubility 2-fold, presumably due to the reduction in lipophilicity (Fig. 2). As **6** was equipotent to ponesimod (pIC₅₀ 7.9) and to maximise compound solubility, subsequent compounds incorporated the *o*-phenol group.

Compounds **11**, **12a–c**, **15a** and **15b** were synthesised using the route shown in Scheme 2. **12a–c** were more soluble than ponesimod, probably due to significant improvements in CHI logD (Table 2). As expected a clear trend can be seen between low lipophilicity and improved solubility. Homologation of the ester group (**15a, b**) showed increased lipophilicity (3.2 and 3.5 respectively) and a commensurate reduction in solubility. However, as with **5a–d**, amide **12a** and esters **12b–c** also showed a lack of metabolism in our human plasma stability assay (Table 2).

As simple esters had shown themselves to be stable to human



Scheme 1. Reagents and conditions for preparation of compounds **4a** and **5a–d**: (a) *n*-PrNH₂ (1 eq), CH₂Cl₂, rt (b) 2-bromoacetyl bromide (1 eq) and pyridine (2 eq), CH₂Cl₂, 0 °C to rt (two steps) (c) aldehyde (1 eq), NaOAc (2 eq), AcOH, 65 °C (d) alcohol (5 eq), HOAt (1.5 eq), EDC (1.5 eq), DIPEA (2.5 eq), CH₂Cl₂, rt. eq: equivalent; rt: room temperature; HOAt: 1-Hydroxy-7-azabenzotriazole; EDC: 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide; DIPEA: *N,N*-Diisopropylethylamine.




Fig. 2. Improvements in aqueous kinetic solubility from introduction of *o*-phenol group.


plasma, we turned our attention to ester functional groups which had been shown to be unstable in human plasma in the literature (half-life < 5min).²⁰ The most successful studies focused on optimising metabolism by the esterases paraoxonase²⁵ and butyrylcholinesterases²⁶, which are mainly expressed in blood plasma.

We synthesised compounds **11** and **12d–f** using the route shown in Scheme 2, which all contain ester groups that are unstable in human plasma when attached to other scaffolds. Although **12d** or **12e** did not show sufficient clearance in human plasma we were pleased to identify compound **12f** containing a (2-oxo-1,3-dioxolan-4-yl)methyl ester which is rapidly degraded in human plasma (Table 3). The half-life of this ester in human plasma was 9 min and metabolite identification confirmed the conversion to the inactive parent acid **11** (Fig. 3).

Unfortunately **12f** was also rapidly metabolised in human skin S9 fraction with a half-life of just 23 min, far faster than the desirable half-life of > 6 h that we target for once daily dosing. In fact, there are suggestions of a correlation between the relative stability of **12d–f** in human plasma and skin S9 fraction. It may be that the esterases responsible for the metabolism of **12f** are present in both human plasma and skin S9 fraction or that skin and plasma esterases have similar

Table 2

Effect of R¹ substitution on kinetic aqueous solubility and human plasma stability.



Compound	n	R ¹	Kinetic solubility (μM) ^a	CHI logD ^b	H Plasma Stability (half-life min) ^c
Ponesimod	–	–	110	3.1	> 180
12a	0	NHCH ₂ CONMe ₂	219	1.9	> 180
12b	0	OCH ₂ CH ₂ CH ₂ OH	183	2.3	> 180
12c	0	OCH ₂ CH(±CH ₂ OH)OH	> 250	1.8	> 180
15a	1	OMe	155	3.2	> 180
15b	2	OMe	79	3.5	> 180


^a The aqueous solubility of the test compounds was measured using laser nephelometry.

^b Reverse-phase HPLC method to determine the chromatographic hydrophobicity index (CHI).

^c Incubated in human plasma at 37 °C.


substrate affinities. Unfortunately, **12f** was also inactive as a S1PR1 modulator (Table 3).




In summary, using a fast follower approach, we identified a series of ester S1PR1 modulators. Further optimisation led to improved solubility and identification of ester **12d** that is plasma unstable.



Scheme 2. Reagents and conditions for preparation of compounds **11**, **12a–f** and **15a**, b: (a) *n*-PrNH₂ (1 eq), CH₂Cl₂, rt (b) 2-bromoacetyl bromide (1 eq) and pyridine (2 eq), CH₂Cl₂, 0 °C to rt (two steps) (c) aldehyde (1 eq), NaOAc (2 eq), AcOH, 85 °C (d) BBr₃ (8 eq), CH₂Cl₂, –78 °C to rt (e) For **12a–c**, amine (3 eq) or alcohol (5 eq), HOAt (1.5 eq), EDC (1.5 eq), DIPEA (2.5 eq), CH₂Cl₂, rt (f) For **12d–f**, alcohol, alkyl chloride or alkyl tosylate (1.1–1.5 eq) K₂CO₃ (1.5 eq), NaI (0.1 eq), DMF, 70 °C (g) NEt₃ (3 eq), EDC (1.1 eq), HOAt (1.1 eq), alcohol (1.5 eq) rt (h) BBr₃ (5 eq), CH₂Cl₂, –78 °C to rt. eq: equivalent; rt: room temperature; HOAt: 1-Hydroxy-7-azabenzotriazole; EDC: 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide; DIPEA: *N,N*-Diisopropylethylamine; DMF: dimethyl formamide.

Table 3
Effect of R¹ substitution on kinetic aqueous solubility and human plasma stability.



Compound	R ¹	H Plasma Stability (half-life min) ^a	Kinetic solubility (μM) ^b	CHI logD ^c	H Skin S9 (half-life min) ^d	S1PR1 pIC ₅₀ ^e
11	OH	> 180	–	–	–	< 6 (0)
12d		180	117	3.0	152	–
12e		61	79	2.8	40	–
12f		9	79	3.3	23	< 6 (0)

^a Incubated in human plasma at 37 °C.

^b The aqueous solubility of the test compounds was measured using laser nephelometry.

^c Reverse-phase HPLC method to determine the chromatographic hydrophobicity index (CHI).

^d Stability measured in skin S9 over 180 mins in the presence of enzymatic cofactors.

^e All pIC₅₀s reported in this table correspond to n of 2, reported as their geometric mean. The range of pIC₅₀ values is provided in brackets.




Fig. 3. Profile of compound 12f being hydrolysed to 11 in human plasma. 12f was incubated in human plasma at 37 °C. The experiment was conducted as an n of one to provide a qualitative analysis.

Unfortunately, the ester is inactive and is metabolised in human skin models. Further efforts will focus on identifying an active, skin stable compound that is rapidly metabolised systemically will be disclosed in due course.

Acknowledgement

The Centre for Dermatology and Genetic Medicine is supported by a Wellcome Trust Strategic Award (reference 098439/Z/12/Z).

References

- Parisi R, Symmons DP, Griffiths CE, Ashcroft DM. *J Invest Dermatol.* 2013;133:377–385.
- Langley RG, Elewski BE, Lebwohl M, et al. *N Engl J Med.* 2014;371:326–338.
- Gordon KB, Blauvelt A, Papp KA, et al. *Engl J Med.* 2016;375:345–356.
- Kimball AB, Gordon KB, Fakharzadeh S, et al. *Br J Dermatol.* 2012;166:861–872.
- Kunkel GT, Maceyka M, Milstien S, Spiegel S. *Nat Rev Drug Discov.* 2013;12:688–702.
- Huang WC, Nagahashi M, Terracina KP, Takabe K. *Biomolecules.* 2013;3.
- Blaho VA, Hla T. *J Lipid Res.* 2014;55:1596–1608.
- Kappos L, Radue EW, O'Connor P, et al. *N Engl J Med.* 2010;362:387–401.
- Vaclavkova A, Chimenti S, Arenberger P, et al. *Lancet.* 2014;384:2036–2045.
- Papp K, Kristian Reich K, Leonardi CL, et al. *J Am Acad Dermatol.* 2015;73:37–49.
- Ports WC, Khan S, Lan S, et al. *Br J Dermatol.* 2013;169:137–145.
- Skvara H, Dawid M, Kleyn E, et al. *J Clin Invest.* 2008;118:3151–3159.
- Reines I, Kietzmann M, Mischke R, et al. *J Invest Dermatol.* 2009;129:1954–1962.
- Griffiths CE, Dearman RJ, Cumberbatch M, Kimber I. *Cytokine.* 2005;32:67–70.
- Shaw FL, Mellody KT, Ogden S, Dearman RJ, Kimber I, Griffiths CE. *J Invest Dermatol.* 2014;134:268–271.
- Bhardwaj YR, Pareek A, Jain V, Kishore D. *Saudi Pharm J.* 2014;22:290–302.
- Bodor N, Buchwald P. *Pure Appl Chem.* 2008;80.
- Felding J, Sorensen MD, Poulsen TD, et al. *J Med Chem.* 2014;57:5893–5903.

19. Roberts JK, Moore CD, Ward RM, Yost GS, Reilly CA. *J Pharmacol Exp Ther.* 2013;345:308–316.
20. Beaumont K, Webster R, Gardner I, Dack K. *Curr Drug Metab.* 2003;4:461–485.
21. Singh P, Roberts MS. *J Pharmacol Exp Ther.* 1994;268:144–151 Skin permeability and local tissue concentrations of nonsteroidal anti-inflammatory drugs after topical application.
22. Bolli MH, Abele S, Binkert C, et al. *J Med Chem.* 2010;53:4198–4211.
23. The PathHunter® CHO-K1 EDG1 β -Arrestin Cell Line (Discoverx) was used to assess agonist activity of the compounds. Briefly, compound at 5 \times desired concentration was added to cells and incubated at 37 °C for 90 min. Assay signal was generated through a single addition of PathHunter Detection reagent cocktail, followed by a one hour incubation at room temperature. Microplates were read following signal generation with a PerkinElmer Envision instrument for chemiluminescent signal detection. Percentage activity was calculated using the following formula: % Activity = 100% \times (mean RLU of test sample – mean RLU of vehicle control)/ (mean MAX control ligand – mean RLU of vehicle control).
24. Mitravotri S, Anissimov YG, Bunge AL, et al. *Int J Pharm.* 2011;418:115–129.
25. Biggadike K, Angell RM, Burgess CM, et al. *J Med Chem.* 2000;43:19–21.
26. Mork N, Bundgaard H. *Pharm Res.* 1992;9:492–496.