



University of Dundee

Mouth Cancer Awareness in General Population

Kawecki, Michal M.; Nedeva, Iva R.; Iloya, Jonathan; Macfarlane, Tatiana

Published in:
Journal of Oral and Maxillofacial Research

DOI:
[10.5037/jomr.2019.10203](https://doi.org/10.5037/jomr.2019.10203)

Publication date:
2019

Licence:
CC BY

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Kawecki, M. M., Nedeva, I. R., Iloya, J., & Macfarlane, T. (2019). Mouth Cancer Awareness in General Population: Results from Grampian Region of Scotland, United Kingdom. *Journal of Oral and Maxillofacial Research*, 10(2), 1-11. [e3]. <https://doi.org/10.5037/jomr.2019.10203>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Mouth Cancer Awareness in General Population: Results from Grampian Region of Scotland, United Kingdom

Michal M. Kawecki¹, Iva R. Nedeva², Jonathan Iloya³, Tatiana V. Macfarlane^{4,5}

¹NHS Wales, Dental Surgery Department, Haverfordwest, United Kingdom.

²University of Ulster, School of Biomedical Sciences, Coleraine, United Kingdom.

³NHS Grampian, Dental Public Health, Aberdeen, United Kingdom.

⁴University of Aberdeen, Epidemiology Group, Aberdeen, United Kingdom.

⁵University of Dundee, Medicines Monitoring Unit (MEMO), Dundee, United Kingdom.

Corresponding Author:

Michal Kawecki

NHS Wales, Hywel Dda University Health Board

Withybush Hospital, Dental Surgery Department

Haverfordwest, SA61 2PZ

United Kingdom

Phone:

Fax:

E-mail: michal.kawecki.07@aberdeen.ac.uk

ABSTRACT

Objectives: The purpose of this project was to determine the level of mouth cancer awareness and to investigate the associated factors in a United Kingdom (UK) general population sample.

Material and Methods: Adult Dental Health Survey (2010) was conducted in a sample of 3,353 adult residents in the Grampian region of the UK (adjusted participation rate 58%). Participants completed a questionnaire consisting of questions on oral health, health behaviour, quality of life and cancer awareness.

Results: Overall, 81% of participants were aware of mouth cancer. This was associated with younger age, higher levels of education and better general health. Current smokers and alcohol drinkers were more aware of mouth cancer. When asked about risk factors for mouth cancer, the following were identified by the respondents: smoking (84%), poor oral hygiene (60%), drinking alcohol heavily (59%), poor diet (37%), stress (15%), being overweight (6%), drinking hot liquids (5%), eating spicy food (3%), using mouthwash (2%) and kissing someone (1%). Smokers were more likely to identify smoking as a risk factor for mouth cancer. Similarly, those who consumed alcohol almost daily were more likely to identify heavy alcohol drinking as a risk factor.

Conclusions: Awareness of mouth cancer is high in respondents from the general population, and participants were able to identify the most important risk factors. Knowledge of tobacco and alcohol as risk factors was highest amongst those exposed to them. The study proposed that the prevention strategies should focus not only on increasing knowledge, but also on changing health behaviour.

Keywords: alcohol drinking; awareness; mouth cancer; risk factors; tobacco.

Accepted for publication: 25 June 2019

To cite this article:

Kawecki MM, Nedeva IR, Iloya J, Macfarlane TV.

Mouth Cancer Awareness in General Population: Results from Grampian Region of Scotland, United Kingdom

J Oral Maxillofac Res 2019;10(2):e3

URL: <http://www.ejomr.org/JOMR/archives/2019/2/e3/v10n2e3.pdf>

doi: [10.5037/jomr.2019.10203](https://doi.org/10.5037/jomr.2019.10203)

INTRODUCTION

Head and neck (HNC) cancers is a collective term which encompasses malignancies of oral cavity, pharynx, larynx, sinuses and salivary glands. Oral and pharyngeal cancer (OPC) is the 8th most common malignancy in Europe and the 11th cause for cancer mortality [1]. It has a heterogeneous distribution and projected incidence of 121,300 new cases in the continent in 2018 [2]. Estimated numbers of deaths from OPC in Europe in 2018 is 41,400 in males and 11,800 in females [2]. In the United Kingdom (UK), there were 12,061 new HNC cases in 2015 and 4,047 related deaths in 2016 [3,4]. Scotland, with 1,289 cases, exhibits the highest HNC incidence rate in the UK at 24.6 (European age-standardised/100,000 population). Values for Northern Ireland, Wales and England have been estimated to be 22.5, 21.9 and 19.2, respectively [3].

Major risk factors for OPC cancer are tobacco consumption, heavy alcohol drinking and poor nutrition - specifically lower fruit and vegetable consumption [5-7]. Human papilloma virus (HPV) infection, poor oral hygiene and dentition, genetic factors and alcohol in mouthwashes have also been suggested to play a causal role [8-11].

To the best of our knowledge there are only two large studies conducted in the UK which have investigated the public awareness of OC and the associated risk factors, and these have yielded conflicting results. While one of the surveys conducted in 1995 by Warnakulasuriya et al. [10] reported an alarming lack of OC awareness with only 56% of the participants being aware, the later survey conducted in 2001 by West et al. [12] reported that 96% of respondents were aware of mouth cancer. Furthermore, the majority (85%) of respondents recognised smoking as a risk factor, however only 19% recognised frequent alcohol consumption as such [12].

The purpose of this project was to determine the level of mouth cancer awareness in a general population sample in Scotland.

MATERIAL AND METHODS

The Grampian Adult Dental Health Survey (GADHS) was conducted in the Grampian region of Scotland between October 2009 and January 2010, and aimed to evaluate current state of oral health, access to dental care and a range of related behaviours [13]. There was a delay in publishing these results due to

study investigators retiring or relocating.

This epidemiological survey was approved by the Grampian National Health Service (NHS) Board. An age-, gender- and area-stratified random sample of 6,000 people aged 25 years and over was selected from the Community Health Index (CHI) database. The CHI database contains details of all Scottish residents registered with a General Medical Practitioner. NHS Grampian had previously undertaken a youth and young people lifestyle survey [14], and therefore 16 to 24-year olds were not sampled.

Each selected individual received a postal questionnaire which consisted of 51 questions on socio-demographic characteristics (age, gender, and education), oral health, health-related behaviours and utilisation of dental services. A follow-up of non-responders was performed initially with a reminder postcard, a further questionnaire, and finally where necessary, a short questionnaire. The short questionnaire consisted of 10 key questions on dental services utilisation and number of teeth. Deprivation was measured using the Scottish Index of Multiple Deprivation (SIMD) 2009 (<http://www.scotland.gov.uk/Topics/Statistics/SIMD/>), which was obtained from the participants' postcodes. The SIMD gives a relative measure of deprivation by providing a relative ranking, with most and least deprived ranked as "1" and "5", respectively.

Participants were asked whether they had heard of eight types of cancer (lung, skin, breast, mouth, cervical, prostate, bowel and throat) and to identify which factors might cause mouth cancer. They were given 10 options: drinking alcohol heavily, poor diet, poor dental hygiene, using mouthwash, being overweight, stress, smoking, drinking hot liquids, kissing someone and eating spicy food.

Information pertaining to the participants' socio-demographic characteristics, general, dental and oral health, smoking, alcohol consumption, dental check-ups and mouthwash use was obtained as a part of the study.

Statistical analysis

Statistical analysis involved descriptive tables, Chi-square test and Cox regression [15]. The magnitude of association between a factor and mouth cancer awareness was described by the relative risk (RR). Data were analysed using Stata 11 for Windows (StataCorp, 2009) and IBM SPSS Statistics version 19 (2010, SPSS INC, an IBM company) statistical packages.

RESULTS

A total of 3,022 full and 331 short questionnaires were returned. Of the remainder, 250 subjects were assumed not to have received the questionnaire, either because notification was received from the occupants or post office that the subject had moved (n = 205), they were severely disabled/terminally ill (n = 2) or had died (n = 30). This resulted in an adjusted participation rate of 58.2%. Participation rate was highest in the age group 55 - 64 (68.5%) and lowest in those aged 25 - 34 (47.3%). The lowest participation rate was among the most deprived participant (SIMD Quintile 1, 44.6%) and the highest in more affluent (60.7% in quintile 4 and 60.4% in quintile 5).

The most commonly identified cancer was cancer of the breast (88.1%) followed by cancer of lung (87.3%), skin (87.3%), bowel (87.3%), prostate (87.2%), cervix (86%), throat (84.6%) and mouth (81%).

Overall, 81% of all the participants were aware of mouth cancer. Mouth cancer awareness was related to age with participants aged 75 years or over being the least aware (67.7%) (test for trend P < 0.001), deprivation (test for trend P = 0.034), better general health (test for trend P < 0.001), self-reported oral health (test for trend P = 0.016) regular dental check-ups (test for trend P < 0.001), mouthwash use (test for trend P = 0.034), higher number of natural teeth (test for trend P < 0.001) and education with participants from primary and secondary school education only being less aware (78.4%) (test for trend P < 0.001). Among 46 people who reported their level of education as only primary school, 26 (56.5%) were male, 37 (80.4%) were aged 65 years or older and 30 (65.2%) lived in remote small towns or rural locations.

On univariate analysis, there was no significant association found with gender (Chi-square test P = 0.065), rurality (test for trend P = 0.3) or geographical area (Chi-square test P = 0.475). Mouth cancer awareness was related to smoking (test for trend P = 0.004) with current smokers being the most aware of mouth cancer (87.2%) and alcohol consumption (test for trend P < 0.001) with participants who did not drink being the least aware (73.5%) (Table 1 and 2).

When factors significant on univariate analysis were considered in multivariate analysis, adjusted for age and gender, the best predictors of mouth cancer awareness were gender, education, smoking, alcohol consumption and number of teeth (Table 3).

When asked about risk factors for mouth cancer the following were identified from the suggested list:

Table 1. Knowledge of mouth cancer by socio-demographic characteristics

| Factor | Total number in group | Aware number (%) | Statistical test (P-value) ^a |
|------------------------------------|-----------------------|------------------|---|
| Age | | | |
| 25 - 34 | 460 | 379 (82.4) | Test for trend (< 0.001) |
| 35 - 44 | 605 | 504 (83.3) | |
| 45 - 54 | 470 | 407 (86.6) | |
| 55 - 64 | 545 | 455 (83.5) | |
| 65 - 74 | 527 | 424 (80.5) | |
| 75+ | 415 | 281 (67.7) | |
| Gender | | | |
| Male | 1438 | 1146 (79.7) | Chi-square test (0.065) |
| Female | 1584 | 1304 (82.3) | |
| Education | | | |
| Primary school | 46 | 23 (50.0) | Test for trend (< 0.001) |
| Secondary school | 1354 | 1074 (79.3) | |
| Technical college | 787 | 671 (85.3) | |
| University/postgraduate | 781 | 647 (82.8) | |
| Deprivation (SIMD Quintile) | | | |
| 1 (most deprived) | 120 | 92 (76.7) | Test for trend (0.034) |
| 2 | 334 | 255 (76.3) | |
| 3 | 764 | 624 (81.7) | |
| 4 | 987 | 809 (82.0) | |
| 5 | 817 | 669 (81.9) | |
| Urban/rural | | | |
| Large urban | 712 | 571 (80.2) | Test for trend (0.3) |
| Other urban | 481 | 393 (81.7) | |
| Accessible small town | 280 | 219 (78.2) | |
| Remote small town | 272 | 218 (80.1) | |
| Accessible rural | 867 | 709 (81.8) | |
| Remote rural | 410 | 339 (82.7) | |
| Geographical area | | | |
| Aberdeen city | 771 | 623 (80.8) | Chi-square test (0.475) |
| Aberdeenshire north | 738 | 585 (79.3) | |
| Aberdeenshire south | 729 | 597 (81.9) | |
| Moray | 784 | 644 (82.1) | |

^aLevel of statistical significance P < 0.05.

SIMD = Scottish Index of Multiple Deprivation.

smoking (84.2%), poor oral hygiene (60%), drinking alcohol heavily (58.8%), poor diet (37.4%), stress (15.2%), being overweight (6.2%), drinking hot liquids (5%), eating spicy food (2.7%), using mouthwash (2.3%) and kissing someone (1.1%) (Table 4). In addition, one respondent identified HPV as a risk factor for mouth cancer (however no dedicated space for open suggestions were provided in the questionnaire).

Table 2. Knowledge of mouth cancer by health-related characteristics

| Factor | Total number in group | Aware number (%) | Statistical test (P-value) ^a |
|-------------------------------------|-----------------------|------------------|---|
| Self-reported general health | | | |
| Excellent | 671 | 559 (83.3) | Test for trend (< 0.001) |
| Good | 1732 | 1440 (83.1) | |
| Fair | 459 | 340 (74.1) | |
| Poor | 83 | 64 (77.1) | |
| Self-reported oral health | | | |
| Excellent | 340 | 288 (84.7) | Test for trend (P = 0.016) |
| Good | 1575 | 1291 (82) | |
| Fair | 741 | 599 (80.8) | |
| Poor | 246 | 189 (76.8) | |
| Dental check-up | | | |
| Less than 1 year ago | 2017 | 1691 (83.4) | Test for trend (< 0.001) |
| 1 - 2 years ago | 283 | 226 (79.9) | |
| 3 - 4 years ago | 164 | 129 (78.7) | |
| 5 - 10 years ago | 197 | 151 (76.7) | |
| Over 10 years ago | 300 | 224 (74.7) | |
| Number of teeth | | | |
| No natural teeth | 356 | 265 (74.4) | Test for trend (< 0.001) |
| < 10 | 225 | 163 (72.4) | |
| 10 - 19 | 522 | 408 (78.2) | |
| 20 or more | 1859 | 1587 (85.4) | |
| Mouthwash use | | | |
| Daily | 646 | 551 (85.3) | Test for trend (P = 0.034) |
| Once every few days | 499 | 415 (83.2) | |
| Less than once a month | 449 | 374 (83.3) | |
| Never | 980 | 795 (81.1) | |
| Smoking | | | |
| Current smoker | 553 | 482 (87.2) | Test for trend (0.004) |
| Used to smoke | 779 | 634 (81.4) | |
| Never smoker | 1530 | 1240 (81) | |
| Alcohol consumption | | | |
| Almost every day | 320 | 275 (85.9) | Test for trend (< 0.001) |
| Once or twice a week | 1045 | 881 (84.3) | |
| Less than once a week | 1014 | 829 (81.8) | |
| Does not drink | 551 | 405 (73.5) | |

^aLevel of statistical significance P < 0.05.

Table 3. Multivariate model of mouth cancer unawareness

| Factor | Rate ratio (95% CI) ^a |
|----------------------------|----------------------------------|
| Gender | |
| Male | 1 |
| Female | 0.8 (0.67; 0.97) |
| Education | |
| University/postgraduate | 1 |
| Technical college | 0.81 (0.62; 1.05) |
| Secondary school | 0.95 (0.75; 1.2) |
| Primary school | 1.83 (1.08; 3.1) |
| Smoking | |
| Never smoker | 1 |
| Used to smoke | 0.91 (0.73; 1.13) |
| Current smoker | 0.6 (0.45; 0.8) |
| Alcohol consumption | |
| Does not drink | 1 |
| Less than once a week | 0.83 (0.64; 1.06) |
| Once or twice a week | 0.7 (0.54; 0.91) |
| Every day | 0.69 (0.48; 1) |
| Number of teeth | |
| 20 or more | 1 |
| 10 - 19 | 1.49 (1.13; 1.97) |
| < 10 | 1.78 (1.25; 2.55) |
| No natural teeth | 1.69 (1.21; 2.35) |

^aRate ratio of being UNAWARE of mouth cancer estimated from Cox regression model adjusted for age. All variables are in the same model. For example, participants with no natural teeth were 1.69 times more likely to be UNAWARE of mouth cancer compared to those who have 20 or more natural teeth. CI = confidence interval.

Table 4. Distribution of participant-identified risk factors for mouth cancer

| Characteristic | % identified characteristic as a risk factor for mouth cancer |
|--------------------------|---|
| Smoking | 84.2 |
| Poor oral hygiene | 60 |
| Drinking alcohol heavily | 58.8 |
| Poor diet | 37.4 |
| Stress | 15.2 |
| Being overweight | 6.2 |
| Drinking hot liquids | 5 |
| Eating spicy food | 2.7 |
| Using mouthwash | 2.3 |
| Kissing someone | 1.1 |

Table 5. Relationship between potential risk factors for mouth cancer and recognising them as risk factors

| Characteristic | % identified characteristic as a risk factor for mouth cancer | Chi-square test P-value ^a |
|----------------------------|---|--------------------------------------|
| Smoking | Smoking is a risk factor | 0.016 |
| Current smoker | 89.3 | |
| Used to smoke | 84.9 | |
| Never smoker | 84.4 | |
| Alcohol consumption | Heavy alcohol consumption is a risk factor | < 0.001 |
| Every day | 66.6 | |
| Once or twice a week | 63.3 | |
| Less than once a week | 56.3 | |
| Mouthwash use | Mouthwash is a risk factor | 0.085 |
| Daily | 1.7 | |
| Once every few days | 1.8 | |
| Less than once a month | 2.2 | |
| Oral Health | Poor oral health is a risk factor | 0.783 |
| Excellent | 61 | |
| Good | 60.2 | |
| Fair | 62.5 | |
| Dental check-up | Poor oral health is a risk factor | 0.033 |
| Less than 1 year ago | 62.1 | |
| 1 - 2 years ago | 63.3 | |
| 3 - 4 years ago | 61 | |
| 5 - 10 years ago | 55.8 | |
| Over 10 years ago | 53.7 | |
| Number of teeth | Poor oral health is a risk factor | < 0.001 |
| No natural teeth | 51.7 | |
| < 10 | 56 | |
| 10 - 19 | 62.6 | |
| 20 or more | 62.8 | |

^aLevel of statistical significance $P < 0.05$.

Current smokers were more likely to identify smoking as a risk factor for mouth cancer ($P = 0.016$). Similarly, those who consumed alcohol almost daily were more likely to identify heavy alcohol drinking as a risk factor ($P < 0.001$). Participants with the most recent dental check-up over 5 years ago and those with less than 10 natural teeth were less likely to identify poor oral health as a risk factor (Table 5).

DISCUSSION

This survey found that awareness of mouth cancer was high in respondents from the general population.

The results are similar to those reported by West et al. [12], and show 25% higher OC awareness than the level (56%) determined by the study of Warnakulasuriya et al. [10] in 1995. It is possible that the level of awareness was higher than reported here. The awareness was assumed if respondents ticked a relevant box in the questionnaire. It is plausible that some participants did not tick all possible boxes rather than being unaware, for example, of lung cancer. In fact, there were 310 participants who did not tick any of the boxes. Assuming they missed the question completely, lung cancer awareness would change from 87.3% to 97.3%.

While the participation rate achieved in the current survey was 58%, the historic rates in this geographical location varied from between 25% and 82% [16-18]. The difference between the previous high response level and the value achieved presently could be attributed to forgetfulness, participants' disinterest in the survey topic [19] or the new requirement to provide prior consent to receive a postal questionnaire [18].

Regular consumption of alcohol and tobacco smoking have been identified as the main, yet preventable, causes for cancers of the upper aerodigestive tract (UADT) [12,20-22]. West et al. [12] reported that members of the general public who were more likely to be unaware of mouth cancer were characterised by: older age, low occupational status, no history of tobacco use or frequent alcohol consumption. In our study current smokers and alcohol users were more aware of mouth cancer. They were also more likely to identify smoking and heavy alcohol drinking as risk factors for mouth cancer. A possible explanation is that people tend to remember information which carries a direct personal relevance. Although even a limited exposure to tobacco products constitutes a risk factor for oral cancer, low doses of alcohol might not necessarily pose a similar threat to individual's health [12]. Future questionnaires may seek to establish not only the frequency, but also the quantity of alcohol intake.

To the best of our knowledge, the present study is the first one to investigate the public perception of potential risk factors for oral cancer such as poor oral hygiene, nutritional deficiencies, high body mass index (BMI), mouthwash use and stress.

Previous research has indicated that there is a positive relationship between poor oral hygiene and cancer of the head and neck, including oral and oropharyngeal squamous cell carcinoma [23-28]. Our study shows that 60% of the participants were aware of the association between poor oral hygiene and oral cancer. This awareness was significantly higher in people with better oral health and dental care behaviour as indicated by the number of natural teeth present ($P = 0.033$) and frequency of dental check-ups ($P < 0.001$) (Table 5). The results may be associated with the oral hygiene education which patients receive during their examination appointments. The current study thereby highlights the important role of the dental team in mouth cancer prevention.

The importance of diet for good health and general well-being is being increasingly recognised. In particular, consumption of fresh fruit and non-starchy vegetables has been associated with decreased cancer risks, while diet rich in red and processed meat has been linked to elevated risks [21,29,30]. The present

study shows that 38% of the respondents indicated that poor diet presents a risk factor for oral cancer. A future survey may include more specific questions relating to different eating habits in order to better understand the public awareness of the association between diet and cancer.

Fifteen percent of participants stated that 'stress' may lead to an increased risk of OC. A physiological theory of 'stress' was developed at the beginning of the 20th century [31]. This provided a potential link between the central nervous system and the release of hormones with inhibitory effects on the immune cells [32]. Although some studies did not support a direct relationship between the adverse life events and the onset of cancer [33], others have indicated that a reduction in stress levels contributed to a decreased rate of metastasis and prolonged survival time [34].

Being overweight was recognised as a malignancy risk factor by 6.2% of the respondents. Rates of weight gain and obesity (high BMI) have increased 2-fold in many developed economies between 1990 and 2005 [21]. Evidence has shown that being obese elevates the risks for type 2 diabetes, hyperlipidaemia, and certain types of cancers [21]. As both the consumption of calorie-dense food and inadequate food intake can result in deficiency of micronutrients, it can be speculated that the actual nutritional status of an individual has an additional bearing on the cancer risk [35,36]. Future studies should integrate individual BMI analysis, biochemical and epidemiological investigations as well as personal nutritional profile.

Our study found that 5% of respondents considered drinking hot liquids to be a risk factor for mouth cancer. In a comprehensive review of the available evidence the World Cancer Research Fund and the American Institute for Cancer Research [21] did not find sufficient evidence to indicate that tea and coffee increase the risk of cancer. However, the consumption of beverages at very high temperature may be associated with such risks [21,37,38]. A notable example is maté, a South American drink, which is drunk traditionally at scalding temperatures through a metal straw, and has been linked to increased risks of oesophageal cancer [21,37,39]. Taking into consideration that hot beverages are consumed daily by the public further research to fully establish the potential role of thermal irritation of the epithelium and UADT cancer is warranted.

A small proportion of the respondents in our study (2.7%) were of the opinion that eating spicy food presents a risk factor for mouth cancer. Certain foods contain phytochemicals which give them distinctive spicy flavour, for instance isothiocyanates in mustard and horseradish, sulphides in garlic and chives, and terpenoids in herbs and spices [21]. There is not much

evidence suggesting that these compounds increase the risks of cancer. Indeed there are data suggesting potentially beneficial effects in the cases of many herbs and spices such as ginger, saffron and turmeric [21,40]. Therefore, it is possible that the participants in our study who considered eating spicy food to be a risk factor for mouth cancer may have simply associated the hot/spicy sensation with potentially harmful effects rather than providing their answer on the basis of prior knowledge or evidence.

Only 2.3% of respondents were of the opinion that using mouthwash presents a risk factor for mouth cancer. At present the research evidence on the issue is controversial. Some studies have reported similar oral cancer risks among mouthwash users and non-users [41-46] while other studies have reported that mouthwash use is associated with significantly increased risks of developing malignancy [11,26,47,48]. It should be noted that positive association between mouthwash use and increased risks of oral cancer appears to be found mainly with alcohol containing mouthwashes [11]. As mouthwash is a product which supplements daily dental hygiene [13,48], it is imperative that the issue of whether mouthwash can increase the risks of oral cancer is resolved.

Infection with HPV has been recognised as a risk factor for oral and oropharyngeal cancer [45]. In a hospital-based case-control study D'Souza et al. [49] demonstrated that oropharyngeal cancer was significantly associated with HPV infection, regardless of the patients' alcohol and tobacco use. Although research evidence suggests that HPV infection is likely to be sexually acquired, the possibility of mouth-to-mouth transmission is also being considered [50]. In our study 1.1% of participants identified "kissing someone" as a potential risk factor for mouth cancer. This highlights the importance of further research to fully establish the association between viral infection and OC. Furthermore, public awareness of the risks should be raised through campaigns similar to those addressing HPV and cervical cancer.

Earlier UK surveys found that public knowledge of initial signs of OC was relatively low [12,51]. A survey conducted in the US indicated that 66% of respondents were not familiar with head and neck cancer, only 55% recognized tobacco smoking as a risk factor and few had knowledge of potential signs and symptoms of OC [52]. Public awareness of risk factors, signs, and symptoms of oral malignancies facilitates early detection and thereby improves the treatment outcomes [53]. This is particularly important for high-risk individuals such as tobacco and alcohol consumers, and people who are not

routinely attending dental examinations. Such individuals might present to a hospital with the disease in an advanced stage which may negatively affect the prognosis and the range of available treatment options [54].

While increasing public awareness of risk factors, signs and symptoms of OC would be beneficial for its early detection, adequate education of health-care providers is critical in this respect [55,56]. A study by Tanriover et al. [57] showed that nearly 30% of primary care physicians did not ask patients about tobacco consumption, and more than 45% did not make enquiries about alcohol consumption. Furthermore, the physicians were of the belief that they did not receive adequate smoking cessation training (69.5%) and alcohol cessation training (79.9%) [57]. A cross-sectional survey conducted by Ahluwalia et al. [58] showed that the majority of GP training programmes in the UK did not include structured training in relation to oral health care and OC detection. Another cross-sectional study conducted in Brazil showed that most primary care dentists were unable to identify potentially malignant lesions and to perform biopsy [53]. Furthermore, Wolever et al. [59] argued that the primary healthcare providers should have a dedicated time and adequate resources in order to concentrate on helping patients to alter their lifestyle. Achieving long lasting behavioural changes requires an ongoing professional support using evidence based interventions.

CONCLUSIONS

Awareness of mouth cancer is high in respondents from the general population and participants were able to identify the most important risk factors. Knowledge of tobacco and alcohol as risk factors was highest amongst those exposed to them. This study suggested that preventative programmes should focus not only on increasing individual's knowledge, but also on changing their health behaviour.

ACKNOWLEDGMENTS AND DISCLOSURE STATEMENTS

The authors are grateful to Angus V, Chaloner F and Wilde K from the University of Aberdeen and Rhodes K, Morgan R and Watkins R from NHS Grampian for help with sampling and data management.

We are grateful to the participants for completing the questionnaire.

The Authors declare that there is no conflict of interest.

REFERENCES

1. Diz P, Meleti M, Diniz-Freitas M, Vescovi P, Warnakulasuriya S, Johnson NW, Kerr AR. Oral and pharyngeal cancer in Europe: Incidence, mortality and trends as presented to the Global Oral Cancer Forum. *Transl Res Oral Oncol*. 2017 Jan;2:1-3. [doi: [10.1177/2057178X17701517](https://doi.org/10.1177/2057178X17701517)]
2. Ferlay J, Colombet M, Soerjomataram I, Dyba T, Randi G, Bettio M, Gavin A, Visser O, Bray F. Cancer incidence and mortality patterns in Europe: Estimates for 40 countries and 25 major cancers in 2018. *Eur J Cancer*. 2018 Nov;103:356-387. [Medline: [30100160](https://pubmed.ncbi.nlm.nih.gov/30100160/)] [doi: [10.1016/j.ejca.2018.07.005](https://doi.org/10.1016/j.ejca.2018.07.005)]
3. Head and neck cancers incidence statistics. Cancer Research UK. 2018. [URL: <https://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type/head-and-neck-cancers/incidence#heading-Zero>]
4. Head and neck cancer incidence by sex and UK country. Cancer Research UK. 2018. [URL: <https://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type/head-and-neck-cancers/mortality>]
5. Cogliano VJ, Baan R, Straif K, Grosse Y, Lauby-Secretan B, El Ghissassi F, Bouvard V, Benbrahim-Tallaa L, Guha N, Freeman C, Galichet L, Wild CP. Preventable exposures associated with human cancers. *J Natl Cancer Inst*. 2011 Dec 21;103(24):1827-39. [Medline: [22158127](https://pubmed.ncbi.nlm.nih.gov/22158127/)] [PMC free article: [3243677](https://pubmed.ncbi.nlm.nih.gov/3243677/)] [doi: [10.1093/jnci/djr483](https://doi.org/10.1093/jnci/djr483)]
6. Ram H, Sarkar J, Kumar H, Konwar R, Bhatt ML, Mohammad S. Oral cancer: risk factors and molecular pathogenesis. *J Maxillofac Oral Surg*. 2011 Jun;10(2):132-7. [Medline: [22654364](https://pubmed.ncbi.nlm.nih.gov/22654364/)] [PMC free article: [3177522](https://pubmed.ncbi.nlm.nih.gov/3177522/)] [doi: [10.1007/s12663-011-0195-z](https://doi.org/10.1007/s12663-011-0195-z)]
7. Chuang SC, Jenab M, Heck JE, Bosetti C, Talamini R, Matsuo K, Castellsague X, Franceschi S, Herrero R, Winn DM, La Vecchia C, Morgenstern H, Zhang ZF, Levi F, Dal Maso L, Kelsey K, McClean MD, Vaughan T, Lazarus P, Muscat J, Ramroth H, Chen C, Schwartz SM, Eluf-Neto J, Hayes RB, Purdue M, Boccia S, Cadoni G, Zaridze D, Koifman S, Curado MP, Ahrens W, Benhamou S, Matos E, Lagiou P, Szeszenia-Dabrowska N, Olshan AF, Fernandez L, Menezes A, Agudo A, Daudt AW, Merletti F, Macfarlane GJ, Kjaerheim K, Mates D, Holcatova I, Schantz S, Yu GP, Simonato L, Brenner H, Mueller H, Conway DI, Thomson P, Fabianova E, Znaor A, Rudnai P, Healy CM, Ferro G, Brennan P, Boffetta P, Hashibe M. Diet and the risk of head and neck cancer: a pooled analysis in the INHANCE consortium. *Cancer Causes Control*. 2012 Jan;23(1):69-88. [Medline: [22037906](https://pubmed.ncbi.nlm.nih.gov/22037906/)] [PMC free article: [3654401](https://pubmed.ncbi.nlm.nih.gov/3654401/)] [doi: [10.1007/s10552-011-9857-x](https://doi.org/10.1007/s10552-011-9857-x)]
8. Lagiou P, Talamini R, Samoli E, Lagiou A, Ahrens W, Pohlabein H, Benhamou S, Bouchardy C, Slamova A, Schejbalova M, Merletti F, Richiardi L, Kjaerheim K, Agudo A, Castellsague X, Macfarlane TV, Macfarlane GJ, Biggs AM, Barzan L, Canova C, Simonato L, Lowry RJ, Conway DI, McKinney PA, Znaor A, McCartan BE, Healy CM, Marron M, Hashibe M, Brennan P. Diet and upper-aerodigestive tract cancer in Europe: the ARCAGE study. *Int J Cancer*. 2009 Jun 1;124(11):2671-6. [Medline: [19230023](https://pubmed.ncbi.nlm.nih.gov/19230023/)] [doi: [10.1002/ijc.24246](https://doi.org/10.1002/ijc.24246)]
9. Franceschi S, Muñoz N, Bosch XF, Snijders PJ, Walboomers JM. Human papillomavirus and cancers of the upper aerodigestive tract: a review of epidemiological and experimental evidence. *Cancer Epidemiol Biomarkers Prev*. 1996 Jul;5(7):567-75. [Medline: [8827362](https://pubmed.ncbi.nlm.nih.gov/8827362/)]
10. Warnakulasuriya KA, Harris CK, Scarrott DM, Watt R, Gelbier S, Peters TJ, Johnson NW. An alarming lack of public awareness towards oral cancer. *Br Dent J*. 1999 Sep 25;187(6):319-22. [Medline: [10589135](https://pubmed.ncbi.nlm.nih.gov/10589135/)] [doi: [10.1038/sj.bdj.4800269a](https://doi.org/10.1038/sj.bdj.4800269a)]
11. McCullough MJ, Farah CS. The role of alcohol in oral carcinogenesis with particular reference to alcohol-containing mouthwashes. *Aust Dent J*. 2008 Dec;53(4):302-5. [Medline: [19133944](https://pubmed.ncbi.nlm.nih.gov/19133944/)] [doi: [10.1111/j.1834-7819.2008.00070.x](https://doi.org/10.1111/j.1834-7819.2008.00070.x)]
12. West R, Alkhatib MN, McNeill A, Bedi R. Awareness of mouth cancer in Great Britain. *Br Dent J*. 2006 Feb 11;200(3):167-9, discussion 151. [Medline: [16474365](https://pubmed.ncbi.nlm.nih.gov/16474365/)] [doi: [10.1038/sj.bdj.4813197](https://doi.org/10.1038/sj.bdj.4813197)]
13. Macfarlane TV, Kawecki MM, Cunningham C, Bovaird I, Morgan R, Rhodes K, Watkins R. Mouthwash use in general population: results from adult dental health survey in grampian, Scotland. *J Oral Maxillofac Res*. 2011 Jan 1;1(4):e2. [Medline: [24421979](https://pubmed.ncbi.nlm.nih.gov/24421979/)] [PMC free article: [3886070](https://pubmed.ncbi.nlm.nih.gov/3886070/)] [doi: [10.5037/jomr.2010.1402](https://doi.org/10.5037/jomr.2010.1402)]
14. Grampian - Youth Lifestyle Survey's 2007. NHS Grampian. [URL: <http://www.nhsgrampian.org>]
15. Cox DR. Regression Models and Life-Tables. *J Roy Stat Soc: Series B (Methodological)*. 1972 Jan;34(2):187-220. [doi: [10.1111/j.2517-6161.1972.tb00899.x](https://doi.org/10.1111/j.2517-6161.1972.tb00899.x)]
16. Elliott AM, Smith BH, Penny KI, Smith WC, Chambers WA. The epidemiology of chronic pain in the community. *Lancet*. 1999 Oct 9;354(9186):1248-52. [Medline: [10520633](https://pubmed.ncbi.nlm.nih.gov/10520633/)] [doi: [10.1016/S0140-6736\(99\)03057-3](https://doi.org/10.1016/S0140-6736(99)03057-3)]
17. Vasianovich A, van Teijlingen ER, Reid G, Scott NW. Key health promotion factors among male members of staff at a higher educational institution: a cross-sectional postal survey. *BMC Public Health*. 2008 Feb 12;8:58. [Medline: [18269744](https://pubmed.ncbi.nlm.nih.gov/18269744/)] [PMC free article: [2270269](https://pubmed.ncbi.nlm.nih.gov/2270269/)] [doi: [10.1186/1471-2458-8-58](https://doi.org/10.1186/1471-2458-8-58)]
18. Angus VC, Entwistle VA, Emslie MJ, Walker KA, Andrew JE. The requirement for prior consent to participate on survey response rates: a population-based survey in Grampian. *BMC Health Serv Res*. 2003 Nov 18;3(1):21. [Medline: [14622444](https://pubmed.ncbi.nlm.nih.gov/14622444/)] [PMC free article: [293468](https://pubmed.ncbi.nlm.nih.gov/293468/)] [doi: [10.1186/1472-6963-3-21](https://doi.org/10.1186/1472-6963-3-21)]
19. Huang JY, Hubbard SM, Mulvey KP. Obtaining valid response rates: considerations beyond the tailored design method. *Eval Program Plann*. 2003 Feb;26(1):91-7. [doi: [10.1016/S0149-7189\(02\)00091-5](https://doi.org/10.1016/S0149-7189(02)00091-5)]
20. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. Tobacco smoke and involuntary smoking. *IARC Monogr Eval Carcinog Risks Hum*. 2004;83:1-1438. [Medline: [15285078](https://pubmed.ncbi.nlm.nih.gov/15285078/)] [PMC free article: [4781536](https://pubmed.ncbi.nlm.nih.gov/4781536/)] [doi: [10.1186/1472-6963-3-21](https://doi.org/10.1186/1472-6963-3-21)]

21. World Cancer Research Fund, American Institute for Cancer Research. Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective. Washington DC: AICR; 2007. [URL: <http://eprints.ucl.ac.uk/4841/1/4841.pdf>]
22. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. Alcohol consumption and ethyl carbamate. IARC Monogr Eval Carcinog Risks Hum. 2010;96:3-1383. [Medline: [21735939](#)] [PMC free article: [4781168](#)]
23. Maier H, Zöller J, Herrmann A, Kreiss M, Heller WD. Dental status and oral hygiene in patients with head and neck cancer. Otolaryngol Head Neck Surg. 1993 Jun;108(6):655-61. [Medline: [8516003](#)] [doi: [10.1177/019459989310800606](#)]
24. Velly AM, Franco EL, Schlecht N, Pintos J, Kowalski LP, Oliveira BV, Curado MP. Relationship between dental factors and risk of upper aerodigestive tract cancer. Oral Oncol. 1998 Jul;34(4):284-91. [Medline: [9813724](#)] [doi: [10.1016/S1368-8375\(98\)80009-2](#)]
25. Rosenquist K, Wennerberg J, Schildt EB, Bladström A, Göran Hansson B, Andersson G. Oral status, oral infections and some lifestyle factors as risk factors for oral and oropharyngeal squamous cell carcinoma. A population-based case-control study in southern Sweden. Acta Otolaryngol. 2005 Dec;125(12):1327-36. [Medline: [16303683](#)] [doi: [10.1080/00016480510012273](#)]
26. Guha N, Boffetta P, Wünsch Filho V, Eluf Neto J, Shangina O, Zaridze D, Curado MP, Koifman S, Matos E, Menezes A, Szeszenia-Dabrowska N, Fernandez L, Mates D, Daudt AW, Lissowska J, Dikshit R, Brennan P. Oral health and risk of squamous cell carcinoma of the head and neck and esophagus: results of two multicentric case-control studies. Am J Epidemiol. 2007 Nov 15;166(10):1159-73. [Medline: [17761691](#)] [doi: [10.1093/aje/kwm193](#)]
27. Abnet CC, Kamangar F, Islami F, Nasrollahzadeh D, Brennan P, Aghcheli K, Merat S, Pourshams A, Marjani HA, Ebadati A, Sotoudeh M, Boffetta P, Malekzadeh R, Dawsey SM. Tooth loss and lack of regular oral hygiene are associated with higher risk of esophageal squamous cell carcinoma. Cancer Epidemiol Biomarkers Prev. 2008 Nov;17(11):3062-8. [Medline: [18990747](#)] [PMC free article: [2586052](#)] [doi: [10.1158/1055-9965.EPI-08-0558](#)]
28. Chang JS, Lo HI, Wong TY, Huang CC, Lee WT, Tsai ST, Chen KC, Yen CJ, Wu YH, Hsueh WT, Yang MW, Wu SY, Chang KY, Chang JY, Ou CY, Wang YH, Weng YL, Yang HC, Wang FT, Lin CL, Huang JS, Hsiao JR. Investigating the association between oral hygiene and head and neck cancer. Oral Oncol. 2013 Oct;49(10):1010-7. [Medline: [23948049](#)] [doi: [10.1016/j.oraloncology.2013.07.004](#)]
29. Fruit and vegetables. In: IARC Handbooks of Cancer Prevention. Vainio H, Bianchini F, editors. Lyon: IARC Press; 2003(8). p.376. [URL: <http://publications.iarc.fr/>]
30. IARC Working Group on the Evaluation of Carcinogenic Risk to Humans. Red Meat and Processed Meat. Lyon (FR): International Agency for Research on Cancer; 2018. [Medline: [29949327](#)]
31. Reiche EM, Nunes SO, Morimoto HK. Stress, depression, the immune system, and cancer. Lancet Oncol. 2004 Oct;5(10):617-25. [Medline: [15465465](#)] [doi: [10.1016/S1470-2045\(04\)01597-9](#)]
32. Haas HS, Schauenstein K. Immunity, hormones, and the brain. Allergy. 2001 Jun;56(6):470-7. [Medline: [11421890](#)] [doi: [10.1034/j.1398-9995.2001.056006470.x](#)]
33. Petticrew M, Fraser JM, Regan MF. Adverse life-events and risk of breast cancer: A meta-analysis. Br J Health Psychol. 1999 Feb;4(1):1-17. [doi: [10.1348/135910799168434](#)]
34. Morgenstern H, Gellert GA, Walter SD, Ostfeld AM, Siegel BS. The impact of a psychosocial support program on survival with breast cancer: the importance of selection bias in program evaluation. J Chronic Dis. 1984;37(4):273-82. [Medline: [6715493](#)] [doi: [10.1016/0021-9681\(84\)90135-8](#)]
35. Franceschi S, Dal Maso L, Levi F, Conti E, Talamini R, La Vecchia C. Leanness as early marker of cancer of the oral cavity and pharynx. Ann Oncol. 2001 Mar;12(3):331-6. [Medline: [11332144](#)] [doi: [10.1023/A:1011191809335](#)]
36. Zheng W, Blot WJ, Shu XO, Diamond EL, Gao YT, Ji BT, Fraumeni JF Jr. Risk factors for oral and pharyngeal cancer in Shanghai, with emphasis on diet. Cancer Epidemiol Biomarkers Prev. 1992 Sep-Oct;1(6):441-8. [Medline: [1302555](#)]
37. Islami F, Pourshams A, Nasrollahzadeh D, Kamangar F, Fahimi S, Shakeri R, Abedi-Ardekani B, Merat S, Vahedi H, Semnani S, Abnet CC, Brennan P, Møller H, Saidi F, Dawsey SM, Malekzadeh R, Boffetta P. Tea drinking habits and oesophageal cancer in a high risk area in northern Iran: population based case-control study. BMJ. 2009 Mar 26;338:b929. [Medline: [19325180](#)] [PMC free article: [3269898](#)] [doi: [10.1136/bmj.b929](#)]
38. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Drinking Coffee, Mate, and Very Hot Beverages. 2016 May;116. [URL: <http://publications.iarc.fr/>]
39. Islami F, Boffetta P, Ren JS, Pedoeim L, Khatib D, Kamangar F. High-temperature beverages and foods and esophageal cancer risk--a systematic review. Int J Cancer. 2009 Aug 1;125(3):491-524. [Medline: [19415743](#)] [PMC free article: [2773211](#)] [doi: [10.1002/ijc.24445](#)]
40. Thomas R, Williams M, Sharma H, Chaudry A, Bellamy P. A double-blind, placebo-controlled randomised trial evaluating the effect of a polyphenol-rich whole food supplement on PSA progression in men with prostate cancer--the U.K. NCRN Pomi-T study. Prostate Cancer Prostatic Dis. 2014 Jun;17(2):180-6. [Medline: [24614693](#)] [PMC free article: [4020278](#)] [doi: [10.1038/pcan.2014.6](#)]
41. Blot WJ, Winn DM, Fraumeni JF Jr. Oral cancer and mouthwash. J Natl Cancer Inst. 1983 Feb;70(2):251-3. [Medline: [6571933](#)]
42. Mashberg A, Barsa P, Grossman ML. A study of the relationship between mouthwash use and oral and pharyngeal cancer. J Am Dent Assoc. 1985 May;110(5):731-4. [Medline: [3859544](#)] [doi: [10.14219/jada.archive.1985.0422](#)]

43. Young TB, Ford CN, Brandenburg JH. An epidemiologic study of oral cancer in a statewide network. *Am J Otolaryngol*. 1986 May-Jun;7(3):200-8. [Medline: [3728828](#)] [doi: [10.1016/S0196-0709\(86\)80007-2](#)]
44. La Vecchia C. Mouthwash and oral cancer risk: an update. *Oral Oncol*. 2009 Mar;45(3):198-200. [Medline: [18952488](#)] [doi: [10.1016/j.oraloncology.2008.08.012](#)]
45. Warnakulasuriya S. Causes of oral cancer--an appraisal of controversies. *Br Dent J*. 2009 Nov 28;207(10):471-5. [Medline: [19946320](#)] [doi: [10.1038/sj.bdj.2009.1009](#)]
46. Gandini S, Negri E, Boffetta P, La Vecchia C, Boyle P. Mouthwash and oral cancer risk quantitative meta-analysis of epidemiologic studies. *Ann Agric Environ Med*. 2012;19(2):173-80. [Medline: [22742785](#)]
47. Winn DM, Blot WJ, McLaughlin JK, Austin DF, Greenberg RS, Preston-Martin S, Schoenberg JB, Fraumeni JF Jr. Mouthwash use and oral conditions in the risk of oral and pharyngeal cancer. *Cancer Res*. 1991 Jun 1;51(11):3044-7. [Medline: [2032242](#)]
48. Ahrens W, Pohlabeln H, Foraita R, Nelis M, Lagiou P, Lagiou A, Bouchardy C, Slamova A, Schejbalova M, Merletti F, Richiardi L, Kjaerheim K, Agudo A, Castellsague X, Macfarlane TV, Macfarlane GJ, Lee YC, Talamini R, Barzan L, Canova C, Simonato L, Thomson P, McKinney PA, McMahon AD, Znaor A, Healy CM, McCartan BE, Metspalu A, Marron M, Hashibe M, Conway DI, Brennan P. Oral health, dental care and mouthwash associated with upper aerodigestive tract cancer risk in Europe: the ARCAGE study. *Oral Oncol*. 2014 Jun;50(6):616-25. [Medline: [24680035](#)] [doi: [10.1016/j.oraloncology.2014.03.001](#)]
49. D'Souza G, Kreimer AR, Viscidi R, Pawlita M, Fakhry C, Koch WM, Westra WH, Gillison ML. Case-control study of human papillomavirus and oropharyngeal cancer. *N Engl J Med*. 2007 May 10;356(19):1944-56. [Medline: [17494927](#)] [doi: [10.1056/NEJMoa065497](#)]
50. Heck JE, Berthiller J, Vaccarella S, Winn DM, Smith EM, Shan'gina O, Schwartz SM, Purdue MP, Pilarska A, Eluf-Neto J, Menezes A, McClean MD, Matos E, Koifman S, Kelsey KT, Herrero R, Hayes RB, Franceschi S, Wünsch-Filho V, Fernández L, Daudt AW, Curado MP, Chen C, Castellsagué X, Ferro G, Brennan P, Boffetta P, Hashibe M. Sexual behaviours and the risk of head and neck cancers: a pooled analysis in the International Head and Neck Cancer Epidemiology (INHANCE) consortium. *Int J Epidemiol*. 2010 Feb;39(1):166-81. [Medline: [20022926](#)] [PMC free article: [2817092](#)] [doi: [10.1093/ije/dyp350](#)]
51. Robb K, Stubbings S, Ramirez A, Macleod U, Austoker J, Waller J, Hiom S, Wardle J. Public awareness of cancer in Britain: a population-based survey of adults. *Br J Cancer*. 2009 Dec 3;101 Suppl 2:S18-23. [Medline: [19956158](#)] [PMC free article: [2790705](#)] [doi: [10.1038/sj.bjc.6605386](#)]
52. Luryi AL, Yarbrough WG, Niccolai LM, Roser S, Reed SG, Nathan CA, Moore MG, Day T, Judson BL. Public awareness of head and neck cancers: a cross-sectional survey. *JAMA Otolaryngol Head Neck Surg*. 2014 Jul;140(7):639-46. [Medline: [24902640](#)] [doi: [10.1001/jamaoto.2014.867](#)]
53. Galvão-Moreira LV, da Cruz MCFN. Screening and early detection of oral cancer: current controversies. *Acta Odontol Scand*. 2017 Jul;75(5):361-365. [Medline: [28447861](#)] [doi: [10.1080/00016357.2017.1316868](#)]
54. Seoane J, Takkouche B, Varela-Centelles P, Tomás I, Seoane-Romero JM. Impact of delay in diagnosis on survival to head and neck carcinomas: a systematic review with meta-analysis. *Clin Otolaryngol*. 2012 Apr;37(2):99-106. [Medline: [22429704](#)] [doi: [10.1111/j.1749-4486.2012.02464.x](#)]
55. Carter LM, Parsonage-Grant S, Marshall A, Achal KS, Kanatas A. Oral cancer teaching of medical students in the UK: time for a new approach? *J Cancer Educ*. 2011 Jun;26(2):308-14. [Medline: [20872262](#)] [doi: [10.1007/s13187-010-0156-0](#)]
56. Sousa FB, Freitas e Silva MR, Fernandes CP, Silva PG, Alves AP. Oral cancer from a health promotion perspective: experience of a diagnosis network in Ceará. *Braz Oral Res*. 2014;28 Spec No. [Medline: [24964281](#)] [doi: [10.1590/1807-3107BOR-2014.vol28.0018](#)]
57. Tanriover O, Hidiroglu S, Save D, Akan H, Ay P, Karavus M, Hayran O. Knowledge of oral cancer, preventive attitudes, and behaviors of primary care physicians in Turkey. *Eur J Cancer Prev*. 2014 Sep;23(5):464-8. [Medline: [24584196](#)] [doi: [10.1097/CEJ.0000000000000020](#)]
58. Ahluwalia A, Crossman T, Smith H. Current training provision and training needs in oral health for UK general practice trainees: survey of General Practitioner Training Programme Directors. *BMC Med Educ*. 2016 May 11;16:142. [Medline: [27169796](#)] [PMC free article: [4863349](#)] [doi: [10.1186/s12909-016-0663-8](#)]
59. Wolever RQ, Caldwell KL, McKernan LC, Hillinger MG. Integrative Medicine Strategies for Changing Health Behaviors: Support for Primary Care. *Prim Care*. 2017 Jun;44(2):229-245. [Medline: [28501227](#)] [doi: [10.1016/j.pop.2017.02.007](#)]

To cite this article:

Kawecki MM, Nedeva IR, Iloya J, Macfarlane TV.

Mouth Cancer Awareness in General Population: Results from Grampian Region of Scotland, United Kingdom

J Oral Maxillofac Res 2019;10(2):e3

URL: <http://www.ejomr.org/JOMR/archives/2019/2/e3/v10n2e3.pdf>

doi: [10.5037/jomr.2019.10203](https://doi.org/10.5037/jomr.2019.10203)

Copyright © Kawecki MM, Nedeva IR, Iloya J, Macfarlane TV. Published in the JOURNAL OF ORAL & MAXILLOFACIAL RESEARCH (<http://www.ejomr.org>), 30 June 2019.

This is an open-access article, first published in the JOURNAL OF ORAL & MAXILLOFACIAL RESEARCH, distributed under the terms of the [Creative Commons Attribution-Noncommercial-No Derivative Works 3.0 Unported License](https://creativecommons.org/licenses/by-nc-nd/3.0/), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work and is properly cited. The copyright, license information and link to the original publication on (<http://www.ejomr.org>) must be included.