

University of Dundee

**Effect of depleted uranium on a soil microcosm fungal community and influence of a plant-ectomycorrhizal association**

Fomina, Marina; Hong, Ji Won; Gadd, Geoffrey Michael

*Published in:*  
Fungal Biology

*DOI:*  
[10.1016/j.funbio.2019.08.001](https://doi.org/10.1016/j.funbio.2019.08.001)

*Publication date:*  
2020

*Document Version*  
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

*Citation for published version (APA):*

Fomina, M., Hong, J. W., & Gadd, G. M. (2020). Effect of depleted uranium on a soil microcosm fungal community and influence of a plant-ectomycorrhizal association. *Fungal Biology*, 124(5), 289-296. <https://doi.org/10.1016/j.funbio.2019.08.001>

**General rights**

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Table 2. Uranium amounts and fractionation in soil-containing microcosms.

Fraction	Microcosm	U (g per jar)	Difference
HNO <sub>3</sub> –fraction in soil	Mycorrhizal	125.77 ± 14.98	a
	Non-mycorrhizal	85.18 ± 9.04	a
	Non-pine	333.50 ± 42.98	b
Acetic acid–fraction in soil	Mycorrhizal	31.67 ± 1.55	b
	Non-mycorrhizal	15.31 ± 1.65	a
	Non-pine	104.32 ± 7.39	c
Water–fraction in soil*	Mycorrhizal	0.011 ± 0.0002	ab*
	Non-mycorrhizal	0.009 ± 0.0002	a*
	Non-pine	0.034 ± 0.0060	b*
HNO <sub>3</sub> –fraction in pine seedling	Mycorrhizal	0.137 ± 0.049	No significant difference
	Non-mycorrhizal	0.343 ± 0.30	
	Non-pine	0	-

± SEM. At least three replicate determinations were made. The different letters denote

significant differences at the 5% level using All Pairwise Multiple Comparison Procedures, the Holm-Sidak method or Tukey Test\* (one-way ANOVA).