

University of Dundee

## Intracellular Transfer of Na<sup>+</sup> in an Active State G Protein Coupled Receptor

Vickery, Owen; Carvalheda Dos Santos, Catarina; Zaidi, Saheem A.; Pislakov, Andrei; Katritch, Vsevolod; Zachariae, Ulrich

*Published in:*  
Structure

*DOI:*  
[10.1016/j.str.2017.11.013](https://doi.org/10.1016/j.str.2017.11.013)

*Publication date:*  
2018

*Document Version*  
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

### *Citation for published version (APA):*

Vickery, O., Carvalheda Dos Santos, C., Zaidi, S. A., Pislakov, A., Katritch, V., & Zachariae, U. (2018). Intracellular Transfer of Na<sup>+</sup> in an Active State G Protein Coupled Receptor. *Structure*, 26(1), 171-180.e2. <https://doi.org/10.1016/j.str.2017.11.013>

### **General rights**

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Intracellular Transfer of Na<sup>+</sup> in an Active State G-Protein Coupled Receptor

Owen N. Vickery<sup>1,2</sup>, Catarina A. Carvalheda<sup>1,2</sup>, Saheem A. Zaidi<sup>3</sup>, Andrei V. Pislakov<sup>1,2</sup>,  
Vsevolod Katritch<sup>3,4</sup>, Ulrich Zachariae<sup>1,2,\*</sup>

<sup>1</sup>School of Life Sciences, University of Dundee, Dundee DD1 5EH, UK.

<sup>2</sup>School of Science and Engineering, University of Dundee DD1 4NH, UK.

<sup>3</sup>Departments of Biological Sciences and <sup>4</sup>Chemistry, Bridge Institute, University of  
Southern California, Los Angeles, CA 90089, USA

\*Corresponding author; email: u.zachariae@dundee.ac.uk

\*Lead Contact; email: u.zachariae@dundee.ac.uk

## ABSTRACT

Playing a central role in cell signaling, GPCRs are the largest superfamily of membrane proteins and form the majority of drug targets in humans. How extracellular agonist binding triggers the activation of GPCRs and associated intracellular effector proteins remains, however, poorly understood. Structural studies have revealed that inactive class-A GPCRs harbor a conserved binding site for Na<sup>+</sup> ions in the center of their transmembrane domain, accessible from the extracellular space. Here, we show that the opening of a conserved hydrated channel in the activated state receptors allows the Na<sup>+</sup> ion to egress from its binding site into the cytosol. Coupled with protonation changes, this ion movement occurs without significant energy barriers, and can be driven by physiological transmembrane ion and voltage gradients. We propose that Na<sup>+</sup> ion exchange with the cytosol is a key step in GPCR activation. Further, we hypothesize that this transition locks receptors in long-lived active-state conformations.

## INTRODUCTION

G-protein coupled receptors (GPCRs) mediate the transfer of external ligand binding information across the plasma membrane to activate a range of intracellular signaling pathways (Pierce et al., 2002). Playing a central role in regulation of vital biological systems, including nervous, cardiovascular, immune, digestive, reproductive etc., they represent the majority of membrane proteins in humans and the largest class of present drug targets (Overington et al., 2006; Rask-Andersen et al., 2014). In recent years, a number of crystal structures have been solved to reveal conformational changes between inactive and active state receptors, including common movement in transmembrane helices and conserved microswitches (Katritch et al., 2013; Venkatakrishnan et al., 2013). However, despite this wealth of structural information, it is still not fully understood how ligand binding leads to activated receptors, which are able to trigger nucleotide exchange in intracellular effector G-protein complexes.

One of the major unknowns is the role of the highly conserved hydrophilic water-filled channel observed in crystal structures of class A GPCRs, which extends along the receptor axis from the external ligand-binding region nearly all the way to the effector binding site. The channel is sealed toward the cytoplasm by a thin layer of hydrophobic residues in inactive state GPCRs (Fig 1 A, B). High resolution X-ray structures of the inactive conformation reveal a Na<sup>+</sup> ion near the floor of this pocket, coordinated by water and three or four conserved residues including an acidic aspartate that is fully conserved in all ligand-sensing class A GPCRs (Christopher et al., 2013; Fenalti et al., 2014; Kruse et al., 2012; Liu et al., 2012; Miller-Gallacher et al., 2014; Pardo et al., 2007; Zhang et al., 2012) (D<sup>2.50</sup>; superscript refers to the Ballesteros and Weinstein residue numbering system) (Isberg et al., 2015). The allosteric effect of monovalent cations, in particular Na<sup>+</sup> ions, for GPCR function has been known for almost half a century (Pert and Synder, 1974), and the bulk of recent evidence shows that these effects are largely mediated by the ion binding at the D<sup>2.50</sup> site at the physiological concentration of Na<sup>+</sup> (140 mM and lower) (Fenalti et al., 2014; Liu et al., 2012; Massink et al., 2015). Due to the highly conserved nature of D<sup>2.50</sup> and other Na<sup>+</sup> ion coordinating residues, Na<sup>+</sup> ion binding at this site is likely to be a ubiquitous feature shared by the vast majority of class A GPCRs (Katritch et al., 2014).

In active receptor conformations, the ion binding site near D<sup>2.50</sup> shows a collapsed state, which is likely not optimal for Na<sup>+</sup> ion binding (Huang et al., 2015; Kruse et al., 2013; Liu et al., 2012; Rasmussen et al., 2011). It was therefore proposed that the Na<sup>+</sup> ion leaves the

hydrophilic pocket upon receptor activation by a ligand or during receptor-G-protein complex formation. However, how this movement is triggered and which pathway is followed by the ion remains unknown.

Here, we investigated the link between ligand-induced receptor activation, the fate of the bound Na<sup>+</sup> ion in class A GPCRs and its implications for transmembrane (TM) signal transduction by equilibrium and non-equilibrium atomistic simulations on the M2 muscarinic receptor (m2r). When one addresses these questions, it is important to take physiologically relevant electrochemical membrane conditions into consideration. Strong TM Na<sup>+</sup> and K<sup>+</sup> gradients produce a sizable voltage across the plasma membrane of up to -100 mV in the resting state of mammalian cells (Kandel et al., 2000). Both the ionic gradients and electric field have been shown to influence the function of GPCRs (Ben-Chaim et al., 2006; Navarro-Polanco et al., 2011; Rinne et al., 2015) and are likely to impact the movement of the Na<sup>+</sup> ion within the membrane region.

Our data reveal that the Na<sup>+</sup> ion observed in the TM domain of class A GPCRs can readily traverse the receptor and, driven by the electrochemical gradients, migrate into the cytoplasm in active receptor conformations. This result implies that a Na<sup>+</sup> ion may be exchanged from the extracellular space to the cytoplasm as an important step in receptor activation. Furthermore, the movement of Na<sup>+</sup> in the receptor, and intracellular egress, are coupled to a protonation change of D<sup>2.50</sup>.

## RESULTS

### *GPCR activation opens a hydrated pathway across the receptor*

We were first interested whether the conformational change from the inactive to active receptor state renders the ion-binding pocket sterically incapable of accommodating a Na<sup>+</sup> ion. The binding site for Na<sup>+</sup> appears to adopt a collapsed conformation in active crystal structures. We started from an inactive state structure of the m2 muscarinic acetylcholine receptor (m2r, PDB ID: 3UON) and, using a targeted molecular dynamics (TMD) approach, gently drove this conformation to the active state of this receptor (PDB ID: 4MQT) (Fig S1). Our simulations show that the active state of m2r initially retains sufficient space for the ion. The electrostatic attraction between the ion and the negatively charged side chain of D69<sup>2.50</sup> keeps the ion bound to this site during and after the transition from the inactive to the active receptor conformation (Fig S1). However, our simulations show a widening of the intracellular portion of the TM helices below the hydrophilic pocket during this conformational change, which subsequently becomes fully hydrated (Fig 1 B, C). The hydrated pathway forms a connection between the orthosteric ligand-binding site, the hydrophilic pocket and the G-protein binding site. The slim hydrophobic layer that delimits the hydrophilic pocket toward the G-protein binding site in the inactive crystal structure undergoes substantial conformational changes, which are especially evident from the sidechain position of Y440<sup>7.53</sup>. Our simulations show two major conformations of the Y440<sup>7.53</sup> sidechain following the transition – an upward state similar to the conformation observed in the inactive crystal structure (PDB: 3UON; Fig 1 D) and a downward configuration, which is also seen in the active crystal structure (PDB: 4MQT; Fig 1 E). The formation of a hydrated pathway connecting the receptor ligand and effector binding sites has been reported in previous simulation studies on the A<sub>2A</sub>R and 5-HT<sub>1A</sub> receptors (Yuan et al., 2014, 2016), however the previous reports did not take the presence of a Na<sup>+</sup> ion into consideration.

### ***The position of the internal Na<sup>+</sup> ion is coupled to protonation of D2.50***

We were next interested in the interplay between the Na<sup>+</sup> ion and the key conserved titratable residue D69<sup>2.50</sup>. A number of computational studies have explored functional implications of the protonation state of D<sup>2.50</sup>, in particular its role in receptor activation, Na<sup>+</sup> ion binding, and interaction with the “ionic lock” motif (DR<sup>3.50</sup>Y) in several class A family GPCRs (Miao et al., 2015; Ranganathan et al., 2014; Vanni et al., 2010). Here, we focused on a potential coupling between the position of the Na<sup>+</sup> ion within the receptor and protonation of D69<sup>2.50</sup>. We carried out pK<sub>a</sub> calculations on D69<sup>2.50</sup> using more than 800 equilibrated frames from simulations of the m2r receptor in a variety of conformations, including both the upward and downward configurations of the Y440<sup>7.53</sup> sidechain. Due to the formation of a hydrated pathway across the receptor from the ligand to the effector binding sites in the active state simulations, we were able to evaluate the effect of the Na<sup>+</sup> ion positional changes on the D69<sup>2.50</sup> pK<sub>a</sub>, where the Na<sup>+</sup> ion was shifted both in the upward (toward the extracellular face) and downward direction.

Figure 2 shows that the pK<sub>a</sub> value and, thus, the most likely protonation state of D69<sup>2.50</sup> are substantially influenced by the Na<sup>+</sup> ion. If the cation is within ~3-5 Å of D69<sup>2.50</sup>, its positive charge strongly stabilizes the negatively charged form of D69<sup>2.50</sup>, leading to a pK<sub>a</sub> value of ~3–4. However, displacement of the Na<sup>+</sup> ion to distances of 5 Å and greater gives rise to a substantial pK<sub>a</sub> shift to values between 8-12. This can be understood given the location of D69<sup>2.50</sup> in the middle of the transmembrane domain, surrounded by many non-polar residues. Transient movements of the internal Na<sup>+</sup> ion from its binding site, facilitated by activation-related conformational changes in the Na<sup>+</sup> pocket, can therefore be sufficient to lead to protonation of D69<sup>2.50</sup>.

For the protonation of D<sup>2.50</sup>, we propose that the most likely proton entry route would be from the extracellular side, along the negative membrane potential gradient. Moreover, in the m2r and other aminergic receptors the proton could be transferred from the conserved D<sup>3.32</sup> in the orthosteric binding pocket via a short chain of water molecules (Isom and Dohlman, 2015). In the apo state, our calculations in m2r indicate that D103<sup>3.32</sup> is generally protonated (pK<sub>a</sub> = 11.2±1.7), whereas upon ligand binding the pK<sub>a</sub> is substantially lowered (pK<sub>a</sub> = 7.6±1.9). A possible protonation change of D<sup>3.32</sup>, induced by ligand binding, could thus facilitate the shuttling of protons to D<sup>2.50</sup>. Furthermore, if a G-protein complex with a receptor is preformed before agonist binding, D<sup>2.50</sup> would be readily accessible for protonation from the extracellular side via a hydrated pathway. In this context, it has further

been argued that bound agonists, but not antagonists, may sustain the hydrated pathway past the ligand which connects the extracellular space with the Na<sup>+</sup> ion binding site upon receptor activation (Yuan et al., 2016). Interestingly, the protonation state of D69<sup>2.50</sup> shows an effect on the stability of the activated receptor state in our simulations. Under equilibrium, the active state remains stable when D69<sup>2.50</sup> is neutral (Fig S2), while it exhibits a greater propensity to revert back to the inactive state when D<sup>2.50</sup> is charged. We obtain similar results for non-equilibrium simulations (Fig S2). This lends further support to an important role of D<sup>2.50</sup> protonation for receptor activation.

### ***Simulations under electrochemical gradient show ion movement to the intracellular face***

Next we conducted atomistic simulations with the Computational Electrophysiology (CompEL) protocol (Kutzner et al., 2016) on the active conformation of m2r. We applied a physiological Na<sup>+</sup> ion gradient of 150:10 mM across the membrane from the extracellular to the intracellular side, in addition to a small ion imbalance evoking a hyperpolarised V<sub>m</sub> at -250 mV. Due to the wide range of pK<sub>a</sub> values that D69<sup>2.50</sup> can adopt, its sidechain was modeled both in charged and neutral forms.

Our simulations at -250 mV show that the Na<sup>+</sup> ion exhibits a substantial degree of mobility even when D69<sup>2.50</sup> is in the charged state (Fig 3 A, B). The Na<sup>+</sup> ion is predominantly coordinated by the residues D69<sup>2.50</sup>, S110<sup>3.39</sup>, N435<sup>7.45</sup> and S433<sup>7.46</sup>. Under a small membrane voltage, a bimodal distribution of distances between the ion and D69<sup>2.50</sup> is observed, where larger distances of 5–6 Å are not uncommon (Fig S1). As our pK<sub>a</sub> calculations showed that moderate excursions of the ion from its original binding site on this scale are likely to have a major impact on the pK<sub>a</sub> and protonation state of the D69<sup>2.50</sup> sidechain (Fig 2), we investigated the effect of a protonation change of D69<sup>2.50</sup> in the active conformation.

Our simulations reveal that, in this receptor conformation, the Na<sup>+</sup> ion readily passes through the hydrated channel into the intracellular solution. When D69<sup>2.50</sup> is neutral, we observe the Na<sup>+</sup> ion to be expelled into the intracellular solution in three out of four simulations at -250 mV (Fig 3 A, C; for a complete list of trajectories see Table S1). At -500 mV the effect is, expectably, even more pronounced and movement into the cytoplasm is seen in all four simulations we conducted (Fig 3 B, D; Table S1). In contrast, when D69<sup>2.50</sup> is in a charged state, such a transition is observed only in one out of eight

simulations, namely at a raised membrane voltage (Fig 3 A, B; Table S1). The observed translocation of  $\text{Na}^+$  to the intracellular side occurs irrespective of the conformation adopted by Y440<sup>7.53</sup> (Fig 1 D, E and 3 C, D).

In our simulations as well as under physiological conditions, both TM ion concentration and voltage gradients drive ion flow across membrane pores. In the case of the  $\text{Na}^+$  ion, both gradients act synergistically in the resting state of the cell, driving the  $\text{Na}^+$  ion toward the cytoplasm. Under the conditions used in the simulations, fast ion motion through the receptor is predominantly voltage-driven. Converted into an effective force, and using a linear approximation to describe the gradient across the membrane (Dill and Bromberg, 2011), the influence of the concentration gradient would be about 10-fold smaller ( $\sim 1.3$  pN) than the driving force caused by the voltage gradient under these conditions ( $\sim 13$  pN). At physiological conditions, both driving forces are likely to be of similar magnitude, such that ion migration could either be induced by the voltage or ion gradients.

### ***Energetics of ion movement to the cytoplasm***

As the initiation of fast ion movement to the intracellular side was initially tested under slightly supra-physiological levels of  $V_m$ , we next evaluated the detailed equilibrium energetics of the  $\text{Na}^+$  ion movement on this pathway (i.e. without applied gradients) to ascertain the physiological relevance of this transition. We calculated the potential-of-mean-force (PMF) for the migration of the cation in four different states. In addition to probing the influence of the D69<sup>2.50</sup> protonation state, we examined the role of the conformation of the Y440<sup>7.53</sup> sidechain, which substantially affects the width and overall shape of the formed hydrated pathway into the cytoplasm (Fig 3 C, D).

When D69<sup>2.50</sup> is charged (Fig 4), the free energy difference between the internal  $\text{Na}^+$  ion binding site and the free intracellular bulk solution is  $\sim 30$  kJ mol<sup>-1</sup>. Accordingly, the active conformation of m2r retains a  $\text{Na}^+$  ion at the allosteric site ( $Z = 7.5\text{-}8$  Å) with relatively high affinity, as long as D69<sup>2.50</sup> remains deprotonated. The major barrier to migration into the cytoplasm is located near the Y440<sup>7.53</sup> sidechain. In its upward state, the free energy barrier amounts to  $\sim 41$  kJ mol<sup>-1</sup>, while it increases to  $\sim 48$  kJ mol<sup>-1</sup> in the downward state (Fig 4).

As our  $pK_a$  calculations showed that even a moderate displacement of the  $\text{Na}^+$  ion away from its binding site at D69<sup>2.50</sup> is likely to lead to a protonation change of the aspartate, we also calculated the PMF for the movement of  $\text{Na}^+$  along the intracellular pathway with neutral D69<sup>2.50</sup>. Importantly, this state no longer shows any affinity for the  $\text{Na}^+$  ion, and ion



movement into the intracellular bulk is not obstructed by any energy barrier significantly larger than the thermal energy ( $kT$ ,  $\sim 2.5$  kJ mol<sup>-1</sup>) in the upward-oriented Y440<sup>7.53</sup> conformation. When Y440<sup>7.53</sup> is oriented downward, a small but readily surmountable energy barrier (on physiologically relevant timescales) of  $\sim 14$  kJ mol<sup>-1</sup> exists for this transition. The downward conformation of Y440<sup>7.53</sup>, in conjunction with the neutral state of D69<sup>2.50</sup> also has a small influence on the shape and configuration of the ion binding site at D69<sup>2.50</sup>, which leads to a reduction of the number of hydrogen bonds formed between the protein and the ion (Fig S3), raising the free energy of binding at this site further by  $\sim 7.5$  kJ mol<sup>-1</sup> (Fig 4 A, B). In the non-equilibrium case, with a physiological  $V_m$  applied, the free energy minima at  $Z = \sim 7.5$  Å will be raised with regard to the intracellular bulk by  $\sim 4.4$  kJ mol<sup>-1</sup> per 100 mV (Fig S4). This means that, in all of these states, the Na<sup>+</sup> ion can readily traverse the receptor and permeate along a hydrated pathway to the intracellular side.

### ***Conservation of the pocket and intracellular exit channel***

Additional support for an important role of intracellular Na<sup>+</sup> egress in the activation of class A GPCRs is provided by an analysis of residue conservation along its exit pathway. As we detailed previously (Katritch et al., 2014), there is a remarkable level of conservation for the 16 residues of the Na<sup>+</sup> binding pocket in class A GPCRs (Figure 5, Table S1), suggesting a conserved functional role of Na<sup>+</sup> in receptor activation mechanism. Interestingly, our analysis of Na<sup>+</sup> contacts along the MD trajectories in this study shows that the residues lining the ion exit path to the intracellular side are also well conserved. Thus, out of the 36 contact residues, 32 are 100% conserved among all five muscarinic receptors, 17 are >90% conserved among all aminergic receptors, and 22 are consensus residues among all class A GPCRs. Most importantly, the discovered exit pathway includes Na<sup>+</sup> contacts with the highly conserved N<sup>1.50</sup> (100% and 98% conserved in aminergic and in all class A respectively), D<sup>3.49</sup> (100% and 64%), Y<sup>5.58</sup> (94% and 73%) and other residue positions generally conserved as polar residues, including N<sup>1.60</sup>, T<sup>2.37</sup> and N<sup>2.39</sup>. Particularly, in the inactive M2 muscarinic receptor and in other inactive state GPCR structures as well, the Y<sup>7.53</sup> residue is directed toward the Na<sup>+</sup> ion-binding pocket, and hence may play a role as first point of polar contact outside the Na<sup>+</sup> ion-binding pocket for the intracellular movement of Na<sup>+</sup>. Na<sup>+</sup> ion passage toward the cytosol may be further facilitated by other conserved polar residues, including D<sup>3.49</sup>, N<sup>2.39</sup>, N<sup>2.40</sup> and T<sup>2.37</sup>. The conservation of the Na<sup>+</sup> ion pocket and the path for intracellular egress of Na<sup>+</sup> suggests that the Na<sup>+</sup> transfer

described in this study can occur in all muscarinic receptors and other class A GPCRs, comprising a key, unidirectional part of the activation mechanism.

## DISCUSSION

The principal role of GPCRs is to transmit information about an extracellular agonist binding event toward the cytoplasm, by catalyzing GDP release from a bound intracellular G-protein complex (Pierce et al., 2002). This is known to involve conformational changes in the receptor, including conserved residue microswitches, and large scale movement of TM helices 6 and 7 on the intracellular side that open the nucleotide binding site of the  $G\alpha$  protein (Dror et al., 2015; Huang et al., 2015; Mahoney and Sunahara, 2016). It has, furthermore, long been recognized that G-protein binding, and stabilization of this conformation on the intracellular side of the receptor, increases agonist affinity on the extracellular face (DeVree et al., 2016; Maguire et al., 1976).

$Na^+$  ions, binding to an internal receptor site between the G-protein and the external ligand binding pockets, are known to act as powerful allosteric modulators of class A GPCRs (Katritch et al., 2014; Pert and Synder, 1974).  $Na^+$  was found to selectively diminish the affinity of agonists, but not antagonists, to GPCRs, which can be interpreted as a structural stabilization of the inactive receptor state by the ions (Miller-Gallacher et al., 2014; Quitterer et al., 1996; Selley et al., 2000). Accordingly, while receptor X-ray structures of sufficient resolution crystallized in the inactive state display a  $Na^+$  ion bound to D<sup>2.50</sup>, this binding site is collapsed in active receptor conformations, and ions are not observed (Huang et al., 2015; Katritch et al., 2014). Mutations around the  $Na^+$  ion binding site have a major impact on receptor function in most class A GPCRs (Liu et al., 2012)(Liu et al., 2012)(Liu et al., 2012)(Liu et al. 2012)(Liu et al. 2012)(Liu et al. 2012)(Liu et al. 2012)(Liu et al. 2012)(Liu et al. 2012)(Liu et al. 2012)(Liu et al. 2012), either completely abolishing G-protein activation, or resulting in constitutive ligand-independent or pathway-biased signaling (Fenalti et al., 2014; Liu et al., 2012; Massink et al., 2015).

Our work shows that the  $Na^+$  ion binding pocket, which is accessible only from the extracellular face in the inactive state (Selent et al., 2010; Vickery et al., 2016a), is transformed into a fully permeable, water-filled channel in the activated receptor conformation of m2r (Fig S5). This channel bridges the extracellular ligand and intracellular G-protein binding sites. Water access from the ligand binding site all the way to the

cytoplasmic side of the receptor has previously also been observed in simulations on the A<sub>2A</sub>R and 5-HT<sub>1A</sub> receptors (Yuan et al., 2014, 2016). We show here that the activated receptor state permits the Na<sup>+</sup> ion to readily cross the receptor toward the cytoplasmic side without experiencing major energy barriers on its pathway. The high hydration level of this pathway in the active state is thereby an important factor in facilitating ion passage. A correlation between hydration level and ion transfer has previously been demonstrated in the case of ion channels (Beckstein et al., 2003; Dong et al., 2013; Zhu and Hummer, 2012). In simulations of the inactive state, by contrast, the application of substantially larger forces seems to be necessary to achieve inward migration of Na<sup>+</sup>, as no continuous hydrated channel is formed (Shang et al., 2014).

The inward motion of the Na<sup>+</sup> ion is likely facilitated by a protonation change of D<sup>2.50</sup> from the negatively charged to the neutral form, which we show to occur even upon small displacements of the ion from its equilibrium binding position. Neutralization of D<sup>2.50</sup> substantially reduces the affinity of the binding site for Na<sup>+</sup> ions. Migration of the ion toward the cytosol is then driven by the negative membrane voltage and by a greater than 10-fold Na<sup>+</sup> gradient across the cytoplasmic membrane under physiological conditions, both strongly attracting Na<sup>+</sup> ions inward. Indeed, we observe that moderately negative membrane voltages allow fast escape of the allosteric Na<sup>+</sup> ion to the cytoplasm on 10–100 ns timescales in our simulations.

According to our results, conformational changes associated with agonist binding from the extracellular side and/or G-protein binding from the cytoplasm alters the Na<sup>+</sup> site conformation and the dynamics of the Na<sup>+</sup>–D<sup>2.50</sup> pair. This, in turn, leads to a protonation change of this residue, and subsequent egress of the Na<sup>+</sup> ion via a hydrated exit channel to the intracellular side.

We therefore suggest that intracellular Na<sup>+</sup> ion transfer, facilitated by the membrane potential and Na<sup>+</sup> gradient, is a pivotal step during receptor activation. We further hypothesize that this transition traps the receptor in the active state (Fig 6). The loss of Na<sup>+</sup> is associated with receptor activation, and it has been shown that, once activated, GPCRs remain in a prolonged active state, capable of signaling even when the receptors are internalized from the cytoplasmic membrane during endocytosis (Irannejad et al., 2013; Thomsen et al., 2016). The crucial role of the Na<sup>+</sup> ion movement within the receptor is reflected by the nearly complete conservation of the Na<sup>+</sup> ion binding site in class A GPCRs, as well as the high conservation level of the exit pathway. The mechanism suggested here is

also consistent with agonist independent basal signaling of GPCRs (Kobilka and Deupi, 2007), explaining this phenomenon as spontaneous protonation of D<sup>2.50</sup> and egress of the bound Na<sup>+</sup> ion on the intracellular pathway, leading to receptor activation. Following arrival on the cytoplasmic side, it is conceivable that the ion induces further conformational transitions through its strong interaction with protein residues, including at the G-protein-receptor interface and the G-protein itself. This region includes a number of charged and polar groups, for example a polar network extending across all G-proteins, similar to the one observed in GPCRs which enables ion movement (Isom and Dohlman, 2015; Isom et al., 2013).

Charge movements within membrane proteins, such as the coupled transfer of Na<sup>+</sup> ions and protons suggested by our MD simulations and pK<sub>a</sub> calculations, should be sensitive to the membrane voltage. Indeed, it has been demonstrated that GPCR signaling is modulated by membrane voltage changes (Ben-Chaim et al., 2006; Mahaut-Smith et al., 2008; Martinez-Pinna et al., 2004; Moreno-Galindo et al., 2016; Rinne et al., 2015; Vickery et al., 2016a). This applies both to the conformation of the receptors as well as their transmitted signal. Our findings are therefore consistent with these observations, as they suggest that movement of ions in the receptors constitute a key element in the receptor activation process. The observed voltage regulation of GPCRs is of particular relevance for receptors expressed in electrically excitable cells (Heifetz et al., 2016). In these cell types, the membrane voltage undergoes large-scale oscillations during action potentials. The transmitted receptor signal could thereby be tuned depending on the specific cell type and its excitation status (Vickery et al., 2016b). Crucially, many GPCR drug targets are located in excitable tissue in the brain or muscle, where voltage regulation and a differential response to drugs may play an important role.

To summarize, our results suggest a model for class A GPCR activation, in which conformational changes induced by G-protein and agonist binding are accompanied by the intracellular transfer of an internally bound Na<sup>+</sup> ion. Importantly, these conformational changes encompass rearrangement of the sidechain of Y<sup>7.53</sup>, a conserved receptor microswitch (Katritch et al., 2013), which in its upward state allows nearly barrier-free intracellular permeation of Na<sup>+</sup> ions. This observation forms a functional link between the major Na<sup>+</sup> binding site D<sup>2.50</sup> and Y<sup>7.53</sup> as the first polar point of contact on the intracellular migration pathway of the Na<sup>+</sup> ion. Translocation of the ion is facilitated by protonation of

the conserved D<sup>2.50</sup> residue (Fig 6) and driven by the physiological membrane Na<sup>+</sup> and voltage gradients. The voltage sensitivity of GPCRs, which has been previously reported for many receptors (Vickery et al., 2016b), would thus be a natural consequence of an activation mechanism incorporating the movement of ions as a key element. The Na<sup>+</sup> free receptors are likely to be trapped in an active state, potentially explaining the prolonged mechanisms of signaling observed in many GPCRs. Our results suggest a link between TM signal transduction by receptor proteins and the voltage and ion-gradient driven permeation of ions across ion channels and pores, forming the basis of electric signal transduction in cells.

Based on our findings, we further speculate that the ligand-induced translocation of an ion across the receptor may reflect a common functional principle, which links microbial 7-transmembrane proteins with the structurally remarkably similar eukaryotic GPCRs. The function of microbial 7-transmembrane proteins, such as bacteriorhodopsin and channelrhodopsins, is to transport protons and ions across the membrane following the absorption of photons (Ernst et al., 2014; Mirzadegan and Benko, 2003).

## **CONTRIBUTIONS**

Conceptualization, V.K., U.Z; Methodology, O.N.V, C.A.C, S.A.Z, A.V.P, V.K, U.Z; Analysis, O.N.V, C.A.C, S.A.Z, A.V.P, V.K, U.Z; Investigation, O.N.V, C.A.C, S.A.Z; Writing – Original draft O.N.V, C.A.C, A.V.P, V.K and U.Z; Writing – Review and editing, O.N.V, C.A.C, S.A.Z, A.V.P, V.K, U.Z; Funding acquisition, V.K. and U.Z; Supervision, V.K. and U.Z.

## **ACKNOWLEDGEMENTS**

This work was supported by the BBSRC (Training Grant BB/J013072/1 to U.Z.) and the Scottish Universities' Physics Alliance (C.A.C, A.V.P. and U.Z.). This research was partially supported by National Institute of Health grant DA035764 to V.K. We thank Salomé Llabrés and Daniel Seeliger for fruitful discussions.

## REFERENCES

- Abraham, M.J., Murtola, T., Schulz, R., Páll, S., Smith, J.C., Hess, B., and Lindahl, E. (2015). GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* 1–2, 19–25.
- Baptista, A.M., Martel, P.J., and Soares, C.M. (1999). Simulation of Electron-Proton Coupling with a Monte Carlo Method: Application to Cytochrome c3 Using Continuum Electrostatics. *Biophys. J.* 76, 2978–2998.
- Bashford, D., and Gerwert, K. (1992). Electrostatic calculations of the pKa values of ionizable groups in bacteriorhodopsin. *J. Mol. Biol.* 224, 473–486.
- Beckstein, O., Biggin, P.C., Bond, P., Bright, J.N., Domene, C., Grottesi, A., Holyoake, J., and Sansom, M.S.P. (2003). Ion channel gating: Insights via molecular simulations. *FEBS Lett.* 555, 85–90.
- Ben-Chaim, Y., Chanda, B., Dascal, N., Bezanilla, F., Parnas, I., and Parnas, H. (2006). Movement of “gating charge” is coupled to ligand binding in a G-protein-coupled receptor. *Nature* 444, 106–109.
- Berendsen, H.J.C., Postma, J.P.M., van Gunsteren, W.F., DiNola, A., and Haak, J.R. (1984). Molecular dynamics with coupling to an external bath. *J. Chem. Phys.* 81, 3684–3690.
- Berendsen, H.J.C., Grigera, J.R., and Straatsma, T.P. (1987). The Missing Term in Effective Pair Potentials. *J. Phys. Chem.* 91, 6269–6271.
- Berger, O., Edholm, O., and Jähnig, F. (1997). Molecular dynamics simulations of a fluid bilayer of dipalmitoylphosphatidylcholine at full hydration, constant pressure, and constant temperature. *Biophys. J.* 72, 2002–2013.
- Bussi, G., Donadio, D., and Parrinello, M. (2007). Canonical sampling through velocity rescaling. *J. Chem. Phys.* 126.
- Case, D., Betz, R., Botello-Smith, W., Cerutti, D.S., Cheatham, T.E., Darden, T.A., Duke, R.E., Giese, T.J., Gohlke, H., Goetz, A.W., et al. (2016). Amber 2016.
- Christopher, J.A., Brown, J., Doré, A.S., Errey, J.C., Koglin, M., Marshall, F.H., Myszka, D.G., Rich, R.L., Tate, C.G., Tehan, B., et al. (2013). Biophysical Fragment Screening of the  $\beta$  1 -Adrenergic Receptor: Identification of High Affinity Arylpiperazine Leads Using Structure-Based Drug Design. *J. Med. Chem.* 56, 3446–3455.
- Cordomí, A., Caltabiano, G., and Pardo, L. (2012). Membrane protein simulations using AMBER force field and Berger lipid parameters. *J. Chem. Theory Comput.* 8, 948–958.
- DeVree, B.T., Mahoney, J.P., Vélez-Ruiz, G.A., Rasmussen, S.G.F., Kuszak, A.J., Edwald, E., Fung, J.-J., Manglik, A., Masureel, M., Du, Y., et al. (2016). Allosteric coupling from G protein to the agonist-binding pocket in GPCRs. *Nature* 535, 182–186.
- Dill, K.A., and Bromberg, S. (2011). Molecular Driving Force.

- Dong, H., Fiorin, G., Carnevale, V., Treptow, W., and Klein, M.L. (2013). Pore waters regulate ion permeation in a calcium release-activated calcium channel. *Proc. Natl. Acad. Sci. U. S. A.* *110*, 17332–17337.
- Dror, R.O., Mildorf, T.J., Hilger, D., Manglik, A., Borhani, D.W., Arlow, D.H., Philippsen, A., Villanueva, N., Yang, Z., Lerch, M.T., et al. (2015). Structural basis for nucleotide exchange in heterotrimeric G proteins. *Science* (80-. ). *348*, 1361–1365.
- Ernst, O.P., Lodowski, D.T., Elstner, M., Hegemann, P., Brown, L.S., and Kandori, H. (2014). Microbial and animal rhodopsins: Structures, functions, and molecular mechanisms. *Chem. Rev.* *114*, 126–163.
- Feenstra, K.A., Hess, B., and Berendsen, H.J.C. (1999). Improving efficiency of large time-scale molecular dynamics simulations of hydrogen-rich systems. *J. Comput. Chem.* *20*, 786–798.
- Fenalti, G., Giguere, P.M., Katritch, V., Huang, X.-P., Thompson, A. a, Cherezov, V., Roth, B.L., and Stevens, R.C. (2014). Molecular control of  $\delta$ -opioid receptor signalling. *Nature* *506*, 191–196.
- Haga, K., Kruse, A.C., Asada, H., Yurugi-Kobayashi, T., Shiroishi, M., Zhang, C., Weis, W.I., Okada, T., Kobilka, B.K., Haga, T., et al. (2012). Structure of the human M2 muscarinic acetylcholine receptor bound to an antagonist. *Nature* *482*, 547–551.
- Heifetz, A., James, T., Morao, I., Bodkin, M.J., and Biggin, P.C. (2016). Guiding lead optimization with GPCR structure modeling and molecular dynamics. *Curr. Opin. Pharmacol.* *30*, 14–21.
- Hess, B., Bekker, H., Berendsen, H.J.C., and Fraaije, J.G.E.M. (1997). LINCS: A linear constraint solver for molecular simulations. *J. Comput. Chem.* *18*, 1463–1472.
- Huang, W., Manglik, A., Venkatakrisnan, A.J., Laeremans, T., Feinberg, E.N., Sanborn, A.L., Kato, H.E., Livingston, K.E., Thorsen, T.S., Kling, R.C., et al. (2015). Structural insights into  $\mu$ -opioid receptor activation. *Nature* *524*, 315–321.
- Irannejad, R., Tomshine, J.C., Tomshine, J.R., Chevalier, M., Mahoney, J.P., Steyaert, J., Rasmussen, S.G.F., Sunahara, R.K., El-Samad, H., Huang, B., et al. (2013). Conformational biosensors reveal GPCR signalling from endosomes. *Nature* *495*, 534–538.
- Isberg, V., de Graaf, C., Bortolato, A., Cherezov, V., Katritch, V., Marshall, F.H., Mordalski, S., Pin, J.-P., Stevens, R.C., Vriend, G., et al. (2015). Generic GPCR residue numbers – aligning topology maps while minding the gaps. *Trends Pharmacol. Sci.* *36*, 22–31.
- Isom, D.G., and Dohlman, H.G. (2015). Buried ionizable networks are an ancient hallmark of G protein-coupled receptor activation. *Proc. Natl. Acad. Sci.* *2015*, 201417888.
- Isom, D., Sridharan, V., Baker, R., Clement, S., Smalley, D., and Dohlman, H. (2013). Protons as second messenger regulators of G protein signaling. *Mol. Cell* *51*, 531–538.



- Kandel, E.R., Schwartz, J.H., and Jessell, T.M. (2000). Principles of Neural Science.
- Katritch, V., Cherezov, V., and Stevens, R.C. (2013). Structure-function of the G protein-coupled receptor superfamily. *Annu. Rev. Pharmacol. Toxicol.* 53, 531–556.
- Katritch, V., Fenalti, G., Abola, E.E., Roth, B.L., Cherezov, V., and Stevens, R.C. (2014). Allosteric sodium in class A GPCR signaling. *Trends Biochem. Sci.* 39, 233–244.
- Kobilka, B.K., and Deupi, X. (2007). Conformational complexity of G-protein-coupled receptors. *Trends Pharmacol. Sci.* 28, 397–406.
- Kruse, A.C., Hu, J., Pan, A.C., Arlow, D.H., Rosenbaum, D.M., Rosemond, E., Green, H.F., Liu, T., Chae, P.S., Dror, R.O., et al. (2012). Structure and dynamics of the M3 muscarinic acetylcholine receptor. *Nature* 482, 552–556.
- Kruse, A.C., Ring, A.M., Manglik, A., Hu, J., Hu, K., Eitel, K., Hübner, H., Pardon, E., Valant, C., Sexton, P.M., et al. (2013). Activation and allosteric modulation of a muscarinic acetylcholine receptor. *Nature* 504, 101–106.
- Kutzner, C., Grubmüller, H., de Groot, B.L., and Zachariae, U. (2011). Computational electrophysiology: the molecular dynamics of ion channel permeation and selectivity in atomistic detail. *Biophys. J.* 101, 809–817.
- Kutzner, C., Köpfer, D.A., Machtens, J., Groot, B.L. De, Song, C., and Zachariae, U. (2016). *Biochimica et Biophysica Acta Insights into the function of ion channels by computational electrophysiology simulations.* BBA - Biomembr.
- Lindorff-Larsen, K., Piana, S., Palmo, K., Maragakis, P., Klepeis, J.L., Dror, R.O., and Shaw, D.E. (2010). Improved side-chain torsion potentials for the Amber ff99SB protein force field. *Proteins* 78, 1950–1958.
- Liu, W., Chun, E., Thompson, A. a, Chubukov, P., Xu, F., Katritch, V., Han, G.W., Roth, C.B., Heitman, L.H., IJzerman, A.P., et al. (2012). Structural basis for allosteric regulation of GPCRs by sodium ions. *Science* 337, 232–236.
- Machtens, J.P., Briones, R., Alleva, C., de Groot, B.L., and Fahlke, C. (2017). Gating Charge Calculations by Computational Electrophysiology Simulations. *Biophys. J.* 112, 1396–1405.
- Maguire, M.E., Van Arsdale, P.M., and Gilman, a G. (1976). An agonist-specific effect of guanine nucleotides on binding to the beta adrenergic receptor. *Mol. Pharmacol.* 12, 335–339.
- Mahaut-Smith, M.P., Martinez-Pinna, J., and Gurung, I.S. (2008). A role for membrane potential in regulating GPCRs? *Trends Pharmacol. Sci.* 29, 421–429.
- Mahoney, J.P., and Sunahara, R.K. (2016). Mechanistic insights into GPCR-G protein interactions. *Curr. Opin. Struct. Biol.* 41, 247–254.
- Martinez-Pinna, J., Tolhurst, G., Gurung, I.S., Vandenberg, J.I., and Mahaut-Smith, M.P.

- (2004). Sensitivity limits for voltage control of P2Y receptor-evoked  $\text{Ca}^{2+}$  mobilization in the rat megakaryocyte. *J. Physiol.* *555*, 61–70.
- Massink, A., Gutierrez-de-Teran, H., Lenselink, E.B., Ortiz Zacarias, N. V., Xia, L., Heitman, L.H., Katritch, V., Stevens, R.C., and IJzerman, A.P. (2015). Sodium Ion Binding Pocket Mutations and Adenosine A2A Receptor Function. *Mol. Pharmacol.* *87*, 305–313.
- Metropolis, N., Rosenbluth, A.W., Rosenbluth, M.N., Teller, A.H., and Teller, E. (1953). Equation of state calculations by fast computing machines. *J. Chem. Phys.* *21*, 1087–1092.
- Miao, Y., Caliman, A.D., and McCammon, J.A. (2015). Allosteric Effects of Sodium Ion Binding on Activation of the M3 Muscarinic G-Protein-Coupled Receptor. *Biophys. J.* *108*, 1796–1806.
- Miller-Gallacher, J.L., Nehmé, R., Warne, T., Edwards, P.C., Schertler, G.F.X., Leslie, A.G.W., and Tate, C.G. (2014). The 2.1 Å Resolution Structure of Cyanopindolol-Bound  $\beta_1$ -Adrenoceptor Identifies an Intramembrane  $\text{Na}^+$  Ion that Stabilises the Ligand-Free Receptor. *PLoS One* *9*, e92727.
- Mirzadegan, T., and Benko, G. (2003). Sequence Analyses of G-Protein-Coupled Receptors: Similarities to Rhodopsin - Corrections. *Biochemistry* *42*, 2759–2767.
- Miyamoto, S., and Kollman, P.A. (1992). SETTLE: an analytical version of the SHAKE and RATTLE algorithm for rigid water models. *J. Comput. Chem.* *13*, 952–962.
- Moreno-Galindo, E.G., Alamilla, J., Sanchez-Chapula, J.A., Tristani-Firouzi, M., and Navarro-Polanco, R.A. (2016). The agonist-specific voltage dependence of M2 muscarinic receptors modulates the deactivation of the acetylcholine-gated  $\text{K}^+$  current ( $I_{\text{KACH}}$ ). *Pflügers Arch. - Eur. J. Physiol.*
- Navarro-Polanco, R. a, Moreno Galindo, E.G., Ferrer-Villada, T., Arias, M., Rigby, J.R., Sánchez-Chapula, J. a, and Tristani-Firouzi, M. (2011). Conformational changes in the M2 muscarinic receptor induced by membrane voltage and agonist binding. *J. Physiol.* *589*, 1741–1753.
- Overington, J.P., Al-Lazikani, B., and Hopkins, A.L. (2006). How many drug targets are there? *Nat. Rev. Drug Discov.* *5*, 993–996.
- Pardo, L., Deupi, X., Dölker, N., López-Rodríguez, M.L., and Campillo, M. (2007). The role of internal water molecules in the structure and function of the rhodopsin family of G protein-coupled receptors. *ChemBioChem* *8*, 19–24.
- Pert, C.B., and Synder, S.H. (1974). Opiate Receptor Binding of Agonists and Antagonists Affected Differentially by Sodium. *Mol. Pharmacol.* *10*, 868–879.
- Pierce, K.L., Premont, R.T., and Lefkowitz, R.J. (2002). Seven-transmembrane receptors. *Nat. Rev. Mol. Cell Biol.* *3*, 639–650.
- Quitterer, U., AbdAlla, S., Jarnagin, K., and Müller-Esterl, W. (1996).  $\text{Na}^+$  ions binding to the bradykinin B2 receptor suppress agonist-independent receptor activation. *Biochemistry*

35, 13368–13377.

Ranganathan, A., Dror, R.O., and Carlsson, J. (2014). Insights into the Role of Asp79<sup>2.50</sup> in  $\beta$ 2 Adrenergic Receptor Activation from Molecular Dynamics Simulations. *Biochemistry* 53, 7283–7296.

Rask-Andersen, M., Masuram, S., and Schiöth, H.B. (2014). The Druggable Genome: Evaluation of Drug Targets in Clinical Trials Suggests Major Shifts in Molecular Class and Indication. *Annu. Rev. Pharmacol. Toxicol.* 54, 9–26.

Rasmussen, S.G.F., DeVree, B.T., Zou, Y., Kruse, A.C., Chung, K.Y., Kobilka, T.S., Thian, F.S., Chae, P.S., Pardon, E., Calinski, D., et al. (2011). Crystal structure of the  $\beta$ 2 adrenergic receptor-Gs protein complex. *Nature* 477, 549–555.

Rinne, A., Mobarec, J.C., Mahaut-Smith, M., Kolb, P., and Bunemann, M. (2015). The mode of agonist binding to a G protein-coupled receptor switches the effect that voltage changes have on signaling. *Sci. Signal.* 8, ra110-ra110.

Šali, A., and Blundell, T.L. (1993). Comparative Protein Modelling by Satisfaction of Spatial Restraints. *J. Mol. Biol.* 234, 779–815.

Selent, J., Sanz, F., Pastor, M., and de Fabritiis, G. (2010). Induced effects of sodium ions on dopaminergic G-protein coupled receptors. *PLoS Comput. Biol.* 6.

Selley, D.E., Cao, C.C., Liu, Q., and Childers, S.R. (2000). Effects of sodium on agonist efficacy for G-protein activation in mu-opioid receptor-transfected CHO cells and rat thalamus. *Br. J. Pharmacol.* 130, 987–996.

Shang, Y., Lerouzic, V., Schneider, S., Bisignano, P., Pasternak, G.W., and Filizola, M. (2014). Mechanistic insights into the allosteric modulation of opioid receptors by sodium ions. *Biochemistry* 53, 5140–5149.

Thomsen, A.R.B., Plouffe, B., Cahill, T.J., Shukla, A.K., Tarrasch, J.T., Dosey, A.M., Kahsai, A.W., Strachan, R.T., Pani, B., Mahoney, J.P., et al. (2016). GPCR-G Protein-B-Arrestin Super-Complex Mediates Sustained G Protein Signaling. *Cell* 166, 907–919.

Vanni, S., Neri, M., Tavernelli, I., and Rothlisberger, U. (2010). A Conserved Protonation-Induced Switch can Trigger “Ionic-Lock” Formation in Adrenergic Receptors. *J. Mol. Biol.* 397, 1339–1349.

Venkatakrishnan, a J., Deupi, X., Lebon, G., Tate, C.G., Schertler, G.F., and Babu, M.M. (2013). Molecular signatures of G-protein-coupled receptors. *Nature* 494, 185–194.

Vickery, O.N., Machtens, J.-P., Tamburrino, G., Seeliger, D., and Zachariae, U. (2016a). Structural Mechanisms of Voltage Sensing in G Protein-Coupled Receptors. *Structure* 24, 997–1007.

Vickery, O.N., Machtens, J.-P., and Zachariae, U. (2016b). Membrane potentials regulating GPCRs: insights from experiments and molecular dynamics simulations. *Curr. Opin. Pharmacol.* 30, 44–50.

Willett, P., and Glen, C. (1995). Molecular Recognition of Receptor Sites using a Genetic Algorithm with a Description of Desolvation. 43–53.

Wolf, M.G., Hoefling, M., Aponte-Santamaría, C., Grubmüller, H., and Groenhof, G. (2010). G-membed: Efficient insertion of a membrane protein into an equilibrated lipid bilayer with minimal perturbation. *J. Comput. Chem.* *31*, 2169–2174.

Yuan, S., Filipek, S., Palczewski, K., and Vogel, H. (2014). Activation of G-protein-coupled receptors correlates with the formation of a continuous internal water pathway. *Nat. Commun.* *5*, 4733.

Yuan, S., Peng, Q., Palczewski, K., Vogel, H., and Filipek, S. (2016). Mechanistic Studies on the Stereoselectivity of the Serotonin 5-HT 1A Receptor. *Angew. Chemie Int. Ed.* *55*, 8661–8665.

Zhang, C., Srinivasan, Y., Arlow, D.H., Fung, J.J., Palmer, D., Zheng, Y., Green, H.F., Pandey, A., Dror, R.O., Shaw, D.E., et al. (2012). High-resolution crystal structure of human protease-activated receptor 1. *Nature* *492*, 387–392.

Zhu, F., and Hummer, G. (2012). Drying transition in the hydrophobic gate of the GLIC channel blocks ion conduction. *Biophys. J.* *103*, 219–227.

## Figure Legends

**Figure 1: Major structural features and internal hydration of class A GPCRs in the inactive and active state as shown by the m2r.** (A) The main structural features of class A GPCRs, as exemplified by m2r, include 7 TM helices (blue), an extracellular ligand binding site, the intracellular effector (G-protein) binding site as well as conserved and functionally important residues termed microswitches (selected ones are highlighted). The vertical axis (Z-coordinate) and all positions stated in the text use the C $\alpha$  atom of D103<sup>3.32</sup> as a reference. (B) Conformation of inactive m2r (PDB: 3UON) during the simulations showing the presence of the hydrophobic layer separating the hydrophilic pocket and effector binding site. (C) After transition to the active state (PDB: 4MQT), and further simulation, m2r displays a continuous water channel connecting the orthosteric ligand binding site, hydrophilic pocket and effector binding site. (D, E) The most populated states of the Y440<sup>7.53</sup> sidechain are demonstrated here in an upward (D) and downward conformation (E); please see Fig S3 for a detailed comparison of the upward and downward tyrosine populations. Water molecules are shown in red (surface representation); the position of the allosteric Na<sup>+</sup> ion, as obtained from our initial simulations, is shown as a green sphere, and residues forming the hydrophobic layer (yellow) as well as the bound ligand (carbachol, light green) are depicted in stick representation.

**Figure 2: Proximity of the Na<sup>+</sup> ion modulates protonation of D69<sup>2.50</sup>.** Continuum electrostatics calculations of the pK<sub>a</sub> of the D69<sup>2.50</sup> sidechain using a multitude of m2r conformations obtained from our atomistic simulations in the carbachol-bound active state, both for Y440<sup>7.53</sup> in the upward (left) and downward (right) conformations. The pK<sub>a</sub> is shown as a function of Z, the separation between the Na<sup>+</sup> ion and the C $\alpha$  atom of D103<sup>3.32</sup>, which marks the orthosteric ligand binding pocket, along the TM axis (see Fig 1A). The data points are in addition coloured according to their distance to the D69<sup>2.50</sup> sidechain. The black continuous line, a smoothed spline fit, indicates the approximate average pK<sub>a</sub> for each separation for illustrative purposes, and the dashed black line shows a pK<sub>a</sub> of 7.

**Figure 3: Migration of the Na<sup>+</sup> ion across the receptor to the intracellular side.**

(A-B) Z-coordinate of the Na<sup>+</sup> ion in m2r under a hyperpolarised V<sub>m</sub> of -250 mV (A) and -500 mV (B). Black and grey lines denote simulations with charged D69<sup>2.50</sup>; purple, green and red lines display simulations with neutral D69<sup>2.50</sup>. (C-D) Trajectories of the Na<sup>+</sup> ion moving from the hydrophilic pocket, accessible from the extracellular space, into the intracellular bulk solution at -250 mV (C) and -500 mV (D). Three example trajectories are shown for each V<sub>m</sub>; please see table S1 for a complete list. The color used to display the Na<sup>+</sup> ion corresponds to the trajectories shown in panels A and B, respectively. Examples of the Y440<sup>7.53</sup> upward and downward conformations are shown in green. The pathways of the ion toward the intracellular side are almost indistinguishable from each other until the ion passes Y440<sup>7.53</sup>. Thereafter, the pathways diverge to some degree due to the widened exit region to the cytoplasm.

**Figure 4: Energetics of Na<sup>+</sup> translocation from the hydrophilic pocket to the intracellular side.**

Equilibrium potential of mean force (PMF) profiles of the energetics of Na<sup>+</sup> translocation along the Z-axis in m2r without any applied voltage or concentration gradients. Four relevant states were considered: (Left) negatively charged D69<sup>2.50</sup> (black) or neutral D69<sup>2.50</sup> (red) with the Y440<sup>7.53</sup> sidechain in an upward conformation; (Right) negatively charged D69<sup>2.50</sup> (black) or neutral D69<sup>2.50</sup> (red) with a downward-oriented Y440<sup>7.53</sup> sidechain. The standard deviation of the PMF, obtained from Bayesian bootstrap analysis, is depicted as shaded area. For each PMF, the intracellular bulk solution was used as a reference, and the range of positions adopted by the Y440<sup>7.53</sup> sidechain is denoted by blue dotted lines.

**Figure 5: Conservation of the intracellular Na<sup>+</sup> ion exit pathway.**

The muscarinic m2 receptor is shown in a blue cartoon representation, along with ball-and-stick representation of residues involved in the egress of the Na<sup>+</sup> ion. The carbon atoms of 17 residues that are >90% conserved among aminergic receptors are shown in green, the carbon atoms of additional 15 residues that are conserved among the muscarinic family of receptors are shown in yellow, the carbon atoms of the 4 non-conserved residues are shown in orange.

**Figure 6: Proposed role of Na<sup>+</sup> translocation in GPCR activation.**

Key checkpoints during the transition from the inactive (A) to active (D) state of the receptor. (A) The initial, inactive receptor conformation shows no bound agonist or G-protein, and displays a Na<sup>+</sup> ion bound in a pocket which is sealed towards the cytosol by a hydrophobic layer around Y<sup>7.53</sup>. (B) G-protein and agonist bind to the receptor (in undetermined order), leading to the formation of a continuous water channel across the GPCR. The increased mobility of the Na<sup>+</sup> ion results in a pK<sub>a</sub> shift and subsequent protonation of D<sup>2.50</sup>. (C) Neutralization of D<sup>2.50</sup> and the presence of the hydrated pathway facilitate transfer of Na<sup>+</sup> to the intracellular side, driven by the transmembrane Na<sup>+</sup> gradient and the negative cytoplasmic membrane voltage. (D) The expulsion of Na<sup>+</sup> towards the cytosol results in a prolonged active state of the receptor.

**STAR METHODS:****CONTACT FOR REAGENT AND RESOURCE SHARING**

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Ulrich Zachariae, [uzachariae@dundee.ac.uk](mailto:uzachariae@dundee.ac.uk)

## **METHOD DETAILS**

### ***System Setup***

The simulation system for the m2r in the inactive state was constructed using the crystal structure (PDB: 3UON)(Haga et al., 2012). Ligands and non-GPCR subunits were removed. The missing loop ICL3 was modelled using Modeller (v9.14)(Šali and Blundell, 1993). All internal water molecules and ions were retained, and a Na<sup>+</sup> ion was positioned into the hydrophilic pocket. The charged N- and C-termini were capped using acetyl and methyl moieties, respectively. All ionisable groups were simulated with default protonation states, unless otherwise mentioned. The receptor was embedded into an equilibrated and hydrated 1,2-palmitoyl-oleoyl-sn-glycero-3-phosphocholine (POPC) lipid bilayer using the GROMACS utility `g_membed` (Wolf et al., 2010) resulting in a system size of ~92 x 88 x 97 Å. A concentration of 150 mM NaCl in the aqueous solution was used for the single bilayer systems. During equilibration, all protein heavy atoms were position-restrained with a force constant of 1000 kJ mol<sup>-1</sup> nm<sup>-2</sup> for 5-10 ns. Due to the low degree of internal hydration and medium resolution of the m2r structure, the equilibration was extended by another 100 ns, now without position restraints, to enable full hydration of the hydrophilic pocket.

### ***Targeted MD***

To study the active structure, the ligand carbachol was parameterised using AMBER16, GAFF2 atom types and AM1-BCC partial charges (Case et al., 2016), and docked into the orthosteric ligand binding site using GOLD (v5.2.2) (Willett and Glen, 1995). We then used a targeted MD (TMD) approach with the RMSD to the protein C $\alpha$  atoms of the active m2r crystal structure (PDB: 4MQT)(Kruse et al., 2013) as a reference, in order to gently enforce the transition from the inactive (PDB: 3UON)(Haga et al., 2012) to the active state, and further equilibrated it for ~250 ns. While the backbone rapidly transitioned toward the active conformation (Fig. S1), the adaptation of sidechains and the increase in hydration of the receptor occurred on a slightly slower timescale, necessitating this simulation time span. The two major conformations of Y440<sup>7.53</sup> observed during this simulation were then probed systematically in the PMF calculations, in which distance restraints between N<sup>1.50</sup>-C $\alpha$  and

D<sup>2.50</sup>-C<sub>α</sub> to Y<sup>7.53</sup>-C<sub>ζ</sub> and dihedral restraints on the sidechain of Y<sup>7.53</sup> were used to maintain the protein in the conformations of interest. To keep the G-protein binding site in an active conformation despite the absence of bound G-protein, we applied, at this interaction site, a minimal set of four distance restraints to the C<sub>α</sub> atoms of the terminal groups of TM helices 2, 5, 6 and 7, namely between residues 2.39-6.33, 2.39-5.61, 2.43-7.54 and 6.36-7.54 (Fig S6).

### ***CompEL setup***

For the CompEL simulations, the aforementioned active state simulation system was duplicated along the Z axis to construct double bilayer systems. A NaCl gradient of 150mM:10mM between the extracellular and intracellular compartments was used, along with an ion imbalance of 1 to 2 Cl<sup>-</sup> ions to generate a V<sub>m</sub> of ~-250 to ~-500 mV, as previously described (Kutzner et al., 2011). The V<sub>m</sub> was determined by the GROMACS utility gmx potential.

### ***Potential of mean force calculations***

To calculate the PMF for Na<sup>+</sup> ion translocation across m2r at neutral V<sub>m</sub>, umbrella sampling calculations were performed in bins of 0.25 Å and analysed with the GROMACS utility gmx wham. We used a simulation time of 50 ns in each window and harmonic potentials of 900–2000 kJ mol<sup>-1</sup> nm<sup>-2</sup> to restrain the Na<sup>+</sup> ion in the Z-direction. The standard deviation of the PMF profiles was estimated by using the Bayesian bootstrap method, as implemented in gmx wham, with 200 runs (See quantification and statistical analysis). The free energy of the Na<sup>+</sup> ion in bulk solution was set to 0. The position of the Na<sup>+</sup> ion (Z-coordinate) is reported relative to the D103<sup>3.32</sup>-C<sub>α</sub> atom (ligand binding site). These calculations were performed for the active conformation in both the apo and carbachol-bound state. In the latter case, D69<sup>2.50</sup> was modelled both in the negatively charged and neutral state with the Y440<sup>7.53</sup> sidechain in an upward or downward conformation.

### ***Gating charge calculations***

To calculate the gating charges, we followed a method previously described in Machtens et al., 2017 and Vickery et al., 2016a). A single bilayer of the active system was duplicated along the Z-axis, with one bilayer inverted (intracellular components of the receptors facing each other). The charge imbalance between compartments was then neutralised by adding ions. All protein atoms except hydrogen atoms were position-restrained using a spring constant of 1000 kJ mol<sup>-1</sup> nm<sup>-2</sup>, whilst the Na<sup>+</sup> ion was restrained with a force constant of

10,000 kJ mol<sup>-1</sup> nm<sup>-2</sup> due to its greater mobility. Bulk Na<sup>+</sup> ions were position-restrained on the Z-axis using a spring constant of 200 kJ mol<sup>-1</sup> nm<sup>-2</sup> to prevent their ingress into the receptor. The systems were calibrated using charge imbalances of -4 to 4; the slopes of the charge imbalance-voltage relationships indicate a near constant capacitance of the membrane/protein system under these conditions. The gating charges were inferred from the voltage differences for each ion position at a given charge imbalance. Errors were derived from the maximum and minimum slopes of the charge imbalance-voltage relationships. The hydrophilic channel was scanned by placing the ion at 2.5 Å intervals from the hydrophilic pocket to the intracellular solution and simulated for 50 ns, with the first 5 ns discarded (Fig S4 E). The gating charge calculated for each interval was taken as a direct measure of the voltage drop within the hydrated channel. This voltage drop, multiplied by the elementary charge  $e$  for a monovalent ion, was added to the equilibrium PMFs obtained by umbrella sampling (Fig S4 A-D), representing the excess free energy.

### ***Forcefield parameters***

For all MD simulations, the amber99sb\_ildn force field was used for the protein (Lindorff-Larsen et al., 2010), Berger parameters for lipids (Berger et al., 1997), which were adapted for use with the amber99sb force field (Cordomí et al., 2012), and the SPC/E model for water molecules (Berendsen et al., 1987). Water bond angles and distances were constrained by SETTLE (Miyamoto and Kollman, 1992) while all other bonds were constrained using the LINCS method (Hess et al., 1997). The temperature and pressure were kept constant throughout the simulations at 310 K and 1 bar, respectively, with the protein, lipids, and water/ions coupled individually to a temperature bath by the v-rescale method using a time constant of 0.2 ps and a semi-isotropic Berendsen barostat (Berendsen et al., 1984; Bussi et al., 2007). Employing a virtual site model for hydrogen atoms (Feenstra et al., 1999) allowed the use of 4-fs time steps during the simulation. All simulations were performed with the GROMACS software package, version 5.1.2 (Abraham et al., 2015).

### ***pK<sub>a</sub> calculations***

The pK<sub>a</sub> calculations were performed using a continuum electrostatics method, namely the Poisson-Boltzmann/Monte Carlo (PB/MC) approach, on multiple snapshots taken at a 2 ns interval from different umbrella sampling simulations in the carbachol-bound active state (both for Y440<sup>7.53</sup> in the upward and downward conformations) and in the apo active state. PB calculations were performed using MEAD (version 2.2.9)(Bashford and Gerwert, 1992)



with a dielectric constant ( $\epsilon_p$ ) of 4 for the protein and 80 for the solvent ( $\epsilon_w$ ), in the presence of an explicit membrane. Both D69<sup>2.50</sup> and D103<sup>3.32</sup> were included as titrating sites, together with Y440<sup>7.53</sup> and other nearby tyrosine residues. The temperature was set to 310 K and the ionic strength to 0.145 M. The same temperature was used for MC calculations ( $10^3$  steps in each calculation), which were performed using MCRP (Baptista et al., 1999). Each MC step consisted of a cycle of random choices of a state for all individual sites and pairs of sites with couplings above 2.0 pK<sub>a</sub> units (Baptista et al., 1999), whose acceptance/rejection followed a Metropolis criterion (Metropolis et al., 1953); tautomeric forms were not included.

### ***Residue conservation***

The GROMACS software package, version 5.0.4 analysis toolkit was used to identify residues with non-hydrogen heavy atoms within 4 Å of the sodium ion path during the simulations. The residue conservation profile of the amino acids was obtained from the GPCRdb server (Isberg et al., 2015).

## **QUANTIFICATION AND STATISTICAL ANALYSIS**

The standard deviation of the PMF profiles was estimated by using the Bayesian bootstrap method, as implemented in gmx wham, with 200 runs. Errors for the gating charge calculation were estimated by deriving the minimum and maximum slopes of the charge imbalance-voltage relationships, giving a mean and standard deviation for the capacitance of the membrane-patch system. This error was then propagated into the determination of the gating charges.

### **KEY RESOURCES TABLE**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
<b>Antibodies</b>		
N/A	N/A	N/A
<b>Bacterial and Virus Strains</b>		
N/A	N/A	N/A
<b>Biological Samples</b>		
N/A	N/A	N/A
<b>Chemicals, Peptides, and Recombinant Proteins</b>		

N/A	N/A	N/A
Critical Commercial Assays		
N/A	N/A	N/A
Deposited Data		
N/A	N/A	N/A
Experimental Models: Cell Lines		
N/A	N/A	N/A
Experimental Models: Organisms/Strains		
N/A	N/A	N/A
Oligonucleotides		
N/A	N/A	N/A
Recombinant DNA		
N/A	N/A	N/A
Software and Algorithms		
GROMACS	(Abraham et al., 2015)	<a href="http://www.gromacs.org/">http://www.gromacs.org/</a>
MEAD	(Bashford and Gerwert, 1992)	<a href="http://www.bisb.uni-bayreuth.de/People/ullmann/">http://www.bisb.uni-bayreuth.de/People/ullmann/</a>
AMBER16	(Case et al., 2016)	<a href="http://ambermd.org/">http://ambermd.org/</a>
GOLD	(Willett and Glen, 1995)	<a href="https://www.ccdc.cam.ac.uk/solutions/csd-discovery/components/gold/">https://www.ccdc.cam.ac.uk/solutions/csd-discovery/components/gold/</a>
MCRP	(Baptista et al., 1999)	<a href="http://www.itqb.unl.pt/labs/molecular-simulation">http://www.itqb.unl.pt/labs/molecular-simulation</a>
Other		
N/A	N/A	N/A