



University of Dundee

Prevalence of Sleep-Disordered Breathing and Associations With Malocclusion in Children

Aroucha Lyra, Maria Carlla; Aguiar, Débora; Paiva, Mabel; Arnaud, Manuela; Alencar Filho, Arnaldo; Rosenblatt, Aronita

Published in:
Journal of Clinical Sleep Medicine

DOI:
[10.5664/jcsm.8370](https://doi.org/10.5664/jcsm.8370)

Publication date:
2020

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Aroucha Lyra, M. C., Aguiar, D., Paiva, M., Arnaud, M., Alencar Filho, A., Rosenblatt, A., Thérèse Innes, N. P., & Heimer, M. V. (2020). Prevalence of Sleep-Disordered Breathing and Associations With Malocclusion in Children. *Journal of Clinical Sleep Medicine*, 16(7), 1007-1012. <https://doi.org/10.5664/jcsm.8370>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Prevalence of sleep-disordered breathing and associations with malocclusion in children

**Maria Carlla Aroucha Lyra^{1,¶,*}, Débora Aguiar^{1,&}, Mabel Paiva^{1,&}, Manuela Arnaud^{1,&},
Arnoldo Alencar Filho^{1,&}, Aronita Rosenblatt^{1,¶} Nicola Patricia Thérèse Innes^{2,¶,#b},
Mônica Vilela Heimer^{1,¶,#a}**

1. Department of Pediatric Dentistry, University of Pernambuco, Camaragibe, Pernambuco, Brazil.
2. School of Dentistry, University of Dundee, Park Place Dundee, DD14HR, Scotland.

*Corresponding author

Email: mcaryllyra@gmail.com (MC)

#a Current Address: Department of Pediatric Dentistry, University of Pernambuco, Camaragibe, Pernambuco, Brazil.

#b Current Address: School of Dentistry, University of Dundee, Park Place Dundee, DD14HR, Scotland

¶ The author and co- authors contributed equally to this work

This article has been accepted for publication in the *Journal of Clinical Sleep Medicine* published by the American Academy of Sleep Medicine [<https://jcs.m.aasm.org/journal/jcs.m>].

ABSTRACT

Objective: This study aimed to determine the prevalence of sleep-disordered breathing (SDB) and its association with malocclusion among children in Recife, Brazil. Methods: 390 children aged seven to eight years took part in the study, comprised by the body mass measurement, orthodontic examination and parent's information required by the Sleep Disturbance Scale for Children. Statistics tools were Pearson's chi-square and Lemeshow test. Results: SDB was found in 33.3% of the children and associated with overjet ($p= 0.007$), anterior open bite ($p=0.008$) and posterior crossbite ($p= 0.001$). There was no association between BMI and SDB. The multivariate logistic regression model indicated that the anterior open bite ($p= 0.002$) and posterior crossbite ($p = 0.014$) have an association with SDB. Conclusions: Results of this study indicated that the prevalence of SDB was high and highly associated with malocclusion; anterior open bite and posterior crossbite are risk factors for SDB.

Keywords: Sleep-disordered Breathing; Child; Prevalence; Malocclusion, Body Weight

INTRODUCTION

Scientific investigation on Sleep disorders usually reports to insomnia or excessive daytime sleepiness in adulthood and its negative impacts on normal physiological functioning and quality of life. However, very little is written on the prevalence of the condition in childhood, even though it impacts for 20 to 30% of problems reported by children's health providers^{1,2}. Inadequate sleep can lead to diverse comorbidities, such as delayed growth as well as cardiovascular, immunological and metabolic conditions, which can affect a child's health and quality of life over the years³. Among the primary risk factors, the body mass index (BMI) is an important predictor of the clinical intensity of sleep disorders³.

Sleep-disordered breathing (SDB) includes a spectrum of upper airway disorders ranging from primary snoring (PS) to obstructive sleep apnea (OSA)⁴. PS is a respiratory noise, in which sleep architecture, alveolar ventilation, and blood oxygen saturation are maintained at normal values whereas OSA is partial or complete obstruction of the upper airways that impairs normal ventilation during sleep and normal sleep patterns⁵. The prevalence of SDB in children ranges from 0.7% to 16.9%; however, in the current literature, PS ranges from 4.3 to 16.9% and OSA from 0.7 to 3%^{6,7,8,9}.

Thus, abnormal craniofacial morphology, such as retrognathia and dental malocclusion contribute to upper airway obstruction and increase the risk of SDB^{6,10}. Therefore, SDB is commonly associated with oro and dentofacial features and may be related to malocclusion. Failure to diagnose SDB in adults may have undesirable health outcomes¹¹ which may require further education for both parents and providers of how occlusal abnormalities associates with SDB^{12,13}.

Sleep-related issues often are not discussed during well-child visits, because of parents not raising the issue and primary care providers not asking about symptoms of sleep disorders. Sleep disorders are underdiagnosed in primary care practices⁷.

Nevertheless, oral screening in children and the use of a validated questionnaire could improve early diagnosis and reduce the under-diagnosis of sleep disorders¹⁴.

This study aims to evaluate the prevalence of SDB and its relationship with malocclusion in Brazilian children.

METHODS

A cross-sectional study, conducted with aged seven and eight year's old children, attending local public schools in the city of Recife, Brazil.

The study was carried out following the Internal research Board of the University of Pernambuco by the record number CAAE 53308915.8.0000.5207 and followed the Resolution 466/2012 of National Commission of Ethics in Research and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards, including Informed consent from all study participants. It also followed the STROBE protocol for observational studies (Additional file 1).

The sample size was calculated using the formula for cross-sectional studies in Epi Info Version 2000 (Atlanta, Georgia, USA), based on Schlarb et al.¹, which is 20% prevalence rate for SDB, 95% confidence interval, 5% margin of error, a 1.5 design effect and the addition of 20% to make up for possible dropouts, a total of 435 children. A two-stage sampling method allowed the representativeness of the sample. The first stage was the randomization of public schools in each administrative district. In the second stage, classes were assigned by random computer number and all children attending those classes were invited to take part in the study.

The physical assessment comprised the measurement of body mass index (BMI), based on height (measured using a stadiometer Tonelli®) with a 2-m capacity and precision of 0.1 cm) and weight (measured using a digital scale Camry®, model EB9013, Brazil) with a capacity for 150 kg and a precision of 100 g). Children in the 85th to 97th percentile for age were classified as overweight, and those above the 97th percentile were classified as obese, as recommended by the World Health Organization (WHO)¹⁵. For the orthodontic assessment, a single experienced orthodontist examined the children¹⁶, as follows.

Sagittal analysis

The molar relationship followed Angle's classification¹⁷, based on the comparison of the anteroposterior diameter of the mandible and maxilla and the relationship between the permanent first molars.

Overjet, the projection of the maxillary incisors beyond the lower incisors on the horizontal plane, measured by using a Community Periodontal Index probe (CPI), placed perpendicular to the occlusal plane, on the buccal surface of the mandibular incisors¹⁸. A

measurement of 0 mm classifies as an edge to edge bite; 1 to 3 mm normal; 4 to 6 mm, moderately increased, and higher than 6 mm, severely increased. In this report, values equal or less than 3 mm, are not classified as overjet; for values higher than are 3mm , classified as overjet. When the maxillary incisors occlude behind the mandibular incisors, negative overjet, then recorded as anterior crossbite ^{19,20}.

Vertical analysis

Overbite is the overlap of the upper teeth along the lower teeth on the vertical plane¹⁶. The distance (in mm) between the incisal edges, measured using a CPI probe¹⁸. A measurement of 0 mm (incisal edges of maxillary incisors in contact with incisal edges of mandibular incisors) classifies as an edge to edge ^{19,20}; 1 to 3 mm classifies as normal; 4 to 6 mm, moderately increased, and higher than 6mm, severely increased. In the present study, values smaller or equal to 3 mm did not indicate overbite, when higher than 3 mm, record overbite. A lack of contact between the incisal edges of the maxillary and mandibular teeth was considered anterior open bite ^{16,19,20}.

Transversal analysis

Posterior crossbite, defined as a labiolingual relationship between the maxillary and mandibular teeth, when out of the normal standards, Inverted.¹⁶ this is classified as present, when involving one or more molars in the posterior region. Posterior crossbite, classified as unilateral (involving only one side) or bilateral (involving both sides)²¹.

The orthodontic chart used in this assessment, also meant to record Angle's classification (class I, II or III), overjet, overbite, anterior open bite, anterior crossbite, and posterior crossbite.

The sociodemographics recorded: sex; marital and employment status of the caregiver, and the Sleep Disturbance Scale for Children (SDSC).

SDB was evaluated using the SDSC (for three to 18 years of age), validated for Brazil²², and filled by parents or caregiver, about the child's sleep habits in the previous six months. The scale has 26 items divided among six subscales: Disorders of Initiating and Maintaining Sleep, Sleep Breathing Disorders, Disorders of Arousal, Sleep-Awake Transition Disorders, Disorders of Excessive Somnolence and Sleep Hyperhydrosis. The SDSC items to detect symptoms of SDB were: (1) "the child has difficulty in breathing at night"; (2) "the child

gasps for breath or is unable to breathe during sleep”; and (3) “the child snores”. Answers, recorded on a Likert scale of 1–5: 1 point for “never”, 2 points for “occasionally” (once or twice per month), 3 points for “sometimes” (once or twice per week), 4 points for “often” (3–5 times per week), and 5 points for “always” (daily). Hence, the sum score for the three questions can be at least 3 and at most 15. According to the total score obtained for these three SDSC items, participants were classified into 3 groups: group 1, at high risk of SDB (12–15 points); group 2, at moderate risk of SDB (7–11 points); and group 3, at low risk of SDB (3–6 points). This variable, dichotomized in children without SDB (group 1 and 2) and children with SDB (group 3)¹⁶. This study also evaluated the BMI of children, since according to Kaditis et al²³ this variable is an important factor for SDB.

Data analysis was performed with the Statistical Package for the Social Sciences (SPSS, version 17.0) by the Pearson’s chi-square test and Lemeshow test, with a 5% margin of error and 95% confidence interval.

RESULTS

A total of 390 out of the 435 students (89.7%) had physical, and orthodontic examination and parents returned the questionnaires. The losses were 45 students (10.3%) that did not complete the questionnaires or did not attend school on the examination days. The majority of the children was seven years of age (n=212; 54.4%), and males accounted for 53.8% (n=210) of the sample. Among the caregivers, the majority was female (n= 363; 93.1%), married or in a stable relationship (n= 238; 61.0%), who had completed high school or university education (n= 214; 54.9%), n = 242; 62.1% were employed. Table 1 shows to the frequency of malocclusion, body mass index and SDB.

Table 1 – Occurrence of malocclusion, SDB and BMI status in participating children

<i>Variable</i>	N	%
TOTAL	390	100.0
Occurrence of malocclusion		
Yes	253	64.9
No	137	35.1
Molar Relationship		
Class I	224	57.4
Class II	91	23.3
Class III	75	19.2
Overjet		
Present	91	23.3
Absent	299	76.7
Overbite		
Present	35	9.0
Absent	355	91.0
Anterior Open bite		
Present	68	17.4
Absent	322	82.6
Posterior Crossbite		
Present	41	10.5
Absent	349	89.5
Anterior Crossbite		
Present	28	7.2
Absent	362	92.8
Sleep-disordered breathing		
Present	130	33.7
Absent	260	66.7
BMI		
Overweight / obese	144	36.9
Ideal range	246	63.1

Among the orthodontic variables, overjet, anterior, and posterior crossbite were significantly associated with the SDB (TABLE 2). A logistic regression model with the variables that showed a significant association up to 20% ($p < 0.20$) in the bivariate study (overjet, anterior open bite. Posterior crossbite) was used to verify which variables influenced

the presence of SDB. In the multivariate logistic regression analysis, posterior crossbite and open bite were considered risk factors for SDB (TABLE 3).

Table 2 – Occurrence of SDB according to malocclusions and BMI status

Variable	Sleep Breathing Disorders				Total		p-value	OR (95% CI)
	Present		Absent					
	N	%	n	%	N	%		
Total Group	130	33.3	260	66.7	390	100.0		
Molar Relationship							p ⁽¹⁾ = 0.559	
Class I	70	31.3	154	68.8	224	100.0		1,00
Class II	34	37.4	57	62.6	91	100.0		1.31 (0.79 a 2.19)
Class III	26	34.7	49	65.3	75	100.0		1.17 (0.67 a 2.03)
Overjet							p ⁽¹⁾ = 0.007*	
Present	41	45.1	50	54.9	91	100.0		1.93 (1.19 a 3.13)
Absent	89	29.8	210	70.2	299	100.0		1.00
Overbite							p ⁽¹⁾ = 0,616	
Present	13	37.1	22	62.9	35	100.0		1.20 (0.58 a 2.47)
Absent	117	33.0	238	67.0	355	100.0		1.00
Anterior Open bite							p ⁽¹⁾ = 0.008*	
Present	32	47.1	36	52.9	68	100.0		2.03 (1.19 a 3.46)
Absent	98	30.4	224	69.6	322	100.0		1.00
Posterior Crossbite							p ⁽¹⁾ = 0.001*	
Present	23	56.1	18	43.9	41	100.0		2.89 (1.59 a 5.58)
Absent	107	30.7	242	69.3	349	100.0		1.00
Anterior Crossbite							p ⁽¹⁾ = 0.488	
Present	11	39.3	17	60.7	28	100.0		1.32 (0.60 a 2.91)
Absent	119	32.9	243	67.1	362	100.0		1.00
BMI							p ⁽¹⁾ = 0.975	
Obese	31	34.1	60	65.9	91	100.0		1.05 (0.63 a 1.75)
Overweight	18	34.0	35	66.0	53	100.0		1.05 (0.56 a 1.96)
Ideal range	81	32.9	165	67.1	246	100.0		1.00

(*) Significant association at 5% level // (1) Pearson's chi-square test

Table 3 - Multivariate logistic regression analysis of the association between SDB and overjet, anterior open bite and posterior crossbite

Variable	Bivariate		Adjusted	
	OR and 95% CI	p - value	OR and 95% CI	p - value
Overjet		0.007*		0.402
Present	1.93 (1.19 a 3.13)		1.29 (0.71 a 2.36)	
Absent	1.00		1.00	
Anterior Open bite		0.008*		0.002*
Present	2.03 (1.19 a 3.46)		2.34 (1.36 a 4.03)	
Absent	1.00		1.00	
Posterior Crossbite		0.001*		0,014*
Present	2.89 (1.59 a 5.58)		2.79 (1.24 a 6.31)	
Absent	1.00		1.00	

(*) Significant association at 5% level // Lemeshow Test

DISCUSSION

Among the 390 children, in this study, 65% presented malocclusion; the result is in agreement with previous studies, which report prevalence rates ranging from 60 to 70%. Angle class I was the most prevalent, followed by classes II and III, which is also in agreement with the previous studies ^{24,25}, conducted in different countries, which report a high prevalence of malocclusion ^{26,27}, that impact the functional, emotional and social aspects of children's life ^{16,28}.

It merits consideration that the present study provides data on the prevalence of malocclusion among children with SDB, which is not a piece of trivial information. According to other studies, untreated SDB in childhood could have a negative impact in adulthood, with

an increase in health-related quality of life^{3,6,14,29}. As described in latter reports, Orthodontic intervention in the mixed dentition may prevent or arrest incipient malocclusions, contributing to the harmonious growth of the basal bones and to the normal development of the occlusion, reducing the chances of deleterious disorders in the permanent dentition.^{3,6,14}

SDB may impact public health throughout the world, once it affects millions of individuals, mainly children³⁰. SDB often more associated with occlusal abnormalities than other sleep disorders, which may be explained by the abnormal development of malocclusions caused by the breathing pattern, facial muscle balance, and skeletal muscles³⁰. This condition can develop into mild to more severe obstructive sleep apnea in a relatively short period in cases of a lack of effective therapy. Adults presenting with SDB seem to have poor dental occlusions and anomalous craniofacial morphology, which may raise a question of whether orthodontic treatment in childhood could prevent SDB in adulthood^{31,32}. Early diagnosis of SDB in childhood and provision of appropriate treatment may not only treat or prevent medium-term complications such as learning difficulties but may potentially prevent long term cardiovascular complications³³.

Among the occlusal features shown in the present study, overjet, posterior crossbite, and open bite significantly associate with SDB, which is in agreement with data in a previous study³⁴. The logistic regression analysis in the present investigation revealed that both posterior crossbite and open bite were risk factors for sleep- disorders breathing. The present findings indicate the importance of the early diagnosis of posterior crossbite, which seems to be the type of malocclusion most associated with sleep-disordered breathing^{24,31,34} followed by anterior open bite^{35,36}. Those may lead to an abnormal breathing pattern, which seems to alter the oral and facial muscular balance and are likely to affect skeletal and occlusal development in children³¹.

In this group of children, 36.9% were overweight/obese. The latter is in agreement with previous studies that report a high prevalence of overweight and obesity among children and adolescents aged two to 19 years³¹. Concerning SDB, in this study, there was no significant association with overweight/obese. Likewise, a previous study also found no association between overweight and SDB and, similar to our findings, individuals with craniofacial abnormalities were more susceptible to developing sleep-disorders breathing than individuals with excess weight³⁷.

Evidence suggests that obesity plays an important role in the pathophysiology of SDB in adults³⁸and, to a lesser extent, in children³⁹. Thus, more knowledge about the interactions between obesity and SDB is relevant when considering the increases in the rates of obesity in childhood^{23,40}.

SDB represents a continuum of manifestations from simple snoring to Obstructive sleep apnea (OSA), and this condition tends to progress from mild SDB to severe OSA over a varying period, which may be surprisingly short in the case of weight gain and the lack of effective treatment^{31,41}.

In children, adenotonsillar hypertrophy, as well as deviations concerning craniofacial morphology and dental occlusion, are risk factors for SDB, which seems to be complex, multifactor pathogenesis in children^{31,37}. Thus, timely orthodontic intervention could prevent SDB in adulthood^{11,16}.

The main limitation of the present report is the study design, which does not allow the determination of cause-and-effect relationships between malocclusion and sleep-disorders breathing. Thus, longitudinal design studies are needed for more precise diagnosis purposes.

CONCLUSIONS

In the study population, the prevalence of SDB was high and associated with malocclusion. Posterior crossbite and anterior open bite were found to be risk factors for SDB.

Acknowledgements

The authors of this research are grateful to the children and their families who have agreed to participate and CAPES, Brazilian Ministry of Education, for funding.

Contributions

MCA Lyra: conceptualization, data curation, writing (original draft preparation) and writing (review & editing) of the manuscript; D Aguiar, MC Paiva, M Arnaud and A Vasconcelos: data curation and visualization; A Rosenblatt, NPT Innes and MV Heimer: conceptualization, supervision and writing (review & editing). All authors read and approved the final manuscript and agree to be accountable for it.

References

1. Schlarb A, Gulewitsch MD, Weltzer V, Ellert U, Enck P. Sleep duration and sleep problems in a representative sample of German children and adolescents. *Health*. 2015; 7:1397-1408.
2. Sadeh A, Mindell J, Rivera L. "My child has a sleep problem": a cross-cultural comparison of parental definitions. *Sleep Med*. 2011; 12: 478-482.
3. Blunden S, Chervin R. Sleep, performance and behaviour in Australian indigenous and non-indigenous children: An exploratory comparison. *J Paediatr Child Health*. 2010; 46: 10-16.
4. Marcus C, Brookers L, Draper K, et al. Diagnosis and management of childhood obstructive sleep apnea syndrome. *Pediatrics*. 2012; 130: 714-755.
5. Lee C, Kang K, Weng W, Lee P, Hsu W. Quality of life after adenotonsillectomy for children with sleep-disordered breathing: a linear mixed model analysis. *Int J Pediatr Otorhinolaryngol*, 2014; 78(8), 1374-1380.
6. Kim E, Choi J, Kim Y, et al. Upper airway changes in severe obstructive sleep apnea: upper airway length and volumetric analyses using 3D MDCT. *Acta Otolaryngol*. 2011; 131: 527-532.
7. Meltzer L, Johnson C, Crosette J, Ramos M, Mindell JA. Prevalence of diagnosed sleep disorders in pediatric primary care practices. *Pediatrics*. 2010;125: 1410-8.
8. Bixler E, Vgontzas A, Lin H, Liao D, Calhoun S, Vela-Bueno A, et al. Sleep disordered breathing in children in a general population sample: prevalence and risk factors. *Sleep*. 2009; 32:731-6.
9. Seo W, Koo B, Kim M, Rho Y, Sung H, Shin J. Preliminary study of children's sleep problems in an elementary school in Daegu. *J Korean Acad Child Adolesc Psychiatry*. 2008;19:156-61.
10. Mallampati S, Gatt S, Gugino L, Desai SP, Waraksa B, Freiburger D, Liu PL. A clinical sign to predict difficult tracheal intubation; a prospective study. *Can Anaesth Soc J*. 1985; 32: 429-434.
11. Fernandes A, Novais A, Feitosa B, et al. Children's sleep: is it given its due importance? *Rev. Assoc Med. Minas Gerais*. 2014; 24: 323-326.
12. Guilleminault C, Lee J, Chan A. Pediatric obstructive sleep apnea syndrome. *Arch Pediatr Adolesc Med*. 2005; 159:775-785.
13. Huynh N, Morton P, Rompré P, Papadakis A, Remise C. Associations between sleep-disordered breathing symptoms and facial and dental morphometry, assessed with screening examinations. *Ajo-do*. 2011; 140: 762-770.
14. De Oliveira H, De Melo A, Couto J, Drummond S, Casali C, Leite R, Sampaio W. Características orofaciais e risco para apneia obstrutiva do sono em hipertensos. *R Enferm Cent O Min*. 2013; 3:706-713.
15. Organização Mundial De Saúde. Orientações para coleta e análise de dados antropométricos em serviços de saúde: norma técnica do sistema de Vigilância Alimentar e Nutricional – SISVAN. IOP; Publishing BVSMS; http://bvsm.s.saude.gov.br/bvs/publicacoes/orientacoes_coleta_analise_dados_antropometricos.pdf; 2011; Accessed 08 June 2018.

16. Carvalho F, Lentini-Oliveira D, Carvalho G, Prado L, Prado G, Carvalho L. Sleep-disordered breathing and orthodontic variables in children—pilot study. *Int J Pediatr Otorhinolaryngol.* 2014; 78: 1965-1969.
17. Moyers R. *Classificação e terminologia da má-oclusão.* Rio de Janeiro, RJ. Guanabara Koogan; 1987.
18. Coordenação Nacional de Saúde Bucal; Projeto SB 2010: Condições de Saúde Bucal da População Brasileira no ano 2010; Manual de Calibração de examinadores. IOP Publishing DAB. http://dab.saude.gov.br/CNSB/sbbrasil/arquivos/SBBrasil2010_Manual_Calibracao.pdf. Accessed 08 August 2017
19. Björk A, Krebs A, Solow B. A Method for Epidemiological Registration of Malocclusion. *Acta Odontol Scand.* 1964; 22: 27-41.
20. Thilander B, Pena L, Infante C, Parada S, Mayorga C. Prevalence of malocclusion and orthodontic treatment need in children and adolescents in Bogota, Colombia. An epidemiological study related to different stages of dental development. *Eur J Orthod.* 2001; 23:153-168.
21. Gandini M, Pinto A, Gandini Junior L, Martins J, Mendes A. Estudo da oclusão dentária de escolares da cidade de Araraquara, na fase da dentadura mista relação inte-arcos, região anterior (overjet e overbite). *Ortodontia.* 2000; 33: 44-49.
22. Ferreira V, Carvalho L, Ruotolo F, Morais J, Prado L, Prado G. Sleep disturbance scale for children: translation, cultural adaptation, and validation. *Sleep Med.* 2009; 10: 457-463.
23. Kaditis AG, Alvarez ML, Boudewyns A, Alexopoulos EI, Ersu R, Joosten K, et al. Obstructive sleep disordered breathing in 2-to 18-year-old children: diagnosis and management. *European Respiratory Journal.* 2016; 47: 69-94.
24. Sauer C, Schluter T, Hinz R, Gesch D. Childhood obstructive sleep apnea syndrome: an interdisciplinary approach: a prospective epidemiological study of 4,318 five-and-a-half-year-old children. *J Orofac Orthop.* 2012; 73:342-358.
25. Garbin A, Perin P, Garbin C, Lolli L. Malocclusion prevalence and comparison between the Angle classification and the Dental Aesthetic Index in scholars in the interior of São Paulo state – Brazil. *Dental Press J Orthod.* 2010; 15: 94-102.
26. Ibrahim A, Abuaffan A. Prevalence of malocclusion and orthodontic treatment needs among down syndrome sudanese individuals. *Braz Dent Sci.* 2015; 18: 95-101.
27. De Freitas C, Souza J, Mendes D, Pordeus I, Jones K, Martins A. Necessidade de tratamento ortodôntico em adolescentes brasileiros: avaliação com base na saúde pública. *Revista Paulista de Pediatria.* 2015; 33: 204-210.
28. Guimarães S, Jorge K, Fontes M, Ramos-Jorge M, Araújo C, et al. Impact of malocclusion on oral health-related quality of life among schoolchildren. *Brazilian oral research.* 2018; 32.
29. Nixon G, Brouillette R. Paediatric obstructive sleep apnoea. *Thorax.* 2005; 60: 511-516.
30. Wei J, Mayo M, Smith H, Reese M, Weatherly R. Improved behavior and sleep after adenotonsillectomy in children with sleep disordered breathing. *Arch Otolaryngol Head Neck Surg.* 2007; 133: 974-979.
31. Ikavalko H, Tuomilehto R, Pahkala T, Tompuri T, Laitinen T, Myllykangas R, Vierola A, Lindi V, Narhi M, Lakka T. Craniofacial morphology but not excess body fat is associated with risk of having sleep-disordered breathing the PANIC Study (a questionnaire-based inquiry in 6–8-year-olds). *Eur J Pediatr.* 2012; 171: 1747-1752.

32. Pahkala R, Puustinen R, Tuomilehto H, Ahlberg J, Seppä J. Risk factors for sleep disordered breathing: the role of craniofacial structure. *Acta Odontol Scand.* 2011; 69:137–143.
33. Stauffer J, Okuji D, Lichty I, Gc B, Whyte F, Miller D, Hussain, J. A review of pediatric obstructive sleep apnea and the role of the dentist. *J Dent Sleep Med,* 2018; 5: 111-130.
34. Pacheco M, Fiorott B, Finck N, Araújo M. Craniofacial changes and symptoms of sleep-disordered breathing in healthy children. *Dental press J of Orthod.* 2015; 20: 80-87.
35. Lofstrand Tidestrom B, Hultcrantz E. The development of snoring and sleep related breathing distress from 4 to 6 years in a cohort of Swedish children. *Int. J. Pediatr. Otorhinolaryngol.* 2007; 71 (7): 1025–1033.
36. Lundeborg I, McAllister A, Graf J, Ericsson E, Hultcrantz E, Oral motor dysfunction in children with adenotonsillar hypertrophy-effects of surgery. *Logoped Phoniatr Vocol.* 2009; 28: 1–6.
37. Hultcrantz E, Tideström B. The development of sleep disordered breathing from 4 to 12 years and dental arch morphology. *Int J Pediatr Otorhinolaryngol.* 2009; 73:1234-1241.
38. Bonsignore MR, McNicholas WT, Montserrat JM, et al. Adipose tissue in obesity and obstructive sleep apnoea. *Eur Respir J.* 2012; 39:746–67.
39. Gozal D, Kheirandish-Gozal L. Childhood obesity and sleep: relatives, partners, or both?--a critical perspective on the evidence. *Ann N Y Acad Sci.* 2012; 1264:135–41.
40. Alonso-Alvarez ML, Cordero-Guevara JA, Teran-Santos J, et al. Obstructive sleep apnea in obese community-dwelling children: the NANOS study. *Sleep.* 2014; 37: 943–949.
41. Berger G, Berger R, Oksenberg A (2009) Progression of snoring and obstructive apnoea: the role of increasing weight and time. *Eur Respir J.* 2009; 33:338–345