



**University of Dundee**

## **NDP-rhamnose biosynthesis and rhamnosyltransferases**

Wagstaff, Ben A.; Zorzoli, Azul; Dorfmueller, Helge C.

*Published in:*  
Biochemical Journal

*DOI:*  
[10.1042/BCJ20200505](https://doi.org/10.1042/BCJ20200505)

*Publication date:*  
2021

*Licence:*  
CC BY

*Document Version*  
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

*Citation for published version (APA):*

Wagstaff, B. A., Zorzoli, A., & Dorfmueller, H. C. (2021). NDP-rhamnose biosynthesis and rhamnosyltransferases: building diverse glycoconjugates in nature. *Biochemical Journal*, 478(4), 685-701. <https://doi.org/10.1042/BCJ20200505>

**General rights**

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# NDP-Rhamnose Biosynthesis and Rhamnosyltransferases - Building Diverse Glycoconjugates in Nature

Wagstaff, B. A.<sup>1,2</sup>, Zorzoli, A.<sup>2</sup>, Dorfmüller, H. C.<sup>2</sup>

From the <sup>1</sup>Manchester Institute of Biotechnology, University of Manchester, 131 Princess Street, Manchester, United Kingdom and the <sup>2</sup>Division of Molecular Microbiology, School of Life Sciences, University of Dundee, Dundee, DD1 5EH, United Kingdom.

E-mail: h.c.z.dorfmüller@dundee.ac.uk

**Keywords:** rhamnose, biosynthesis, glycosyltransferase, glycoproteins, polysaccharides, pathogens

## Abstract

Rhamnose is an important 6-deoxy sugar present in many natural products, glycoproteins, and structural polysaccharides. Whilst predominantly found as the L-enantiomer, instances of D-rhamnose are also found in nature, particularly in the Pseudomonads bacteria. Interestingly, rhamnose is notably absent from humans and other animals, which poses unique opportunities for drug discovery targeted towards rhamnose utilising enzymes from pathogenic bacteria. Whilst the biosynthesis of nucleotide-activated rhamnose (NDP-rhamnose) is well studied, the study of rhamnosyltransferases that synthesise rhamnose-containing glycoconjugates is the current focus amongst the scientific community.

In this review, we describe where rhamnose has been found in nature, as well as what is known about TDP-β-L-rhamnose, UDP-β-L-rhamnose, and GDP-α-D-rhamnose biosynthesis. We then focus on examples of rhamnosyltransferases that have been characterized using both *in vivo* and *in vitro* approaches from plants and bacteria, highlighting enzymes where 3D structures have been obtained. The ongoing study of rhamnose and rhamnosyltransferases, in particular in pathogenic organisms, is important to inform future drug discovery projects and vaccine development.

## Background

Rhamnose (Rha) is a 6-deoxy hexose sugar found widely throughout nature in plant and microbial natural products (1), including glycoproteins (2) and structural polysaccharides (3). The importance of these glycoconjugates in biological processes leads to rhamnose often being an essential sugar for cell viability in many organisms. Although the L-enantiomer of rhamnose is by far the most abundant throughout nature, D-Rha has been found extensively throughout the *Pseudomonas* genus (4) as a constituent of their surface exposed lipopolysaccharide (5). Animal production of biological compounds based on L-rhamnose have not been found. Thus, the microbial biosynthesis pathway for L-Rha production has

previously been targeted as a potential therapeutic drug target to fight bacterial infections (3, 6).

The biosynthesis pathways of the nucleotide-activated forms of L- and D-Rha differ depending on the enantiomer and the domain of life. L-Rha is biosynthesized as two variants, thymidine 5'-diphospho- $\beta$ -L-rhamnose (TDP- $\beta$ -L-rhamnose) or uridine 5'-diphospho- $\beta$ -L-rhamnose (UDP- $\beta$ -L-rhamnose). D-Rha is much less common and is biosynthesized as GDP- $\alpha$ -D-rhamnose, using a different set of enzymes to those utilised in the NDP- $\beta$ -L-rhamnose biosynthesis pathways. Bacteria (7), some microalgae (8), and archaea (9), have been shown to primarily produce TDP- $\beta$ -L-rhamnose, whilst the biosynthesis of UDP- $\beta$ -L-rhamnose is mainly observed in plants or other eukaryotes such as fungi and algae, on large, multifunctional enzymes (10, 11). GDP- $\alpha$ -D-rhamnose has been observed in the *Pseudomonas* genus (12), *Aneurinibacillus thermoaerophilus* (13), and also the giant algal-infecting virus, *Paramecium bursaria Chlorella virus 1* (14).

Following nucleotidyl-activation, NDP-Rha becomes a substrate for diverse rhamnosyltransferases that transfer rhamnose onto their glycoconjugate products. Many of the functionally characterized rhamnosyltransferases are responsible for the transfer of L-Rha to bacterial LPS (15-18), but other rhamnosyltransferases that transfer rhamnose to plant natural products have also been described (19-21). In this work, we review the presence of rhamnose in glycoconjugates across the domains of life and viruses, with a particular focus on Gram-negative and Gram-positive pathogenic bacteria, which are classified by their rhamnose-rich surface carbohydrate virulence factors, where rhamnose is abundant and found with a variety of linkages. We also review the rhamnosyltransferases that have been characterised structurally and highlight some limitations that currently hinder the progress towards the elucidation of other rhamnosyltransferases from a functional and structural perspective.

## **Presence of Rhamnose in Nature**

### **Eukaryotes**

Although notably absent from animals and mammals, rhamnose plays a crucial role for plants and a number of lower eukaryotes. Arguably, the most prominent rhamnose glycoconjugate found in land plants is the pectic polysaccharide rhamnogalacturonan-I (RG-I). In this glycoconjugate,  $\alpha$ -L-Rha forms the backbone repeating unit of 4)- $\alpha$ -D-GalpA-(1,2)- $\alpha$ -L-Rhap-(1, and is the branching point for the sidechains (Figure 1). Rhamnose is therefore an essential sugar to the polysaccharide structure. Pectic polysaccharides like RG-I, along with other polysaccharides, constitute the major component of the plant cell wall, and are crucial for cell wall flexibility and stability, but also essential for growth and plant development (22). As well as RG-I, rhamnose frequently decorates flavonoids such as kaempferol and quercetin, often found in the flowering tissues of plants (23). The glycosylation of these flavonoids is believed to prevent self-toxicity; importantly, when flavonol glycosylation is genetically blocked, some plants down-regulate flavonol aglycone biosynthesis (24).

The presence of rhamnose in fungal glycoconjugates is less common, but the filamentous human pathogen *Pseudallescheria boydii*, as well as the plant pathogen, *Rhynchosporium secalis* (25), produce a distinct rhamnose-containing cell wall polysaccharide termed 'peptido-rhamno-mannan', which has been proposed as a potential diagnostic marker due to its rarity (26). Other eukaryotes such as algae are known to biosynthesise nucleotide-activated rhamnose (8) and produce prominent rhamnose-containing polysaccharides. Worth mentioning is the sulphated structural polysaccharide, ulvan, from the green 'sea lettuce' of the *Ulva* genus. It consists of either 4)- $\beta$ -D-GlcA-(1-4)- $\alpha$ -L-Rha-3S-(1 or alternatively 4)- $\alpha$ -D-IdoA-(1-4)- $\alpha$ -L-Rha-3S-(1 repeating units and has been found to display useful pharmaceutical or nutraceutical properties (27). It has even been recommended as a food source to improve the egg laying ability of hens (28). Rhamnose has also been identified to be a major component of other algal polysaccharides, but their chemical structures and carbohydrate compositions have not been yet elucidated (29, 30). Rhamnose and rhamnose biosynthesis genes and subsequent enzymes are absent in higher eukaryotes, including mammals. This uniqueness suggests that these family of enzymes that are mechanistically distinct to any proteins found in the hosts are potential drug targets in human and veterinary pathogens.

## Prokaryotes

Whilst less common in eukaryotes, rhamnose appears to be ubiquitously incorporated into surface carbohydrates amongst both Gram-negative and Gram-positive bacteria. The Gram-negative O-antigen is often the exclusive component that incorporates rhamnose, with structural diversity arising from the linkage types, repeating units, and the substitution of the core backbone. As well as being found in the *E. coli* O-antigens, L-Rha makes up the major component of the surface carbohydrates in the pathogenic *Shigella* O-antigens. *Shigella* are composed of four subgroups (*S. flexneri*, *S. sonnei*, *S. dysenteriae*, and *S. boydii*), which are further categorised into serotypes; these serotypes are identified based on the carbohydrate structure of the lipopolysaccharide repeat units of the O-antigen. Rhamnose is often the major component of these repeating units and forms the backbone of them, with carbohydrate diversity arising in the glycosidic linkage type and substitutions (31) (Figure 1).

Like *Shigella*, the Streptococcus genus is grouped based on the carbohydrate composition of the antigens found on their cell wall and its ability to trigger unique immune responses. This grouping, first described by Rebecca Lancefield in 1933 (32), has prompted the study of the chemical structure and composition of numerous Streptococcal Lancefield antigens, which have since been found to contain rhamnose as their major substituent. These rhamnose-rich surface polysaccharides are currently being studied as a potential vaccine candidate against Group A Streptococcus (GAS), both by academics and the pharmaceutical industry (33-35). As well as species from GAS such as *Streptococcus pyogenes*, the streptococcus genus contains important pathogens to humans and animals, with GAS being the causative agent for a variety of diseases such as streptococcal pharyngitis (strep throat), scarlet fever and autoimmune disease such as rheumatic heart disease (36). Group B Streptococcus (GBS) is an important human pathogen; although mostly harmless in adults it can be fatal if passed from a pregnant mother to a newborn baby (37). Importantly, GBS is also a bovine

pathogen, similarly to other Lancefield bacteria from groups C, D, F, and G, which are both human and animal pathogens (38). The Group A Carbohydrate (GAC) was the first cell wall carbohydrate reported as a virulence determinant for Group A Streptococcus (33), and similar studies on other Group carbohydrates are eagerly awaited. Some species such as the oral pathogen *Streptococcus mutans* and other Streptococci from the viridans group do not belong to a conventional Lancefield group due to the lack of a group specific antigen. Nonetheless, these strains also produce cell wall polysaccharides similar to the Lancefield Streptococci that are rich in rhamnose and have been found to be vital for pathogenesis (39) (Figure 1). In addition to the observations made for *S. mutans*, previous work has emphasized the crucial role of the rhamnose-containing Lancefield antigens from GAS and GBS in pathogenesis (33, 40). Work is ongoing to unravel the complex structures behind some of these carbohydrates, but GAS has previously been shown to contain a rhamnose backbone of alternating  $\alpha$ 1-2,  $\alpha$ 1-3 linkages, periodically decorated with N-acetylglucosamine residues (41, 42); more recently it has been found to contain periodic decorations of glycerol-phosphate on the side chains of the carbohydrate GlcNAc (43) (Figure 1). The same negative charge decoration has also been observed in SCC and was proposed to be present in many more Streptococci (41). The Group B Carbohydrate (GBC) is unique from other streptococcal Lancefield antigens as it forms a multi-antenna branched structure, with an adapter region to link it to the peptidoglycan layer (44, 45). Importantly, the GBC, like other Lancefield antigens, are rich in rhamnose in both the adapter region and repeat unit of the carbohydrate (Figure 1). Moreover, it was recently shown that the majority of streptococcal rhamnose polysaccharides from almost all Lancefield groups are synthesised via a conserved GlcNAc-rhamnose initiator disaccharide (17). Most annotated rhamnosyltransferase genes and the TDP-rhamnose biosynthesis genes have been reported as being essential in Group A, B and C Streptococcus (33, 46).

Another genus of bacteria found to contain rich rhamnose cell wall polysaccharides is the gram-positive Lactobacillus genus. Famous for their ability to convert sugars into lactic acid, the Lactobacilli spp. form a major component of a healthy gut microbiota (47). *Lactobacillus casei* BL23, a species used extensively for fermentation purposes, has recently been found to contain a rhamnose rich cell wall polysaccharide structurally related in parts to Streptococcal Lancefield antigens, however synthesised in heptasaccharide repeat units, which are branched on the rhamnose sugars. The GalNAc-rhamnose backbone is decorated with glucose or n-acetylglucosamine side chains forming the core heptasaccharide repeating unit (48). *L. lactis* has also been found to produce a cell wall polysaccharide containing rhamnose (49), and interestingly the biosynthesis gene cluster for cell wall polysaccharide biogenesis in *L. lactis* has been shown to share similarities with the Lancefield antigen gene clusters from Streptococcus (3). Each of these gene clusters starts with one or more of the biosynthetic genes for TDP- $\alpha$ -L-rhamnose biosynthesis, highlighting the importance of rhamnose in these diverse cell wall polysaccharides (3). The most recently characterised rhamnose containing polysaccharide is the enterococcal polysaccharide antigen from vancomycin resistant *Enterococcus faecalis* (50, 51). The reported gene clusters contain in fact three pathways, i) TDP-Rha biosynthesis, ii) the rhamnan backbone biosynthesis cluster and iii) repeat unit biosynthesis.

Several bacteria contain the rhamnose-rich S-layer, an array of abundant proteins or glycoproteins found on the surface (S-layer) of walled bacteria and archaeal envelopes. As

with other post-translational modifications, the proteins found in the S-layer are glycosylated (typically N-glycosylation) to add a layer of further complexity to the proteinaceous structure. Rhamnose has been found as a component of several bacterial S-layers, for instance as polyrhamnan in the S-layer of *Geobacillus stearotherophilus* (2, 52). Another example is that rhamnose is incorporated into the complex N-glycan structure from *Paenibacillus alvei* (53) (Figure 1). Although less widespread, rhamnose has also been found in the S-layer of some archaea as a response to different growth conditions (54).

Whilst L-Rha is much more common across all domains of life than the D-enantiomer, the Gram-negative plant, animal, and human pathogen, *Pseudomonas aeruginosa*, produces a common polysaccharide antigen on its LPS made up solely of D-Rha residues (55). These antigenic regions have been found to contain an average of ~70 D-Rha residues per molecule, making them a major component of the *P. aeruginosa* cell surface. Previous work has shown that *P. aeruginosa* mutants that lack the ability to produce GDP-D-rhamnose are significantly less efficient at adhering to human bronchial epithelial cells, suggesting that these large D-Rha rich antigenic common polysaccharide antigens may play an important role in *P. aeruginosa* adherence to tissues (56).

## Viruses

Although the glycobiology of viruses is less understood than eukaryotes or bacteria, with many viruses hijacking host glycosylation machinery for capsid glycosylation, it has recently been shown that viruses do produce rhamnose glycoconjugates. The giant Chlorella virus, *Paramecium bursaria* chlorovirus PBCV-1, synthesises highly complex N-glycans decorated on the major capsid protein that contain both L- and D-Rha (57, 58) and are the first reported viral species that produce both enantiomers (Figure 1). Mimivirus has also been found to contain high amounts of L-Rha in its fibers (59). Interestingly, unlike other viruses, giant viruses such as mimivirus and PBCV-1 appear to encode most of the glycosylation machinery in their large genomes for autonomous glycosylation. PBCV-1 has been found to encode a functional GDP-L-fucose biosynthesis pathway, which also produces the rare GDP-D-rhamnose as a major product (14), and mimivirus has been shown to contain a biosynthesis pathway for TDP-L-rhamnose (60).

## NDP-Rhamnose biosynthesis

The biosynthesis of NDP- $\beta$ -L-rhamnose has been studied extensively in bacteria and plants, and more recently microalgae and giant viruses have even had their NDP- $\beta$ -L-rhamnose biosynthesis pathways discovered (8, 60). Production of nucleotide-activated L-Rha arises from either the Rml pathway (commonly found in bacteria) or the NRS/ER or RHM pathways (commonly found in plants and other eukaryotes) (Fig. 2). Whilst there are examples of some enzyme promiscuity *in vitro* in each pathway (8, 10, 61), the Rml pathway primarily produces TDP- $\beta$ -L-rhamnose and the NRS/ER or RHM pathway primarily produces UDP- $\beta$ -L-rhamnose (Figure 2).

### **TDP- $\beta$ -L-rhamnose biosynthesis**

TDP- $\beta$ -L-rhamnose is the most commonly found rhamnose nucleotide sugar and is ubiquitous amongst gram-negative and gram-positive bacteria, where it serves as a common component of the O-antigen of lipopolysaccharides, as previously discussed. Biosynthesis of TDP- $\beta$ -L-rhamnose commonly follows a four-step route, starting with the production of TDP- $\alpha$ -D-glucose by the thymidyltransferase RmlA. The three other chemical modifications of the substrate TDP- $\alpha$ -D-glucose include 1) dehydration (RmlB – 4,6 dehydratase), 2) epimerization (RmlC – 3,5-epimerase), and subsequent 3) reduction (RmlD – 4-reductase) to produce TDP- $\beta$ -L-rhamnose. Interestingly, the four Rml genes (*rmlA-D*) that encode the complete pathway are often found clustered in bacteria. Conversely, the Streptococci from more than 40 different species, including Group A, C and G Streptococci, do not cluster *rmlA-D* (62). *Streptococcus pyogenes* contains RmlD (GacA) in a separate cluster to the rest of the Rml pathway, at the start of the cluster responsible for Lancefield antigen biosynthesis (62, 63). Van der Beek *et al.*, showed that this novel monomeric homolog of RmlD is essential for Group A Streptococcus, and thus represents a good therapeutic target for drug discovery (63). Homologs of RmlB, RmlC and RmlD have been studied for many years, with all having had their crystal structures solved (64-66). Van der Beek also targeted recombinantly expressed and purified RmlBC and GacA (RmlD) from *Streptococcus pyogenes* in a small compound inhibitor screen, highlighting one compound that inhibited both the growth of *S. pyogenes* but also the rhamnose-dependant *Mycobacterium tuberculosis* (62). Alphey *et al.*, have previously targeted purified RmlA from *Pseudomonas aeruginosa* for inhibitor identification, using a combination of high-throughput screening, and subsequent optimization using protein crystallography, *in silico* screening, and chemical synthesis. This vigorous approach yielded nanomolar inhibitors of *P. aeruginosa* RmlA (67).

### **UDP- $\beta$ -L-rhamnose biosynthesis**

Research into UDP- $\beta$ -L-rhamnose biosynthesis in eukaryotes is more recent, but the multidomain *Arabidopsis thaliana* enzyme, RHM, has drawn particular biotechnological attention for its ability to convert as a single protein UDP- $\alpha$ -D-glucose to UDP- $\beta$ -L-rhamnose, via the three common enzymatic steps: dehydration, epimerization, and reduction of the Rml pathway (11, 68) (Figure 2). Pei *et al.*, fused an RHM homolog from *Vitis vinifera* to a bifunctional NRS/ER enzyme from *A. thaliana*, resulting in a self-sufficient NADPH-independent enzyme able to convert UDP- $\alpha$ -D-glucose to UDP- $\beta$ -L-rhamnose (68). As well as the RHM pathway, *A. thaliana* encodes for the NRS/ER pathway, where the three conserved reaction steps are conducted by two enzymes only (10). This pathway is also found in other eukaryotes such as fungi (61), where it is essential for pathogenicity in some species (69). The NRS/ER pathway is also present in giant viruses (60), where both L- and D- enantiomers of rhamnose are found in complex N-glycans (57). Whilst most species of algae appear to encode for either the RHM or NRS/ER pathway, depending on their eukaryotic origin, Wagstaff *et al.* recently reported a novel RmlC/RmlD fusion protein in the Haptophyceae and Alveolata phyla. The authors suggested that this chimera was likely the result of horizontal and endosymbiotic gene transfer events during plastid evolution (8).

### **GDP- $\alpha$ -D-rhamnose biosynthesis**

Unlike L-Rha which is activated with either uridine or thymidine, D-Rha is biosynthesized as guanidine 5'-diphospho- $\alpha$ -D-rhamnose (GDP- $\alpha$ -D-rhamnose) by the actions of a GDP- $\alpha$ -D-mannose 4,6-dehydratase (GMD) (14, 70) and a GDP-6-deoxy-D-lyxo-4-hexulose reductase (13) (Figure 2). The presence of D-Rha and the responsible biosynthetic pathway are much less common across all domains of life. Like the L-Rha pathways, the sugar ring is first dehydrated by the enzyme GDP-mannose 4,6-dehydratase, before undergoing subsequent stereoselective reduction at position 4 to produce GDP- $\alpha$ -D-rhamnose. This pathway has evolved to show some level of promiscuity in the end product, with reduction to the alternative stereoisomer resulting in GDP- $\alpha$ -D-talose, or an additional 3,5-epimerisation step before reduction resulting in GDP- $\beta$ -L-fucose. GDP- $\alpha$ -D-mannose 4,6-dehydratases (GMD) have been found in many organisms due to their ability to produce the precursors for the biosynthesis of GDP- $\alpha$ -D-rhamnose, GDP- $\alpha$ -D-talose, and GDP- $\beta$ -L-fucose. However, the characterized GDP-4-keto-rhamnose reductases (Rmd) that produces specifically GDP- $\alpha$ -D-rhamnose are less conserved but have been reported in *Aneurinibacillus thermoaerophilus* and *Pseudomonas aeruginosa* (12, 13), which produce D-Rha as a major substituent of their respective LPS, as discussed previously. Interestingly, a GMD enzyme from *Paramecium bursaria* *Chlorella* virus 1 has been shown to have both 4,6-dehydratase and 4-reductase activity, leading to a promiscuous production of GDP- $\alpha$ -D-rhamnose and GDP- $\beta$ -L-fucose from the same enzymatic pathway (14).

## Rhamnosyltransferases

Rhamnosyltransferases are the catalytic enzymes that are responsible for the transfer of rhamnose to the target glycoconjugate. Using nucleotide-activated rhamnose (NDP-Rha) as a sugar donor, rhamnosyltransferases can transfer one or multiple rhamnose residues to a diverse range of acceptor molecules. These range from small natural products such as plant flavonoids, to much larger polysaccharides within bacterial LPS (Table 1). Recombinant production and purification of glycosyltransferases are often challenging, due to their nature of being commonly membrane associated, multidomain proteins. Furthermore, the acceptor substrates are often chemically unusual, and the enzymes require elaborate substrates that cannot be easily mimicked or synthesised for *in vitro* studies. Therefore, a lot of work that has been reported on functionally characterized rhamnosyltransferases involves the genetic manipulation of the host organism, and subsequent metabolite profiling to identify changes in levels of rhamnosylation of target products.

## Eukaryotic rhamnosyltransferases characterized using genetic manipulation

The pectic polysaccharide RG-I is arguably the most notorious rhamnose containing cell wall polysaccharide in land plants. Recently, Takenaka *et al.*, were able to functionally characterize the rhamnosyltransferases from *Arabidopsis thaliana* that are responsible for the transfer of rhamnose from UDP- $\beta$ -L-rhamnose to RG-I oligosaccharides (RRTs) (94). In this work, the authors found that RRT1, which is one of four rhamnosyltransferases responsible for generation of RG-I, is a single-spanning transmembrane protein that is highly expressed during the formation of seed coat mucilage – a specialized cell wall component



abundant with RG-I. In agreement with its proposed function, inactive mutants of RRT1 resulted in a reduction in levels of RG-I in seed coat mucilage.

Other rhamnosyltransferases from plants that have been analysed using genetic manipulation approaches include the  $\beta$ -solanine/ $\beta$ -chaconine rhamnosyltransferase, SGT3, from *Solanum tuberosum* (20). Steroidal glycoalkaloids are often harmful speciality metabolites found in *Solanaceous* plants and include  $\alpha$ -solanine and  $\alpha$ -chaconine, two tri-glycosylated alkaloids that accumulate in potato tubers. Using antisense downregulation of *Sgt3*, McCue *et al.*, were able to show that SGT3 is responsible for the transfer of rhamnose to the first galactose or glucose residue of  $\beta$ -solanine or  $\beta$ -chaconine, respectively (20). Downregulation of *Sgt3* resulted in accumulation of both  $\beta$ -solanine and  $\beta$ -chaconine in tubers, with a concomitant reduction in levels of  $\alpha$ -solanine and  $\alpha$ -chaconine.

### **Prokaryotic rhamnosyltransferases characterized using genetic manipulation**

*Pseudomonas* produce glycoconjugates containing both L- and D-Rha, including virulence factors such as extracellular rhamnolipids that contain one (mono-rhamnolipid) or two (di-rhamnolipid) L-Rha residues. Rhamnosyltransferase 1 (RhlAB) is known to transfer the first rhamnose residue from TDP- $\beta$ -L-rhamnose to  $\beta$ -hydroxydecanoyl- $\beta$ -hydroxydecanoate. Rahim *et al.* characterized the function of the *rhlC* gene product as the second rhamnosyltransferase, responsible for the generation of di-rhamnolipid (85). In this work the authors were able to generate a chromosomal mutant of *rhlC* which was subsequently unable to produce di-rhamnolipid as judged by TLC analysis of mono- and di-rhamnolipid in the culture supernatants. The authors also showed that this mutant did not affect the levels of mono-rhamnolipid, but the *rhlA* null mutant was unable to produce both mono- and di-rhamnolipid. Interestingly, levels of the di-rhamnolipid appeared to increase in the stationary phase of growth, suggesting a possible role for it in allelopathy.

The majority of *Streptococci* produce rhamnose rich lipo-oligosaccharides on their cell-surface as either non-capsular Lancefield antigens or capsule polysaccharides (32). Although recent work has begun to biochemically characterize enzymes responsible for the biosynthesis of these rhamnose-rich carbohydrates (17, 43), the identity of three important rhamnosyltransferases was uncovered over a decade ago for *Streptococcus mutans* (87). In this work, Shibata *et al.*, were able to analyse LPS profiles of *E. coli* strains harbouring various combinations of genes from the *S. mutans* rhamnose-glucose polysaccharide (RGP) pathway. Using Tricine-SDS-PAGE analysis, they showed that an *E. coli* transformant harbouring *rgpA* was able to produce a lipooligosaccharide with an additional rhamnose residue and were also able to use similar experiments to show that *rgpB* and *rgpF* can further rhamnosylate these lipooligosaccharide samples. Based on these early findings, Shibata *et al.*, were first able to propose that RgpA, RgpB, and RgpF, utilize TDP-rhamnose for rhamnose polysaccharide biosynthesis.

### **Eukaryotic rhamnosyltransferases characterized *in vitro***

Possibly the first example of successful expression, purification, and biochemical characterization of a rhamnosyltransferase was reported in 1991. Bar-Peled *et al.* homogenously purified a flavanone-7-O-glucoside-2''-O-rhamnosyltransferase responsible

for the rhamnosylation of the bitter flavanone glucosides, naringin and neo-hesperidin (97). In this work, Bar-Peled *et al.* underwent a thorough column chromatography strategy to purify the rhamnosyltransferase >2700 fold from extracts of young pumelo leaves. The purified enzyme was able to selectively transfer rhamnose from UDP- $\beta$ -L-rhamnose to flavonoid-7-O-glucosides, but interestingly not to 5-O-glucosides or flavonoid aglycones. The authors then performed a detailed biochemical analysis, showing that the enzyme only rhamnosylates with the donor substrate UDP- $\beta$ -L-rhamnose, and detailing the relative activities for a number of other acceptor substrates. Further work from Bar-Peled and co-workers, published in 2004 with improved genomic data, allowed the identification of the *Cm1,2RhaT* gene from pumelo that was responsible for this rhamnosyltransferase activity observed first in 1991 (19). In this work, Frydman *et al.* showed that the 1,2-rhamnosyltransferase enzyme responsible for the bitter taste of species such as grapefruit and pumelo, is only found in neohesperidoside-producing citrus species. Using RT-PCR they revealed that the expression of this gene is significantly upregulated in young tissue, which is in agreement with the developmental pattern of flavanone-neohesperidosides accumulating in younger tissues.

Flavanol glycosides are one of the most prominent natural product classes found in the model plant species, *Arabidopsis thaliana*. Despite this abundant present in nature, no flavonoid glycosyltransferases had been biochemically characterized from *A. thaliana* until 2003. Jones *et al.* conducted an *in planta* and *in vitro* approach, to characterize both a UDP-rhamnose:flavonol-3-O-rhamnosyltransferase and UDP-glucose:flavonol-3-O-glycoside-7-O-glucosyltransferase (80). In this work, the authors used LC-MS to analyse the presence or absence of rhamnosylated/glucosylated glycosides in the mutant lines *ugt78D1* and *ugt73C6*, proposing that *ugt78D1* was a rhamnosyltransferase responsible for the formation of quercetin-3-O-rhamnoside. The authors also recombinantly expressed UGT78D1 in *E. coli* as a GST-fusion, but this enzyme lost activity rapidly. Therefore, subsequent assays were performed with crude extracts of *E. coli* harbouring the rhamnosyltransferase-containing plasmid. Mixtures of recombinant UGT78D1 combined with a UDP-rhamnose preparation, and the acceptor compound quercetin resulted in production of quercetin-3-O-rhamnoside, in agreement with their *in planta* data that UGT78D1 is flavonol-3-O-rhamnosyltransferase.

Soybean (*Glycine max*) is another plant containing an abundance of flavonol glycosides. Like Jones *et al.*, in 2014 Rodas *et al.* took both an *in planta* and *in vitro* approach in order to functionally characterize a flavonol 3-O-glucoside-1,6-rhamnosyltransferase (82). In this work the authors used HPLC to investigate the flavonol glycosides produced by a number of recombinant inbred lines, which were developed by crossing Kitakomachi and Koganejiro cultivars of soybean. The authors found that the flavonol glycosides of Koganejiro were rhamnosylated, whereas flavonol glycosides of Kitakomachi did not contain rhamnose. Using this methodology, the authors identified a candidate gene for this specific rhamnosyltransferase activity, designated GmF3G6''Rt (UGT79A6). An *in vitro* assay was developed for recombinantly produced UGT79A6 and subsequent HPLC and MS analysis of the reaction products confirmed that UGT79A6 was a rhamnosyltransferase, which produced kaempferol 3-O-rutinoside (rhamnosylated) utilising kaempferol 3-O-glucoside and UDP-rhamnose. The authors showed that the activity was specific for UDP-rhamnose, as UGT79A6 could not utilise UDP-glucose, UDP-galactose, or UDP-glucuronic acid as sugar-donors.

To date, only a single rhamnosyltransferase from eukaryotes has been crystallised; the flavonol  $\alpha$ -7-O-L-rhamnosyltransferase, UGT89C1, from *A. thaliana*. Zong *et al.*, recently reported the crystal structure of this plant rhamnosyltransferase in complex with the sugar-donor, UDP-rhamnose, and the flavonol acceptor, quercetin (98). Having previously been characterized as a flavonol-7-O-L-rhamnosyltransferase based on a combination of *in planta* and *in vitro* work (21), Zong and co-workers used mutagenesis studies to shed detailed insight into key amino acid residues for sugar donor recognition and specificity for UDP-rhamnose. The mutant H357Q, in particular, was shown to exhibit activity with both UDP-rhamnose and UDP-glucose. Discovery of key active site amino acids, like this histidine sidechain, are important for the bioinformatic-based prediction of whether a glycosyltransferase may be a rhamnosyltransferase or have other sugar-donor specificity. Interestingly, of the six rhamnosyltransferases that have been crystallised to date, all display a GT-B fold (Table 1, Figure 3).

### **Prokaryotic rhamnosyltransferases characterized *in vitro***

The first crystal structure of a rhamnosyltransferase was reported by Steiner *et al.*, in 2010, when they successfully solved the structure of the  $\beta$ -1,2-rhamnosyltransferase WsaF from *Geobacillus stearothermophilus* (52) (Figure 3). WsaF is responsible, along with WsaE, for the generation of a trirhamnan repeating unit of the polyrhamnan that decorates proteins in the S-layer in *G. stearothermophilus*. Steiner *et al.* revealed through recombinant expression in 2008, that WsaE is a trifunctional rhamnosyltransferase responsible for both methylation and generation of independent  $\alpha$ -1,2 and  $\alpha$ -1,3-linked rhamnose residues in the growing polyrhamnan chain (88). Using a clever design of truncated enzyme constructs that included or excluded these individual domains, the authors were able to use TLC analysis with an octyl-linked oligosaccharide acceptor to show that both glycosyltransferase domains were needed for the full activity of the enzyme, and that the C-terminal glycosyltransferase was responsible for the  $\alpha$ -1,2-rhamnosyltransferase activity. In the same work, the authors also recombinantly expressed and purified WsaF. Using an octyl-linked oligosaccharide and TLC analysis, they showed that WsaF is a rhamnosyltransferase that adds a single rhamnose to the growing chain. Using NMR spectroscopy, the authors were able to show that the new rhamnose introduced by WsaF was of a  $\beta$ -1,2-linkage, and that the additional rhamnosyltransferase domain of WsaE was transferring rhamnose in an  $\alpha$ -1,3-linkage. Alongside WsaE and WsaF, Steiner *et al.* used a similar approach to characterize two further rhamnosyltransferases, WsaC and WsaD. The authors found that these two  $\alpha$ -1,3-rhamnosyltransferases were responsible for the addition of a single rhamnose residue each to the adaptor saccharide region (88).

Just two years after the first crystal structure of WsaF was published, the crystal structure of another rhamnosyltransferase was revealed (99). SpnG is a rhamnosyltransferase responsible for the final tailoring of the spinosyns A and D. The spinosyns are complex polyketides produced by the actinomycete *Saccharopolyspora spinose* and make up the active ingredient of pesticides. SpnG crystallised as a homodimer, as it has been observed for other glycosyltransferases belonging to the GT1 family (102, 103). This crystal structure

highlighted the importance of residues H13 and D316 in SpnG activity, in agreement to previous reports in 2009 (76). The crystal structure also highlighted how SpnG was selective for TDP-sugars over UDP-sugars; the amine of the Asn202 sidechain forms a hydrogen bond with the 3-OH from the thymidine moiety whilst also preventing ribose-containing nucleotides from binding due to clashes with 2'-OH.

N-linked glycosylation is a common posttranslational modification of proteins across all domains of life, which contributes to the correct folding, localization, and signalling of cellular proteins. Typically, proteins are glycosylated with either glucose, or more commonly N-acetylglucosamine. Recently, a novel form of arginine glycosylation with rhamnose has been discovered in *Pseudomonas aeruginosa*, *Pseudomonas putida*, *Shewanella oneidensis*, and *Neisseria meningitidis* (90-93). Known as EarP enzymes, these rhamnosyltransferases have been shown to rhamnosylate translation elongation factor P. This translational modification has been shown to be crucial for the rescue of polyproline-stalled ribosomes, and for the pathogenicity of organisms such as *Pseudomonas aeruginosa* (104). A structural basis for this arginine rhamnosylation was uncovered by Krafczyk and coworkers in 2017 when they crystallised EarP from *Pseudomonas putida* (91). This work has been crucial in identifying important active site residues and catalytic residues such as Asp20. This conserved residue was also described by Sengoku *et al.* in their report of the crystal structure of the EarP homolog from *Neisseria meningitidis* in 2018 (100). The latest of these EarP rhamnosyltransferases to be crystallised was EarP from *Pseudomonas aeruginosa* by He and co-workers (101). Interestingly, EarP from *Pseudomonas aeruginosa* was shown to display important conformational changes upon TDP-rhamnose and translation elongation factor P binding; with sugar-donor binding enhancing acceptor binding. The continued study of these novel rhamnosyltransferases and future development of drug discovery screens should be a key focus for the upcoming years.

Only the WsaF enzyme from *Geobacillus stearothermophilus* has a retaining mechanism of action, whilst all other reported structural studies reported an inverting mechanism of catalysis, alternating the stereochemistry of the nucleotide sugar donor during catalysis to generate the product (Figure 3). Four of the reported structures are functional dimers, with only WsaF and SpnG, from *Saccharopolyspora spinosa*, being functional homodimers. A major structural difference between the retaining WsaF enzyme and the inverting SpnG, from *Saccharopolyspora spinosa* is that the dimerization interface appears to be on the opposite site (Figure 3). Further structural and functional characterisation is required to reveal if this is a common tertiary structure feature for retaining and inverting rhamnosyltransferase.

In *Mycobacterium tuberculosis*, the conserved gene *wbbl* encodes an inverting rhamnosyltransferase responsible for the transfer of  $\alpha$ -1,3-rhamnose from TDP- $\beta$ -L-rhamnose to the GlcNAc moiety of decaprenyldiphosphoryl- $\alpha$ -D-GlcNAc. This growing lipooligosaccharide chain is then further elongated with galactofuranose and arabinofuranose units before being transferred onto peptidoglycan. Grzegorzewicz *et al.* were able to recombinantly assay WbbL's activity by overexpression in *E. coli* and purifying the WbbL-containing membrane fractions (105). Using this methodology, the authors created a plate-based assay using the natural substrate, GlcNAc-PP-decaprenyl, and were able to characterize pH optima, metal dependency, and kinetic parameters for WbbL.

An enzyme with a similar rhamnosyltransferase activity, GacB, has been characterized from the human exclusive pathogen *Streptococcus pyogenes* (Group A Streptococcus). Zorzoli, Meyer *et al.* reported that GacB, like WbbL, is responsible for the rhamnosylation of a GlcNAc-PP-undecaprenyl substrate *in vivo* (17). In Group A Streptococcus, this primed lipodissacharide is then further elongated with  $\alpha$ -1,3- and  $\alpha$ -1,2-rhamnose residues, before having GlcNAc side-chains incorporated that are further decorated with glycerol-phosphate modifications to build the Group A Carbohydrate virulence factor (43). Zorzoli *et al.* genetically incorporated streptococcal polyrhamnose biosynthesis gene clusters into *E. coli* using parts of the gene cluster from *Streptococcus mutans* and used antibodies against the polyrhamnose product as a marker to investigate the effects of manipulating individual genes. Using this strategy, GacB was shown to be essential for polyrhamnose biosynthesis (RhaPS). Using radiolabelling experiments, GacB was proposed to transfer a single rhamnose residue to the growing lipid-linked oligosaccharide, which was confirmed *in vitro* by detergent extracting the GacB enzyme from *E. coli* membrane fractions and assaying using MALDI-TOF. This *in vitro* assay was also used to show that GacB is metal independent, and NMR analysis of the reaction product confirmed that GacB is a novel retaining rhamnosyltransferase producing a  $\beta$ -1,4 linkage, unlike WbbL, which is an  $\alpha$ -1,3-rhamnosyltransferase (105). Strikingly, the authors revealed that the GacB enzyme produces the first committed step not only in Group A Streptococcus and *S. mutans*, but also is conserved in the RhaPS biosynthesis gene cluster of Group B, Group C, and Group G Lancefield Streptococci. This conservation and the essentiality of the *gacB* gene and its homologs has significant implications for the future development of novel streptococcal drug candidates.

### **Viral rhamnosyltransferases characterized *in vitro***

Very recently, Speciale *et al.* biochemically characterized the rhamnosyltransferase activity of a multifunctional enzyme from the algal-infecting giant virus, PBCV-1 (58). In this work, the authors showed that A064R contains 3 domains; 2 of which have  $\beta$ -1,4- and  $\alpha$ -1,2-rhamnosyltransferase activity, with the third responsible for 2-*O*-methyltransferase activity which installs the 2-*OMe* group on the terminal rhamnose residue (58). In this work, the authors used several synthetic lipo-oligosaccharides and constructs of A064R varying in length, together with HPLC analysis, to characterize the activity of each domain. Both rhamnosyltransferases domains were shown to use UDP- $\beta$ -L-rhamnose as the L-rhamnose nucleotide sugar donor. A064R-D1 was shown to encode the  $\beta$ -1,4-rhamnosyltransferase activity required to install the first rhamnose at the 4'-OH position of the xylose residue. A064R-D2 was shown to encode an  $\alpha$ -1,2-rhamnosyltransferase responsible for installing the terminal rhamnose residue onto the previously installed L-rhamnose. After both rhamnose residues are installed, A064R-D3 was shown to methylate the terminal rhamnose at the 2'-OH position. This breakthrough work is, to our knowledge, the first example of a biochemically characterized viral rhamnosyltransferase, and sheds light on how giant viruses have evolved to perform N-glycosylation.

### **Concluding remarks and perspectives**

Rhamnose, in both the L- and D-form, is found widespread in plant and microbial natural products and surface glycans. It is particularly prevalent in lipopolysaccharides and surface

carbohydrate in Gram-positive and Gram-negative bacteria. Strikingly, rhamnose and any nucleotide-rhamnose biosynthesis enzymes are entirely absent from animals and higher eukaryotes. This distinction has often led to the proposal to screen for small molecule inhibitors to target those enzymes that synthesise or utilise rhamnose-nucleotide sugars, which are ideal drug targets as they are often essential in pathogenic bacteria and absent in humans. A number of projects have focused on bacterial enzymes involved in the biosynthesis of TDP- $\beta$ -L-rhamnose, including inhibitor screens and optimization to target successfully the first enzyme of the pathway, RmlA (62, 67). However, it is also noteworthy to highlight that there is no report of any drug discovery programme operated by academia or industry to develop these small molecule inhibitors into lead compounds for drug discovery pipelines. It is commonly known that the financial costs to develop novel antimicrobial drugs is hindering those developments. Researchers and private as well as governmental funding bodies need to continue working together to increase the opportunities to successfully develop a generation of new antimicrobial drugs, including those targeting the human and veterinary absent sugars L- and D-Rha.

The absence of any rhamnose in humans has also been exploited for vaccine development to prevent infections from pathogenic bacteria, including *Shigella* and the *Streptococcus* genus. A particularly well investigated glycoconjugate vaccine candidate has been developed over the past two decades for *Shigella dysenteriae* serotype 1. This vaccine candidate is composed of a carrier protein that carries recombinantly produced *Shigella dysenteriae* serotype 1 surface polysaccharides, which contain a rhamnose repeat unit (Fig 1). Clinical trials were found safe and immunogenic in a phase I clinical trial (106). Preclinical trials in animals have shown that the Group A Carbohydrate backbone is also a strong immunogenic vaccine candidate and is currently exploited by several research groups and industry (33). This unique opportunity will hopefully benefit the vaccine development of a number of pathogenic bacteria, that produce similar, rhamnose-containing glycoconjugates on their surface.

The biological activity of many rhamnosyltransferases has been revealed and the CAZy database continues to be an invaluable resource for the cataloguing of glycosyltransferases. The current difficulties in characterizing the rhamnosyltransferase function *in vitro* are often linked to the biological and physiochemical properties of the enzymes, in particular the membrane association due to the lipid-linked nature of their acceptor substrates. These features significantly reduce the chances to crystallize a full length or functional rhamnosyltransferase. Another major limitation is the access to affordable quantities of TDP- or UDP-rhamnose. In addition, radiolabelled derivatives of TDP- or UDP-rhamnose are not available – in comparison to other nucleotide sugars - challenging the researchers to develop their bespoke analytical systems and tools. The huge diversity of the acceptor substrates, in particular with regards to the lipid linked carbohydrates, is a significant obstacle to reveal functional insights into the enzymes under study. These tools need to be continuously developed and made widely accessible to study rhamnosyltransferases. These current constraints result in fewer studies conducted on these classes of enzymes and the majority of studies has relied on genetic manipulation of host organisms or semi-crude purifications of the target enzyme. Only half a dozen of rhamnosyltransferases have been purified to homogeneity and crystallised. As more structures of rhamnosyltransferase families become available, detailed insights into their biochemistry will be revealed and can

be exploited to develop better tools to study related rhamnosyltransferases. This progress and development will ultimately benefit the discovery of the first inhibitors to target pathogenic bacteria via their rhamnosyltransferases.

## References

1. Moses T, Papadopoulou KK, Osbourn A. Metabolic and functional diversity of saponins, biosynthetic intermediates and semi-synthetic derivatives. *Crit Rev Biochem Mol Biol.* 2014;49(6):439-62.
2. Steiner K, Pohlentz G, Dreisewerd K, Berkenkamp S, Messner P, Peter-Katalinić J, et al. New insights into the glycosylation of the surface layer protein SgsE from *Geobacillus stearothermophilus* NRS 2004/3a. *J Bacteriol.* 2006;188(22):7914-21.
3. Mistou M-Y, Sutcliffe IC, van Sorge NM. Bacterial glycobiology: rhamnose-containing cell wall polysaccharides in Gram-positive bacteria. *FEMS Microbiol Rev.* 2016;40(4):464-79.
4. Mishra T, Balaji PV. Highly clade-specific biosynthesis of rhamnose: present in all plants and in only 42% of prokaryotes. Only *Pseudomonas* uses both D-and L-rhamnose. *bioRxiv.* 2019:854612.
5. Wang S, Hao Y, Lam JS, Vlahakis JZ, Szarek WA, Vinnikova A, et al. Biosynthesis of the common polysaccharide antigen of *Pseudomonas aeruginosa* PAO1: characterization and role of GDP-d-rhamnose: GlcNAc/GalNAc-diphosphate-lipid  $\alpha$ 1, 3-d-rhamnosyltransferase WbpZ. *J Bacteriol.* 2015;197(12):2012-9.
6. Yamashita Y, Shibata Y, Nakano Y, Tsuda H, Kido N, Ohta M, et al. A Novel Gene Required for Rhamnose-Glucose Polysaccharide Synthesis in *Streptococcus mutans*. *J Bacteriol.* 1999;181(20):6556-9.
7. Giraud MF, Naismith JH. The rhamnose pathway. *Curr Opin Struct Biol.* 2000;10(6):687-96.
8. Wagstaff BA, Rejzek M, Kuhaudomlarp S, Hill L, Mascia I, Nepogodiev SA, et al. Discovery of an RmlC/D fusion protein in the microalga *Prymnesium parvum* and its implications for NDP- $\beta$ -l-rhamnose biosynthesis in microalgae. *J Biol Chem.* 2019;294(23):9172-85.
9. Kaminski L, Eichler J. *Haloferax volcanii* N-Glycosylation: Delineating the Pathway of dTDP-rhamnose Biosynthesis. *PLOS ONE.* 2014;9(5):e97441.
10. Han X, Qian L, Zhang L, Liu X. Structural and biochemical insights into nucleotide-rhamnose synthase/epimerase-reductase from *Arabidopsis thaliana*. *Biochim Biophys Acta.* 2015;1854(10, Part A):1476-86.
11. Oka T, Nemoto T, Jigami Y. Functional analysis of *Arabidopsis thaliana* RHM2/MUM4, a multidomain protein involved in UDP-D-glucose to UDP-L-rhamnose conversion. *J Biol Chem.* 2007;282(8):5389-403.
12. Mäki M, Järvinen N, Rabinä J, Roos C, Maaheimo H, Renkonen R. Functional expression of *Pseudomonas aeruginosa* GDP-4-keto-6-deoxy-D-mannose reductase which synthesizes GDP-rhamnose. *Eur J Biochem.* 2002;269(2):593-601.
13. Kneidinger B, Graninger M, Adam G, Puchberger M, Kosma P, Zayni S, et al. Identification of two GDP-6-deoxy-D-lyxo-4-hexulose reductases synthesizing GDP-D-rhamnose in *Aneurinibacillus thermoaerophilus* L420-91T. *J Biol Chem.* 2001;276(8):5577-83.

14. Tonetti M, Zanardi D, Gurnon JR, Fruscione F, Armirotti A, Damonte G, et al. *Paramecium bursaria* Chlorella virus 1 encodes two enzymes involved in the biosynthesis of GDP-L-fucose and GDP-D-rhamnose. *J Biol Chem*. 2003;278(24):21559-65.
15. Rocchetta HL, Burrows LL, Pacan JC, Lam JS. Three rhamnosyltransferases responsible for assembly of the A-band D-rhamnan polysaccharide in *Pseudomonas aeruginosa*: a fourth transferase, WbpL, is required for the initiation of both A-band and B-band lipopolysaccharide synthesis. *Mol Microbiol*. 1998;28(6):1103-19.
16. Poon KK, Westman EL, Vinogradov E, Jin S, Lam JS. Functional characterization of MigA and WapR: putative rhamnosyltransferases involved in outer core oligosaccharide biosynthesis of *Pseudomonas aeruginosa*. *J Bacteriol*. 2008;190(6):1857-65.
17. Zorzoli A, Meyer BH, Adair E, Torgov VI, Veselovsky VV, Danilov LL, et al. Group A, B, C, and G *Streptococcus* Lancefield antigen biosynthesis is initiated by a conserved  $\alpha$ -D-GlcNAc- $\beta$ -1, 4-l-rhamnosyltransferase. *J Biol Chem*. 2019;294(42):15237-56.
18. Eckstein TM, Silbaq FS, Chatterjee D, Kelly NJ, Brennan PJ, Belisle JT. Identification and Recombinant Expression of a *Mycobacterium avium* Rhamnosyltransferase Gene (*rtfA*) Involved in Glycopeptidolipid Biosynthesis. *J Bacteriol*. 1998;180(21):5567-73.
19. Frydman A, Weisshaus O, Bar-Peled M, Huhman DV, Sumner LW, Marin FR, et al. Citrus fruit bitter flavors: isolation and functional characterization of the gene *Cm1*, 2RhaT encoding a 1, 2 rhamnosyltransferase, a key enzyme in the biosynthesis of the bitter flavonoids of citrus. *The Plant Journal*. 2004;40(1):88-100.
20. McCue KF, Allen PV, Shepherd LV, Blake A, Maccree MM, Rockhold DR, et al. Potato glycoesterol rhamnosyltransferase, the terminal step in triose side-chain biosynthesis. *Phytochemistry*. 2007;68(3):327-34.
21. Yonekura-Sakakibara K, Tohge T, Niida R, Saito K. Identification of a flavonol 7-O-rhamnosyltransferase gene determining flavonoid pattern in *Arabidopsis* by transcriptome coexpression analysis and reverse genetics. *J Biol Chem*. 2007;282(20):14932-41.
22. Caffall KH, Mohnen D. The structure, function, and biosynthesis of plant cell wall pectic polysaccharides. *Carbohydr Res*. 2009;344(14):1879-900.
23. Saffer AM, Irish VF. Flavonol rhamnosylation indirectly modifies the cell wall defects of RHAMNOSE BIOSYNTHESIS1 mutants by altering rhamnose flux. *The Plant Journal*. 2018;94(4):649-60.
24. Yin R, Messner B, Faus-Kessler T, Hoffmann T, Schwab W, Hajirezaei M-R, et al. Feedback inhibition of the general phenylpropanoid and flavonol biosynthetic pathways upon a compromised flavonol-3-O-glycosylation. *J Exp Bot*. 2012;63(7):2465-78.
25. Pettolino F, Sasaki I, Turbic A, Wilson SM, Bacic A, Hrmova M, et al. Hyphal cell walls from the plant pathogen *Rhynchosporium secalis* contain (1,3/1,6)- $\beta$ -D-glucans, galacto- and rhamnomannans, (1,3;1,4)- $\beta$ -D-glucans and chitin. *FEBS J*. 2009;276(14):3698-709.
26. Pinto MR, Mulloy B, Haido RMT, Travassos LR, Barreto Bergter E. A peptidorhamnomannan from the mycelium of *Pseudallescheria boydii* is a potential diagnostic antigen of this emerging human pathogen. *Microbiology*. 2001;147(Pt 6):1499-506.
27. Lahaye M, Robic A. Structure and Functional Properties of Ulvan, a Polysaccharide from Green Seaweeds. *Biomacromolecules*. 2007;8(6):1765-74.
28. Li Q, Luo J, Wang C, Tai W, Wang H, Zhang X, et al. Ulvan extracted from green seaweeds as new natural additives in diets for laying hens. *J Appl Phycol*. 2018;30(3):2017-27.



29. Ogawa K, Yamaura M, Maruyama I. Isolation and identification of 2-O-methyl-L-rhamnose and 3-O-methyl-L-rhamnose as constituents of an acidic polysaccharide of *Chlorella vulgaris*. *Biosci, Biotechnol, Biochem*. 1997;61(3):539-40.
30. Cogburn JN, Schiff JA. PURIFICATION AND PROPERTIES OF THE MUCUS OF *EUGLENA GRACILIS* (EUGLENOPHYCEAE)1. *J Phycol*. 1984;20(4):533-44.
31. Anderson M, Sansonetti PJ, Marteyn BS. *Shigella* Diversity and Changing Landscape: Insights for the Twenty-First Century. *Frontiers in Cellular and Infection Microbiology*. 2016;6(45).
32. Lancefield RC. A SEROLOGICAL DIFFERENTIATION OF HUMAN AND OTHER GROUPS OF HEMOLYTIC STREPTOCOCCI. *The Journal of experimental medicine*. 1933;57(4):571-95.
33. Van Sorge NM, Cole JN, Kuipers K, Henningham A, Aziz RK, Kasirer-Friede A, et al. The classical lancefield antigen of group a *Streptococcus* is a virulence determinant with implications for vaccine design. *Cell host & microbe*. 2014;15(6):729-40.
34. Di Benedetto R, Mancini F, Carducci M, Gasperini G, Moriel DG, Saul A, et al. Rational Design of a Glycoconjugate Vaccine against Group A *Streptococcus*. *International Journal of Molecular Sciences*. 2020;21(22):8558.
35. Gao NJ, Uchiyama S, Pill L, Dahesh S, Olson J, Bautista L, et al. Site-Specific Conjugation of Cell Wall Polyrrhamnose to Protein SpyAD Envisioning a Safe Universal Group A *Streptococcal* Vaccine. *Infectious Microbes & Diseases*. 2020;Latest Articles.
36. Spellerberg B, Brandt C, Ferretti J, Stevens D, Fischetti V. *Streptococcus pyogenes: Basic Biology to Clinical Manifestations* [Internet]. University of Oklahoma Health Sciences Center Oklahoma City; 2016.
37. Anthony BF, Okada DM, Hobel CJ. Epidemiology of group B *Streptococcus*: longitudinal observations during pregnancy. *J Infect Dis*. 1978;137(5):524-30.
38. Patterson MJ. *Streptococcus*. *Medical Microbiology* 4th edition: University of Texas Medical Branch at Galveston; 1996.
39. Kovacs CJ, Faustoferri RC, Bischer AP, Quivey Jr RG. *Streptococcus mutans* requires mature rhamnose-glucose polysaccharides for proper pathophysiology, morphogenesis and cellular division. *Mol Microbiol*. 2019;112(3):944-59.
40. Caliot E, Dramsi S, Chapot-Chartier M-P, Courtin P, Kulakauskas S, Pechoux C, et al. Role of the Group B antigen of *Streptococcus agalactiae*: a peptidoglycan-anchored polysaccharide involved in cell wall biogenesis. *PLoS Path*. 2012;8(6).
41. Pritchard DG, Coligan JE, Geckle JM, Evanochko WT. High-resolution 1H-and 13C-nmr spectra of the group A-variant streptococcal polysaccharide. *Carbohydr Res*. 1982;110(2):315-9.
42. Huang DH, Krishna NR, Pritchard DG. Characterization of the group A streptococcal polysaccharide by two-dimensional 1H-nuclear-magnetic-resonance spectroscopy. *Carbohydr Res*. 1986;155:193-9.
43. Edgar RJ, van Hensbergen VP, Ruda A, Turner AG, Deng P, Le Breton Y, et al. Discovery of glycerol phosphate modification on streptococcal rhamnose polysaccharides. *Nat Chem Biol*. 2019;15(5):463-71.
44. Michon F, Katzenellenbogen E, Kasper DL, Jennings HJ. Structure of the complex group-specific polysaccharide of group B *Streptococcus*. *Biochemistry*. 1987;26(2):476-86.
45. Michon F, Brisson JR, Dell A, Kasper DL, Jennings HJ. Multiantennary group-specific polysaccharide of group B *Streptococcus*. *Biochemistry*. 1988;27(14):5341-51.
46. Charbonneau ARL, Forman OP, Cain AK, Newland G, Robinson C, Boursnell M, et al. Defining the ABC of gene essentiality in streptococci. *BMC Genomics*. 2017;18(1):426.

47. Walter J. Ecological Role of Lactobacilli in the Gastrointestinal Tract: Implications for Fundamental and Biomedical Research. *Appl Environ Microbiol.* 2008;74(16):4985.
48. Vinogradov E, Sadovskaya I, Grard T, Chapot-Chartier M-P. Structural studies of the rhamnose-rich cell wall polysaccharide of *Lactobacillus casei* BL23. *Carbohydr Res.* 2016;435:156-61.
49. Chapot-Chartier M-P, Vinogradov E, Sadovskaya I, Andre G, Mistou M-Y, Trieu-Cuot P, et al. Cell surface of *Lactococcus lactis* is covered by a protective polysaccharide pellicle. *The Journal of biological chemistry.* 2010;285(14):10464-71.
50. Guerardel Y, Sadovskaya I, Maes E, Furlan S, Chapot-Chartier M-P, Mesnage S, et al. Complete Structure of the Enterococcal Polysaccharide Antigen (EPA) of Vancomycin-Resistant *Enterococcus faecalis* V583 Reveals that EPA Decorations Are Teichoic Acids Covalently Linked to a Rhamnopolysaccharide Backbone. *mBio.* 2020;11(2):e00277-20.
51. Smith RE, Salamaga B, Szkuta P, Hajdamowicz N, Prajsnar TK, Bulmer GS, et al. Decoration of the enterococcal polysaccharide antigen EPA is essential for virulence, cell surface charge and interaction with effectors of the innate immune system. *PLoS Path.* 2019;15(5):e1007730.
52. Steiner K, Hagelueken G, Messner P, Schäffer C, Naismith JH. Structural basis of substrate binding in WsaF, a rhamnosyltransferase from *Geobacillus stearothermophilus*. *J Mol Biol.* 2010;397(2):436-47.
53. Ristl R, Steiner K, Zarschler K, Zayni S, Messner P, Schäffer C. The S-layer glycome—adding to the sugar coat of bacteria. *International journal of microbiology.* 2010;2011.
54. Guan Z, Naparstek S, Calo D, Eichler J. Protein glycosylation as an adaptive response in Archaea: growth at different salt concentrations leads to alterations in *Haloferax volcanii* S-layer glycoprotein N-glycosylation. *Environ Microbiol.* 2012;14(3):743-53.
55. Lam JS, Taylor VL, Islam ST, Hao Y, Kocíncová D. Genetic and functional diversity of *Pseudomonas aeruginosa* lipopolysaccharide. *Frontiers in microbiology.* 2011;2:118.
56. Microbiology UoGDo, Matewish JM. The functional role of lipopolysaccharide in the cell envelope and surface proteins of *Pseudomonas aeruginosa*: University of Guelph; 2004.
57. De Castro C, Molinaro A, Piacente F, Gurnon JR, Sturiale L, Palmigiano A, et al. Structure of N-linked oligosaccharides attached to chlorovirus PBCV-1 major capsid protein reveals unusual class of complex N-glycans. *Proceedings of the National Academy of Sciences.* 2013;110(34):13956-60.
58. Speciale I, Laugieri ME, Noel E, Lin S, Lowary TL, Molinaro A, et al. Chlorovirus PBCV-1 protein A064R has three of the transferase activities necessary to synthesize its capsid protein N-linked glycans. *Proceedings of the National Academy of Sciences.* 2020;117(46):28735.
59. Piacente F, Gaglianone M, Laugieri M, Tonetti M. The Autonomous Glycosylation of Large DNA Viruses. *International Journal of Molecular Sciences.* 2015;16(12):26169.
60. Chothi MP, Duncan GA, Armirotti A, Abergel C, Gurnon JR, Van Etten JL, et al. Identification of an L-rhamnose synthetic pathway in two nucleocytoplasmic large DNA viruses. *J Virol.* 2010;84(17):8829-38.
61. Martinez V, Ingwers M, Smith J, Glushka J, Yang T, Bar-Peled M. Biosynthesis of UDP-4-keto-6-deoxyglucose and UDP-rhamnose in pathogenic fungi *Magnaporthe grisea* and *Botryotinia fuckeliana*. *J Biol Chem.* 2012;287(2):879-92.

62. van der Beek SL, Zorzoli A, Çanak E, Chapman RN, Lucas K, Meyer BH, et al. Streptococcal dTDP-L-rhamnose biosynthesis enzymes: functional characterization and lead compound identification. *Mol Microbiol.* 2019;111(4):951-64.
63. van der Beek SL, Le Breton Y, Ferenbach AT, Chapman RN, van Aalten DM, Navratilova I, et al. GacA is essential for Group A streptococcus and defines a new class of monomeric dTDP-4-dehydrorhamnose reductases (RmlD). *Mol Microbiol.* 2015;98(5):946-62.
64. Allard ST, Giraud MF, Whitfield C, Graninger M, Messner P, Naismith JH. The crystal structure of dTDP-D-Glucose 4,6-dehydratase (RmlB) from *Salmonella enterica* serovar Typhimurium, the second enzyme in the dTDP-l-rhamnose pathway. *J Mol Biol.* 2001;307(1):283-95.
65. Blankenfeldt W, Kerr ID, Giraud MF, McMiken HJ, Leonard GA, Whitfield C, et al. Variation on a theme of SDR: dTDP-6-deoxy-L-lyxo-4-hexulose reductase (RmlD) shows a new Mg<sup>2+</sup>-dependent dimerisation mode in a well-known enzyme family. *Structure.* 2002;10:773-86.
66. Dong C, Major LL, Srikannathasan V, Errey JC, Giraud MF, Lam JS, et al. RmlC, a C3' and C5' carbohydrate epimerase, appears to operate via an intermediate with an unusual twist boat conformation. *J Mol Biol.* 2007;365(1):146-59.
67. Alphey MS, Pirrie L, Torrie LS, Boulkeroua WA, Gardiner M, Sarkar A, et al. Allosteric Competitive Inhibitors of the Glucose-1-phosphate Thymidyltransferase (RmlA) from *Pseudomonas aeruginosa*. *ACS Chemical Biology.* 2013;8(2):387-96.
68. Pei J, Chen A, Sun Q, Zhao L, Cao F, Tang F. Construction of a novel UDP-rhamnose regeneration system by a two-enzyme reaction system and application in glycosylation of flavonoid. *Biochem Eng J.* 2018;139:33-42.
69. Santhanam P, Boshoven JC, Salas O, Bowler K, Islam MT, Saber MK, et al. Rhamnose synthase activity is required for pathogenicity of the vascular wilt fungus *Verticillium dahliae*. *Mol Plant Pathol.* 2017;18(3):347-62.
70. King JD, Poon KK, Webb NA, Anderson EM, McNally DJ, Brisson JR, et al. The structural basis for catalytic function of GMD and RMD, two closely related enzymes from the GDP-d-rhamnose biosynthesis pathway. *The FEBS journal.* 2009;276(10):2686-700.
71. McCaughey LC, Grinter R, Josts I, Roszak AW, Waløen KI, Cogdell RJ, et al. Lectin-like bacteriocins from *Pseudomonas* spp. utilise D-rhamnose containing lipopolysaccharide as a cellular receptor. *PLoS Pathog.* 2014;10(2):e1003898.
72. Xiao Y, Li S, Niu S, Ma L, Zhang G, Zhang H, et al. Characterization of tiacumicin B biosynthetic gene cluster affording diversified tiacumicin analogues and revealing a tailoring dihalogenase. *J Am Chem Soc.* 2011;133(4):1092-105.
73. Miyamoto Y, Mukai T, Nakata N, Maeda Y, Kai M, Naka T, et al. Identification and characterization of the genes involved in glycosylation pathways of mycobacterial glycopeptidolipid biosynthesis. *J Bacteriol.* 2006;188(1):86-95.
74. Pérez E, Constant P, Lemassu A, Laval F, Daffé M, Guilhot C. Characterization of three glycosyltransferases involved in the biosynthesis of the phenolic glycolipid antigens from the *Mycobacterium tuberculosis* complex. *J Biol Chem.* 2004;279(41):42574-83.
75. Ochsner UA, Fiechter A, Reiser J. Isolation, characterization, and expression in *Escherichia coli* of the *Pseudomonas aeruginosa* rhlAB genes encoding a rhamnosyltransferase involved in rhamnolipid biosurfactant synthesis. *J Biol Chem.* 1994;269(31):19787-95.

76. Chen YL, Chen YH, Lin YC, Tsai KC, Chiu HT. Functional characterization and substrate specificity of spinosyn rhamnosyltransferase by in vitro reconstitution of spinosyn biosynthetic enzymes. *J Biol Chem.* 2009;284(11):7352-63.
77. Luzhetskyy A, Mayer A, Hoffmann J, Pelzer S, Holzenkämper M, Schmitt B, et al. Cloning and heterologous expression of the aranciamycin biosynthetic gene cluster revealed a new flexible glycosyltransferase. *ChemBioChem.* 2007;8(6):599-602.
78. Blanco G, Patallo EP, Braña AF, Trefzer A, Bechthold A, Rohr J, et al. Identification of a sugar flexible glycosyltransferase from *Streptomyces olivaceus*, the producer of the antitumor polyketide elloramycin. *Chem Biol.* 2001;8(3):253-63.
79. Olano C, Abdelfattah MS, Gullón S, Braña AF, Rohr J, Méndez C, et al. Glycosylated derivatives of steffimycin: insights into the role of the sugar moieties for the biological activity. *ChemBioChem.* 2008;9(4):624-33.
80. Jones P, Messner B, Nakajima J, Schäffner AR, Saito K. UGT73C6 and UGT78D1, glycosyltransferases involved in flavonol glycoside biosynthesis in *Arabidopsis thaliana*. *J Biol Chem.* 2003;278(45):43910-8.
81. Chen J, Yuan Z, Zhang H, Li W, Shi M, Peng Z, et al. Cit1,2RhaT and two novel CitdGlcTs participate in flavor-related flavonoid metabolism during citrus fruit development. *J Exp Bot.* 2019;70(10):2759-71.
82. Rojas Rodas F, Rodriguez TO, Murai Y, Iwashina T, Sugawara S, Suzuki M, et al. Linkage mapping, molecular cloning and functional analysis of soybean gene Fg2 encoding flavonol 3-O-glucoside (1 → 6) rhamnosyltransferase. *Plant Mol Biol.* 2014;84(3):287-300.
83. Kroon J, Souer E, De Graaff A, Xue Y, Mol J, Koes R. Cloning and structural analysis of the anthocyanin pigmentation locus Rt of *Petunia hybrida*: characterization of insertion sequences in two mutant alleles. *The Plant Journal.* 1994;5(1):69-80.
84. Mills JA, Motichka K, Jucker M, Wu HP, Uhlik BC, Stern RJ, et al. Inactivation of the mycobacterial rhamnosyltransferase, which is needed for the formation of the arabinogalactan-peptidoglycan linker, leads to irreversible loss of viability. *J Biol Chem.* 2004;279(42):43540-6.
85. Rahim R, Ochsner UA, Olvera C, Graninger M, Messner P, Lam JS, et al. Cloning and functional characterization of the *Pseudomonas aeruginosa* rhIC gene that encodes rhamnosyltransferase 2, an enzyme responsible for di-rhamnolipid biosynthesis. *Mol Microbiol.* 2001;40(3):708-18.
86. Morona R, Mavris M, Fallarino A, Manning PA. Characterization of the rfc region of *Shigella flexneri*. *J Bacteriol.* 1994;176(3):733-47.
87. Shibata Y, Yamashita Y, Ozaki K, Nakano Y, Koga T. Expression and characterization of streptococcal rgp genes required for rhamnan synthesis in *Escherichia coli*. *Infect Immun.* 2002;70(6):2891-8.
88. Steiner K, Novotny R, Werz DB, Zarschler K, Seeberger PH, Hofinger A, et al. Molecular Basis of S-layer Glycoprotein Glycan Biosynthesis in *Geobacillus stearothermophilus*. *J Biol Chem.* 2008;283(30):21120-33.
89. Williams DM, Ovchinnikova OG, Koizumi A, Mainprize IL, Kimber MS, Lowary TL, et al. Single polysaccharide assembly protein that integrates polymerization, termination, and chain-length quality control. *Proceedings of the National Academy of Sciences of the United States of America.* 2017;114(7):E1215-e23.
90. Rajkovic A, Erickson S, Witzky A, Branson OE, Seo J, Gafken PR, et al. Cyclic Rhamnosylated Elongation Factor P Establishes Antibiotic Resistance in *Pseudomonas aeruginosa*. *mBio.* 2015;6(3):e00823.

91. Krafczyk R, Macošek J, Jagtap PKA, Gast D, Wunder S, Mitra P, et al. Structural Basis for EarP-Mediated Arginine Glycosylation of Translation Elongation Factor EF-P. *mBio*. 2017;8(5).
92. Lassak J, Keilhauer EC, Fürst M, Wuichet K, Gödeke J, Starosta AL, et al. Arginine-rhamnosylation as new strategy to activate translation elongation factor P. *Nat Chem Biol*. 2015;11(4):266-70.
93. Yanagisawa T, Takahashi H, Suzuki T, Masuda A, Dohmae N, Yokoyama S. *Neisseria meningitidis* Translation Elongation Factor P and Its Active-Site Arginine Residue Are Essential for Cell Viability. *PLoS One*. 2016;11(2):e0147907.
94. Takenaka Y, Kato K, Ogawa-Ohnishi M, Tsuruhama K, Kajiura H, Yagyu K, et al. Pectin RG-I rhamnosyltransferases represent a novel plant-specific glycosyltransferase family. *Nature Plants*. 2018;4(9):669-76.
95. Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B. The carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Res*. 2014;42(Database issue):D490-5.
96. Cantarel BL, Coutinho PM, Rancurel C, Bernard T, Lombard V, Henrissat B. The Carbohydrate-Active EnZymes database (CAZy): an expert resource for Glycogenomics. *Nucleic Acids Res*. 2009;37(Database issue):D233-D8.
97. Bar-Peled M, Lewinsohn E, Fluhr R, Gressel J. UDP-rhamnose:flavanone-7-O-glucoside-2''-O-rhamnosyltransferase. Purification and characterization of an enzyme catalyzing the production of bitter compounds in citrus. *J Biol Chem*. 1991;266(31):20953-9.
98. Zong G, Fei S, Liu X, Li J, Gao Y, Yang X, et al. Crystal structures of rhamnosyltransferase UGT89C1 from *Arabidopsis thaliana* reveal the molecular basis of sugar donor specificity for UDP- $\beta$ -l-rhamnose and rhamnosylation mechanism. *The Plant Journal*. 2019;99(2):257-69.
99. Isiorho EA, Liu H-w, Keatinge-Clay AT. Structural Studies of the Spinosyn Rhamnosyltransferase, SpnG. *Biochemistry*. 2012;51(6):1213-22.
100. Sengoku T, Suzuki T, Dohmae N, Watanabe C, Honma T, Hikida Y, et al. Structural basis of protein arginine rhamnosylation by glycosyltransferase EarP. *Nat Chem Biol*. 2018;14(4):368-74.
101. He C, Liu N, Li F, Jia X, Peng H, Liu Y, et al. Complex Structure of *Pseudomonas aeruginosa*; Arginine Rhamnosyltransferase EarP with Its Acceptor Elongation Factor P. *J Bacteriol*. 2019;201(13):e00128-19.
102. Mittler M, Bechthold A, Schulz GE. Structure and action of the C-C bond-forming glycosyltransferase UrdGT2 involved in the biosynthesis of the antibiotic urdamycin. *J Mol Biol*. 2007;372(1):67-76.
103. Zhang C, Bitto E, Goff RD, Singh S, Bingman CA, Griffith BR, et al. Biochemical and structural insights of the early glycosylation steps in calicheamicin biosynthesis. *Chem Biol*. 2008;15(8):842-53.
104. Pan X, Luo J, Li S. Bacteria-Catalyzed Arginine Glycosylation in Pathogens and Host. *Front Cell Infect Microbiol*. 2020;10:185.
105. Grzegorzewicz AE, Ma Y, Jones V, Crick D, Liav A, McNeil MR. Development of a microtitre plate-based assay for lipid-linked glycosyltransferase products using the mycobacterial cell wall rhamnosyltransferase WbbL. *Microbiology (Reading)*. 2008;154(Pt 12):3724-30.
106. Hatz CF, Bally B, Rohrer S, Steffen R, Kramme S, Siegrist CA, et al. Safety and immunogenicity of a candidate bioconjugate vaccine against *Shigella dysenteriae* type 1

administered to healthy adults: A single blind, partially randomized Phase I study. Vaccine. 2015;33(36):4594-601.





Figure 1 – Examples of structurally characterised rhamnose-containing glycoconjugates found across nature. The pectic polysaccharide in plant cell walls, RG-I; O-antigens of *Shigella dysenteriae* Type I and *Shigella flexneri* 5a (RU = repeat units); Polyramnan from the S-layer glycan of *G. stearo-thermophilus* NRS 2004/3a and an S-layer glycan of *Paenibacillus alvei* CCM 2051; Lancefield antigens from Group A and B *Streptococcus*, *Streptococcus mutans* serotype C; N-glycan from *Chlorovirus* PBCV-1.

### Classical NDP-β-L-rhamnose biosynthesis



### GDP-α-D-rhamnose biosynthesis




Figure 2 – Biosynthesis pathways of nucleotide activated L- and D-Rha. NDP-β-L-rhamnose is the most common form of activated rhamnose in nature. Successive dehydration, epimerization, and reduction steps are either conducted via three enzymes (bacteria, archaea, phages and some microalgae - RmIB to RmID), or by a single or two enzymes (plants, fungi, viruses, and other eukaryotic algae - NRS/ER pathway or RHM). GDP-α-D-rhamnose biosynthesis is less prevalent and has only been reported in *Aneurinibacillus*, chlorovirus PBCV-1, and *Pseudomonas*, where D-Rha is widespread throughout the genus (71).






Figure 3 – Timeline of rhamnosyltransferase crystal structures deposited to the PDB database with their PDB IDs. Dark domains refer to the N-terminal sequence, whilst bright coloured domains highlight the C-terminal domain. (a) crystal structure of the  $\beta$ -1,2-rhamnosyltransferase, WsaF, from *Geobacillus stearothermophilus* (52); (b) crystal structure of Spinosyn 9-O- $\alpha$ -L-rhamnosyltransferase, SpnG, from *Saccharopolyspora spinosa* (99); (c) crystal structure of arginine  $\alpha$ -L-rhamnosyltransferase, EarP, from *Pseudomonas putida* (91); (d) crystal structure of arginine  $\alpha$ -L-rhamnosyltransferase, EarP, from *Neisseria meningitidis* (100); (e) crystal structure of arginine  $\alpha$ -L-rhamnosyltransferase, EarP, from *Pseudomonas aeruginosa* (101); (f) crystal structure of the flavonol  $\alpha$ -7-O-L-rhamnosyltransferase, UGT89C1, from *Arabidopsis thaliana* (98).

|            | Family     | Name    | Activity  | Host organism  | Genbank                           | Reference  |
|------------|------------|---------|---|--|-----------------------------------|------------|
| PROKARYOTE | GT1        | TiaG1   | tiacumicin 5-C-methyl-D-rhamnosyltransferase  | <i>Dactylosporangium aurantiacum</i>                         | ADU86026.1                        | (72)       |
|            | GT1        | TiaG2   | tiacumicin 2-O-methyl-D-rhamnosyltransferase  | <i>Dactylosporangium aurantiacum</i>                         | ADU85989.1                        | (72)       |
|            | GT1        | RtfA    | GPL/6-deoxytalose $\alpha$ -L-1,2-rhamnosyltransferase A                                | <i>Mycobacterium avium</i>                                   | AA44209.1                         | (18)       |
|            | GT1        | GtfA    | GPL/3,4-di-O-Me-rhamnose $\alpha$ -L-1,2-(3-O-methyl)-rhamnosyltransferase              | <i>Mycobacterium smegmatis</i>                               | AAN28688.1                        | (73)       |
|            | GT1        | Rv2962c | <i>p</i> -hydroxybenzoic acid ethyl ester rhamnosyltransferase                          | <i>Mycobacterium tuberculosis</i>                            | CAB05415.1                        | (74)       |
|            | GT1        | RhlB    | rhamnosyltransferase  | <i>Pseudomonas aeruginosa</i>                                | AAG06866.1                        | (75)       |
|            | GT1        | SpnG    | spinosyn 9-O- $\alpha$ -L-rhamnosyltransferase  | <i>Saccharopolyspora spinosa</i>                             | AAG23268.1                        | (76)       |
|            | GT1        | AraGT   | $\alpha$ -L-rhamnosyltransferase (aranciamycin synthase)                                | <i>Streptomyces echinatus</i>                                | ABL09968.1                        | (77)       |
|            | GT1        | ElmGT   | 8-demethyl-tetracenomycin C $\alpha$ -L-rhamnosyltransferase                            | <i>Streptomyces olivaceus</i>                                | CAC16413.2                        | (78)       |
|            | GT1        | StfG    | Steffimycin L-rhamnosyltransferase  | <i>Streptomyces steffisburgensis</i>                         | CAJ42338.1                        | (79)       |
| EUKARYOTE  | GT1        | UGT78D1 | flavonol-3-O-rhamnosyltransferase   | <i>Arabidopsis thaliana</i>                                  | OAP13716.1                        | (80)       |
|            | GT1        | UGT89C1 | flavonol $\alpha$ -7-O-L-rhamnosyltransferase   | <i>Arabidopsis thaliana</i>                                  | AAF80123.1                        | (21)       |
|            | GT1        | C12RT1  | flavonoid $\alpha$ -1,2-L-rhamnosyltransferase  | <i>Citrus maxima</i>   | AAU99997.1                        | (81)       |
|            | GT1        | UGT79A6 | flavonol 3-O-glucoside $\alpha$ -1,6-L-rhamnosyltransferase                             | <i>Glycine max</i>   | BAN91401.1                        | (82)       |
|            | GT1        | UGT72   | anthocyanidin-3-glucoside $\alpha$ -L-rhamnosyltransferase                              | <i>Petunia x hybrida</i>                                     | CAA50376.1                        | (83)       |
|            | GT1        | Sgt3    | $\beta$ -solanine/ $\beta$ -chaconine rhamnosyltransferase                              | <i>Solanum tuberosum</i>                                     | ABB84472.1                        | (20)       |
|            | PROKARYOTE | GT2     | WbbL  | $\alpha$ -GlcNAc-PP-C5O $\alpha$ -1,3-L-rhamnosyltransferase | <i>Mycobacterium tuberculosis</i> | AFN51271.1 |
| GT2        |            | WapR    | LPS $\alpha$ -1,3-L-rhamnosyltransferase  | <i>Pseudomonas aeruginosa</i>                                | AAG08385.1                        | (16)       |
| GT2        |            | RhIC    | L-Rha- $\beta$ -hydroxydecanoate- $\beta$ -rhamnosyltransferase                         | <i>Pseudomonas aeruginosa</i>                                | AAG04519.1                        | (85)       |
| GT2        |            | MigA    | LPS $\alpha$ -1,6-L-rhamnosyltransferase  | <i>Pseudomonas aeruginosa</i>                                | AAG04094.1                        | (16)       |
| GT2        |            | RfbF    | L-Rha $\alpha$ -1,2-rhamnosyltransferase  | <i>Shigella flexneri</i> O2A                                 | CAA50772.1                        | (86)       |
| GT2        |            | RgpB    | $\alpha$ -1,3-rhamnosyltransferase*   | <i>Streptococcus mutans</i>                                  | Q840W4                            | (87)       |
| GT2        |            | WsaC    | $\alpha$ -1,3-rhamnosyltransferase  | <i>Geobacillus stearothermophilus</i>                        | Q7BG54                            | (88)       |
| GT2        |            | WsaD    | $\alpha$ -1,3-rhamnosyltransferase  | <i>Geobacillus stearothermophilus</i>                        | Q7BG45                            | (88)       |
| GT2        |            | WsaE    | Multifunctional $\alpha$ -1,3- and $\alpha$ -1,2-rhamnosyltransferase                   | <i>Geobacillus stearothermophilus</i>                        | Q7BG51                            | (88)       |
| PROKARYOTE |            | GT4     | RgpA  | $\alpha$ -D-GlcNAc- $\beta$ -1,4-rhamnosyltransferase*       | <i>Streptococcus mutans</i>       | Q840W5     |
|            | GT4        | WsaF    | $\beta$ -1,2-rhamnosyltransferase   | <i>Geobacillus stearothermophilus</i>                        | Q7BG50                            | (88)       |
|            | GT4        | GacB    | $\alpha$ -D-GlcNAc- $\beta$ -1,4-rhamnosyltransferase                                   | <i>Streptococcus pyogenes</i>                                | N/A                               | (17)       |
| PROKARYOTE | GT102      | WbbB    | Multimodular glycosyltransferase catalyzing glycan polymerization and chain termination | <i>Raoultella terrigena</i>                                  | AAQ82931.1                        | (89)       |
| PROKARYOTE | GT104      | EarP    | arginine $\alpha$ -L-rhamnosyltransferase   | <i>Pseudomonas aeruginosa</i>                                | AAG06240.1                        | (90)       |
|            | GT104      | EarP    | arginine $\alpha$ -L-rhamnosyltransferase   | <i>Pseudomonas putida</i>                                    | AAN67476.1                        | (91)       |
|            | GT104      | EarP    | arginine $\alpha$ -L-rhamnosyltransferase   | <i>Shewanella oneidensis</i>                                 | AAN55363.1                        | (92)       |

|                   |         |       |  |                               |            |      |
|-------------------|---------|-------|--|-------------------------------|------------|------|
|                   | GT104   | EarP  | arginine $\alpha$ -L-rhamnosyltransferase                          | <i>Neisseria meningitidis</i> | BAU19337.1 | (93) |
| <b>EUKARYOTE</b>  | GT106   | RRT1  | rhamnogalacturonan-l rhamnosyltransferase                          | <i>Arabidopsis thaliana</i>   | CAC01773.1 | (94) |
|                   | GT106   | RRT2  | rhamnogalacturonan-l rhamnosyltransferase                          | <i>Arabidopsis thaliana</i>   | AEE73784.1 | (94) |
|                   | GT106   | RRT3  | rhamnogalacturonan-l rhamnosyltransferase                          | <i>Arabidopsis thaliana</i>   | AEC05682.1 | (94) |
|                   | GT106   | RRT4  | rhamnogalacturonan-l rhamnosyltransferase                          | <i>Arabidopsis thaliana</i>   | AEE29099.1 | (94) |
| <b>PROKARYOTE</b> | Unknown | RgpF  | Multidomain $\alpha$ -1,3- and $\alpha$ -1,2-rhamnosyltransferase* | <i>Streptococcus mutans</i>   | Q840W3     | (87) |
| <b>VIRAL</b>      | Unknown | A064R | Multidomain $\beta$ -1,4- and $\alpha$ -1,2-rhamnosyltransferase   | PBCV-1                        | Q89399     | (58) |