



University of Dundee

A mechanistic study of the influence of nitrogen and energy availability on the NH_4^+ sensitivity of nitrogen assimilation in *Synechococcus*

Giordano, Mario; Goodman, Charles A.; Huang, Fengying; Raven, John A.; Ruan, Zuoxi

Published in:
Journal of Experimental Botany

DOI:
[10.1093/jxb/erac219](https://doi.org/10.1093/jxb/erac219)

Publication date:
2022

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Giordano, M., Goodman, C. A., Huang, F., Raven, J. A., & Ruan, Z. (2022). A mechanistic study of the influence of nitrogen and energy availability on the NH_4^+ sensitivity of nitrogen assimilation in *Synechococcus*. *Journal of Experimental Botany*, 73(16), 5596-5611. <https://doi.org/10.1093/jxb/erac219>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

A mechanistic study of the influence of nitrogen and energy availability on the NH_4^+ sensitivity of nitrogen assimilation in *Synechococcus*

Mario Giordano^{#1,2,3,4,5}, Charles A. Goodman³, Fengying Huang¹, John A. Raven^{6,7,8} and Zuoxi Ruan^{1,2,*}

¹STU-UNIVPM Joint Algal Research Center, Marine Biology Institute, Shantou University, Shantou, Guangdong 515063, China;

²Dipartimento di Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, Ancona 60131, Italy;

³CMNS-Cell Biology and Molecular Genetics, University of Maryland, 2107 Bioscience Research Building, College Park, MD 20742-4407, USA;

⁴Institute of Microbiology ASCR, Algatech, Trebon, Czech Republic;

⁵National Research Council, Institute of Marine Science, Venezia, Italy;

⁶Division of Plant Sciences, University of Dundee at the James Hutton Institute, Invergowrie, Dundee DD2 5 DA, UK;

⁷Plant Functional Biology and Climate Change Cluster, University of Technology Sydney, Ultimo NSW 2007, Australia;

⁸School of Biological Sciences, University of Western Australia, 35 Stirling Highway, Crawley, WA 6009.

* Corresponding author: Zuoxi Ruan; email: run@stu.edu.cn; tel.: +86 754 8650 3159

Deceased

Highlights

Cyanobacterial NO_3^- assimilation is inhibited by NH_4^+ only when light is growth limiting, exploiting the lower energy cost of NH_4^+ utilization, and not exacerbating N-shortage when N supply limits growth.

Accepted Manuscript

Abstract

In most algae, NO_3^- assimilation is tightly controlled and is often inhibited by the presence of NH_4^+ . In the marine, non-colonial, non-diazotrophic cyanobacterium *Synechococcus* UTEX 2380, NO_3^- assimilation is sensitive to NH_4^+ only when N does not limit growth. We sequenced the genome of *Synechococcus* UTEX 2380, studied the genetic organization of the Nitrogen Assimilation Related (NAR) genes, and investigated expression and kinetics of the main NAR enzymes, under N or light limitation. We found that *Synechococcus* UTEX 2380 is a β -cyanobacterium with a full complement of N uptake and assimilation genes and NAR regulatory elements. The nitrate reductase (NR) of our strain showed biphasic kinetics, previously observed only in freshwater or soil diazotrophic *Synechococcus* strains. Nitrite reductase and glutamine synthetase showed little response to our growth treatments, and their activity was usually much higher than that of nitrate reductase. NH_4^+ insensitivity of NAR genes may be associated with the stimulation of the binding of the regulator NtcA to NAR genes promoters by the high 2-oxoglutarate concentrations produced under N-limitation. NH_4^+ -sensitivity in energy-limited cells fits with the fact that, under these conditions, the use of NH_4^+ rather than NO_3^- decreases N-assimilation cost, whereas it would exacerbate N-shortage under N-limitation.

Keywords

ammonium, cyanobacteria, nitrate reductase, nitrite reductase, glutamine synthetase, NtcA regulation, N metabolism, limitation

Abbreviations

NAR genes, nitrate assimilation related genes

Rubisco, Ribulose-1,5-bisphosphate carboxylase/oxygenase

RbcL, Rubisco large subunit

NrtP, nitrate/nitrite permease

nrtP, NrtP gene

NR; nitrate reductase

narB, NR gene

NiR, nitrite reductase

nirA, NiR gene

GSI, type I glutamine synthetase

glnA, GSI gene

NtcA, global nitrogen regulator

ntcA, NtcA gene

NtcB, activator of NAR genes

ntcB, NtcB gene

Pil, nitrogen regulation protein

glnB, Pil gene

PipX, Pil interacting protein

pipX, PipX gene

NrtS1 and NrtS2 (NrtS1/S2), nitrate/nitrite bispecific transporter

nrtS1 and *nrtS2*, NrtS1 and NrtS2 gene

GOGAT, glutamate synthase

2OG, 2-oxoglutarate

Glu, glutamate

Gln, glutamine

Accepted Manuscript

Introduction

Nitrate and ammonium are the most common chemical forms of combined inorganic N in nature (Giordano and Raven, 2014); the energetic costs for their assimilation are very different due to the different requirement of reductants for their assimilation into organic matter. The assimilation of one NO_3^- into glutamate requires the oxidation of 10 NADPH and the hydrolysis of 2 ATP. Assuming that the energy obtained for the oxidation of 1 NADPH is equal to that released by the hydrolysis of 3 ATP (Nitschmann and Peschek, 1986; Scherer *et al.*, 1988), this process has a cost of 32 ATP equivalents. The assimilation of 1 NH_4^+ requires 2 NADPH and 1 ATP, corresponding to about 7 ATP equivalents; i.e., 78% fewer ATP equivalents are needed for the assimilation of NH_4^+ than for that of NO_3^- (Lee and Erickson, 1987; Noctor and Foyer, 1998), although ammonium assimilation may have additional costs associated with minimizing its potential toxicity (Dai *et al.*, 2008; Collos and Harrison, 2014). Assuming a cell C:N:P:S stoichiometry of 124:16:1:1.3 (Giordano, 2013), NH_4^+ -grown cells, other things being equal, consume 13% less energy than NO_3^- -grown cells to produce the same amount of biomass (Ruan and Giordano, 2017). This becomes relevant when energy is in short supply (Ruan *et al.*, 2017).

In algae (including cyanobacteria; Giordano and Raven, 2014), the concomitant presence of NO_3^- and NH_4^+ very often leads to the shutdown of NO_3^- assimilation (Dortch, 1990; L'Helguen *et al.*, 2008). However the underlying mechanisms of this regulation appear to be different in cyanobacteria and eukaryotic algae (Fernandez and Galvan, 2008; Ohashi *et al.*, 2011). In *Chlamydomonas reinhardtii*, NH_4^+ interferes with gene activation via an impediment on the activity of the transcription factor NIT2. Cyanobacteria, commonly divided into two groups: α -cyanobacteria with Form 1A Rubisco and β cyanobacteria with Form 1B Rubisco, contain a rather complex and not fully elucidated mechanism controlling the negative regulation of NO_3^- assimilation by NH_4^+ (Ohashi *et al.*, 2011). Although β -cyanobacteria have the 2-oxoglutarate decarboxylase bypass (Zhang and Bryant, 2011; Zhang *et al.*, 2016; Wan *et al.*, 2017), the carbon flux from 2-oxoglutarate to glutamate, catalyzed by the GS/GOGAT cycle, is considered to be the dominant route (Forchhammer and Selim, 2020), especially since cyanobacteria seem to preferentially perform a noncyclic

tricarboxylic acid cycle during the light period (Esteves-Ferreira *et al.*, 2018). 2-oxoglutarate is still accepted as the metabolic signal superintending the expression of nitrate assimilation related (NAR) genes in cyanobacteria: When 2-oxoglutarate is abundant, such as when the cell experiences a shortage of reduced N, it triggers the activation of NAR genes via the transcriptional regulator NtcA (Tanigawa *et al.*, 2002). Conversely, the presence of NH_4^+ leads to a depletion of the cell pool of 2-oxoglutarate and to the interruption of the activation of NAR genes via NtcA (Muro-Pastor *et al.*, 2005; Ll acer *et al.*, 2010; Herrero and Flores, 2018). The NtcA mediated regulation of NAR genes is favored by the interaction of NtcA with the PII interacting protein (PipX), a coactivator of NtcA, which appears to stabilize the active form of NtcA (Ll acer *et al.*, 2010). When 2-oxoglutarate is low, the PipX protein preferentially binds to the PII protein; this prevents the formation of the NtcA-PipX complex and leads to a down-regulation of NAR gene expression (Espinosa *et al.*, 2014; Forcada-Nadal *et al.*, 2018). In β -cyanobacteria, when NO_3^- is scarce, the activation of some of the NAR genes is controlled via both NO_2^- and NtcA; the NO_2^- -dependent regulation of NAR gene is mediated by the transcription factor NtcB (Aichi *et al.*, 2004; Ohashi *et al.*, 2011).

The negative effect of NH_4^+ on NO_3^- assimilation can also be exerted at the level of NO_3^- transport. Cyanobacteria are equipped with an ABC-NRT type bispecific nitrate/nitrite transporter, present in most freshwater β -cyanobacteria (*Nostoc punctiforme* ATCC29133 is the only known exception), an NRT2 type nitrate/nitrite permease (NrtP), present in α -cyanobacteria apart from *Prochlorococcus* (Scanlan *et al.*, 2009; Ohashi *et al.*, 2011), or a newly found transporter NrtS, which has been found in both α and β -cyanobacteria (Maeda *et al.*, 2019). A few strains of β -cyanobacteria of the genera *Cyanothece* and *Arthrospira* possess both ABC-NRT and NrtP (see table 1 in Ohashi *et al.*, 2011). It has been shown that ABC-NRT are post-translationally regulated by NH_4^+ , via the ATP binding subunit NrtC and NrtD (Watzer *et al.*, 2019), whose long C-terminal extension is the site to which an NH_4^+ -induced negative effector binds (Kobayashi *et al.*, 1997; Kobayashi *et al.*, 2005). On the contrary, NrtP is probably NH_4^+ -insensitive, as observed for *Nostoc punctiforme* ATCC29133 (Aichi *et al.*, 2006). In this strain, nitrate reductase (NR) is NH_4^+ -sensitive and thus the NH_4^+ -inhibitory effect on NO_3^- assimilation is retained. It is not clear to what extent this fact can be generalized. No information is available for the regulation of NrtS (Maeda *et al.*, 2019).

To elucidate the mechanisms behind the inhibitory effect of NH_4^+ on NO_3^- assimilation (or the lack of it), and to attain a better understanding of the influence of energy availability on N assimilation, we cultured *Synechococcus* sp. UTEX 2380, originally isolated from coastal waters in Western Caroline Islands, Palau (see details of inorganic N, pH etc. in Ruan *et al.*, 2017), under either N or energy (light) limitation, in the presence of either NO_3^- and NH_4^+ , and conducted a genomic, enzymological, and expression-based study on the main Nitrogen Assimilation Related (NAR) genes, with special attention to nitrate reductase.

Materials and Methods

1. Culture conditions

The unicellular marine non-diazotrophic cyanobacterium *Synechococcus* UTEX LB 2380 was cultured semicontinuously as in Ruan and Giordano (2017), under a 12 h light : 12 h dark photoperiod and an irradiance of $100 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ (Ruan *et al.*, 2018). The Artificial Multipurpose Complement for the Nutrition of Algae (AMCONA) medium (Fanesi *et al.*, 2014) was modified to have two concentrations (22 and 550 μM) of either NO_3^- or NH_4^+ . At the lowest N concentration, growth was limited by N, cells grew faster if N concentration increased (up to 550 μM); at the highest N concentration, growth was limited by light (energy), when cultures were moved to an irradiance of $200 \mu\text{mol photons m}^{-2} \text{s}^{-1}$, a higher growth rate was elicited regardless of N chemical form, however this was not true for low N grown cultures (Ruan and Giordano, 2017; Ruan *et al.*, 2017). The semi-continuous cultures were diluted at 0.40 d^{-1} (22 μM nitrate), 0.35 d^{-1} (22 μM ammonium), 0.60 d^{-1} (550 μM nitrate), and 0.67 d^{-1} (550 μM ammonium) (Ruan and Giordano, 2017). These dilution rates corresponded to the maximum specific growth rates measured in batch cultures under the same culture conditions. The cultures were maintained under these conditions for at least 16 days (≥ 8 generations) prior to any measurement. Measurements were done on samples collected 3 hours after the onset of the light period.

2. Growth, protein abundance, C and N cell quotas, C and N net assimilation rates

The methods used to determine these parameters were reported in Ruan and Giordano (2017). In brief, cell numbers (from which specific growth rates were derived) and cell volume were measured with an automatic cell counter (Casy TT, Innovatis AG, Reutlingen, Germany). The protein cell content was determined according to Peterson (1977). C and N cell quotas were assessed with an elemental analyzer (EA1108, Carlo Erba Instruments, Milan, Italy). C and N assimilation rates were calculated by multiplying the cell quotas by the specific growth rates.

3. Enzyme activities and kinetics

Cells were collected by centrifugation at 16,000 g for 20 min with a Beckman J2-MC centrifuge (Beckman Coulter, Brea, USA); they were then washed twice in a solution containing 25 mM HEPES-NaOH (pH 7.0) and 1 mM KCl, and resuspended in the same solution (Herrero *et al.*, 1981; Sakamoto and Bryant, 1999).

Nitrate reductase (NR) activity assay - Prior to NR activity measurements, 1 mL of cell suspension containing $5-8 \times 10^9$ cells was whirl mixed in the presence of 20 μ L of toluene for 2 minutes to permeabilize the cells (Herrero *et al.*, 1981). The cells were assayed within 60 min after this procedure and kept on ice until used. To avoid temperature variation in the course of the assay, a 60 μ L aliquot of permeabilized cells suspension was added to a 2 mL reaction mixture pre-incubated at the assay temperature (30°C), in a thermostated bath. The temperature of this mix was allowed to equilibrate for 5 min. The reaction mixture contained: 0.1 M Na_2CO_3 - NaHCO_3 buffer (pH 10.5), 1 mM KCl, 4 mM methyl viologen, and NaNO_3 at a concentration varying from 0 to 80 mM (see Fig. 3). The reaction was initiated by the addition of 20 μ L of a 1.0 M $\text{Na}_2\text{S}_2\text{O}_4$ solution. The $\text{Na}_2\text{S}_2\text{O}_4$ solution was freshly prepared in N_2 -sparged 0.3 M NaHCO_3 . The addition of this solution caused the reduction of methyl viologen, which acted as electron provider to NR. Aliquots of 0.7 mL of the reaction mix were sampled after 1 and 4 minutes (linearity in this time range was preliminarily checked) and shaken vigorously until the blue color associated to the reduced form of methyl viologen disappeared. Samples were then centrifuged for 5 min at 13,700 g, and 0.5 mL of

the supernatant was transferred to a new Eppendorf tube. To this solution, 0.5 mL of 1% (w/v) sulfanilamide in 3 M HCl was added, followed by 0.5 mL of 0.02% (w/v) *N*-(1-naphthyl)ethylenediamine (Nicholas and Nason, 1957). The samples were incubated in the presence of these reagents for 15 minutes. Then the absorbance at 540 nm, due to interaction of the reagents with NO_2^- , was determined with a Beckman DU 640 spectrophotometer (Beckman Coulter, Brea, USA). The NO_2^- concentration was calculated by interpolating the absorbance values into a standard curve generated using 0 to 50 μM NaNO_2 .

Nitrite reductase (NiR) activity assay - NiR activity was assayed on toluene permeabilized cells, as for the NR assay. An aliquot of 0.15 mL of cell suspension containing $0.5\text{--}1.3 \times 10^9$ permeabilized cells was incubated for 5 min at 30°C in 5 ml of reaction mix, before initiating the assay. The reaction mixture contained 25 mM HEPES buffer (pH 7.0), 1 mM KCl, 20 mM NaNO_2 and 5 mM methyl viologen. The reaction was started by the addition of 100 μL of a 2 M NaNO_2 stock. The NaNO_2 concentration in our reaction mixture was much higher than that used by other authors, e.g. 100 μM (Sakamoto and Bryant, 1999), because in preliminary experiments we noticed that the NiR of our *Synechococcus* strain was saturated above 30 mM NaNO_2 . One hundred micromoles of sodium dithionite solution in 0.1 ml N_2 -sparged 0.3 M NaHCO_3 were added to start the reaction. Aliquots of 0.4 mL were taken from the reaction mixture at time 0 and after 10 min (linearity was checked preliminarily; the enzyme activity was verified to be stable for at least 15 min). The reaction was terminated by vigorous shaking to oxidize the reduced methyl viologen. The samples were then centrifuged (13,700 g for 5 min) and the supernatant was diluted 1 in 400 (v : v). Aliquots (0.5 mL) of the diluted supernatant were used for the quantification of nitrite concentration (see above *Nitrate reductase (NR) activity assay*).

Glutamine synthetase (GS) activity assay - For the GS assay, alkyltrimethylammonium bromide (MTA, Chemical Abstract Service registry NO. 8044-71-1) at a final concentration of 0.025 % (w/v) was used as the permeabilizing agent, since toluene interferes with the GS assay (Marqués *et al.*, 1989).

The assay was carried out according to Marqués *et al.* (1989), except for the killing solution and the substrate: The killing solution contained 2.5% FeCl₃ (w/v), 5% TCA (w/v) and 1.5 N HCl. NH₂OH was used as the inorganic substrate, according to Kaffes *et al.* (2010) and Oaks *et al.* (1980), substituting for the NH₄Cl used by Marqués *et al.* (1989). This was done because NH₂OH, being less volatile than the NH₃ in equilibrium with NH₄⁺, ensures a greater control of substrate concentration. The absorbance of the supernatant at 540 nm due to the reaction product (L-glutamic acid γ -monohydroxamate) was determined with a Beckman DU-64 spectrophotometer (Beckman Coulter, Brea, USA).

An aliquot of 50 μ L of permeabilized cell suspension (containing $2.5-4 \times 10^8$ cells) was added to a reaction mixture containing 50 mM HEPES (pH 7.5), 50 mM MgCl₂, 100 mM monosodium glutamate (pH 7.5), 2 mM NH₂OH·HCl (pH 7.5), 7.5 mM ATP (pH \sim 7.4), in a total volume of 1 mL. The reaction mixture was carefully prepared to make sure that the pH was always 7.5, at which ATP is stable and the enzyme activity is high (close to optimum pH) (Gawronski and Benson, 2004). The reaction took place in a water bath at 30°C and was initiated by ATP addition. After 15 min, the reaction was stopped by adding 200 μ L of killing solution to the assay mixture. The control assays were stopped at time 0. After the termination of the reaction, cells were pelleted by centrifugation at 13,700 g for 5 min, in a Heraeus *fresco* microcentrifuge (Thermo Fisher Scientific, Langenselbold, Germany). The absorbance of the supernatant at 540 nm was determined with a Beckman DU-64 spectrophotometer (Beckman Coulter, Brea, USA). The amount of L-glutamic acid- γ -monohydroxamate generated over the 15 min of the assay was determined by interpolation of absorbance in a standard curve constructed with γ -monohydroxamate (0 to 250 μ M).

GS kinetics was studied by varying both the inorganic (NH₂OH) and the organic (monosodium glutamate) substrates. One of the substrates was added to the reaction mix at saturating concentration; the other substrate was added in incremental amounts to obtain a substrate concentration response. When glutamate was the “variable” substrate, its concentration went from 2 to 200 mM. For the NH₂OH response curve, the substrate concentration range was between 60 and 3000 μ M.

4. Expression of nitrate reductase gene (*narB*)

Thirty-four amino acid sequences of NR from marine *Synechococcus* strains were downloaded from NCBI's publicly available database and were aligned using DNAMAN V6 (Lynnon Biosoft, San Raman, USA). Two highly conserved regions were identified: NNFDANSRLCMSSA and GPFSLTGQPNAMGGREAGGL. An alignment of 27 *Synechococcus narB* sequences obtained with DNAMAN was examined for codon preference. Five pairs of degenerate primers were designed according to CODEHOP (COnsensus-DEgenerate Hybrid Oligonucleotide Primer) (Rose et al., 2003). CODEHOP primers consists of a short 3' degenerate core region with three to four highly conserved amino acid residues and a longer 5' consensus clamp region that permits stable annealing during PCR amplification. Primer efficiencies were verified on serial dilutions of cDNA. LightCycler 480 Software (F. Hoffmann-La Roche Ltd, Basel, Switzerland) was used to analyze the data. Analyses of the denaturation curves for the reactions showed a single peak in all cases and no amplification products of primer-dimer formations were observed. The primer pair shown in Table S1 was used for template amplification in a Block Incubator (ASTEC BI-525A, Fukuoka, Japan), according to the protocol described in table S2. The target segment was retrieved and purified with Midi Purification Kit (DP209, Tiangen, Beijing, China) following the manufacturer's specifications and amplified again to increase the amount of product. 2 μL of the final PCR product were electrophoresed on 1.0% agarose gel to verify fragment size and quality. An additional 2 μL aliquot of PCR product was used to determine the product concentration in a Microvolume Nanodrop Spectrophotometer (NanoDrop 2000, Thermo Fisher scientific, Waltham, USA); the concentration was $> 30 \text{ ng } \mu\text{L}^{-1}$. The target product was sequenced with Illumina Hiseq (Illumina, San Diego, USA). Based on this sequence, a new pair of "confirmation" primers (Table S1) was designed with Primer Premier V5.00 (Premier Biosoft International, Palo Alto, USA). PCR products obtained with the confirmation primers were sequenced to ensure no replication error had occurred during amplification. A partial sequence of UTEX 2380 *narB* was submitted to NCBI Genebank (accession number: MH684745). This sequence was used in PrimerQuest (Integrated DAN Technology, Skokie, USA) to design RT-qPCR primers.

Total RNA was extracted from 8×10^7 cells ground in a mortar with TRNzol Universal reagent (Catalogue No. DP424, Tiangen, Beijing, China) following the manufacturer's instructions. Total RNA was resuspended in 20 μ L of DEPC water and quantified in a Nanodrop Spectrophotometer (Thermo Fisher Scientific, Waltham, USA); protein and carbohydrate contamination was assessed by determining the OD_{260}/OD_{280} and OD_{260}/OD_{230} ratio respectively; a value in the range of 1.8 ~ 2.0 for the former and a value >2.0 for the latter were considered acceptable for further processing. Samples of the extracted RNA (5 μ L) were mixed with 1 μ L of 6 \times Loading Buffer containing 0.05% bromophenol blue and 0.05% xylene cyanol FF (Tiangen, Beijing, China) and subject to electrophoresis on 1.0% agarose gels. After completion of the electrophoretic run, the gel was inspected with a Gel-Imaging System (BIO-RAD EOBR/01438, Hercules, USA) to check for RNA quality. The template RNA was mixed with 2 μ L of 5 \times gDNA buffer and RNase-free ddH₂O at 42°C to remove residual genomic DNA, then stored on ice. Reverse transcription was carried out using the FastQuant RT Kit (Cat. No. KR106, Tiangen, Beijing, China). The reverse transcription mix was composed of 10 \times Fast RT Buffer (2 μ L), RT Enzyme Mix (1 μ L), FQ-RT Primer Mix (2 μ L), and RNase-free ddH₂O (5 μ L). This mix was added to the template RNA mix, at 42°C for 15 min, and then incubated at 95°C for 3 min. For RT-qPCR, the SuperReal PreMix Plus (SYBR green) kit (Catalogue No. FP205, Tiangen, Beijing, China) was used. The RT-qPCR was conducted in a Roche LightCycler 480II (Hoffmann-La Roche Ltd, Basel, Swiss), in a reaction mix composed of 12.5 μ L of 2 \times SuperReal PreMix Plus, 0.75 μ L of forward and reverse primers (20 μ M), 5.0 μ L of cDNA (30 ng/ μ L) and 6.75 μ L of RNase-free ddH₂O. Samples were subject to 15 min pre-denaturation at 95°C, followed by forty 10-s cycles of denaturation at 95°C, annealing at 55°C for 32 s, and elongation at 72°C for 20 s. Fluorescence was measured after the elongation step. The RNase P gene (*rnpB*) (Paerl *et al.*, 2012) was used as the internal control gene, after verifying that its expression was stable under all our experimental conditions (Table S3).

5. Genome sequencing and annotation

Observing that NR activity in UTEX 2380 was insensitive to NH₄⁺ inhibition while energy was sufficient for growth, and that the UTEX 2380 enzyme showed biphasic kinetics prompted us to look for peculiarities in the sequences of N transport and assimilation genes and in their regulatory elements. We thus sequenced genomic DNA from *Synechococcus* UTEX 2380

using Illumina Hiseq XTen and Miseq platforms (Illumina, San Diego, USA). Sequencing, assembly, and annotation were carried out with the assistance of Sangon Biotech Shanghai Co. Ltd. (Shanghai, China; contract No.: GEN1815830SZ).

DNA for sequencing was prepared using a Rapid Bacterial Genomic DNA Isolation Kit (Cat. No.: B518225, Sangon Biotech, Shanghai, China) according to manufacturer instructions. DNA integrity was checked by agarose electrophoresis (0.7%, 20 V for 15 h), and concentration was determined via fluorometric assay using a Qubit 4.0 Fluorometer (Thermo Fisher Scientific, Waltham, USA). DNA was then fragmented into 300-500 bp segments with a S220 Focused-Ultrasonicator (Covaris, Woburn, USA). A genomic paired-end library was prepared with a NEB Next[®] Ultra™ DNA Library Prep Kit for Illumina (Catalogue No.: NEB#E6177, New England Biolabs, Ipswich, USA). Adaptor-ligated fragments were then purified and enriched for PCR with Agencourt AMPure XP Clean Beads. Library volume was checked by 2% agarose gel electrophoresis, and concentration was determined via Qubit 4.0 fluorometric assay. The range of DNA fragments size was determined using a 2100 DNA 1000 Kit (Recorder No.: 5067-1504, Agilent Technologies, Waldbronn, Germany).

Sequencing was completed using both Illumina Hiseq Xten and Illumina Miseq platforms. Data quality was assessed with FastQC 0.11.2 (Babraham Bioinformatics, Cambridge, UK), and artifact primer sequence and read pairs below quality threshold (where Q val.<20 and/or length <35nt) were trimmed with Trimmomatic 0.36 (The Usadel Lab of RWTH Aachen University, Aachen, Germany) (Bolger et al., 2014). Filtered reads were assembled with SPAdes 3.5.0 (Center for Algorithmic Biotechnology, St. Petersburg State University, Russia). Contig gaps were filled using GapFiller 1.11 (Boetzer and Pirovano, 2012), and then PrInSeS-G 1.0.0 (Massouras et al., 2010) was utilized to detect and correct structural variants incurred during assembly. Putative gene annotation was accomplished with the bacterial annotation pipeline Prokka 1.10 (Torsten Seemann, The University of Melbourne, Australia).

Specific genes involved in N acquisition and metabolism (*narB*, *nrtP*, *nrtS1*, *nrtS2*, *nirA*, *focA*, *glnA*, *glnN*, *glsF*, *glnB*, *ntcA*, *ntcB*, *glnB*, *pipX*) were identified by BLASTing known genes from *Synechococcus* sp. PCC 7002 (phylogenetically close to UTEX 2380, Fig. 1) against our assembly of *Synechococcus* UTEX 2380 (Altschul et al., 1990). Multiple sequence alignment was used to visually assess BLAST hit similarities. Putative matches were verified through BLASTing the NCBI Conserved Domain Database, where the genes of interest (*narB*, *nrtP*, *nrtS1*, *nrtS2*, *nirA*, *focA*, *glnA*, *glnN*, *glsF*, *glnB*, *ntcA*, *ntcB*, *glnB*, *pipX*) were found to contain the conserved domains common to other cyanobacteria. The NH₄⁺-sensitive regulatory motifs in the genome were identified by blasting the NtcA or NtcB motif against the upstream of identified N-related gene sequences. The query sequences used were GTA-N₈-TAC-N_{22/23}-TA-N₃-T (the consensus binding sites for NtcA-motif) and ATGC-N₇-GCAT-N₇-ATGC-N₇-GCAT (consensus binding sites for NtcB-motif) (Jiang et al., 2000; Ohashi et al., 2011).

For *nrtS* and *pipX*, a simple BLAST failed to determine matching sequence. In these cases, we obtained 67 and 42 homologous sequences (Table S4; Table S5), respectively. The 67 homologs of *nrtS* were derived from Maeda et al. (2019) supplementary data. The 42 homologs of *pipX* were obtained by BLASTing a known *pipX*, UniProtKB entry Q7X386_SYNE7 (PipX, *Synechococcus elongatus* strain PCC 7942), against NCBI's conserved Domain Database to identify Conserved Protein Domain Family DUF3539. All family members were obtained as a multiple sequence alignment. For each set, Hidden Markov Models were built using HMMER's hmmbuild function with standard parameters (<http://hmmer.org>). We then used HMMER's hmmsearch function to identify putative matches in our translated dataset. The *nrtS* search returned exactly two matches, with scores of 108.8 and 101.9, with E-values < 1e-30. The *pipX* search returned exactly one match, with a score of 100.2, E-value < 1e-29.

6. Classification of *Synechococcus* UTEX 2380

We sought to determine whether *Synechococcus* UTEX 2380 should be classified as an α -cyanobacterium or a β -cyanobacterium. Typically, the attribution to α - or β -cyanobacteria is based on the type of Rubisco and/or carboxysome proteins (Badger and Bek, 2008). Given

the comparatively larger availability of Rubisco sequence data relative to those of carboxysome protein, we proceeded to compare the sequence of Rubisco large subunits (Rbcl) in our genome with orthologues from a large number of strains of both α - and β -cyanobacteria. Seventy-six non-redundant cyanobacterial and two terrestrial bacterial Rbcl peptide sequences were obtained from NCBI's publicly available sequence database (Table S6). Sequence orientation was manually confirmed, and sequences were subsequently aligned using Clustal Omega, with standard parameters (Sievers *et al.*, 2011). The resulting fasta-format multiple sequence alignment was converted to phylip format via Fasta2Phylip (<https://github.com/jvollme/fasta2phylip>). The final maximum likelihood tree was generated using RAxML version 8 (Stamatakis, 2014), employing the GTR gamma substitution model with 100 bootstrap replicates.

7. Statistical methods

The programs and statistical packages used for the analysis of the genome are given above. For all other measurements, data were expressed as means and standard deviations of three biological replicates (i.e. three distinct cultures). The significance of variance was assessed by two-tailed t-test using the Origin 7.0 SR0 software (OriginLab Corporation, Northampton, USA) or by two-way ANOVA analysis followed by Student-Newman-Keuls (SNK) multiple comparison test using GMAV 5.0 for Windows XP (Underwood and Chapman, Institute of Marine Ecology, University of Sydney, Australia). In the two-way ANOVA, the significance of homogeneity of variance was always checked by Cochran's test, before analysis; if it was significant, the original data were transformed until the homogeneity of variance was not significant. The level of significance was always set at 95%.

Results

1. Growth, protein abundance, C and N cell quotas, C and N net assimilation rates

In Table S7, we summarized the information on basic cell parameters. These data were already published in (Ruan and Giordano, 2017), but are essential for the comprehension of this dataset and were thus deemed necessary to include herein. In brief, the cells cultured under energy limitation (550 μ M N) grew faster, were larger, contained more protein and had higher net rate of C and N assimilation than the N-limited cells. The C cell quota was

instead similar under the two limitations. Overall, a greater allocation of C to protein was observed in the cells grown under energy limitation (which also had more N available). The N chemical form had an impact on the various parameters in Table S7 (growth, protein abundance, C and N cell quotas, C and N net assimilation) only when energy was limiting, with higher values for all parameters in the NH_4^+ grown cells.

2. Phylogeny of *Synechococcus* UTEX 2380

Our phylogenetic analysis of the sequence of the Rubisco large subunit gene of *Synechococcus* UTEX 2380 (GenBank accession number: MN373284) conclusively places this organism among β -cyanobacteria (Tabita *et al.*, 2008) and within a single clade containing all other unicellular β -cyanobacteria examined herein (Fig. 1), with the branch dividing α and β cyanobacteria shows 100% bootstrap support. The α -cyanobacteria can be considered a monophyletic group, which previously reported multi-gene phylogenies have indicated as being nested within paraphyletic β -cyanobacteria, with *Gloeobacter* placed as sister to all other β -cyanobacteria – though a consensus has not been reached for finer details of the topology (Rae *et al.*, 2013; Sánchez-Baracaldo *et al.*, 2014; Moore *et al.*, 2019). Within β -cyanobacteria, the unicellular and the multicellular (filamentous or colonial) species generally group separately; however, the relationships between these β subsets are not sufficiently resolved in the present analysis to be described as strictly distinct clades. Four outlier species within the unicellular β subset preclude this observation: *Chroococciopsis thermalis* and *Dactylococcopsis salina* are both rooted outside the greater clade of unicellular β -cyanobacteria, with bootstrap support of 22 and 51, respectively. These values are too low to be certain of these species' placement in the presented topology. More deeply within the unicellular β -cyanobacteria, we observed *Bacillus subtilis* (Ashida *et al.*, 2003), a terrestrial non-autotrophic bacterium, placed with 100% support. *Bacillus* Rubisco-like proteins (RLP) have been reported to be involved in methionine metabolism (Sekowska *et al.*, 2019 and references therein), and it has been long recognized that Rubisco phylogenies can topologically differ from those based on other loci due to differential retention of duplicated gene copies, or as may be the case with *Bacillus*, a horizontal gene transfer (Delwiche and Palmer, 1996). Perhaps surprising is the highly supported presence of a filamentous *Leptolyngbya* within the unicellular β -cyanobacteria.

Among the included filamentous and colonial β -cyanobacteria, and excepting the aforementioned *Leptolyngbya* sp., both *Prochlororothix hollandica* and *Arthrospira* sp. are rooted outside the greater clade, with respective bootstrap values of 75 and 22. We consider these values inconclusive, and cannot be certain of their true placement at this time.

3. Nitrate assimilation genes in *Synechococcus* UTEX 2380

The Illumina HiSeq platform sequencing effort provided 2,482,535 paired-end reads, totaling 360,847,093 base pairs (bp), with an average read length of 143.1 bp. The MiSeq platform effort provided 982,685 paired-end reads totaling 237,484,177 bp in length, with reads averaging at 241.7 bp. Our *de novo* assembly was resolved to a total concatenated length of 3,478,287 bp across 95 contigs, with a mean GC-content of 49%. Contigs averaged 36,613.3 bp in length, and the current draft genome has a N50 of 80,905 base pairs. A total of 3360 protein coding genes were identified across all available databases and ranged 37 bp to 8163 bp in length. At the date of submission, 97.52% and 97.03% of identified protein coding genes have maintained accession records in the NCBI NR and TrEMBL databases, respectively. Total coding bases in the draft genome amounted to 3,035,878 bp, for a coding ratio of 87.28%. We identified 130 repeated regions, totaling 8672 bp, with an overall ratio of 0.25% repeated sequence.

Among our genes of interest, two pairs of genes were found to co-occur in their given contigs of the current assembly. The first pair included *narB* (GenBank accession No.: MN373283) and *narP* (MN373275); the second pair was constituted by *nrtS1* (MN603758) and *nrtS2* (MN603759). The remaining nitrate assimilation genes, *focA* (MN603757), *nirA* (MN373276), *glnA* (MN373277), *glnN* (MN373278), *glsF* (MN373279), *ntcA* (MN373280), *ntcB* (MN373281), *glnB* (MN373282), *pipX* (MN793312), were distributed in independent contigs. Further sequencing and analysis are required to resolve sequence gaps and determine operon structure and organization in UTEX2380.

Nitrate transporters - In the genome of *Synechococcus* UTEX 2380, we found one *nrtP* gene; no ABC type transporter was found. Upstream of *nrtP*, one NtcA and one NtcB binding motifs are present (Fig. 2). Two nitrate/nitrite bispecific transporter genes *nrtS1* and *nrtS2* (Table S4), possibly forming heteromeric transporter complexes (Maeda *et al.*, 2019), were closely related (96-100%) to the *nrtS1* and *nrtS2* of *Synechococcus* strains PCC7117, PCC7002, PCC8807, and PCC7003. In UTEX 2380, these were found to occur in tandem, sharing a reading frame and separated by 14 only amino acid residues. No typical NtcA or NtcB motif was found upstream of these two genes. In *Synechococcus* PCC7003 and NIES-970, we found only a single *nrtS* gene; the similarity of this gene to the *nrtS* of our experimental strain is much lower than that of the aforementioned *Synechococcus* strains and appreciably higher with *nrtS1* than with *nrtS2* (Table S8).

Nitrate reductase – Immediately downstream to the *nrtP* gene, a gene coding for NR was identified (*narB*), with 81-97% similarity to *narB* genes of other *Synechococcus* strains (Table S8). No other NR encoding gene was identified in the genome. Both *nrtP* and *narB* reside in the same orientation, however they are separated by a 208-nucleotide intergenic region, suggesting they occupy different reading frames. A putative NtcA-binding motif (GTA-N₉-TAC) was found between these two genes which contained one base more than normal (NtcA-binding motifs are GTA-N₈-TAC). Furthermore, a -10 promoter box is usually located 22 or 23 bases after the NtcA-binding motif, but is not present in *Synechococcus* UTEX 2380 (Fig. 2). This could indicate the presence of a pseudogenic or nonfunctional NtcA-binding motif. We were unable to identify any other putative promoter sequence in the intergenic region: Thus, *narB* and *nrtP* could potentially share an operon (Fig. 2).

Nitrite reductase – A single *nirA* gene was identified. The similarity of this gene to those of other *Synechococcus* is between 91 and 99% (Table S8). The *nirA* sequence is preceded by NtcB and NtcA binding sites (Fig. 2). Only one of the two NtcB-binding sites described by Ohashi *et al.* (2011) is present upstream of the *nirA* gene.

Glutamine synthetase (GS) - Genes encoding for type I (*glnA*) and III (*glnN*) GS were identified in the genome of *Synechococcus* UTEX 2380. Their similarity with cyanobacterial orthologs was between 84 and 99% (Table S8). A typical NtcA-binding site is present upstream of GSI, but not of GSIII (Fig. 2).

Glutamine 2-oxoglutarate aminotransferase (GOGAT) – A *glsF* gene, encoding a ferredoxin-dependent glutamate synthase (Fd-GOGAT), was identified in *Synechococcus* UTEX 2380 genome; its identity to other Fd-GOGAT of cyanobacteria ranged in 77-98 % (Table S8). No NADP-dependent GOGAT was found in the genome. No canonical NtcA and NtcB binding motifs were found upstream of the gene encoding GOGAT.

Nitrogenase – The genome analysis confirmed that this strain of *Synechococcus* does not contain genes encoding nitrogenases and is thus unable to fix N₂.

N metabolism regulators – In *Synechococcus* UTEX 2380, the sequence of *glnB*, encoding the PII protein, was almost identical to that of *Synechococcus* PCC 7117, PCC 73109, PCC 7002, and PCC 8807 (99%). The NtcA and NtcB sequences were also highly similar (99% and 98%, respectively) to those of the above-mentioned *Synechococcus* strains (Table S8). A putative NtcA binding motif was found upstream of the *ntcA* and *pipX* gene (Fig. 2; Table S5). Upstream of the *ntcB* gene, an NtcA and two NtcB binding sites were found, and the -10 promoter box overlapped with one of the NtcB binding sites.

4. Enzyme activities and kinetics

Nitrate reductase (NR)- The kinetics of NR activity was studied for NO₃⁻-grown cells (Fig. 3). NR activity showed biphasic kinetics. The first phase encompassed nitrate concentration between 0 and 1 mM; the second phase occurred between 1 and 80 mM NO₃⁻ (Fig. 3). The NR V_{max} (Fig. 3; Table S9), expressed on per cell basis, was independent of the nitrate growth concentration in phase I, but was 2.7-fold higher at 550 μM nitrate (energy limited cells) in

phase II. If V_{\max} was expressed on a per protein basis, however, the value for phase II was about 3-fold higher in N-limited cells than that in their energy-limited counterparts. The $K_m(\text{NO}_3^-)$ was lower at low N (22 μM NO_3^- , nitrogen-limited), with values that were about 75% and 48% those of 550 μM NO_3^- -grown cells, in phase I and phase II respectively (Fig 3; Table S9). Due to resource constraints, we were unable to study the kinetics of NR from N-limited NH_4^+ -grown cells.

Interestingly, NR activity was present in NH_4^+ -grown cells under N-limitation (22 μM NH_4^+); the typical inhibitory effect of NH_4^+ on NR expression/activity was thus not observed in *Synechococcus* UTEX 2380 under N limitation. The NR V_{\max} of N-limited NH_4^+ -grown cells was about half of that measured in the same conditions for NO_3^- -grown cells (Fig. 3; Table S9). Under 550 μM NH_4^+ (and energy limitation), no NR activity was detected in the presence of NH_4^+ .

Nitrite reductase (NiR) - NiR activity was two to three orders of magnitude higher than that of NR, under all conditions (Fig. 4), with very low affinity for NO_2^- (in the mM range) and very high saturating concentration (>40 mM) (data not shown). This may imply that the intracellular NO_2^- concentration could be very low to allow NiR activity to match that of NR. NiR V_{\max} , whether expressed on a per cell or a per protein basis, had similar rates in all treatments, with only slight differences (Fig. 4; table S9).

Glutamine synthetase (GS) – GS response to substrate concentration was studied for both glutamate (Fig. 5) and NH_2OH (Fig. S1). When glutamate concentration varied and NH_2OH was saturating, the V_{\max} of GS responded to the N concentration, but not to the N chemical form. Only when V_{\max} was expressed on a protein basis, it was significantly higher in cells cultured in the presence of NO_3^- , under energy limitation and high N (Fig. 5; Table S9). The V_{\max} of low N grown cells was 2.0- (NO_3^-) and 1.7- (NH_4^+) fold higher than at high N on a per cell basis, and 2.4- (NO_3^-) to 3.7- (NH_4^+) fold on a protein basis (Table S9). The $K_m(\text{Glu})$ of GS was not affected by either the N chemical form (NO_3^- or NH_4^+) or the N concentration (22 or

550 μM). The same trends for both V_{max} and K_m were observed, except for K_m between N-limited and energy-limited cultures grown on NH_4^+ , when the NH_2OH concentration was varied and glutamate was maintained constant at saturating concentrations (Fig. S1; Table S9).

5. Nitrate reductase expression

The presence of NR activity on NH_4^+ -grown cells prompted us to check if the expression of the *narB* gene, encoding nitrate reductase, followed the same trend (Fig. 6). At low N, no obvious difference in *narB* gene expression was shown between NO_3^- - and NH_4^+ -grown cells. At high N, *narB* expression in NH_4^+ -grown cells is indistinguishable from zero; in NO_3^- -grown cells, *narB* expression is much higher at high N than at low N.

Discussion

The inhibitory role of even small amounts of NH_4^+ on nitrate acquisition and assimilation in cyanobacteria has been known for many years (e.g. Dortch, 1990; Mérida *et al.*, 1991; Luque *et al.*, 1994). This is also the case for *Synechococcus* sp. UTEX 2380, under energy limitation. When, however, this strain is subject to N-limitation (Table S7), nitrate is reduced even in the presence of NH_4^+ (Fig. 4). Our work was aimed at understanding the mechanisms that give rise to this phenotype.

Are transporters involved in the NH_4^+ insensitivity of nitrate reductase?

From a metabolic point of view, the production and consumption of 2-oxoglutarate are key control points in N assimilation (Fig. 7): when reduced N is low within the cell, the 2-oxoglutarate pool is large and it favors the binding of NtcA to the promoters of most NAR operons, inducing their expression. When reduced N is generated in high amount, 2-oxoglutarate is aminated to glutamate and the 2-oxoglutarate pool size decreases and NtcA binding to NAR genes regulatory sites is downregulated (Herrero *et al.*, 2001; Forchhammer, 2008; Herrero and Flores, 2018; Forchhammer and Selim, 2020). In some organisms, the 2-oxoglutarate pool size is also at the basis of post-translational modification of ABC-type NRT transport systems, whose activity is blocked in the presence of NH_4^+ (Ohashi *et al.*, 2011).

Some NRT2 transporters (NrtP) found in cyanobacteria do not appear to be sensitive to NH_4^+ . Our biochemical analysis did not allow us to directly test whether transport was affected by the presence of NH_4^+ , since our activity assays were performed on permeabilized cells. The analysis of the genome showed that one *nrtP* gene and no ABC-type NRT genes encoding nitrate transporters are present in *Synechococcus* UTEX 2380. This is in agreement with previous reports for marine cyanobacteria (Ohashi *et al.*, 2011). If the *Synechococcus* UTEX 2380 NrtP transporter behaves like a typical cyanobacterial NRT2, then NH_4^+ should not have any effect on transport. It is worth noticing, however, that Sakamoto *et al.* (1999) reported on the NH_4^+ sensitivity of *nrtP* gene expression in *Synechococcus* PCC 7002, a strain that our Rubisco phylogeny places close to *Synechococcus* UTEX 2380. In *Synechococcus* UTEX 2380, *nrtP* is likely to be in the same operon with *narB*; the promoter region upstream of *nrtP* has typical NtcA and NtcB binding motifs, while the intergenic region separating it from *narB* contains only an atypical NtcA-binding motif that lacks a -10 box; we thus might expect *narB* and *nrtP* to show a similar expression pattern controlled primarily by the NtcA and NtcB systems. However, in *Nostoc punctiforme* ATCC29133, in which *nrtP* and *narB* share an operon with *nirA* (Ohashi *et al.*, 2011), *nrtP* is NH_4^+ insensitive, whereas *narB* appears to be down-regulated or inhibited in the presence of 3.75 mM NH_4^+ ; unfortunately, no data are available on *N. punctiforme* at limiting N (Aichi *et al.*, 2006). In the case of UTEX 2380, NR is expressed and active in the presence of NH_4^+ in N-limited cells, but not at high N when cells are energy limited. We currently lack data on *nrtP* expression, thus cannot verify whether NH_4^+ sensitivity of *nrtP* expression is concentration-dependent.

Does the NR of Synechococcus UTEX 2380 have structural or regulatory anomalies?

The insensitivity of NR to NH_4^+ in cyanobacteria has only been reported in rare cases, such as those of the freshwater β -cyanobacteria *Synechocystis* PCC 6803 (Kobayashi *et al.*, 2005) and *Cyanothece* PCC8801 (previously classified as *Synechococcus* sp. RF-1 in Wang *et al.*, 2003). It was proposed that these NRs are structurally different from the NH_4^+ sensitive NRs (Kobayashi *et al.*, 2005; Ohashi *et al.*, 2011). In our strain, only one *narB* gene was found. Its derived NR protein sequence had only 63% similarity with the NR of *Synechocystis* sp. PCC 6803 (although it shares with it biphasic kinetics; Martin-Nieto *et al.*, 1992), whereas its protein sequence was identical for 99% to that of the NR of *Synechococcus* sp. PCC 7002

(Table S8), which is sensitive to NH_4^+ (Sakamoto and Bryant, 1999). It is therefore unlikely that the NH_4^+ insensitivity of *Synechococcus* UTEX 2380 NR is associated with structural differences of the enzymes.

As we mentioned earlier, the *nrtP* promoter region, upstream of *narB*, contains typical NtcA and NtcB binding motifs. This is not *per se* sufficient proof of NH_4^+ sensitivity, since the binding of NtcA to regulatory sites is mediated by 2-oxoglutarate through PII and PipX proteins (Fig. 7, Ll acer *et al.*, 2010; Herrero and Flores, 2018). Abundant 2-oxoglutarate interferes with the interaction of PipX with PII and favors the binding of PipX to NtcA protein increasing NtcA activity as gene inducer (Espinosa *et al.*, 2006; Espinosa *et al.*, 2007; Ll acer *et al.*, 2010; Zhao *et al.*, 2010; Espinosa *et al.*, 2014). It has been reported that in the freshwater *Synechococcus elongatus* PCC7942, when the *glnB* gene, which produces the PII protein, is inactivated, NH_4^+ sensitivity disappears (Lee *et al.*, 1998). We may thus expect that a non-functional *glnB* gene may contribute to generating the phenotype we observed. However, no obvious defect was observed in the *glnB* gene of *Synechococcus* sp. UTEX 2380.

Proposed mechanism for metabolic regulation of N assimilation

Although no obvious peculiarities could be identified in the NAR genes of *Synechococcus* UTEX 2380, NH_4^+ sensitivity of NR in this cyanobacterium is suppressed under N limitation (Fig. 4; Fig. 6). The high 2-oxoglutarate concentration attained under N-limitation (Jiang *et al.*, 2000; Ludwig and Bryant, 2012) appears to favor *narB* gene induction, even in the presence of NH_4^+ . The fact that a NtcA binding site is present upstream of the *ntcA* gene itself is suggestive of an autocatalytic effect of NtcA on its own expression (Jiang *et al.*, 2000) that may amplify the induction of this gene under high 2-oxoglutarate. It is noteworthy, however, that NtcA binding to its target motif is under the control of redox status, only when the thiol groups of NtcA is in the reduced state (e.g. cells in N-limitation), this autocatalysis of NtcA may function (Alfonso *et al.*, 2001). Besides, we found that PipX, a coactivator of *ntcA*, is also promoted by NtcA. Consequently, these would enhance *narB* (and possibly other NAR genes) expression (Luque *et al.*, 1994; Luque *et al.*, 2004). It is only when the reduced N in the cell is high (see Giordano and Raven, 2014 for a detailed discussion on N algal metabolism) and the level of 2-oxoglutarate falls, that the interference

of 2-oxoglutarate with the interaction of PipX with NtcA would downregulate, and possibly block, the expression of *narB* (Espinosa *et al.*, 2014). High NO_3^- availability would not have the same effect as high NH_4^+ concentration due to the presence, in addition of the NtcA regulatory system, of a NO_2^- -dependent induction system, based on the NtcB protein (Aichi and Omata, 1997; Aichi *et al.*, 2001), which would maintain the expression of *narB* when NO_2^- is produced from NO_3^- reduction. NtcB binding motifs are present in the promoter region of both the *nrtP-narB* operon and the *nirA* operon, in *Synechococcus* UTEX 2380. In the case of the *nirA* operon of *Synechococcus elongatus* PCC 7942, it was shown that it ceases to respond to NO_2^- if the NtcB system is disrupted (Aichi and Omata, 1997); however, a complete functional NtcA system is required for NtcB function (Maeda *et al.*, 1998; Ohashi *et al.*, 2011 and references therein).

The regulatory mechanism proposed above (Fig. 7) makes ecological sense, because it would allow cells not to discriminate for the N- NO_3^- in the presence of NH_4^+ under severe N-limitation. N-limitation is frequent in the open ocean, where NH_4^+ must be present at appreciable concentrations (although low), as it is demonstrated by the fact that many oceanic *Prochlorococcus* strains are unable to use NO_3^- and are only able to use NH_4^+ (Moore *et al.*, 2002). In such an environment, an NH_4^+ -triggered impediment to NO_3^- assimilation would most likely lead to N starvation. Furthermore, under these conditions, the use of NH_4^+ rather than NO_3^- would bring no advantage, since energy would not be controlling growth (Ruan and Giordano, 2017; Ruan *et al.*, 2017; Ruan *et al.*, 2018). An NH_4^+ -mediated control only becomes relevant in N-rich and energy-limiting environments (Ruan and Giordano, 2017; Ruan *et al.*, 2017).

Impact of energy limitation on N metabolism

NR was not expressed at high NH_4^+ concentration in *Synechococcus* UTEX 2380. However, under energy limitation NiR and GS were always present and active, though their activity was modulated by growth conditions (Fig. 4 and 5; Table S9). It is known that NiR (especially) and GS activity are often constitutively high to allow the prompt removal from the intracellular environment of the potentially harmful NO_2^- and NH_4^+ , and are little

responsive to the chemical form in which N is available outside the cell (Bird and Wyman, 2003; Wyman and Bird, 2007). The fact that the *nirA* gene is preceded in the genome by NtcA and NtcB binding sites suggests that it occupies a distinct operon from *narB*; this may play a role in the finding that, at high N, NiR does not show NH_4^+ sensitivity. In support of this idea (although, admittedly, not conclusively), in organisms in which the genes encoding NR and NiR are in the same operon, such as *Synechococcus elongatus* PCC 7942, both are NH_4^+ sensitive (Forchhammer and de Marsac, 1995; Ohashi *et al.*, 2011); on the contrary NR and NiR appear to respond independently to the N-source in *Synechococcus* WH 8103, where the two genes are located in distinct operons (Bird and Wyman, 2003). In the case of glutamine synthetase, the situation is complicated by the presence of two genes, *glnA* and *glnN*. *GlnA* appears to be under the control of the NtcA system, but not under that of the NtcB system, whereas neither *ntcA* nor *ntcB* binding motifs were found upstream of the *glnN* gene (see Fig. 2; Jiang *et al.*, 2000).

The nitrate reductase of Synechococcus sp. UTEX 2380 shows biphasic kinetics

At the end of this paper, it seems useful to spend a few words on the biphasic (multimodal sensu Nissen and Martín-Nieto, 1998) kinetics of *Synechococcus* sp. UTEX 2380 NR. Biphasic NRs have already been reported for cyanobacteria, but always in freshwater or soil nitrogen fixers (Martín-Nieto *et al.*, 1992; Wang *et al.*, 2003; Wang *et al.*, 2011). Up to this point in time, *Synechococcus* sp. UTEX 2380 is the only non-diazotrophic and marine cyanobacterium with a biphasic NR. The K_m for nitrate in phase I ($K_{mI}(\text{NO}_3^-) = 24\text{-}31 \mu\text{M}$) and phase II ($K_{mII}(\text{NO}_3^-) = 2.14\text{-}4.44 \text{ mM}$) (Table S9) were lower than those reported for other biphasic NRs ($K_{mI} = 50\text{-}550 \mu\text{M}$; $K_{mII} = 4.2\text{-}22 \text{ mM}$) (Martín-Nieto *et al.*, 1992; Nissen and Martín-Nieto, 1998; Wang *et al.*, 2003; Wang *et al.*, 2011), and cells cultured under N-limitation had a higher affinity for NO_3^- than cells cultured in N-replete conditions (Table S9). *Synechococcus* sp. UTEX 2380 possesses only one gene encoding NR (*narB*). The biphasic kinetics must therefore be due to allosteric regulation. This seems to be the case for all the biphasic cyanobacterial NR studied so far in sufficient detail (Martín-Nieto *et al.*, 1992; Wang *et al.*, 2003; Wang *et al.*, 2011). The biphasic NR of *Cyanothece* sp. PCC8801 (previously recognized as *Synechococcus* sp. RF-1; Wang *et al.*, 2003) and *Synechocystis* PCC 6803 (Kobayashi *et al.*, 2005) are NH_4^+ -insensitive at high N (4 mM N), unlike the NR of our

Synechococcus sp. UTEX 2380, which is insensitive only under N-limitation. Whether strains with biphasic kinetics have a common phylogeny is unclear. Nissen and Martín-Nieto (1998) pointed out that the strains with biphasic NR they analyzed were all heterocystous N₂ fixers and phylogenetically related. For each of the Martín-Nieto *et al.* (1992) species with an available Rubisco sequence, our phylogenetic analysis indicates that biphasic species – both filamentous and unicellular – are placed among the β -cyanobacteria, while the single monophasic species was an α -cyanobacterium. *Synechococcus* sp. UTEX2380, identified herein as a β -cyanobacterium, however, is not a N₂ fixer and does not have heterocysts. Therefore, if biphasic NRs follow a common phylogenetic trajectory, *Synechococcus* sp. UTEX 2380 may have lost its *nif* genes.

Beta-cyanobacteria include freshwater, coastal and terrestrial species (e.g. in desert crusts and in lichens), while the α -cyanobacteria include only marine strains of *Synechococcus* and *Prochlorococcus*. This is relevant to our work because α - and β -cyanobacteria also differ for features related to N acquisition and assimilation, although some characteristics (e.g. the type of NO₃⁻ transporter) appears to be more influenced by ecology than by phylogeny, i.e. marine strain of both α - and β -cyanobacteria tend to have NrtP transporters only, whereas the freshwater strains are usually equipped with ABC-type transporters, with rare exceptions (*Cyanothece* sp. and *Arthrospira platensis*) (Ohashi *et al.*, 2011 and references therein). This however requires further confirmation by increasing the number of species analyzed; this work also provides a contribution in this direction. It is noteworthy that different from α -cyanobacteria with NtcA as the sole transcription activator for NAR genes, β -cyanobacteria own a dual activation mechanism involving both NtcA and NtcB. NtcA *per se* can not fully activate NAR genes (Aichi and Omata, 1997; Maeda *et al.*, 1998; Aichi *et al.*, 2001); the coactivator NtcB play an essential role, nitrate assimilation capability of its mutant may decrease by 90%, e.g. in *Synechococcus* sp. PCC7002 (Sakamoto *et al.*, 2008). However, whether and how much this regulator affects NH₄⁺ sensitivity is not clear. Since NtcB is also subjected to NtcA regulation, we may thus expect that the control of NH₄⁺ over nitrate acquisition and assimilation in β -cyanobacteria are mainly via NtcA as described above, albeit different organization of NAR genes. Besides, there are relatively few related

researches available on α -cyanobacteria (Bird and Wyman, 2003; Wyman and Bird, 2007), it is still very hard to tell the difference of NH_4^+ sensitivity between α - and β -cyanobacteria.

The ecological relevance of biphasic kinetics is not obvious: millimolar concentrations of NO_3^- are unlikely to occur in the environment, except in rare cases (Bruland *et al.*, 2008). This may become relevant if NR operates in an intracellular environment where NO_3^- is accumulated at high concentration. There is evidence that some cyanobacteria are capable of accumulating nitrate intracellularly at concentration as high as 0.2-1.5 mM (Lara *et al.*, 1987; Kobayashi *et al.*, 1997), a range compatible with the phase II kinetics of the biphasic NRs. Data are insufficient to say whether organisms with biphasic NR are capable of high NO_3^- accumulation.

Accepted Manuscript

Acknowledgements

The University of Dundee is a registered Scottish charity, NO 015096.

Mario Giordano died on December 28, 2019. Mario was an innovative scientist, a mentor who always had time for junior colleagues, and a generous and supportive colleague. His premature death is a loss to science, and especially a loss to his colleagues and friends.

Authors contribution

MG formulated the hypothesis, made the experimental design, analyzed the results, wrote/revised the paper, and provided funding for this work; CAG analyzed and interpreted the genomic data, contributed to write/revise the manuscript; FH carried out the RT-qPCR measurements; JAR contributed to the analysis and revised the manuscript; ZR carried out the experiments, analyzed the results and contributed to write/revise the manuscript.

Conflict of interest

The authors declare no conflict of interest.

Funding

MG was funded by GAČR 16-16343S, Guangdong Province Universities and Colleges Pearl River Scholar Funded Scheme (2017). ZR was funded by a PhD scholarship awarded by Università Politecnica delle Marche, National natural Science Foundation of China (42076206) and Guangdong Basic and Applied Basic Research Foundation (2020A1515011073). CAG was funded by NIH NIGMS T32-GMN080201-09.

Data availability

The data supporting the findings of this study are available from the corresponding author Zuoxi Ruan upon request.

References

- Aichi M, Maeda S-I, Ichikawa K, Omata T.** 2004. Nitrite-responsive activation of the nitrate assimilation operon in cyanobacteria plays an essential role in up-regulation of nitrate assimilation activities under nitrate-limited growth conditions. *Journal of Bacteriology* **186**, 3224-3229.
- Aichi M, Omata T.** 1997. Involvement of NtcB, a LysR family transcription factor, in nitrite activation of the nitrate assimilation operon in the cyanobacterium *Synechococcus* sp. strain PCC 7942. *Journal of Bacteriology* **179**, 4671-4675.
- Aichi M, Takatani N, Omata T.** 2001. Role of NtcB in activation of nitrate assimilation genes in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *Journal of Bacteriology* **183**, 5840-5847.
- Aichi M, Yoshihara S, Yamashita M, Maeda S, Nagai K, Omata T.** 2006. Characterization of the nitrate-nitrite transporter of the major facilitator superfamily (the *nrtP* gene product) from the cyanobacterium *Nostoc punctiforme* strain ATCC 29133. *Bioscience, Biotechnology, and Biochemistry* **70**, 2682-2689.
- Alfonso M, Perewoska I, Kirilovsky D.** 2001. Redox control of *ntcA* gene expression in *Synechocystis* sp. PCC 6803. Nitrogen availability and electron transport regulate the levels of the NtcA protein. *Plant Physiology* **125**, 969-981.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ.** 1990. Basic local alignment search tool. *Journal of molecular biology* **215**, 403-410.
- Ashida H, Saito Y, Kojima C, Kobayashi K, Ogasawara N, Yokota A.** 2003. A functional link between RuBisCO-like protein of *Bacillus* and photosynthetic RuBisCO. *Science* **302**, 286-290.
- Badger MR, Bek EJ.** 2008. Multiple Rubisco forms in proteobacteria: their functional significance in relation to CO₂ acquisition by the CBB cycle. *Journal of Experimental Botany* **59**, 1525-1541.
- Bird C, Wyman M.** 2003. Nitrate/nitrite assimilation system of the marine picoplanktonic cyanobacterium *Synechococcus* sp. strain WH 8103: effect of nitrogen source and availability on gene expression. *Applied and Environmental Microbiology* **69**, 7009-7018.
- Boetzer M, Pirovano W.** 2012. Toward almost closed genomes with GapFiller. *Genome biology* **13**, R56.

- Bolger AM, Lohse M, Usadel B.** 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114-2120.
- Bruland KW, Lohan MC, Aguilar - Islas AM, Smith GJ, Sohst B, Baptista A.** 2008. Factors influencing the chemistry of the near-field Columbia River plume: Nitrate, silicic acid, dissolved Fe, and dissolved Mn. *Journal of Geophysical Research: Oceans* **113**, doi:10.1029/2007JC004702.
- Collos Y, Harrison PJ.** 2014. Acclimation and toxicity of high ammonium concentrations to unicellular algae. *Marine Pollution Bulletin* **80**, 8-23.
- Dai G, Deblois CP, Liu S, Juneau P, Qiu B.** 2008. Differential sensitivity of five cyanobacterial strains to ammonium toxicity and its inhibitory mechanism on the photosynthesis of rice-field cyanobacterium Ge-Xian-Mi (*Nostoc*). *Aquatic Toxicology* **89**, 113-121.
- Delwiche CF, Palmer JD.** 1996. Rampant horizontal transfer and duplication of Rubisco genes in eubacteria and plastids. *Molecular Biology and Evolution* **13**, 873-882.
- Dortch Q.** 1990. The interaction between ammonium and nitrate uptake in phytoplankton. *Marine Ecology Progress Series* **61**, 183-201.
- Espinosa J, Forchhammer K, Burillo S, Contreras A.** 2006. Interaction network in cyanobacterial nitrogen regulation: PipX, a protein that interacts in a 2-oxoglutarate dependent manner with PII and NtcA. *Molecular Microbiology* **61**, 457-469.
- Espinosa J, Forchhammer K, Contreras A.** 2007. Role of the *Synechococcus* PCC 7942 nitrogen regulator protein PipX in NtcA-controlled processes. *Microbiology* **153**, 711-718.
- Espinosa J, Rodríguez-Mateos F, Salinas P, Lanza VF, Dixon R, de la Cruz F, Contreras A.** 2014. PipX, the coactivator of NtcA, is a global regulator in cyanobacteria. *Proceedings of the National Academy of Sciences* **111**, E2423-E2430.
- Esteves-Ferreira AA, Inaba M, Fort A, Araújo WL, Sulpice R.** 2018. Nitrogen metabolism in cyanobacteria: metabolic and molecular control, growth consequences and biotechnological applications. *Critical Reviews in Microbiology* **44**, 541-560.
- Fanesi A, Raven JA, Giordano M.** 2014. Growth rate affects the responses of the green alga *Tetraselmis suecica* to external perturbations. *Plant, Cell and Environment* **37**, 512-519.

- Fernandez E, Galvan A.** 2008. Nitrate assimilation in *Chlamydomonas*. *Eukaryotic Cell* **7**, 555-559.
- Forcada-Nadal A, Llácer JL, Contreras A, Marco-Marín C, Rubio V.** 2018. The PII-NAGK-PipX-NtcA regulatory axis of cyanobacteria: a tale of changing partners, allosteric effectors and non-covalent interactions. *Frontiers in Molecular Biosciences* **5**, doi: 10.3389/fmolb.2018.00091.
- Forchhammer K.** 2008. PII signal transducers: novel functional and structural insights. *Trends in microbiology* **16**, 65-72.
- Forchhammer K, de Marsac NT.** 1995. Functional analysis of the phosphoprotein PII (*glnB* gene product) in the cyanobacterium *Synechococcus* sp. strain PCC 7942. *Journal of Bacteriology* **177**, 2033-2040.
- Forchhammer K, Selim KA.** 2020. Carbon/nitrogen homeostasis control in cyanobacteria. *FEMS Microbiology Reviews* **44**, 33-53.
- Gawronski JD, Benson DR.** 2004. Microtiter assay for glutamine synthetase biosynthetic activity using inorganic phosphate detection. *Analytical biochemistry* **327**, 114-118.
- Giordano M.** 2013. Homeostasis: an underestimated focal point of ecology and evolution. *Plant Science* **211**, 92-101.
- Giordano M, Raven JA.** 2014. Nitrogen and sulfur assimilation in plants and algae. *Aquatic Botany* **118**, 45-61.
- Herrero A, Flores E.** 2018. Genetic responses to carbon and nitrogen availability in *Anabaena*. *Environmental Microbiology* **21**, 1-17.
- Herrero A, Flores E, Guerrero MG.** 1981. Regulation of nitrate reductase levels in the cyanobacteria *Anacystis nidulans*, *Anabaena* sp. strain 7119, and *Nostoc* sp. strain 6719. *Journal of Bacteriology* **145**, 175-180.
- Herrero A, Muro-Pastor AM, Flores E.** 2001. Nitrogen control in cyanobacteria. *Journal of Bacteriology* **183**, 411-425.
- Jiang F, Wisén S, Widersten M, Bergman B, Mannervik B.** 2000. Examination of the transcription factor NtcA-binding motif by in vitro selection of DNA sequences from a random library. *Journal of Molecular Biology* **301**, 783-793.
- Kaffes A, Thoms S, Trimborn S, Rost B, Langer G, Richter KU, Köhler A, Norici A, Giordano M.** 2010. Carbon and nitrogen fluxes in the marine coccolithophore *Emiliana huxleyi* grown under different nitrate concentrations. *Journal of Experimental Marine Biology and Ecology* **393**, 1-8.

- Kobayashi M, Rodríguez R, Lara C, Omata T.** 1997. Involvement of the C-terminal domain of an ATP-binding subunit in the regulation of the ABC-type nitrate/nitrite transporter of the cyanobacterium *Synechococcus* sp. strain PCC 7942. *Journal of Biological Chemistry* **272**, 27197-27201.
- Kobayashi M, Takatani N, Tanigawa M, Omata T.** 2005. Posttranslational regulation of nitrate assimilation in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *Journal of Bacteriology* **187**, 498-506.
- L'Helguen S, Maguer J-F, Caradec J.** 2008. Inhibition kinetics of nitrate uptake by ammonium in size-fractionated oceanic phytoplankton communities: implications for new production and f-ratio estimates. *Journal of Plankton Research* **30**, 1179-1188.
- Lara C, Romero JM, Guerrero MG.** 1987. Regulated nitrate transport in the cyanobacterium *Anacystis nidulans*. *Journal of Bacteriology* **169**, 4376-4378.
- Lee H, Erickson L.** 1987. Theoretical and experimental yields for photoautotrophic, mixotrophic, and photoheterotrophic growth. *Biotechnology and Bioengineering* **29**, 476-481.
- Lee HM, Flores E, Herrero A, Houmard J, Tandeau de Marsac N.** 1998. A role for the signal transduction protein PII in the control of nitrate/nitrite uptake in a cyanobacterium. *FEBS Letters* **427**, 291-295.
- Llácer JL, Espinosa J, Castells MA, Contreras A, Forchhammer K, Rubio V.** 2010. Structural basis for the regulation of NtcA-dependent transcription by proteins PipX and PII. *Proceedings of the National Academy of Sciences* **107**, 15397-15402.
- Ludwig M, Bryant DA.** 2012. Acclimation of the global transcriptome of the cyanobacterium *Synechococcus* sp. strain PCC 7002 to nutrient limitations and different nitrogen sources. *Frontiers in Microbiology* **145**, 1-15.
- Luque I, Flores E, Herrero A.** 1994. Molecular mechanism for the operation of nitrogen control in cyanobacteria. *The EMBO journal* **13**, 2862-2869.
- Luque I, Vázquez-Bermúdez MF, Paz-Yepes J, Flores E, Herrero A.** 2004. In vivo activity of the nitrogen control transcription factor NtcA is subjected to metabolic regulation in *Synechococcus* sp. strain PCC 7942. *FEMS Microbiology Letters* **236**, 47-52.
- Mérida A, Candau P, Florencio FJ.** 1991. Regulation of glutamine synthetase activity in the unicellular cyanobacterium *Synechocystis* sp. strain PCC 6803 by the nitrogen source: effect of ammonium. *Journal of Bacteriology* **173**, 4095-4100.

- Maeda S-I, Aoba R, Nishino Y, Omata T.** 2019. A novel bacterial nitrate transporter composed of small transmembrane proteins. *Plant and Cell Physiology* **60**, 2180-2192.
- Maeda S-I, Kawaguchi Y, Ohe T-A, Omata T.** 1998. Cis-acting sequences required for NtcB-dependent, nitrite-responsive positive regulation of the nitrate assimilation operon in the cyanobacterium *Synechococcus* sp. strain PCC 7942. *Journal of Bacteriology* **180**, 4080-4088.
- Marqués S, Florencio FJ, Candau P.** 1989. Ammonia assimilating enzymes from cyanobacteria: In situ and in vitro assay using high-performance liquid chromatography. *Analytical Biochemistry* **180**, 152-157.
- Martin-Nieto J, Flores E, Herrero A.** 1992. Biphasic kinetic behavior of nitrate reductase from heterocystous, nitrogen-fixing cyanobacteria. *Plant Physiology* **100**, 157-163.
- Massouras A, Hens K, Gubelmann C, Uplekar S, Decouttere F, Rougemont J, Cole ST, Deplancke B.** 2010. Primer-initiated sequence synthesis to detect and assemble structural variants. *Nature methods* **7**, 485-486.
- Moore KR, Magnabosco C, Momper LM, Gold DA, Bosak T, Fournier GP.** 2019. An expanded ribosomal phylogeny of cyanobacteria supports a deep placement of plastids. *Frontiers in Microbiology* **10**, 1612.
- Moore LR, Post AF, Rocap G, Chisholm SW.** 2002. Utilization of different nitrogen sources by the marine cyanobacteria *Prochlorococcus* and *Synechococcus*. *Limnology and Oceanography* **47**, 989-996.
- Muro-Pastor MI, Reyes JC, Florencio FJ.** 2005. Ammonium assimilation in cyanobacteria. *Photosynthesis Research* **83**, 135-150.
- Nicholas D, Nason A.** 1957. Determination of nitrate and nitrite. *Methods in Enzymology* **3**, 981-984.
- Nissen P, Martín-Nieto J.** 1998. 'Multimodal' kinetics: Cyanobacterial nitrate reductase and other enzyme, transport and binding systems. *Physiologia Plantarum* **104**, 503-511.
- Nitschmann WH, Peschek GA.** 1986. Oxidative phosphorylation and energy buffering in cyanobacteria. *Journal of Bacteriology* **168**, 1205-1211.
- Noctor G, Foyer CH.** 1998. A re-evaluation of the ATP: NADPH budget during C3 photosynthesis: a contribution from nitrate assimilation and its associated respiratory activity? *Journal of Experimental Botany* **49**, 1895.

- Oaks A, Stulen I, Jones K, Winspear MJ, Misra S, Boesel IL.** 1980. Enzymes of nitrogen assimilation in maize roots. *Planta* **148**, 477-484.
- Ohashi Y, Shi W, Takatani N, Aichi M, Maeda S, Watanabe S, Yoshikawa H, Omata T.** 2011. Regulation of nitrate assimilation in cyanobacteria. *Journal of Experimental Botany* **62**, 1411-1424.
- Paerl RW, Tozzi S, Kolber ZS, Zehr JP.** 2012. Variation in the abundance of *Synechococcus* sp. cc9311 narB mRNA relative to changes in light, nitrogen growth conditions and nitrate assimilation. *Journal of Phycology* **48**, 1028-1039.
- Peterson GL.** 1977. A simplification of the protein assay method of Lowry *et al.* which is more generally applicable. *Analytical Biochemistry* **83**, 346-356.
- Rae BD, Long BM, Badger MR, Price GD.** 2013. Functions, compositions, and evolution of the two types of carboxysomes: Polyhedral microcompartments that facilitate CO₂ fixation in cyanobacteria and some proteobacteria. *Microbiology and Molecular Biology Reviews* **77**, 357-379.
- Rose TM, Henikoff JG, Henikoff S.** 2003. CODEHOP (COnsensus-DEgenerate hybrid oligonucleotide primer) PCR primer design. *Nucleic Acids Research* **31**, 3763-3766.
- Ruan Z, Giordano M.** 2017. The use of NH₄⁺ rather than NO₃⁻ affects cell stoichiometry, C allocation, photosynthesis and growth in the cyanobacterium *Synechococcus* sp. UTEX LB 2380, only when energy is limiting. *Plant, Cell and Environment* **40**, 227-236.
- Ruan Z, Prášil O, Giordano M.** 2018. The phycobilisomes of *Synechococcus* sp. are constructed to minimize nitrogen use in nitrogen-limited cells and to maximize energy capture in energy-limited cells. *Environmental and Experimental Botany* **150**, 152-160.
- Ruan Z, Raven JA, Giordano M.** 2017. In *Synechococcus* sp. competition for energy between assimilation and acquisition of C and those of N only occurs when growth is light limited. *Journal of Experimental Botany* **68**, 3829-3839.
- Sánchez-Baracaldo P, Ridgwell A, Raven JA.** 2014. A neoproterozoic transition in the marine nitrogen cycle. *Current Biology* **24**, 652-657.
- Sakamoto T, Bryant DA.** 1999. Nitrate transport and not photoinhibition limits growth of the freshwater cyanobacterium *Synechococcus* species PCC 6301 at low temperature. *Plant Physiology* **119**, 785-794.

- Sakamoto T, Inoue-Sakamoto K, Bryant DA.** 1999. A novel nitrate/nitrite permease in the marine cyanobacterium *Synechococcus* sp. strain PCC 7002. *Journal of Bacteriology* **181**, 7363-7372.
- Sakamoto T, Inoue - Sakamoto K, Persson S, Bryant DA.** 2008. Transcription factor NtcB specifically controls the nitrate assimilation genes in the marine cyanobacterium *Synechococcus* sp. strain PCC 7002. *Phycological research* **56**, 223-237.
- Scanlan DJ, Ostrowski M, Mazard S, Dufresne A, Garczarek L, Hess WR, Post AF, Hagemann M, Paulsen I, Partensky F.** 2009. Ecological genomics of marine picocyanobacteria. *Microbiology and Molecular Biology Reviews* **73**, 249-299.
- Scherer S, Häfele U, Krüger GH, Böger P.** 1988. Respiration, cyanide - insensitive oxygen uptake and oxidative phosphorylation in cyanobacteria. *Physiologia Plantarum* **72**, 379-384.
- Sekowska A, Ashida H, Danchin A.** 2019. Revisiting the methionine salvage pathway and its paralogues. *Microbial Biotechnology* **12**, 77-97.
- Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Söding J.** 2011. Fast, scalable generation of high - quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems Biology* **7**, 539.
- Stamatakis A.** 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312-1313.
- Tabita FR, Satagopan S, Hanson TE, Kreel NE, Scott SS.** 2008. Distinct form I, II, III, and IV Rubisco proteins from the three kingdoms of life provide clues about Rubisco evolution and structure/function relationships. *Journal of Experimental Botany* **59**, 1515-1524.
- Tanigawa R, Shirokane M, Maeda S-i, Omata T, Tanaka K, Takahashi H.** 2002. Transcriptional activation of NtcA-dependent promoters of *Synechococcus* sp. PCC 7942 by 2-oxoglutarate *in vitro*. *Proceedings of the National Academy of Sciences* **99**, 4251-4255.
- Wan N, DeLorenzo DM, He L, You L, Immethun CM, Wang G, Baidoo EE, Hollinshead W, Keasling JD, Moon TS.** 2017. Cyanobacterial carbon metabolism: Fluxome plasticity and oxygen dependence. *Biotechnology and Bioengineering* **114**, 1593-1602.

- Wang TH, Chen YH, Huang JY, Liu KC, Ke SC, Chu HA.** 2011. Enzyme kinetics, inhibitors, mutagenesis and electron paramagnetic resonance analysis of dual-affinity nitrate reductase in unicellular N₂-fixing cyanobacterium *Cyanothece* sp. PCC 8801. *Plant Physiology and Biochemistry* **49**, 1369-1376.
- Wang TH, Fu H, Shieh YJ.** 2003. Monomeric NarB is a dual-affinity nitrate reductase, and its activity is regulated differently from that of nitrate uptake in the unicellular diazotrophic cyanobacterium *Synechococcus* sp. strain RF-1. *Journal of Bacteriology* **185**, 5838-5846.
- Watzer B, Spät P, Neumann N, Koch M, Sobotka R, Macek B, Hennrich O, Forchhammer K.** 2019. The signal transduction protein Pii controls ammonium, nitrate and urea uptake in cyanobacteria. *Frontiers in Microbiology* **10**, 1428.
- Wyman M, Bird C.** 2007. Lack of control of nitrite assimilation by ammonium in an oceanic picocyanobacterium, *Synechococcus* sp. strain WH 8103. *Applied and Environmental Microbiology* **73**, 3028-3033.
- Zhang S, Bryant DA.** 2011. The tricarboxylic acid cycle in cyanobacteria. *Science* **334**, 1551-1553.
- Zhang S, Qian X, Chang S, Dismukes GC, Bryant DA.** 2016. Natural and synthetic variants of the tricarboxylic acid cycle in cyanobacteria: introduction of the GABA shunt into *Synechococcus* sp. PCC 7002. *Frontiers in microbiology* **7**, 1972.
- Zhao M-X, Jiang Y-L, He Y-X, Chen Y-F, Teng Y-B, Chen Y, Zhang C-C, Zhou C-Z.** 2010. Structural basis for the allosteric control of the global transcription factor NtcA by the nitrogen starvation signal 2-oxoglutarate. *Proceedings of the National Academy of Sciences* **107**, 12487-12492.

Figure legends

Fig. 1 Phylogenetic tree based on the sequences of Rubisco large subunit.

Fig. 2 Putative promoters of nitrate assimilation genes and their regulatory genes.

Fig. 3 Nitrate reductase (NR) kinetics for *Synechococcus* UTEX 2380 cells cultured under either N (\square) or energy (\circ) limitation. The NR activity is shown as a function of NO_3^- concentration on a per cell (A, C) and per mg of protein (B, D) basis. Panels C and D are enlarged view of the first phase of the curves in A and B. The error bars indicate the standard deviations ($n=3$). **Fig. 4** V_{\max} of *Synechococcus* sp. UTEX 2380 nitrate reductase (NR; panels A, B for kinetic phase I; C, D for kinetic phase II), nitrite reductase (NiR; panels E, F), and glutamine synthetase (GS; panels G, H). The data are expressed on a per cell basis in panels A, C, E, G, and on a protein basis in panels B, D, F, H. Different letters in the superscripts indicate significantly different means across all treatments (i.e. between different N sources and between N- and energy-limited cells)

Fig. 5 Glutamine synthetase (GS) activity of NO_3^- (A, C) and NH_4^+ -grown (B, D) *Synechococcus* sp. UTEX 2380 cells as a function of sodium glutamate concentration. The activities were normalized by the number of cell (A, C) or by the amount of protein (B, D). The error bars indicate the standard deviations ($n=3$).

Fig. 6 Expression of the nitrate reductase gene (*narB*) of *Synechococcus* sp. UTEX 2380 cultured either under N or energy limitation. Values are reported relative to the *narB* expression in energy limited cells grown in NO_3^- at 550 μM . The error bars show the standard deviation ($n=3$). Different letters in the superscripts indicate significantly different means across all treatments (i.e. between different N sources and between N- and energy-limited cells) ($p<0.05$).

Fig. 7 Nitrate assimilation in *Synechococcus* UTEX 2830 is regulated indirectly via 2-oxoglutarate (2OG), the carbon skeleton for glutamate synthesis, depending on intracellular NH_4^+ . In the presence of high concentration of NH_4^+ (cell growth is limited by energy), 2OG is consumed, leading to the segregation of PipX-NtcA-2OG complex and the consequent inactivation of NtcA. This inactivation downregulates or

even complete blocks nitrate assimilation related (NAR) genes expression and nitrate utilization. On the contrary, when the growth is limited by N (irrespective of N source, energy is sufficient), intracellular 2OG accumulates and tend to bind to PII, inducing the release of PipX from PII-PipX complex. PipX together with OG form PipX-NtcA-2OG complex and activated NtcA, the active NtcA promotes NAR genes expression (*nrtP*, *narB*, *nirA* also require NtcB) and nitrate assimilation. The internal NH_4^+ is between these two extremes (high NH_4^+ and N limitation) with sufficient NO_3^- present in the culture media. NrtP, nitrate/nitrite permease; *nrtP*, NrtP gene; NR; nitrate reductase; *narB*, NR gene; NiR, nitrite reductase; *nirA*, NiR gene; GSI, type I glutamine synthetase; *glnA*, GSI gene; NtcA; global nitrogen regulator; *ntcA*, NtcA gene; NtcB, activator of NAR genes; *ntcB*, NtcB gene; PII, nitrogen regulation protein; *glnB*, PII gene; PipX, interacting protein; *pipX*, PipX gene; NrtS1 and NrtS2 (NrtS1/S2), nitrate/nitrite bispecific transporter; *nrtS1* and *nrtS2*, NrtS1 and *nrtS2* gene; GOGAT, glutamate synthase; 2OG, 2-oxoglutarate; Glu, glutamate; Gln, glutamine.

Accepted Manuscript




Fig. 1 Phylogenetic tree based on the sequences of Rubisco large subunit.

ntcA-binding motif GTA-N8-TAC_(-10 box)N22/23-TA-N3-T

ntcB-binding motif ATGC-N7-GCAT-N7-ATGC-N7-GCAT

Exact match **BOLD**

nrtP 5'-GG**ATGC**AGGATTAAAGCCCTTGT**ATCCCAAAGCGCAT**CGATCCCCATAAAG**AGTACAGCGGTAC**SAATTAGCGAAGAAGAAT**GTGATT**CTTTATCACAA [-233bp] -3'

nirA 5'-GA**ATGCGCGCATCGCAT**ATCAAT**ATTCTGATTTCGCA**AAAGCCAGCCAGGCC**GTAGTTAACACTAC**AAAATCCCTGACTGAATG**TACCTT**GGTCAAATC [-60bp] -3'

ntcA 5'-TATGCGATCGTGTGAATTCCTAAAGGTAGGTTAGAGCAGACATGGATCGCC**GTAATAAGGATT**GTGAGTAGATAAAAACGCCCTGG**TAACTT**CATAAAGCC [-134bp] -3'

glnA 5'-TAAATCTAACCTACCTGACAGTAAACAGTTCCTGGGCCGCAACCAAGTGGAGAT**GTGACCCAGACTAC**AACGAAGTGCCTAACTCTT**TAGGAT**EAAAGCGAAC [-75bp] -3'

glnB 5'-CGGGCGCTGGGTCAGTCGGCTCCCCACTGGATCGATTGACGA TTGAGAA**GTTAAAGAGTTTAC**TTGGTAGGGGCTGGAGCAG**CATCGT**ATTTTTGC [-611bp] -3'

pipX 5'-GATCACGCTATCCAGGGGAGCCTTGCCTATATACTATGTCTAATATTTTGTATG**GTA**TTTTG**ATTAC**AGCAAAGCTAACCTCTGATTTT**TGTGGT**CAGAAACCGA [-87bp] -3'

ntcB 5'-TT**GTAACATCCGGAA**CAAAAAGCCCCAAAATCA**ATGCATTA**ACT**AT**TGGGCAC**ATG**CACAGGAA**AAACAT**TTAAACCAAGGGGATGACCC [-0bp] -3'

Fig. 2 Putative promoters of nitrate assimilation genes and their regulatory genes. *nrtP*, nitrate/nitrite permease gene; *nirA*, nitrite reductase gene; *ntcA*, global nitrogen regulator gene; *glnA*, type I glutamine synthetase gene; *glnB*, nitrogen regulation protein PII gene; *ntcB*, activator gene of nitrate assimilation related genes; *pipX*, interacting protein PipX gene.

Accepted Manuscript




Fig. 3 Nitrate reductase (NR) kinetics for *Synechococcus* UTEX 2380 cells cultured under either N (\square) or energy (\circ) limitation. The NR activity is shown as a function of NO_3^- concentration on a per cell (A, C) and per mg of protein (B, D) basis. Panels C and D are enlarged view of the first phase of the curves in A and B. The error bars indicate the standard deviations ($n=3$).




Fig. 4 V_{max} of *Synechococcus* sp. UTEX 2380 nitrate reductase (NR; panels A, B for kinetic phase I; C, D for kinetic phase II), nitrite reductase (NiR; panels E, F), and glutamine synthetase (GS; panels G, H). The data are expressed on a per cell basis in panels A, C, E, G, and on a protein basis in panels B, D, F, H. Different letters in the superscripts indicate significantly different means across all treatments (i.e. between different N sources and between N- and energy-limited cells)




Fig. 5 Glutamine synthetase (GS) activity of NO₃⁻ (A, C) and NH₄⁺-grown (B, D) *Synechococcus* sp. UTEX 2380 cells as a function of sodium glutamate concentration. The activities were normalized by the number of cell (A, C) or by the amount of protein (B, D). The error bars indicate the standard deviations (n=3).

Accepted Manuscript




Fig. 6 Expression of the nitrate reductase gene (*narB*) of *Synechococcus* sp. UTEX 2380 cultured either under N or energy limitation. Values are reported relative to the *narB* expression in energy limited cells grown in NO₃⁻ at 550 μM. The error bars show the standard deviation (n=3). Different letters in the superscripts indicate significantly different means across all treatments (i.e. between different N sources and between N- and energy-limited cells) (p<0.05).

Accepted Article




Fig. 7 Nitrate assimilation in *Synechococcus* UTEX 2830 is regulated indirectly via 2-oxoglutarate (2OG), the carbon skeleton for glutamate synthesis, depending on intracellular NH_4^+ . In the presence of high concentration of NH_4^+ (cell growth is limited by energy), 2OG is consumed, leading to the segregation of PipX-NtcA-2OG complex and the consequent inactivation of NtcA. This inactivation downregulates or even complete blocks nitrate assimilation related (NAR) genes expression and nitrate utilization. On the contrary, when the growth is limited by N (irrespective of N source, energy is sufficient), intracellular 2OG accumulates and tends to bind to PII, inducing the release of PipX from PII-PipX complex. PipX together with OG form PipX-NtcA-2OG complex and activate NtcA, the active NtcA promotes NAR genes expression (*nrtP*, *narB*, *nirA* also require NtcB) and nitrate assimilation. The internal NH_4^+ is between these two extremes (high NH_4^+ and N limitation) with sufficient NO_3^- present in the culture media. NrtP, nitrate/nitrite permease; *nrtP*, NrtP gene; NR; nitrate reductase; *narB*, NR gene; NiR, nitrite reductase; *nirA*, NiR gene; GSI, type I glutamine synthetase; *glnA*, GSI gene; NtcA; global nitrogen regulator; *ntcA*, NtcA gene; NtcB, activator of NAR genes; *ntcB*, NtcB gene; PII, nitrogen regulation protein; *glnB*, PII gene; PipX, interacting protein; *pipX*, PipX gene; NrtS1 and NrtS2 (NrtS1/S2), nitrate/nitrite bispecific transporter; *nrtS1* and *nrtS2*, NrtS1 and nrtS2 gene; GOGAT, glutamate synthase; 2OG, 2-oxoglutarate; Glu, glutamate; Gln, glutamine.