

University of Dundee

Phosphoenolpyruvate carboxykinase, pyruvate orthophosphate dikinase and isocitrate lyase in both tomato fruits and leaves, and in the flesh of peach and some other fruits

Famiani, Franco; Paoletti, Andrea; Battistelli, Alberto; Moscatello, Stefano; Chen, Zhi Hui; Leegood, Richard C.

Published in:
Journal of Plant Physiology

DOI:
[10.1016/j.jplph.2016.07.003](https://doi.org/10.1016/j.jplph.2016.07.003)

Publication date:
2016

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Famiani, F., Paoletti, A., Battistelli, A., Moscatello, S., Chen, Z. H., Leegood, R. C., & Walker, R. P. (2016). Phosphoenolpyruvate carboxykinase, pyruvate orthophosphate dikinase and isocitrate lyase in both tomato fruits and leaves, and in the flesh of peach and some other fruits. *Journal of Plant Physiology*, 202, 34-44. <https://doi.org/10.1016/j.jplph.2016.07.003>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Manuscript Number: JPLPH-D-16-00294

Title: Phosphoenolpyruvate carboxykinase, pyruvate orthophosphate dikinase and isocitrate lyase in both tomato fruits and leaves, and in the flesh of peach and some other fruits

Article Type: Research Paper

Section/Category: Biochemistry

Keywords: gluconeogenesis, isocitrate lyase, malate, pyruvate, pyruvate orthophosphate dikinase, senescence

Corresponding Author: Prof. Franco Famiani,

Corresponding Author's Institution: Università degli Studi di Perugia

First Author: Franco Famiani

Order of Authors: Franco Famiani; Andrea Paoletti; Alberto Battistelli; Stefano Moscatello; Richard C Leegood; Robert P Walker

Abstract: In this study the occurrence of a number of enzymes involved in gluconeogenesis was investigated in both tomato fruits and leaves during their development and senescence and in some other fruits. The enzymes studied were phosphoenolpyruvate carboxykinase (PEPCK), pyruvate orthophosphate dikinase (PPDK) and glyoxysomal isocitrate lyase (ICL). PPDK was detected in the ripe flesh of tomato, and much smaller amounts were detected in the flesh of both peach and pepper, whereas it was not detected (not present or at very low abundance) in the other fruits which were investigated (apricot, aubergine, blackberry, blueberry, cherry, grape, plum, raspberry, red current and tomato). By contrast PEPCK was present in the flesh of all the fruits investigated. Very small amounts of ICL were detected in ripe tomato flesh. PEPCK was present in the skin, flesh, locular gel and columella of tomato fruit, and in these its abundance increased greatly during ripening. PPDK showed a similar distribution, however, its abundance did not increase during ripening. PEPCK was not detected in tomato leaves at any stage of their development or senescence. The content of PPDK g⁻¹ fresh weight (FW) increased in tomato leaves as they matured, however, it declined during their senescence. In tomato leaves the content of ICL g⁻¹ FW increased until the mid-stage of development, then decreased as the leaf matured, and then increased during the latter stages of senescence. In the flesh of tomato fruits the contents of PPDK and PEPCK g⁻¹ FW decreased during senescence.

The results suggest that in fruits other than tomato the bulk of any gluconeogenic flux proceeds via PEPCK, whereas in tomato both PEPCK and PPDK could potentially be utilised. Further, the results indicate that the conversion of pyruvate/acetyl-CoA to malate by the glyoxylate cycle, for which ICL is necessary, is not a major pathway utilised by gluconeogenesis in fruits under normal conditions of growth. Finally, the results contribute to our understanding of the role of several enzymes in the senescence of both leaves and fruits.

1 **Phosphoenolpyruvate carboxykinase, pyruvate orthophosphate dikinase and**
2 **isocitrate lyase in both tomato fruits and leaves, and in the flesh of peach and some**
3 **other fruits**

4
5
6 **Franco Famiani^{1,*}, Andrea Paoletti¹, Alberto Battistelli², Stefano Moscatello², Richard C. Leegood³ and Robert**
7 **P. Walker^{1,*}**

8
9 ¹ Dipartimento di Scienze Agrarie, Alimentari e Ambientali, Università degli Studi di Perugia, Borgo XX Giugno, 74,
10 06121, Perugia, Italy

11 ² Istituto di Biologia Agroambientale e Forestale, CNR, Viale Marconi, 2, 05010, Porano (TR), Italy

12 ³ Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2 TN, UK

13
14 * To whom correspondence should be addressed: E-mail: rob.walker@talktalk.net; franco.famiani@unipg.it

15
16 Abbreviations: isocitrate lyase, ICL; malate synthase, MS; malate dehydrogenase, MDH; malic enzyme, ME;
17 oxaloacetate, OAA; phosphoenolpyruvate, PEP; phosphoenolpyruvate carboxylase, PEPC; phosphoenolpyruvate
18 carboxykinase, PEPCK; pyruvate kinase, PK; pyruvate orthophosphate dikinase, PPDK.

19
20 **Abstract**

21 **In this study the occurrence of a number of enzymes involved in gluconeogenesis was investigated in both**
22 **tomato fruits and leaves during their development and senescence and in some other fruits. The enzymes**
23 **studied were phosphoenolpyruvate carboxykinase (PEPCK), pyruvate orthophosphate dikinase (PPDK) and**
24 **glyoxysomal isocitrate lyase (ICL). PPDK was detected in the ripe flesh of tomato, and much smaller amounts**

25 were detected in the flesh of both peach and pepper, whereas it was not detected (not present or at very low
26 abundance) in the other fruits which were investigated (apricot, aubergine, blackberry, blueberry, cherry,
27 grape, plum, raspberry, red current and tomato). By contrast PEPCK was present in the flesh of all the fruits
28 investigated. Very small amounts of ICL were detected in ripe tomato flesh. PEPCK was present in the skin,
29 flesh, locular gel and columella of tomato fruit, and in these its abundance increased greatly during ripening.
30 PPDK showed a similar distribution, however, its abundance did not increase during ripening. PEPCK was
31 not detected in tomato leaves at any stage of their development or senescence. The content of PPDK g^{-1} fresh
32 weight (FW) increased in tomato leaves as they matured, however, it declined during their senescence. In
33 tomato leaves the content of ICL g^{-1} FW increased until the mid-stage of development, then decreased as the
34 leaf matured, and then increased during the latter stages of senescence. In the flesh of tomato fruits the
35 contents of PPDK and PEPCK g^{-1} FW decreased during senescence.

36 The results suggest that in fruits other than tomato the bulk of any gluconeogenic flux proceeds via PEPCK,
37 whereas in tomato both PEPCK and PPDK could potentially be utilised. Further, the results indicate that the
38 conversion of pyruvate/acetyl-CoA to malate by the glyoxylate cycle, for which ICL is necessary, is not a major
39 pathway utilised by gluconeogenesis in fruits under normal conditions of growth. Finally, the results
40 contribute to our understanding of the role of several enzymes in the senescence of both leaves and fruits.

41

42 Key words: fruits, gluconeogenesis, isocitrate lyase, leaves, malate, peach, phosphoenolpyruvate carboxykinase,
43 pyruvate, pyruvate orthophosphate dikinase, senescence

44

45

46 **1. Introduction**

47 Gluconeogenesis from malate occurs in the flesh of ripening grape, tomato and cherry fruits (Farineau and Laval-
48 Martin, 1977; Halinska and Frenkel, 1991; Huang et al., 2015a, 2015b; Leegood and Walker, 1999; Osorio et al.,
49 2013; Ruffner, 1982). In plants gluconeogenesis from malate can occur by two alternate pathways. One pathway

50 utilises malate dehydrogenase (MDH) in conjunction with phosphoenolpyruvate carboxykinase (PEPCK). The other
51 pathway utilises malic enzyme (ME) in conjunction with pyruvate orthophosphate dikinase (PPDK) (Famiani et al.,
52 2015, 2014b; Leegood and Walker; 2003). In the fruits of both cherry and grape it appears that the PEPCK pathway
53 is used in gluconeogenesis, and this is because PPDK is not present (or is at very low abundance) (Famiani et al.,
54 2014b; Walker et al., 2011a). PEPCK is also present in ripening tomato flesh in which it catalyses a gluconeogenic
55 flux from malate/citrate (Bahrami et al., 2001; Huang et al., 2015a, 2015b; Osorio et al., 2013). In tomato,
56 radiolabelling experiments have shown that gluconeogenesis from pyruvate occurs, and this requires either PPDK or
57 PEPCK (Farineau and Laval-Martin, 1977). If PEPCK is utilised, pyruvate can be converted to malate by either the
58 Krebs cycle or the glyoxylate cycle. Isocitrate lyase (ICL) is an essential component of the glyoxylate cycle
59 (Eastmond and Graham, 2001). In both cucumber flesh and ripening banana flesh very low amounts of ICL are
60 present (Liu et al., 2004; Yang et al., 1998). However, glyoxysomal ICL was not detected in either grape flesh or that
61 of some soft fruits (Famiani et al., 2014b, 2005). In tomato fruits the occurrence of both PPDK and the glyoxylate
62 cycle is uncertain. The first aim of the present study was to investigate the occurrence of PEPCK, PPDK and
63 glyoxysomal ICL in tomato and other fruits. There is contradictory information regarding the occurrence of PPDK in
64 senescing leaves (Chen et al., 2000; Taylor et al., 2010), and very little is known about the occurrence of either
65 PEPCK or PPDK in most fruits during their senescence. The second aim of this study was to determine whether the
66 abundance of PEPCK and PPDK increases in tomato fruits and leaves during their senescence.

67

68 **2. Materials and methods**

69

70 *2.1. Plant material*

71 Tomato plants (*Solanum lycopersicum*) were grown in a greenhouse in Perugia and fruits were collected at four
72 stages of development. These were: 1, small-green (20% final FW); 2, medium-green (50% final FW); 3, breaker
73 (turning colour); 4, full colouration (red). Plants of sweet pepper (*Capsicum annuum*, cv. World Beater), aubergine
74 (*Solanum melongena*, cv. Black Enorma), *Hoya carnososa* and maize were grown in pots of garden soil in a greenhouse

75 in Perugia. In 2004, fruits of both peach (*Prunus persica* L. Batsch) cultivar Adriatica and other fruits were collected
76 from mature trees growing in the experimental orchard of the Department of Agricultural, Food and Environmental
77 Sciences of the University of Perugia, in Deruta (PG), central Italy. The fruits of all species were harvested from
78 several positions on the plant. Maize seedlings (*Zea mays*) were germinated and grown in trays of perlite in a
79 greenhouse in Perugia, Italy, and they were then fed NH₄Cl as previously described (Walker et al., 2001). Cucumber
80 cotyledons (*Cucumis sativus*) were obtained from seeds germinated in perlite at 25°C under darkness for 4 d. For both
81 tomato leaf and fruit senescence experiments mature leaves or ripe fruit (development stage 5) were detached and
82 placed on moist filter paper in petri dishes. These were then placed in an incubator under darkness at 25°C.

83

84 2.2. Preparation of a nitrogen powder

85 To ensure that the sample was representative of the tissue a nitrogen powder was prepared. Tissues were frozen in
86 liquid nitrogen. When required for analysis, the tissue was removed from the liquid nitrogen. For fruits three samples
87 of the components of the pericarp (skin, flesh, locular gel and placenta), each composed of subsamples of 5-10 fruits,
88 were used. Tissues were then ground together in a mortar containing liquid nitrogen and the resulting powder was
89 used either immediately or after storage at -80°C. This powder was used for electrophoretic analysis.

90

91 2.3. Enzyme assays

92 Two hundred mg of frozen powder of fruit flesh was ground in a mortar containing 800 µL of ice cold 200 mM
93 Bicine-KOH (pH 9.0), 50 mM DTT and clarified by centrifugation at 12 000 g for 5 min. PEPCK activity in the
94 supernatant was measured, using an enzyme coupled method, in the carboxylation direction as described by Walker
95 et al. (1999) and Famiani et al. (2005). For the measurement of PPDK activity plant material was extracted and
96 enzyme activity determined in the forward direction (PEP formation) using an enzyme-coupled spectrophotometric
97 method as described previously (Ashton et al., 1990). For both PEPCK and PPDK one unit of activity is that which
98 produces 1 µmol product min⁻¹ at 25 °C.

99

100 2.4. SDS-PAGE and immunoblotting

101 For pericarp 500 mg of frozen powder was added to 500 µl electrophoresis buffer (62.5 mM Tris-HCl (pH 6.8), 10%
102 (v/v) glycerol, 5% (w/v) SDS, 50 mM ascorbate, 5% (v/v) 2-mercaptoethanol and 0.002% (w/v) bromophenol blue)
103 contained in a mortar, and ground with a pestle. Other tissues were extracted in the same way except that for leaves
104 100 mg, for roots 250 mg and for either developing or germinating seeds 50 mg of frozen powder was used. If the
105 extract became yellow, several microlitres of 20% (w/v) NaOH were added until it just became blue. The suspension
106 was immediately poured into an Eppendorf tube, which was then incubated at 100 °C for 5 min and then centrifuged
107 at 12 000 g for 5 min. The supernatant was separated from the pellet and stored at –20 °C until required. After
108 centrifugation at 10 000 g for 5 min, 4 µl of extract was loaded onto each track of SDS-PAGE gels. SDS-PAGE,
109 staining of gels with Coomassie Brilliant Blue dye and immunoblotting were done as described in Walker and
110 Leegood (1996). Protein measurement was done using the Lowry method as previously described (Walker et al.,
111 1995). Briefly SDS-PAGE was performed in a Hoefer mini-gel apparatus (SE 250 Mighty Small II; Hoefer Scientific
112 Instruments, San Francisco, USA) and western transfer done using a Pharmacia Multiphor device (Multiphor II
113 Electrophoresis System; Pharmacia Biotech, Uppsala, Sweden) in conjunction with Millipore Immobilon-P
114 membrane (Millipore, Billerica, Massachusetts, USA). Anti-rabbit peroxidase (diluted 1/1000) was used in conjunction
115 with an ECL kit (GE Healthcare, Little Chalfont, UK) to visualize immunoreactive polypeptides. The antisera to
116 aldolase, PEPC, phosphoenolpyruvate carboxylase (PEPC) and ribulose-1,5-bisphosphate carboxylase/oxygenase
117 (Rubisco) were produced using as antigens the enzymes that had been purified from *Panicum maximum* leaves
118 (Walker et al., 2002; R.P. Walker, unpublished work). The antisera to PPDK was that used in a previous study of
119 grape (Chastain et al., 2002; Famiani et al., 2014b). ICL antiserum was raised against the enzyme from castor bean
120 endosperm (Maeshima et al., 1988). The glutamine synthetase (GS); antiserum was raised against the plastidic
121 enzyme from *Sinapsis alba* (Höpfner et al., 1990).

122

123 3. Results and discussion

124

125 3.1. The occurrence of PPDK and PEPCK in the ripe flesh of both tomato and some other fruits

126 Many studies have established the specificity of both the PEPCK and PPDK antibodies used in the present study for
127 the enzymes from a wide range of plant species (Chastain et al., 2006, 2002; Famiani et al., 2014b, 2005; Taylor et
128 al., 2010; Walker et al., 2011a, 2011b, 2011c). PPDK was detected in both tomato flesh and leaves, and was of the
129 same molecular mass as PPDK from both maize and *Hoya carnososa* leaves (Fig. 1A). PEPCK was detected in tomato
130 flesh, *H. carnososa* leaves, maize leaves but not tomato leaves (Fig. 1A). The occurrence of PEPCK and PPDK in both
131 maize and *H. carnososa* leaves is consistent with previous studies (Black et al., 1996; Leegood and Walker, 1996;
132 Sugiyama, 1973; Walker et al., 1997). The specific activity ([enzyme activity $\mu\text{mol product min}^{-1}$]/[mg total protein])
133 of PPDK in tomato leaves is 0.0006 (Hocking and Anderson, 1986) and in maize leaves 0.1-0.19 (Aoyagi and
134 Bassham, 1983; Hocking and Anderson, 1986; Sugiyama, 1973), and these values give an indication of the
135 abundance of PPDK in tomato tissues.

136 The occurrence of PPDK in the flesh of a number of other fruits was investigated (Fig. 2). All these fruits are known
137 to contain PEPCK (Baldicchi et al., 2015; Famiani et al., 2012, 2005; Walker and Chen, 2002; Walker et al., 2015,
138 2011a). Under the conditions used for the immunoblot (antibody dilution 1/10 000) PPDK was only detected in
139 tomato (Fig. 2). Similar amounts of PPDK polypeptide were also detected in the ripe flesh of a wide range of tomato
140 cultivars (data not shown). PPDK is also present in the peel of cactus pear fruits, however, in these PPDK functions
141 in Crassulacean acid metabolism (CAM) associated with photosynthesis (Walker et al., 2011c). In keeping with
142 previous work, PPDK polypeptide was not detected in extracts of the ripe flesh of several soft fruits, cherry, plum or
143 grape (Fig. 2; Famiani and Walker, 2009; Famiani et al., 2014b, 2012, 2005; Walker et al., 2011a). To ensure that
144 PPDK polypeptide was not lost after extraction, the flesh of each fruit shown in Fig. 2 was individually co-extracted
145 with one twentieth the FW of maize leaf and then subjected to SDS-PAGE and immunoblotting. In extracts of all
146 fruits co-extracted with maize leaf PPDK was detected, and this shows that PPDK was unlikely to be lost from the
147 fruit tissues (data not shown).

148 In addition to tomatoes, other solanaceous fruits of commercial importance are aubergines and peppers, and the
149 abundance of both PEPCK and PPDK was determined in their ripening flesh (Fig. 1B). PEPCK was not detected in

150 sweet pepper, however, it was present in aubergine in which its abundance was very low (Fig. 1B; a barely visible
151 band of the same molecular mass as PEPCK was present in the extract of the flesh of red pepper, this band was so
152 faint that it is not visible in the figure). Previously, very low amounts of PEPCK activity were measured in extracts of
153 aubergine fruits (0.002-0.005 U g⁻¹ FW) (Blanke et al., 1988), and this is about 10-fold less than in tomato flesh
154 (Bahrami et al., 2001; Huang et al., 2015a, 2015b; Osorio et al., 2013). PPDK was not detected in aubergine flesh,
155 however, a very long exposure of the immunoblot showed that very low amounts of PPDK were present in the fruit
156 wall of pepper at both stages of development studied (Fig. 1B). However, the band was so faint that it was not
157 possible to reliably determine whether its abundance was different at the two stages of development studied.

158 We attempted to measure PPDK activity in extracts of the fruits. In peach flesh although PPDK polypeptide has been
159 detected by immunoblotting its activity was not determined (Borsani et al., 2009; Lara et al., 2010, 2009). We failed
160 to measure PPDK activity in the flesh of any of these fruits including tomato (the tissue was harvested and extracted
161 around noon). Previously we failed to measure PPDK in extracts of tomato flesh, and based on this it was stated that
162 there was no evidence for the presence of PPDK in this tissue (Bahrami et al., 2001). To investigate this further we
163 co-extracted ripe fruit flesh with one twentieth its FW of maize leaf (also harvested around noon). This was done for
164 each fruit, and for all co-extractions we could measure a similar amount of PPDK activity as that which was present
165 in the maize leaf extracted alone. Hence it was unlikely that PPDK activity was lost after extraction of the fruit
166 tissues. An explanation for the failure to measure PPDK activity is that in many plant tissues a large proportion of the
167 enzyme is often present as a phosphorylated inactive form (Chastain et al., 2006). For example, in the leaves of the
168 CAM plant *Kalanchoë fedtschenkoi* PPDK is only active at certain times during the day-night cycle, and for most of
169 the day-night cycle although PPDK protein can be detected its activity cannot be measured (Dever et al., 2015).

170 Therefore, in tomato flesh it is possible that PPDK is often present in an inactive phosphorylated form, and is only
171 active at certain times or under certain conditions.

172 To have an approximate idea of the abundance of the PPDK polypeptide in tomato flesh different amounts of extracts
173 of either maize leaf or tomato flesh were subjected to SDS-PAGE and immunoblotting. Based on this the amount of
174 PPDK polypeptide mg⁻¹ total protein in tomato flesh was about 3-12% of that in maize leaves and on a g⁻¹ FW basis

175 0.2-0.8 % of maize leaves (data not shown). In maize leaves the activity of PPDK is of the order of $1.7 \text{ U g}^{-1} \text{ FW}$
176 (Aoyagi and Bassham, 1984). Thus the amount of PPDK polypeptide in tomato flesh corresponds to a maximum
177 activity of about $0.003\text{-}0.014 \text{ U g}^{-1} \text{ FW}$. In ripening tomato flesh the activity of PEPCK is about $0.04 \text{ U g}^{-1} \text{ FW}$
178 (Bahrami et al., 2001; Huang et al., 2015b; Osorio et al., 2013).

179 In contrast to the present study (Fig. 2), PPDK polypeptide was detected in extracts of ripe peach flesh by
180 immunoblotting, and this was using the same antibody as employed in the present study (Borsani et al., 2009; Lara et
181 al., 2010, 2009). An exposure of a western blot probed with the PPDK antibody at a dilution of 1/10 000 detected a
182 large band of PPDK in maize leaf but nothing in ripe peach flesh (Fig. 3A). By contrast PEPCK was detected in both
183 tissues (Fig. 3A). A western blot was done of extracts of peach flesh (from different stages of its development) and
184 maize leaves (Fig. 3B). An exposure of the western blot, using PPDK antibody at a dilution of 1/10 000, detected a
185 large amount of PPDK in maize leaves but none in peach flesh (Fig. 3B). The blot was reprobed using the PPDK
186 antibody at 1/1000, and a very long exposure showed that PPDK was present throughout the ripening of peach flesh
187 (Fig 5). This shows that in peach flesh PPDK is at low abundance and must be at the very most 10-times less
188 abundant than in tomato flesh. This would give an activity of $0.0003\text{-}0.0014 \text{ U g}^{-1} \text{ FW}$. By contrast PEPCK is quite
189 abundant in peach flesh throughout ripening (about $0.18 \text{ U g}^{-1} \text{ FW}$; Famiani et al., 2016a). Similar considerations
190 apply to the flesh of cherry, plum and apricot. Thus PEPCK activity in the flesh of these stone fruits is likely over
191 100-times higher than that of PPDK. Previous studies of both grape and some soft fruits did not detect PPDK in their
192 flesh, and it was concluded that the enzyme was either not present or at low abundance (Famiani et al., 2014b, 2005).
193 The results of the present study suggest that if PPDK is present in the flesh of these fruits, then at most its abundance
194 is similar to that in peach flesh. Thus in the flesh of these PEPCK activity should be at least 35-times higher than that
195 of PPDK. Therefore, as previously suggested in the flesh of all the aforementioned fruits, apart from tomato, the bulk
196 of the gluconeogenic flux should utilise PEPCK and not PPDK. Of course this does not rule out the possibility that in
197 the flesh of these fruits under certain conditions (eg low O_2) the abundance of PPDK increases (Lara et al., 2010).

198

199 *3.2. The occurrence of PEPCK and PPDK in tomato fruits*

200 PPDK was present at a somewhat similar abundance g^{-1} FW (from stage 1 to stage 4 of development) in the flesh,
201 locular gel and columella of tomato fruits, however, its abundance was lower in the skin of green fruits (Figs. 4, 5A).
202 The abundance of PPDK g^{-1} FW was lower in very young flesh (fruit FW 10% of that of ripe fruit) than in the flesh
203 of fruits later in development (Fig. 5B). Except for in the skin of green fruits, in which its abundance was lower, the
204 abundance of PPDK g^{-1} FW was comparable in the different tissues of both green (developmental stage 2) and ripe
205 tomatoes (developmental stage 4) (Fig. 5A). PEPCK was present in comparable amounts g^{-1} FW in the flesh, locular
206 gel and columella (Fig. 5A). In the skin of green fruits the abundance of PEPCK g^{-1} FW was lower than in the other
207 tissues, and in red fruits slightly higher (Fig. 5A). In the flesh of red fruit the abundance of PEPCK was lower than in
208 the other tissues (Fig. 5A). However, this observation was not representative of the situation because in other
209 experiments the abundance of PEPCK in the flesh was found to be similar g^{-1} FW in the different tissues of the ripe
210 fruit (data not shown). In all the tissues of the fruit the abundance of PEPCK g^{-1} FW increased greatly at the onset of
211 ripening (Fig. 4). In each of these tissues the amount of PEPCK was 0.003 U g^{-1} FW (SE = 0.002) (developmental
212 stages 1,2) and 0.04 U g^{-1} FW (SE = 0.005) (developmental stages 3,4). In blueberry, cherry, plum, grape and peach
213 PEPCK is also present in the different tissues of the fruit (Famiani et al., 2016a, 2012, 2005; Walker et al., 2015,
214 2011a, 2011b). A previous study also found that there was a large increase in abundance of PEPCK in tomato flesh
215 during ripening, however, it was not detected before ripening in the flesh, or in the locular gel during ripening
216 (Bahrami et al., 2001). It is possible that these contradictions are a result of either difficulties in detecting PEPCK or
217 differences in growth conditions or variety. The presence of PEPCK before ripening is consistent with the presence of
218 PEPCK mRNA in tomato flesh before ripening (Saito et al., 2008; Yin et al., 2010). In several soft fruits, cherry,
219 plum, grape and peach the abundance of PEPCK is also higher during ripening (Famiani and Walker, 2009; Famiani
220 et al., 2016a, 2012, 2005; Walker et al., 2015, 2011a). Nevertheless, in tomato fruits the presence of PEPCK before
221 ripening suggests that the enzyme does not simply function in the dissimulation of malate/citrate during ripening.
222 Ripe tomato fruits were detached from the plant and incubated under darkness at 25°C for 3-9 d. In the flesh of these
223 fruits the content of both PPDK and PEPCK g^{-1} FW decreased throughout the incubation (Fig. 6).

224

225 *3.3. PPDK and PEPCK abundance in tomato leaves during their development and senescence*

226 PPDK has been detected in the leaves of some C₃ plants but not in those of others (Bailey and Leegood, 2016;
227 Famiani and Walker, 2009; Hocking and Anderson, 1986; Walker et al., 2011a). In extracts of tomato leaves both
228 PPDK activity and polypeptide have been detected (Famiani et al., 2005; Hocking and Anderson, 1986). PPDK has
229 been found to increase in abundance during the senescence of some leaves, but not in others (Chen et al., 2000;
230 Taylor et al., 2010). We therefore investigated whether PPDK polypeptide content increased in tomato leaves during
231 their senescence. PPDK polypeptide abundance and not PPDK activity was determined, because in the darkened
232 leaves of C₃ plants PPDK is present largely as an inactive phosphorylated form (Chastain et al., 2002). The
233 abundance of PPDK polypeptide mg⁻¹ total protein in mature tomato leaves was less than in the fruit (Figs. 7, 8). This
234 comparison was done on a protein basis and not on a FW basis. This was because most of the volume of the cells of
235 the fruit is occupied by a large vacuole and this contains little protein, therefore, the amount of cytoplasm g⁻¹ FW
236 (where PPDK, PEPCK are located) in the fruit is less than in the leaf (Famiani et al., 2012). The abundance of PPDK
237 g⁻¹ FW increased as the leaf matured (Fig. 8). Similarly the abundance of PPDK increases during the development of
238 both wheat and Arabidopsis leaves (Aoyagi and Bassham, 1984; Taylor et al., 2010). The function of PPDK in
239 mature leaves is uncertain, although as proposed for senescing leaves one function could be in amino acid
240 metabolism (Chastain et al., 2011; Taylor et al., 2010). A preliminary study indicated that the abundance of PPDK g⁻¹
241 FW did not increase in senescing leaves of tomato that were attached to the plant (data not shown), and similarly
242 PPDK g⁻¹ FW did not increase in senescing cucumber cotyledons that were attached to the plant (Chen et al., 2000).
243 In tomato leaves the content of PPDK g⁻¹ FW declined greatly in leaves detached and incubated under darkness (Fig.
244 8). Hence in tomato leaves the content of PPDK g⁻¹ FW was highest in mature non-senescing leaves. Similarly, when
245 either barley leaves or mature cucumber cotyledons are detached and incubated under darkness, PPDK polypeptide g⁻¹
246 FW declines (Chen et al., 2000). By contrast there is an increase in PPDK abundance mg⁻¹ total protein during the
247 senescence of Arabidopsis leaves attached to the plant (Taylor et al., 2010). It is possible that this contradiction is a
248 result of differences between species. An alternative explanation is that in the present study each track on the gel was
249 loaded with the protein content of an equal amount FW of tissue, whereas, in the study of Arabidopsis gels were

250 loaded with an equal amount of protein (Taylor et al., 2010), and during senescence there is often a large decrease in
251 the total protein content of the tissue (Chen et al., 2000; Fig. 8). Several studies have arrived at the conclusion that
252 PPDK RNA abundance increases during senescence, however, there are potentially difficulties in interpreting these
253 experiments. This is because the decrease in total RNA content of the tissue during senescence was often not
254 considered. It is necessary to show that the abundance of PPDK RNA g^{-1} FW increases: because, as shown by Chen
255 et al. (2000), studies of PEPCK RNA content in senescing tissues which did not do this led to the false conclusion
256 that PEPCK was associated with senescence. Therefore, further experiments to determine whether PPDK protein g^{-1}
257 FW increases during the senescence of leaves from both different species and different tissues would be informative.
258 Nevertheless, these considerations do not affect the proposal that PPDK functions in amino acid metabolism in some
259 senescing leaves (Taylor et al., 2010), and this is because irrespective of whether PPDK g^{-1} FW increases during
260 senescence it is still present.

261 In tomato leaves, detached from the plant and incubated under darkness, as in barley leaves and cucumber cotyledons
262 (Chen et al., 2000), there was a large decrease in the content of total polypeptides and in the abundance of PEPC and
263 Rubisco (Fig. 8). In tomato leaves there was a large decrease in the larger form of GS (Fig. 8). Two forms of GS were
264 detected and these correspond to the plastidic (larger polypeptide) and cytosolic (smaller polypeptide) GS in tomato
265 (Bortolotti et al., 2003; Fig. 8). In maize root a still smaller form of GS was also present, and this is consistent with
266 previous work which showed that this is a cytosolic GS (Sakakibara et al., 1992). There was a large decrease in the
267 smaller form of aldolase during the senescence of tomato leaves (Fig. 8). A larger form of aldolase was present in
268 maize roots and tomato fruits (Fig. 8). The larger aldolase is the cytosolic enzyme and the smaller one the plastidic
269 enzyme (Famiani and Walker, 2009; Famiani et al., 2014b). Thus the abundance of PEPC, Rubisco, plastidic GS and
270 plastidic aldolase changed in a similar way to that of PPDK (Fig. 8). PEPCK was not detected in tomato leaves at any
271 stage of their development or senescence, and this is consistent with studies of PEPCK RNA content (Fig. 8; Bahrami
272 et al., 2001). Previous studies have shown that PEPCK is unlikely to play a role in leaf senescence (Chen et al., 2000;
273 Taylor et al., 2010). As previously reported (Nieri et al., 1997) small amounts of ICL were present in immature
274 tomato leaves, and in the mature leaf its abundance was much less (Fig. 8). In immature tomato leaves the function of

275 ICL is unknown (Nieri et al., 1997). In detached leaves of many plants incubated under darkness, there is an
276 induction of ICL and the glyoxylate cycle, and this could potentially function in the anaplerotic replenishment of the
277 Krebs cycle during starvation (Chen et al., 2000; Eastmond and Graham, 2001; Smith, 2002). In detached tomato
278 leaves there was also an induction of ICL (Fig. 8).

279

280 *3.4. The occurrence of ICL in tomato fruit*

281 In tomato flesh radiolabelling experiments have shown that gluconeogenesis from pyruvate occurs (Farineau and
282 Laval-Martin, 1977). This process can potentially utilise PEPCK in conjunction with the glyoxylate cycle. The
283 enzymes ICL and malate synthase (MS) are essential components of the glyoxylate cycle (Eastmond and Graham,
284 2001). An early radiolabelling study indicated that MS was present in tomato flesh (Doyle et al., 1960). Both MS
285 RNA and ICL activity are present in the flesh of ripening banana fruits (Liu et al., 2004; Pua et al., 2003). Further,
286 ICL activity has been detected in cucumber flesh (Yang et al., 1998), and in this tissue there is a very low expression
287 of the MS gene (Graham et al., 1992). Glyoxysomal ICL polypeptide and activity were not detected in pumpkin flesh
288 at commercial harvest, however, they subsequently appeared in slices of the flesh incubated under darkness (Pistelli
289 et al., 1996). Glyoxysomal ICL polypeptide was either absent or at very low abundance in the flesh of grape and that
290 of some soft fruits (Famiani et al., 2014b, 2005). In fruits in which ICL activity has been measured it is usually very
291 low: in ripening banana flesh around $0.0001 \text{ U g}^{-1} \text{ FW}$ (Liu et al., 2004) and in cucumber flesh about $0.008 \text{ U g}^{-1} \text{ FW}$
292 (Yang et al., 1998).

293 The occurrence of glyoxysomal ICL polypeptide was investigated in tomato flesh using immunoblotting, and the
294 specificity of the antibody employed has been established in previous studies (Chen et al., 2000; Famiani et al.,
295 2014b, 2005). ICL polypeptide was detected in ripe tomato flesh (development stage 4), however, in this it was at a
296 much lower abundance than in the leaf (Fig. 9). Previous studies have shown that ICL polypeptide is present at very
297 low abundance in young tomato leaves (Nieri et al., 1997), and hence it must be at extremely low abundance in
298 tomato flesh. We could not reliably determine how the abundance of ICL polypeptide changed in the different tissues
299 of tomato fruits during their development and senescence, and this was because of difficulties in detecting ICL on

300 immunoblots. The very low abundance of glyoxysomal ICL in both tomato flesh and that of other fruits, suggests that
301 in these, the conversion of pyruvate/acetyl-CoA to malate by the glyoxylate cycle is not a major pathway utilised by
302 gluconeogenesis under normal conditions of growth. However, recent studies have shown that plants also possess a
303 cytosolic form of ICL with limited sequence similarity to the glyoxysomal form (Eprintsev et al., 2015). This enzyme
304 is thought to function in organic acid interconversions (eg oxalate synthesis in some leaves) and not in the glyoxylate
305 cycle (Eprintsev et al., 2015). Our results do not preclude the presence of this enzyme in fruits, and indeed there is
306 evidence for the occurrence of this enzyme in banana flesh (Eprintsev et al., 2015).

307

308 *3.5. Gluconeogenesis in tomato fruits*

309 Radiolabelling experiments have shown that gluconeogenesis from malate occurs in tomato flesh (Farineau and Laval
310 Martin, 1977; Halinska and Frenkel, 1991). It is likely that PEPCK is utilised in catalysing this gluconeogenic flux
311 (Bahrami et al. 2001; Huang et al., 2015a, 2015b; Osorio et al., 2013). The bulk of the malate/citrate content of
312 tomato flesh is located in the vacuole (Farineau and Laval-Martin, 1977; Knee and Finger, 1992; Rolin et al., 2000),
313 and this is the most likely source of this malate. Nevertheless, some malate could arise from the metabolism of amino
314 acids and amides, and evidence has been provided that the metabolism of vacuolar malate/citrate could be associated
315 with nitrogen metabolism in the flesh of fruits (Famiani et al., 2016a).

316 In the flesh of both ripening grape and peach, citrate and/or malate accumulated before ripening can provide only a
317 small amount of substrate utilised by metabolism (Famiani et al., 2016a, 2016b, 2014a). For tomato flesh, by
318 comparing the amounts of CO₂ released from the fruit with the decrease in malate/citrate contents (Biais et al., 2014;
319 Campbell et al., 1990; Chalmers and Rowan, 1971; Knee and Finger 1992), it can be deduced that the situation is
320 similar. A comparison of the amount of glycolytic flux and the amount of CO₂ released from tomato fruits (Campbell
321 et al., 1990; Carrari et al., 2006; Chalmers and Rowan, 1971) indicates that a large proportion of pyruvate so
322 produced is completely oxidised to CO₂ by the Krebs cycle. This is similar to the situation in the ripening flesh of
323 both grape and peach (Famiani et al., 2016a, 2016b, 2014a). Therefore, in tomato flesh glycolysis from sugars is
324 necessary throughout ripening, and this raises the question as to why gluconeogenesis (reversal of glycolysis) occurs.

325 For both grape and peach flesh one explanation is: the actual rate of malate/citrate dissimilation is not that shown by
326 the changes in their content as determined by two measurements done several days apart. Rather there can be short-
327 term effluxes of malic/citric acid from the vacuole, and this results in a much higher rate of their dissimilation for
328 short periods of time (Famiani et al., 2016a, 2016b; Walker et al., 2015). That is there are times when the amount of
329 malate/citrate released from the vacuole is sufficient to increase the cytosolic malate concentration to a level that
330 brings about gluconeogenesis, and during these periods malate/citrate and not sugars provides the substrate for
331 metabolism (Famiani et al., 2016a, 2016b; Walker et al., 2015). There is evidence to support this proposal in tomato.
332 Firstly, in tomato fruits it appears that there are small diurnal changes in malate content (Farineau and Laval-Martin,
333 1977). Secondly, reducing the amount of PEPCK activity by around 20% in ripening tomato fruits has an inhibitory
334 effect on gluconeogenesis (Osorio et al., 2013). This is not consistent with the observation that activity of PEPCK is
335 80-fold higher than the rate of long-term malate dissimilation (i.e. that measured over a period of several days)
336 (Bahrami et al., 2001). However, this is consistent with there being short-term effluxes of malate/citrate from the
337 vacuole, and that during these periods flux through PEPCK is high.

338 Radiolabelling experiments have shown that gluconeogenesis from pyruvate occurs in tomato flesh (Farineau and
339 Laval Martin, 1977). In the leaves of some CAM plants, PPDK is utilised in gluconeogenesis from pyruvate that is
340 derived from stored vacuolar malate via the action of either cytosolic NADP-ME or mitochondrial NAD-ME
341 (Christopher and Holtum, 1996; Dever et al., 2015; Dittrich, 1976; Häusler et al, 2000). PPDK was present in tomato
342 flesh (Figs. 1A and B, 2, 4), and this raises the possibility that PPDK functions in gluconeogenesis from
343 pyruvate/malate in this tissue. In plants there are both cytosolic and plastidic forms of NADP-ME and a
344 mitochondrial NAD-ME (Famiani et al., 2005). All three enzymes are present in tomato flesh, and their activities
345 appear to be comparable (Bahrami et al., 2001; Jeffery et al., 1986; Knee and Finger, 1992; Knee et al., 1996; Osorio
346 et al., 2013). Tomato flesh contains about 50-times more NAD-ME than does the flesh of either grape or peach (Biais
347 et al., 2014; Borsani et al., 2009; Osorio et al., 2013; Sweetman et al., 2014). The higher abundance of both PPDK
348 and NAD-ME in tomato flesh, compared to that in the flesh of these other fruits, raises the possibility that they can
349 act in concert to catalyse gluconeogenesis from malate/citrate. A recent study showed that pyruvate can be converted

350 to sugars using the PEPCK pathway in ripening tomato pericarp (Osorio et al., 2013). Hence, it appears that a
351 proportion of pyruvate enters the Krebs cycle and is converted to OAA/malate. A proportion of this OAA/malate then
352 equilibrates with the cytosolic pool of these, and a proportion of this cytosolic pool is then utilised in gluconeogenesis
353 using PEPCK. In plants containing reduced amounts of PEPCK gluconeogenesis from pyruvate was almost abolished
354 (Osorio et al., 2013). However, this does not preclude the utilisation of PPDK in gluconeogenesis because it is
355 possible that PPDK is only used at certain times or under certain conditions. In addition, it is also possible that as in
356 some other tissues pyruvate utilised by PPDK could be generated by amino acid metabolism (Chastain et al., 2011;
357 Taylor et al., 2010). Nevertheless, determining whether PPDK contributes to gluconeogenesis in tomato fruits will
358 require further detailed studies. A scheme depicting the potential pathways utilised in gluconeogenesis in tomato
359 flesh is shown in Fig. 10.

360

361 **5. Concluding remarks**

362 In the ripe flesh of a range of fruits PPDK polypeptide was only quite abundant in tomato. Smaller amounts of PPDK
363 polypeptide were detected in the flesh of both peach and pepper. By contrast PEPCK was present in the flesh of all the
364 fruits investigated. A very small amount of glyoxysomal ICL was detected in ripe tomato flesh. In tomato leaves the
365 content g^{-1} FW of PPDK did not increase during their senescence, and this content was highest when the leaf was
366 mature but not senescent. Both PEPCK and PPDK polypeptides decreased in the flesh of tomato during its
367 senescence. The results suggest that in fruits other than tomato the bulk of any gluconeogenic flux proceeds via
368 PEPCK, whereas in tomato both PEPCK and PPDK could potentially be utilised. Further, the results indicate that the
369 conversion of pyruvate/acetyl-CoA to malate by the glyoxylate cycle, for which ICL is necessary, is not a major
370 pathway utilised by gluconeogenesis in fruits under normal conditions of growth. Finally, the results contribute to our
371 understanding of the role of several enzymes in the senescence of both leaves and fruits.

372

373 **Acknowledgements**

374 The research was funded by the “Dipartimento di Scienze Agrarie, Alimentari e Ambientali – Università degli Studi
375 di Perugia - Fondo Ricerca di Base 2014”, Perugia, Italy.

376

377 **References**

- 378 Aoyagi, K., Bassham, J.A., 1983. Pyruvate orthophosphate dikinase in wheat leaves. *Plant Physiol.* 73, 853–854.
- 379 Aoyagi, K., Bassham, J.A., 1984. Pyruvate orthophosphate dikinase of C₃ seeds and leaves as compared to the
380 enzyme from maize. *Plant Physiol.* 75, 387–392.
- 381 Ashton, A.R., Burnell, J.N., Furbank, R.T., Jenkins, C.L.D., Hatch, M.D., 1990. The enzymes in C₄ photosynthesis,
382 in: Lea, P.J., Harborne, J.B. (Eds.), *Enzymes of Primary Metabolism. Methods in Plant Biochemistry*, 3.
383 Academic Press, London, pp. 39–72.
- 384 Bahrami, A.R., Chen, Z.H., Walker, R.P., Leegood, R.C., Gray, J.E., 2001. Ripening-related occurrence of
385 phosphoenolpyruvate carboxykinase in tomato fruit. *Plant Mol. Biol.* 47, 499–506.
- 386 Bailey, K.J., Leegood, R.C., 2016. Nitrogen recycling from the xylem in rice leaves: dependence upon metabolism
387 and associated changes in xylem hydraulics. *J. Exp. Bot.* doi: 10.1093/jxb/erw132.
- 388 Baldicchi, A., Farinelli, D., Micheli, M., Di Vaio, C., Moscatello, S., Battistelli, A., Walker, R.P., Famiani, F., 2015.
389 Analysis of seed growth, fruit growth and composition and phosphoenolpyruvate carboxykinase (PEPCK)
390 occurrence in apricot (*Prunus armeniaca* L.). *Sci. Hortic.* 186, 38–46.
- 391 Biais, B., Bénard, C., Beauvoit, B., Colombié, S., Prodhomme, D., Ménard, G., Bernillon, S., Gehl, B., Gautier, H.,
392 Ballias, P., Mazat, J.P., Sweetlove, L., Génard, M., Gibon, Y., 2014. Remarkable reproducibility of enzyme
393 activity profiles in tomato fruits grown under contrasting environments provides a roadmap for studies of fruit
394 metabolism. *Plant Physiol.* 164, 1204–1221.
- 395 Black, C.C., Chen, J.Q., Doong, R.L., Angelov, M.N., Sung, S.J.S., 1996. Alternative carbohydrate reserves used in
396 the daily cycle of crassulacean acid metabolism, in: Winter, K., Smith, J.A.C. (Eds.), *Crassulacean acid*
397 *metabolism (Ecological studies vol. 114)*. Springer, Heidelberg, pp. 31–45.
- 398 Blanke, M.M., Hucklesby, D.P., Notton, B.A., 1988. Phosphoenolpyruvate carboxykinase in aubergine, kiwi and
399 apple fruit. *Die Gartenbauwissenschaft* 53, 65–70.
- 400 Borsani, J., Budde, C.O., Porrini, L., Lauxmann, M.A., Lombardo, V.A., Murray, R., Andreo, C.S., Drincovich, M.F.,
401 Lara, M.V., 2009. Carbon metabolism of peach fruit after harvest: changes in enzymes involved in organic acid
402 and sugar level modifications. *J. Exp. Bot.* 60, 1823–1837.
- 403 Bortolotti, S., Boggio, S.B., Delgado, L., Orellano, E.G., Valle, E.M., 2003. Different induction patterns of glutamate
404 metabolizing enzymes in ripening fruits of the tomato mutant green flesh. *Physiol. Plant.* 119, 384–391.
- 405 Campbell, A.D., Huysamer, M., Stotz, H.U., Greve, L.C., Labavitch, J.M., 1990. Comparison of ripening processes
406 in intact tomato fruit and excised pericarp discs. *Plant Physiol.* 94, 1582–1589.
- 407 Carrari, F., Baxter, C., Usadel, B., Urbanczyk-Wochniak, E., Zanon, M.I., Nunes-Nesi, A., Nikiforova, V., Centero,
408 D., Ratzka, A., Pauly, M., Sweetlove, L.J., Fernie, A.R., 2006. Integrated analysis of metabolite and transcript
409 levels reveals the metabolic shifts that underlie tomato fruit development and highlight regulatory aspects of
410 metabolic network behavior. *Plant Physiol.* 142, 1380–1396.
- 411 Chalmers, D.J., Rowan, K.S., 1971. The climacteric in ripening tomato fruit. *Plant Physiol.* 48, 235–240.
- 412 Chastain, C.J., Failing, C.J., Manandhar, L., Zimmerman, M.A., Lakner, M.M., Nguyen, T.H.T., 2011. Functional
413 evolution of C₄ pyruvate, orthophosphate dikinase. *J. Exp. Bot.* 62, 3083–3091.
- 414 Chastain, C.J., Heck, J.W., Colquhoun, T.A., Voge, D.G., Gu, X.Y., 2006. Posttranslational regulation of pyruvate,
415 orthophosphate dikinase in developing rice (*Oryza sativa*) seeds. *Planta* 224, 924–934.
- 416 Chastain, C.J., Fries, J.P., Vogel, J.A., Randklev, C.L., Vossen, A.P., Dittmer, S.K., Watkins, E.E., Fiedler, L.J.,
417 Wacker, S.A., Meinhover, K.C., Sarath, G., Chollet, R., 2002. Pyruvate orthophosphate dikinase in leaves and
418 chloroplasts of C₃ plants undergoes light-/dark-induced reversible phosphorylation. *Plant Physiol.* 128, 1368–
419 1378.

420 Chen, Z.H., Walker, R.P., Acheson, R.M., Técsi, L.I., Wingler, A., Lea, P.J., Leegood, R.C., 2000. Are isocitrate
421 lyase and phosphoenolpyruvate carboxykinase involved in gluconeogenesis during senescence of barley leaves
422 and cucumber cotyledons? *Plant Cell Physiol.* 41, 960–967.

423 Christopher, J.T., Holtum, J.A.M., 1996. Patterns of carbon partitioning in leaves of crassulacean acid metabolism
424 species during deacidification. *Plant Physiol.* 111, 393–399.

425 Dever, L.V., Boxall, S.F., Knežova, J., Hartwell, J., 2015. Transgenic perturbation of the decarboxylation phase of
426 crassulacean acid metabolism alters physiology and metabolism but has only a small effect on growth. *Plant*
427 *Physiol.* 167, 44–59.

428 Dittrich, P., 1976. Nicotinamide adenine dinucleotide-specific “malic” enzyme in *Kalanchoë daigremontiana* and
429 other plants exhibiting crassulacean acid metabolism. *Plant Physiol.* 57, 310–314.

430 Doyle, W.P., Huff, R., Wang, C.H., 1960. Role of glyoxylate biosynthesis of acids in tomato fruit. *Plant Physiol.* 35,
431 745–750.

432 Eastmond, P.J., Graham, I.A., 2001. Re-examining the role of the glyoxylate cycle in oilseeds. *Trends Plant Sci.* 6,
433 72–78.

434 Eprintsev, A., Fedorin, D.N., Salnikov, A.V., Igamberdiev, A.U., 2015. Expression and properties of the glyoxysomal
435 and cytosolic forms of isocitrate lyase in *Amaranthus caudatus* L. *J. Plant. Physiol.* 181, 1–8.

436 Famiani, F., Battistelli, A., Moscatello, S., Cruz-Castillo, J.G., Walker, R.P., 2015a. The organic acids that are
437 accumulated in the flesh of fruits: occurrence, metabolism and factors affecting their contents - a review. *Rev.*
438 *Chapingo Ser. Hortic.* 21, 97–128.

439 Famiani, F., Casulli, V., Baldicchi, A., Battistelli, A., Moscatello, S., Walker, R.P., 2012. Development and
440 metabolism of the fruit and seed of the Japanese plum Ozark Premier (Rosaceae). *J. Plant Physiol.* 169, 551–560.

441 Famiani, F., Cultrera, N., Battistelli, A., Casulli, V., Proietti, P., Standardi, A., Chen, Z.H., Leegood, R.C., Walker,
442 R.P., 2005. Phosphoenolpyruvate carboxykinase and its potential role in the catabolism of organic acids in the
443 flesh of soft fruit during ripening. *J. Exp. Bot.* 56, 2959–2969.

444 Famiani, F., Farinelli, D., Moscatello, S., Battistelli, A., Leegood, R.C., Walker, R.P., 2016a. The contribution of
445 stored malate and citrate to the substrate requirements of metabolism of ripening peach (*Prunus persica* L.
446 Batsch) flesh is negligible. Implications for the occurrence of phosphoenolpyruvate carboxykinase and
447 gluconeogenesis. *Plant Physiol. Biochem.* 101, 33–42.

448 Famiani, F., Moscatello, S., Ferradini, N., Gardi, T., Battistelli, A., Walker, R.P., 2014b. Occurrence of a number of
449 enzymes involved in either gluconeogenesis or other processes in the pericarp of three cultivars of grape (*Vitis*
450 *vinifera* L.) during development. *Plant Physiol. Biochem.* 84, 261–270.

451 Famiani, F., Walker, R.P., 2009. Changes in abundance of enzymes involved in organic acid, amino acid and sugar
452 metabolism, and photosynthesis during the ripening of blackberry fruit. *J. Am. Soc. Hortic. Sci.* 134, 167–175.

453 Famiani, F., Farinelli, D., Palliotti, A., Moscatello, S., Battistelli, A., Walker, R.P., 2014a. Is stored malate the
454 quantitatively most important substrate utilised by respiration and ethanolic fermentation in grape berry pericarp
455 during ripening? *Plant Physiol. Biochem.* 76, 52–57.

456 Famiani, F., Farinelli, D., Frioni, T., Palliotti, A., Battistelli, A., Moscatello, S., Walker, R.P., 2016b. Malate as a
457 substrate for catabolism and gluconeogenesis during ripening in the pericarp of different grape cultivars. *Biol.*
458 *Plant.* 60, 155–162.

459 Farineau, J., Laval-Martin, D., 1977. Light versus dark carbon metabolism in cherry tomato fruits. II. Relationship
460 between malate metabolism and photosynthetic activity. *Plant Physiol.* 60, 877–880.

461 Graham, I.A., Leaver, C.J., Smith, S.M., 1992. Induction of malate synthase gene expression in senescent and
462 detached organs of cucumber. *The Plant Cell* 4, 349–357.

463 Halinska, A., Frenkel, C., 1991. Acetaldehyde stimulation of net gluconeogenic carbon movement from applied malic
464 acid in tomato fruit pericarp tissue. *Plant Physiol.* 95, 954–960.

465 Häusler, R.E., Baur, B., Scharte, J., Teichmann, T., Eicks, M., Fischer, K.L., Flügge, U.I., Schubert, S., Weber, A.,
466 Fischer, K., 2000. Plastidic metabolite transporters and their physiological functions in the inducible crassulacean
467 acid metabolism plant *Mesembryanthemum crystallinum*. *Plant J.* 24, 285–296.

468 Hocking, C.G., Anderson, J.W., 1986. Survey of pyruvate, phosphate dikinase activity of plants in relation to the C₃,
469 C₄ and CAM mechanisms of CO₂ assimilation. *Phytochemistry* 25, 1537–1543.

470 Höpfner, M., Ochs, G., Wild, A., 1990. Glutamine synthetase of green and etiolated leaves of *Sinapsis alba*. Evidence
471 of the identity of the respective enzyme proteins. *Planta* 181, 155–161.

472 Huang, Y.X., Goto, Y., Nonaka, S., Fukuda, N., Ezura, H., Matsukura, C., 2015a. Overexpression of the
473 phosphoenolpyruvate carboxykinase gene (SIPEPCK) promotes soluble sugar accumulation in fruit and post-
474 germination growth of tomato (*Solanum lycopersicum* L.). *Plant Biotechnol.* 32, 281–289.

475 Huang, Y.X., Yin, Y.G., Sanuki, A., Fukuda, N., Ezura, H., Matsukura, C., 2015b. Phosphoenolpyruvate
476 carboxykinase (PEPCK) deficiency affects the germination, growth and fruit sugar content in tomato (*Solanum*
477 *lycopersicum* L.). *Plant Physiol. Biochem.* 96, 417–425.

478 Jeffery, D., Goodenough, P.W., Weitzman, P.D., 1986. Enzyme activities in mitochondria isolated from tomato fruit.
479 *Planta* 168, 390–394.

480 Knee, M., Finger, F.L., 1992. NADP⁺-malic enzyme and organic acid levels in developing tomato fruits. *J. Am. Soc.*
481 *Hortic. Sci.* 117, 799–801.

482 Knee, M., Finger, F.L., Lagrimini, L.M., 1996. Evidence for a cytosolic NADP-malic enzyme in tomato.
483 *Phytochemistry* 42, 11–16.

484 Lara, M.V., Borsani, J., Budde, C.O., Lauxmann, M.A., Lombardo, V.A., Murray, R., Andreo, C.S., Drincovich,
485 M.F., 2009. Biochemical and proteomic analysis of ‘Dixiland’ peach fruit (*Prunus persica*) upon heat treatment.
486 *J. Exp. Bot.* 60, 4315–4333.

487 Lara, M.V., Budde, C.O., Porrini, L., Borsani, J., Murray, R., Andreo, C.S., Drincovich, M.F., 2010. Peach (*Prunus*
488 *persica*) fruit response to anoxia: reversible ripening delay and biochemical changes. *Plant Cell Physiol.* 52, 392–
489 403.

490 Leegood, R.C., Walker, R.P., 2003. Regulation and roles of phosphoenolpyruvate carboxykinase in plants. *Arch.*
491 *Biochem. Biophys.* 414, 204–210.

492 Leegood, R.C., Walker, R.P., 1999. Phosphoenolpyruvate carboxykinase in plants: its role and regulation, in: Bryant,
493 J.A., Burrell, M.M., Kruger, N.J. (Eds.), *Plant carbohydrate biochemistry*. BIOS Scientific Publishers Ltd,
494 Oxford, pp. 201–211.

495 Leegood, R.C., Walker, R.P., 1996. Regulation of C₄ pathway, in: Sage, R.F., Monson, R.K. (Eds.), *C₄ Plant biology*.
496 Academic Press, London, pp. 89–132.

497 Liu, S., Yang, Y., Murayama, H., Taira, S., Fukushima, T., 2004. Effects of CO₂ on respiratory metabolism in
498 ripening banana fruit. *Postharvest Biol. Technol.* 33, 27–34.

499 Maeshima, M., Yokoi, H., Asahi, T., 1988. Evidence for no proteolytic processing during transport of isocitrate lyase
500 into glyoxysomes in castor bean endosperm. *Plant Cell Physiol.* 29, 381–384.

501 Nieri, B., Ciurli, A., Pistelli, L., Smith, S.M., Alpi, A., De Bellis, L., 1997. Glyoxylate cycle enzymes in seedlings
502 and in mature plants of tomato (*Lycopersicon esculentum* Mill.). *Plant Sci.* 129, 39–47.

503 Osorio, S., Vallarino, J.G., Szecowka, M., Ufaz, S., Tzin, V., Angelovici, R., Galili, G., Fernie, A.R., 2013.
504 Alteration of the interconversion of pyruvate and malate in the plastid or cytosol of ripening tomato fruit invokes
505 diverse consequences on sugar but similar effects on cellular organic acid, metabolism, and transitory starch
506 accumulation. *Plant Physiol.* 161, 628–643.

507 Pistelli, L., Nieri, B., Smith, S.M., Alpi, A., De Bellis, L., 1996. Glyoxylate cycle enzyme activities are induced in
508 senescent pumpkin fruits. *Plant Sci.* 119, 23–29.

509 Pua, E.C., Chandramouli, S., Han, P., Liu, P., 2003. Malate synthase gene expression during fruit ripening of
510 Cavendish banana (*Musa acuminata* cv. Williams). *J. Exp. Bot.* 54, 309–316.

511 Rolin, D., Baldet, P., Just, D., Chevalier, C., Biran, M., Raymond, P., 2000. NMR study of low subcellular pH during
512 the development of cherry tomato fruit. *Aust. J. Plant Physiol.* 27, 61–69.

513 Ruffner, H.P., 1982. Metabolism of tartaric and malic acid in *Vitis*: a review-Part B. *Vitis* 21, 346–358.

514 Saito, T., Matsukura, C., Ban, Y., Shoji, K., Sugiyama, M., Fukuda, N., Nishimura, S., 2008. Salinity stress affects
515 assimilate metabolism at the gene-expression level during fruit development and improves fruit quality in tomato
516 (*Solanum lycopersicum* L.). *J. Jpn. Soc. Hortic. Sci.* 77, 61–68.

517 Sakakibara, H., Kawabata, S., Takahashi, H., Hase, T., Sugiyama, T., 1992. Molecular cloning of the family
518 glutamine synthetase genes from maize: expression of genes from glutamine synthetase and ferredoxin-
519 dependent glutamate synthase in photosynthetic and non-photosynthetic tissues. *Plant Cell Physiol.* 33, 49–58.

520 Smith, S.M., 2002. Does the glyoxylate cycle have an anaplerotic function in plants? *Trends Plant Sci.* **7**, 12–13.

521 Sugiyama, T., 1973. Purification, molecular, and catalytic properties of pyruvate phosphate dikinase from the maize
522 leaf. *Biochemistry* **12**, 2862–2868.

523 Sweetman, C., Sadras, V.O., Hancock, R.D., Soole, K.L., Ford, C.M., 2014. Metabolic effects of elevated
524 temperature on organic acid degradation in ripening *Vitis vinifera* fruit. *J. Exp. Bot.* **65**, 5975–5988.

525 Taylor, L., Nunes-Nesi, A., Parsley, K., Leiss, A., Leach, G., Coates, S., Wingler, A., Fernie, A.R., Hibberd, J.M.,
526 2010. Cytosolic pyruvate, orthophosphate dikinase functions in nitrogen remobilization during leaf senescence
527 and limits individual seed growth and nitrogen content. *Plant J.* **62**, 641–652.

528 Walker, R.P., Acheson, R.M., Técsi, L.I., Leegood, R.C., 1997. Phosphoenolpyruvate carboxykinase in C4 plants: its
529 role and regulation. *Aust. J. Plant Physiol.* **24**, 459–468.

530 Walker, R.P., Battistelli, A., Moscatello, S., Chen, Z.H., Leegood, R.C., Famiani, F., 2011a. Phosphoenolpyruvate
531 carboxykinase in cherry (*Prunus avium* L.) fruit during development. *J. Exp. Bot.* **62**, 5357–5365.

532 Walker, R.P., Battistelli, A., Moscatello, S., Chen, Z.H., Leegood, R.C., Famiani, F., 2011b. Metabolism of the seed
533 and endocarp of cherry (*Prunus avium* L.) during development. *Plant Physiol. Biochem.* **49**, 923–930.

534 Walker, R.P., Battistelli, A., Moscatello, S., Técsi, L., Leegood, R.C., Famiani, F., 2015. Phosphoenolpyruvate
535 carboxykinase and gluconeogenesis in grape pericarp. *Plant Physiol. Biochem.* **97**, 62–69.

536 Walker, R.P., Chen, Z.H., 2002. Phosphoenolpyruvate carboxykinase: structure, function and regulation. *Adv. Bot.*
537 *Res.* **38**, 93–189.

538 Walker, R.P., Chen, Z.H., Acheson, R.M., Leegood, R.C., 2002. Effects of phosphorylation on phosphoenolpyruvate
539 carboxykinase from the C4 plant, Guinea grass. *Plant Physiol.* **128**, 165–172.

540 Walker, R.P., Chen, Z.H., Johnson, K.E., Famiani, F., Técsi, L., Leegood, R.C., 2001. Using immunohistochemistry
541 to study plant metabolism: the examples of its use in the localization of amino acids in plant tissues, and of
542 phosphoenolpyruvate carboxykinase and its possible role in pH regulation. *J. Exp. Bot.* **52**, 565–576.

543 Walker, R.P., Chen, Z.H., Técsi, L.I., Famiani, F., Lea, P.J., Leegood, R.C., 1999. Phosphoenolpyruvate
544 carboxykinase plays a role in interactions of carbon and nitrogen metabolism during grape seed development.
545 *Planta* **210**, 9–18.

546 Walker, R.P., Famiani, F., Baldicchi, A., Cruz-Castillo, J.G., Inglese, P., 2011c. Changes in enzymes involved in
547 photosynthesis and other metabolic processes in the fruit of *Opuntia ficus-indica* during growth and ripening.
548 *Sci. Hortic.* **128**, 213–219.

549 Walker, R.P., Leegood, R.C., 1996. Phosphorylation of phosphoenolpyruvate carboxykinase in plants: studies in
550 plants with C4 photosynthesis and Crassulacean acid metabolism and in germinating seeds. *Biochem. J.* **317**,
551 653–658.

552 Walker, R.P., Trevanion, S.J., Leegood, R.C., 1995. Phosphoenolpyruvate carboxykinase from higher plants:
553 purification from cucumber and evidence of rapid proteolytic cleavage in extracts from a range of plant tissues.
554 *Planta* **195**, 58–63.

555 Yang, Y., Murayama, H., Fukushima, T., 1998. Activation of glyoxylate cycle enzymes in cucumber fruits exposed
556 to CO₂. *Plant Cell Physiol.* **39**, 533–539.

557 Yin, Y.G., Tominaga, T., Iijima, Y., Aoki, K., Shibata, D., Ashihara, H., Nishimura, S., Ezura, H., Matsukura, C.,
558 2010. Metabolic alterations in organic acids and γ -aminobutyric acid in developing tomato (*Solanum*
559 *lycopersicum* L.) fruits. *Plant Cell Physiol.* **51**, 1300–1314.

560 **Figure legends**

561

562 **Fig. 1.** A) PEPCK and PPDK in ripe tomato flesh and other tissues. Extracts containing the protein content of either 5
563 mg FW ripe tomato flesh (development stage 4), 0.7 mg FW mature Hoya carnosia leaf, 0.4 mg FW mature tomato
564 leaf or 0.4 mg FW mature maize leaf, were subjected to SDS-PAGE. Polypeptides were then either stained in the gel
565 using Coomassie Brilliant Blue dye or transferred to Immobilon-P membrane and enzymes detected using specific
566 antisera. B) PEPCK and PPDK abundance in solanaceous fruits. Extracts containing 5 µg total protein were subjected
567 to SDS-PAGE. Polypeptides were then either stained in the gel using Coomassie Brilliant Blue dye or transferred to
568 Immobilon-P membrane and enzymes detected using specific antisera.

569

570 **Fig. 2.** PPDK abundance in the ripe flesh of tomato (development stage 4) and that of other fruits. Extracts containing
571 the protein content of 4 mg FW of ripe fruit flesh were subjected to SDS-PAGE. Polypeptides were then transferred
572 to Immobilon-P membrane and PPDK detected using a specific antiserum.

573

574 **Fig. 3.** A) Occurrence of PEPCK and PPDK in peach flesh. Extracts containing the protein content of either 5 mg FW
575 ripe peach flesh or 0.4 mg FW mature maize leaf, were subjected to SDS-PAGE. Polypeptides were then transferred
576 to Immobilon-P membrane and PPDK detected using a specific antiserum. The dilution of PPDK antiserum was 1/10
577 000. B) Occurrence of PEPCK and PPDK in peach flesh. Extracts of either 5 mg of peach flesh (from peach fruits at
578 different stages of development; these are the same extracts as used in Famiani et al. 2016a) or 0.4 mg FW mature
579 maize leaf were were subjected to SDS-PAGE. Polypeptides were then transferred to Immobilon-P membrane and
580 PPDK detected using a specific antiserum. The dilution of PPDK antiserum was 1/10 000 (short exposure) and
581 1/1000 (long exposure).

582

583 **Fig. 4.** PEPCK and PPDK abundance in the flesh, locular gel or columella of tomato during development. Extracts
584 containing the protein content of either 5 mg FW tissue (flesh and columella) or 15 mg FW tissue (locular gel) were

585 subjected to SDS-PAGE. Polypeptides were then transferred to Immobilon-P membrane and enzymes detected using
586 specific antisera. The developmental stage of the fruits was based on their size and/or colour: 1, small-green; 2,
587 medium-green; 3, half colouration; 4, full colouration.

588

589 **Fig. 5.** A) PEPCK and PPDK abundance in different parts of tomatoes (green fruits = development stage 2, red fruits
590 = developmental stage 4). Extracts containing 5 µg total protein were subjected to SDS-PAGE. Polypeptides were
591 then transferred to Immobilon-P membrane and enzymes detected using specific antisera. B) PEPCK and PPDK in
592 the flesh of either very young tomato fruit (fruit FW 10% of that of ripe fruit) or ripe fruit (developmental stage 4).
593 Extracts containing the protein content of 5 mg FW were subjected to SDS-PAGE. Polypeptides were then
594 transferred to Immobilon-P membrane and detected using specific antisera.

595

596 **Fig. 6.** PEPCK and PPDK abundance in ripe tomato pericarp post-harvest. Tomato fruits were detached from the
597 plant and incubated under darkness at 25°C for either 3, 6 or 9 days. Extracts corresponding to 4 mg FW of tissue
598 were subjected to SDS-PAGE. Polypeptides were then either stained in the gel using Coomassie Brilliant Blue dye or
599 transferred to Immobilon-P membrane and enzymes detected using specific antisera.

600

601 **Fig. 7.** The abundance of polypeptides in tomato leaves, at different stages of development and senescence, and in
602 other tissues. Extracts corresponding to either 1 mg FW of leaf or 4 mg FW of either fruit or root were subjected to
603 SDS-PAGE. Polypeptides were then stained in the gel using Coomassie Brilliant Blue dye.

604

605 **Fig. 8.** The abundance of PEPCK, PPDK, ICL and other enzymes in tomato leaves, at different stages of
606 development and senescence, and in other tissues. Extracts corresponding to either 1 mg FW of leaf or 4 mg FW of
607 either fruit or root were subjected to SDS-PAGE. Polypeptides were then transferred to Immobilon-P membrane and
608 enzymes detected using specific antisera.

609

610 **Fig. 9.** ICL abundance in young tomato leaves and the flesh of ripe tomato fruits (development stage 4). Extracts
611 containing 5 µg total protein were subjected to SDS-PAGE. Polypeptides were then transferred to Immobilon-P
612 membrane and ICL detected using a specific antiserum.

613

614 **Fig. 10.** Simplified metabolic scheme showing potential pathways utilised by gluconeogenesis in tomato flesh. NAD-
615 ME = NAD-malic enzyme, NADP-ME = cytosolic NADP-malic enzyme, OAA = oxaloacetate, PEP =
616 phosphoenolpyruvate, PEPCK = phosphoenolpyruvate carboxykinase, PDK = pyruvate, orthophosphate dikinase.
617 Note that the bulk of the malate and citrate in the vacuole is almost certainly synthesised from sugars. Nevertheless a
618 small proportion could arise from the carbon skeletons of amino acids and amides.

Figure 1
[Click here to download high resolution image](#)





Figure 2
[Click here to download high resolution image](#)




Figure 3
[Click here to download high resolution image](#)




Figure 4
[Click here to download high resolution image](#)




Figure 5
[Click here to download high resolution image](#)






Figure 6
[Click here to download high resolution image](#)



SDS-PAGE gel



Young tomato leaf

Mid-stage tomato leaf

Mature tomato leaf

Detached mature tomato leaf
after 3 d of darkness (25 °C)

Detached mature tomato leaf
after 6 d of darkness (25 °C)

Detached mature tomato leaf
after 9 d of darkness (25 °C)


Root of maize seedling fed NH₄Cl
In buffer at pH 5.5 for 3 d

Mature maize leaf

Flesh of ripe tomato

Flesh of ripe tomato detached from
plant and incubated at 25 °C for 3 d

Figure 8
[Click here to download high resolution image](#)



Immunoblot

ICL



Young tomato leaf

Flesh of ripe tomato fruit

Figure 10
[Click here to download high resolution image](#)

